

Super-Ehlers in any dimension

Original

Super-Ehlers in any dimension / Sergio, Ferrara; Alessio, Marrani; Trigiante, Mario. - In: JOURNAL OF HIGH ENERGY PHYSICS. - ISSN 1029-8479. - STAMPA. - 2012:11(2012). [10.1007/JHEP11(2012)068]

Availability:

This version is available at: 11583/2504965 since: 2024-07-09T07:23:11Z

Publisher:

Springer

Published

DOI:10.1007/JHEP11(2012)068

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Super-Ehlers in any dimension

Sergio Ferrara,^{a,b,c} Alessio Marrani^a and Mario Trigiante^d

^a*Physics Department, Theory Unit, CERN,
CH 1211, Geneva 23, Switzerland*

^b*INFN - Laboratori Nazionali di Frascati,
Via Enrico Fermi 40, I-00044 Frascati, Italy*

^c*Department of Physics and Astronomy, University of California,
Los Angeles, CA U.S.A.*

^d*Laboratory of Theoretical Physics, Dipartimento di Scienze Applicate e Tecnologia, Politecnico di
Torino, C.so Duca degli Abruzzi, I-10129 Torino, Italy*

E-mail: sergio.ferrara@cern.ch, alessio.marrani@cern.ch,
mario.trigiante@polito.it

ABSTRACT: We classify the enhanced helicity symmetry of the Ehlers group to extended supergravity theories in any dimension. The vanishing character of the pseudo-Riemannian cosets occurring in this analysis is explained in terms of Poincaré duality. The latter resides in the nature of regularly embedded quotient subgroups which are noncompact rank preserving.

KEYWORDS: Supergravity Models, Extended Supersymmetry

ARXIV EPRINT: [1206.1255](https://arxiv.org/abs/1206.1255)

Contents

1	Introduction	1
2	Clifford algebras and “pure” theories	4
2.1	$N = 16$ (<i>maximal</i>)	5
2.2	$N = 12$	12
2.3	$N = 10$	12
3	$N = 8, 6$ matter coupled theories	12
3.1	$N = 8$	12
3.2	$N = 6$	14
4	$N = 4$ matter coupled <i>symmetric</i> theories	14
4.1	<i>Minimal coupling</i> infinite sequence and “ <i>pure</i> ” $D = 4$ supergravity	17
4.2	“ <i>Pure</i> ” $D = 5, 6$ supergravity and T^3 and ST^2 models in $D = 4$	17
4.2.1	$D = 5$	17
4.2.2	$D = 6$	18
4.3	The Jordan symmetric infinite sequence	19
4.3.1	$D = 6$	19
4.3.2	$D = 5$	19
4.3.3	$D = 4$	20
4.4	<i>Magical</i> models	20
4.4.1	$D = 4$	20
4.4.2	$D = 5$	21
4.4.3	$D = 6$	21
5	Cosets with $\chi = 0$ and <i>Poincaré duality</i>	22
5.1	The cosets	26
5.1.1	$N = 16$	26
5.1.2	$N = 12$	26
5.1.3	$N = 10$	28
5.1.4	$N = 8$	28
5.1.5	$N = 6$	28
5.1.6	$N = 4$ <i>symmetric</i>	28
5.2	<i>Poincaré duality</i>	32
5.2.1	$N = 16$	32
5.2.2	$N = 4$ <i>Magical</i> models	39
5.3	Hodge involution and coset cohomology	42
6	Conclusion	46

A	Embeddings	46
A.1	The embeddings $SL(D - 2, \mathbb{R}) \times E_{11-D(11-D)} \subset E_{8(8)}$	47
A.2	Other embeddings	50
A.3	General features	53
B	$\mathfrak{so}(8, 8)$ outer automorphisms and dual subalgebras of $\mathfrak{e}_{8(8)}$	54
C	Poincaré duality and level decomposition	58

1 Introduction

Three decades ago it was shown [1] that the D -dimensional *Ehlers group* $SL(D - 2, \mathbb{R})$ is a symmetry of D -dimensional Einstein gravity, provided that the theory is formulated in the light-cone gauge. For any $D \geq 4$ -dimensional Lorentzian space-time, this results enables to identify the graviton degrees of freedom with the Riemannian coset

$$\mathcal{M}_{\text{grav}} = \frac{SL(D - 2, \mathbb{R})}{SO(D - 2)}, \tag{1.1}$$

even if the action of the theory is not simply the sigma model action on this coset (with the exception of a theory reduced to $D = 3$). In $D = 4$, this statement reduces to the well known fact that the massless graviton described by the Einstein-Hilbert action with two degrees of freedom of ± 2 helicity has an enhanced symmetry $SO(2) \rightarrow SL(2, \mathbb{R})$.

In \mathcal{N} -extended supergravity in D dimensions, U -duality¹ symmetries play an important role to uncover, in terms of geometrical constructions, the non-linear structure of the theories, whose most symmetric one is the theory with maximal supersymmetry ($2N = 32$ supersymmetries). Furthermore, U -duality symmetries get unified with the Ehlers space-time symmetry if one descends to $D = 3$ [5, 6]. In the maximal case, the $D = 3$ U -duality group is $E_{8(8)}$, with *maximal compact subgroup (mcs)* $SO(16)$, which is also the underlying Clifford algebra for massless supermultiplets of maximal supersymmetry. As a consequence, the bosonic sector of the theory is described by the sigma model $E_{8(8)}/SO(16)$ [7–9].

Following these preliminaries, it comes as no surprise that it was further discovered that in light-cone Hamiltonian formulation maximal supergravity exhibits $E_{7(7)}$ symmetry in $D = 4$ [11] and $E_{8(8)}$ symmetry in $D = 3$ [12] (for the $D = 11$ theory, see [10]). Indeed, in any space-time dimension D and for any number of supersymmetries $\mathcal{N} = 2N$, it is known that the $D = 3$ U -duality group G_N^3 [13] embeds (through a rank-preserving embedding; for some basic definitions, see the start of appendix A) the Ehlers group $SL(D - 2, \mathbb{R})$ as a commutant of the U -duality group G_N^D [16, 17]:

$$G_N^3 \supset G_N^D \times SL(D - 2, \mathbb{R}). \tag{1.2}$$

¹Here U -duality is referred to as the “continuous” symmetries of [2, 3]. Their discrete versions are the U -duality non-perturbative string theory symmetries introduced by Hull and Townsend [4].

It is then natural to conjecture that in a suitable light-cone formulation of any \mathcal{N} -extended supergravity theories $G_N^D \times \text{SL}(D-2, \mathbb{R})$ (which we dub *super-Ehlers group*) is a manifest symmetry of the theory. Even if the super-Ehlers group is a bosonic extension of the Ehlers group itself, the presence of the U -duality commutant G_N^D in (1.2) is closely related to supersymmetry. It is intriguing to notice that the super-Ehlers symmetries, which we classify below in any dimension, sometimes exhibit an “*enhancement*” into some larger group;² this occurs whenever the embedding (1.2) is *non-maximal*, and in $D = 10$ type IIB supergravity. Furthermore, it sometimes occurs that the embedding (1.2) is maximal but *non-symmetric*, as in $D = 11$ supergravity.

At any rate, we will show that the common features of the embedding (1.2) are *at least* two (*cfr.* the start of appendix A):

- It is *regular* and preserves the rank of the group. Indeed, it generally holds that

$$\text{rank}(G_N^3) = \text{rank}(G_N^D) + \text{rank}(\text{SL}(D-2, \mathbb{R})) = \text{rank}(G_N^D) + D - 3. \quad (1.3)$$

The same relation holds for the *non-compact rank* of these groups, namely the rank of the corresponding symmetric Riemannian manifolds of which the groups encode the isometries:

$$\text{rank}\left(\frac{G_N^3}{H_N^3}\right) = \text{rank}\left(\frac{G_N^D}{H_N^D}\right) + \text{rank}\left(\frac{\text{SL}(D-2, \mathbb{R})}{\text{SO}(D-2)}\right) = \text{rank}\left(\frac{G_N^D}{H_N^D}\right) + D - 3, \quad (1.4)$$

where H_N^3 and H_N^D are the maximal compact subgroups of G_N^3 and G_N^D , respectively. As mentioned above, this does not imply the embeddings to be in general maximal nor symmetric.

- The pseudo-Riemannian coset resulting from (1.2) has always zero *character* [14, 15], namely it has the same number of compact and non-compact generators. We will show that this latter property is related to the *Poincaré duality* (*alias* electric-magnetic duality) of the spectrum of massless $p > 0$ forms of the theory, which can essentially be traced back to the existence of an *Hodge involution* in the cohomology of the scalar manifold, singling out only the physical forms and their duals in the cohomology of the $(D-2)$ -dimensional transverse space. This property also follows from the regularity of the embedding of $G_N^D \times \text{SL}(D-2)$ inside G_N^3 , the semisimplicity of the two groups and properties (1.3), (1.4), as it will be shown in appendix A.3.

There is also another aspect of interest in the present analysis: the role played by exceptional Lie groups and their relation to Jordan algebras and Freudenthal triple systems [19–21]. In particular, a mathematical construction, called *Jordan pairs* (see e.g. [22] for a recent treatment, and a list of refs.) corresponds to the maximal non-symmetric embedding

$$E_{8(8)} \supset E_{6(6)} \times \text{SL}(3, \mathbb{R}), \quad (1.5)$$

²For enhancement to infinite symmetries, see [18].

which is nothing but (1.2) specified for maximal supersymmetry ($N = 16$) and $D = 5$. We point out that the *Jordan pairs* relevant for supergravity theories always pertain to suitable non-compact real forms of Lie algebras, differently e.g. from the treatment given in [22].

Moreover, it is worth observing that in $D = 11$ supergravity G_{16}^{11} is empty, and thus (1.2) is the following maximal non-symmetric embedding [5]:

$$E_{8(8)} \supset \text{SL}(9, \mathbb{R}), \tag{1.6}$$

which in fact was used long time ago [23] in order to construct the gravity multiplet of this theory [24]. For maximal supergravity ($N = N_{\text{max}} = 16$), (1.2) reads³

$$E_{8(8)} \supset E_{11-D(11-D)} \times \text{SL}(D - 2, \mathbb{R}), \tag{1.7}$$

where $G_{16}^D = E_{11-D(11-D)}$ denotes the so-called *Cremmer-Julia sequence* [8, 9]. The unique exception is provided by type IIB chiral $D = 10$ supergravity, in which (1.2) is given by a two-step chain of maximal embeddings:⁴

$$E_{8(8)} \supset_s \text{SL}(2, \mathbb{R}) \times E_{7(7)} \supset_s \text{SL}(2, \mathbb{R}) \times \text{SL}(8, \mathbb{R}), \tag{1.8}$$

which preserves the group rank.

The plan of the paper is as follows.

In section 2 we start by recalling some basic facts on $\text{SO}(N)$ Clifford algebras relevant for the classification of massless multiplets of \mathcal{N} -extended supersymmetry in any dimension. Here $\mathcal{N} = 2N$ denotes the number of supersymmetries, regardless of the dimension D . Thus, for instance maximal supergravity corresponds to $\mathcal{N} = 32$ (8 spinor supercharges in $D = 4$), whereas the minimal supergravity we consider has $\mathcal{N} = 8$ (2 spinor supercharges in $D = 4$). We then proceed to considering the embedding (1.7) pertaining to maximal supergravity in any dimension $D \geq 4$ (in $D = 10$ both IIA and IIB theories are considered). The embedding (1.2), which can be regarded as the “*non-compact enhancement*” of Nahm’s analysis [23], in all cases consistently provides the massless spectrum of the corresponding theory with the correct spin-statistics content; illustrative analysis is worked out for $D = 11$ and $D = 10$ maximal theories. Other theories which do not exhibit matter coupling are also considered, namely $N = 10, 12$ in $D = 4$ and $N = 12$ in $D = 5$.

In section 3 we consider half-maximal supergravity theories ($N = 8$), which can be matter coupled and exist in all $D \leq 10$ dimensions; for $D = 6$ we consider both inequivalent theories, namely the chiral (2,0) (type IIB) and the non-chiral (1,1) (type IIA) ones. Theories with $N = 6$, living in $D = 4$, are also considered.

Then, in section 4 we consider quarter-maximal theories ($N = 4$), which live in $D = 4, 5, 6$ and admit two different kinds of matter multiplets. We confine ourselves to theories with symmetric scalar manifolds, which (apart from the *minimally coupled* models in $D = 4$

³This embedding was considered, but not proved, in [16]. A proof is presented in appendix A of the present paper.

⁴“*s*” and “*ns*” stand for “symmetric” and “non-symmetric” (embedding), respectively.

and the *non-Jordan* symmetric sequence in $D = 5$) admit an interpretation in terms of Euclidean Jordan algebras.

Pseudo-Riemannian cosets associated to the maximal-rank embeddings (1.2) are then analyzed in section 5. All such cosets enjoy the property of having the same number of compact and non-compact generators. This is also proven, using general group theoretical arguments, in appendix A.3. In subsection 5.2 this property is related to the invariance of the spectrum of massless bosonic $p > 0$ forms under *Poincaré-duality*, or equivalently in subsection 5.3 in terms of an *Hodge involution* acting on the coset cohomology.

Final remarks and outlook are given in section 6.

Three appendices conclude the paper. In appendix A, some embeddings of non-compact, real forms relevant for our analysis are rigorously proved, while in appendix B the issue of inequivalent “dual pairs” of subalgebras of the U -duality algebra is discussed (see also [25]). The related notions of T -dualities as $\mathfrak{so}(8, 8)$ outer-automorphisms are also dealt with. In appendix C the issue of Poincaré duality is revisited with an explicit algebraic construction which makes use of appropriate *level decompositions*.

2 Clifford algebras and “pure” theories

In the seminal paper by Nahm [23], it was shown how massless multiplets of supergravity are built in terms of irreps. of $SO(D - 2)$, the little group (*spin*) of massless particles in D dimensions. The number of supersymmetries $2N$ is encoded in the Clifford algebra of $SO(N)$, and therefore the supermultiplets can be regarded as $SO(N)$ spinors decomposed into $SO(D - 2)$ irreps (for theories with particles with spin $s \leq 2$, which we consider throughout, $N_{\max} = 16$). Bosons and fermions thus correspond to the two semi-spinors (or chiral spinors) of⁵ $SO(N)$.

In any dimension $D \geq 4$, $SO(N)$ exhibits a certain commuting factor with the massless little group $SO(D - 2)$. For “pure” supergravities, in which only the gravity multiplet is present, such a commuting factor is the so-called \mathcal{R} -symmetry of the theory. Then the question arises as to which is the non-compact group commuting with the $SL(D - 2, \mathbb{R})$ Ehlers group (which thus extends the massless little group including the \mathcal{R} -symmetry), and furthermore which is the non-compact group which extends the $SO(N)$ of the N -dimensional Clifford algebra pertaining to $2N$ local supersymmetries.

In describing massless multiplets of theories with $\mathcal{N} = 2N$ local supersymmetries, one consider the the rest-frame supersymmetry algebra without central extension. Since the momentum squares to zero ($P^\mu P_\mu = 0$), only half of the supersymmetry charges survive, and the creation operators of N charges describe an $SO(N)$ Clifford algebra. Moreover, due to the fact that in $D \geq 4$ spinors always have real dimension multiple of 4, N is always even: $N = 4, 6, 8, 10, 12, 16$ (we do not consider here $N = 2$ at $D = 4$, namely minimal 4-dimensional supergravity with 1 spinor supercharge). It thus comes as no surprise that

⁵Note that N is always even, since for $D \geq 4$ spinor charges have real dimensions multiples of 4.

Furthermore, it should be remarked that the cases $D = 4$, $N = 2$ and $D = 10$, $N = 8$ are somewhat particular, because $N = D - 2$, so the two Clifford spinors directly provide bosonic and fermionic supermultiplets’ representations.

U -duality groups G_N^3 in $D = 3$ (in which there is only distinction between bosons and fermions, but no spin is present for massless states) contain in their mcs the Clifford algebra symmetry $SO(N)$.

Supersymmetry dictates that massless bosons and fermions are simply the two (chiral, semi-) spinor irreps. of $SO(N)$, while their spin s content in D space-time dimensions is obtained by suitably branching such irreps. into $SO(D - 2)$, which is the little group (spin) for massless particles in D dimensions.

In the present section we consider “pure” theories in which the matter coupling is not allowed; they include *maximally supersymmetric* ($N = 16$) theories in any dimension $D \leq 11$, as well as *non-maximal* theories with $N = 10, 12$ in $D = 4$ and $N = 12$ in $D = 5$. For such theories, the Clifford algebra $SO(N)$ is nothing but the mcs of the U -duality group G_N^3 in $D = 3$; for *non-maximal* theories ($N < 16$), this is true up to the presence of the so-called *Clifford vacuum* factor group, which expresses further degeneracy of the Clifford algebra symmetry. Moreover, the group $H_N^D = mcs(G_N^D)$ which commutes with $SO(D - 2)$ inside $SO(N)$ is the \mathcal{R} -symmetry, providing the degeneracy of the spin s representations in the decomposition of the chiral spinors under the embedding⁶ [23]

$$SO(N) \supset H_N^D \times SO(D - 2)_J \tag{2.1}$$

which is the (not necessarily maximal-rank, nor maximal nor symmetric) counterpart of (1.2) at the level of mcs . The subscript “ J ” denotes the spin group throughout.

2.1 $N = 16$ (*maximal*)

For *maximal* ($N = 16$) supergravity theories with massless particles, the $D = 3$ U -duality group is $G_3^{16} = E_{8(8)}$, with mcs $SO(16)$, which is the Clifford algebra for massless particles with $\mathcal{N} = 32$ supersymmetries. (1.7) provides the rank-preserving embedding of D -dimensional Ehlers group $SL(D - 2, \mathbb{R})$ into $E_{8(8)}$. The group commuting with $SL(D - 2, \mathbb{R})$ inside $E_{8(8)}$ is nothing but the D -dimensional U -duality group $G_{16}^D = E_{11-D(11-D)}$, belonging to the so-called the Cremmer-Julia sequence. All cases in $4 \leq D \leq 11$ dimensions are reported in table 1 (non-compact level (1.2)–(1.7)) and in table 2 (mcs level (2.1)). In particular, in table 2 also the decomposition of the vector irrep. **16** of the Clifford algebra $SO(16) = mcs(E_{8(8)})$ of maximal ($N = 16 \rightarrow \mathcal{N} = 32$) supersymmetry is reported for the embedding (2.1) pertaining to this case, namely [23] (see also [26]):

$$SO(16) \supset \mathcal{R}_D^{16} \times SO(D - 2)_J, \tag{2.2}$$

where, as mentioned above, $\mathcal{R}_D^{16} \equiv mcs(G_D^{16}) \equiv H_D^{16}$ is the \mathcal{R} -symmetry of the maximal supergravity in D (Lorentzian) space-time dimensions. Note that the irrep. of $SO(D - 2)$ occurring in the branching of the **16** along (2.2) are all spinors, and the \mathcal{R} -symmetry \mathcal{R}_D^{16} is real, pseudo-real (quaternionic), complex, depending on whether such spinor irrep. is real, pseudo-real or complex, respectively.

⁶Further commuting factor group occurs in the l.h.s. of (2.1) in non-maximal ($N \leq 16$) theories; see analysis below.

D	$E_{8(8)} \supset E_{11-D(11-D)} \times \text{SL}(D-2, \mathbb{R})$	type
11	$E_{8(8)} \supset \text{SL}(9, \mathbb{R})$	<i>max, ns</i>
10, <i>IIA</i>	$E_{8(8)} \supset \text{SO}(1, 1) \times \text{SL}(8, \mathbb{R})$	<i>nm, ns</i>
10, <i>IIB</i>	$E_{8(8)} \supset \text{SL}(2, \mathbb{R}) \times \text{SL}(8, \mathbb{R})$	<i>nm, ns</i>
9	$E_{8(8)} \supset \text{GL}(2, \mathbb{R}) \times \text{SL}(7, \mathbb{R})$	<i>nm, ns</i>
8	$E_{8(8)} \supset \text{SL}(2, \mathbb{R}) \times \text{SL}(3, \mathbb{R}) \times \text{SL}(6, \mathbb{R})$	<i>nm, ns</i>
7	$E_{8(8)} \supset \text{SL}(5, \mathbb{R}) \times \text{SL}(5, \mathbb{R})$	<i>max, ns</i>
6	$E_{8(8)} \supset \text{SO}(5, 5) \times \text{SL}(4, \mathbb{R})$	<i>nm, ns</i>
5	$E_{8(8)} \supset E_{6(6)} \times \text{SL}(3, \mathbb{R})$	<i>max, ns</i>
4	$E_{8(8)} \supset E_{7(7)} \times \text{SL}(2, \mathbb{R})$	<i>max, s</i>

Table 1. Embedding $G_N^3 \supset G_N^D \times \text{SL}(D-2, \mathbb{R})$ (1.2) for *maximal* supergravity theories ($N = 16$) in $11 \geq D \geq 4$ Lorentzian space-time dimensions [16, 17]. G_N^D is the U -duality group in D dimensions for the theory with $\mathcal{N} = 2N$ supersymmetries. $\text{SL}(D-2, \mathbb{R})$ is the Ehlers group in D dimensions. For $N = 16$, $G_{16}^3 = E_{8(8)}$, and $G_N^D = E_{11-D(11-D)}$ belongs to the Cremmer-Julia sequence; thus, (1.7) is obtained. The type (*max*(imal), *n*(ext-to-)*m*(aximal), *s*(ymmetric), *n*(on-)*s*(ymmetric)) of embedding is indicated. Explicit proofs are given in appendix A.

Let us scan them briefly (as anticipated, for $D = 11$ and $D = 10$ the massless spectrum analysis is also worked out, as an example of the consistence of the embeddings with the massless spectrum of the corresponding theory). For convenience of the reader, we anticipate that the embeddings (1.2) and (2.1) are *maximal* in $D = 11, 7, 5$ (non-symmetric)

D	$SO(16) \supset H_{16}^D \times SO(D-2)_J$	type
11	$SO(16) \supset SO(9)$ $\mathbf{16} = \mathbf{16}$	<i>max, ns</i>
10, <i>IIA</i>	$SO(16) \supset SO(8)$ $\mathbf{16} = \mathbf{8}_s + \mathbf{8}_c$	<i>nm, ns</i>
10, <i>IIB</i>	$SO(16) \supset SO(2) \times SO(8)$ $\mathbf{16} = (\mathbf{2}, \mathbf{8}_s)$	<i>nm, ns</i>
9	$SO(16) \supset SO(2) \times SO(7)$ $\mathbf{16} = (\mathbf{2}, \mathbf{8})$	<i>nm, ns</i>
8	$SO(16) \supset U(1) \times SU(2) \times SU(4)$ $\mathbf{16} = (\mathbf{2}, \mathbf{4}) + (\bar{\mathbf{2}}, \bar{\mathbf{4}})$	<i>nm, ns</i>
7	$SO(16) \supset USp(4) \times USp(4)$ $\mathbf{16} = (\mathbf{4}, \mathbf{4})$	<i>max, ns</i>
6	$SO(16) \supset USp(4)_L \times USp(4)_R \times SU(2)_L \times SU(2)_R$ $\mathbf{16} = (\mathbf{4}, \mathbf{1}, \mathbf{2}, \mathbf{1}) + (\mathbf{1}, \mathbf{4}, \mathbf{1}, \mathbf{2})$	<i>nm, ns</i>
5	$SO(16) \supset USp(8) \times SU(2)$ $\mathbf{16} = (\mathbf{8}, \mathbf{2})$	<i>max, ns</i>
4	$SO(16) \supset SU(8) \times U(1)$ $\mathbf{16} = \mathbf{8}_1 + \bar{\mathbf{8}}_{-1}$	<i>max, s</i>

Table 2. Embedding $H_N^3 \supset H_N^D \times SO(D-2)$ (2.1) [23] for *maximal* supergravity theories ($N = 16$) in $11 \geq D \geq 4$ Lorentzian space-time dimensions. In this case, as for all “*pure*” theories, H_N^D is the \mathcal{R} -symmetry for the theory with $\mathcal{N} = 2N$ supersymmetries. $SO(D-2)$ is the little group (spin group) for massless particles. In this case, $H_{16}^3 = SO(16)$ is the Clifford algebra of maximal supersymmetry.

and 4 (symmetric), while they are *next-to-maximal* in $D = 10, 9, 8, 6$; in these latter cases, an “*enhancement*” of $E_{11-D(11-D)} \times SL(D-2, \mathbb{R})$ occurs (see analysis below).

1. $D = 11$ (*M*-theory). There is no continuous U -duality (and thus \mathcal{R} -symmetry) group, and (1.7) specifies to (1.6), namely the maximal non-symmetric embedding of

the Ehlers group $SL(9, \mathbb{R})$ only:

$$\begin{aligned} E_{8(8)} \supset_{ns} SL(9, \mathbb{R}); \\ \mathbf{248} = \mathbf{80} + \mathbf{84} + \mathbf{84}', \end{aligned} \tag{2.3}$$

where $\mathbf{84}$ and $\mathbf{84}'$ are the 3-fold antisymmetric of $SL(9, \mathbb{R})$ and its dual; they correspond to gauge fields coupling to $M2$ branes and $M5$ branes, respectively. The corresponding *mcs* level is given by the specification of (2.1) to the following non-symmetric embedding of the massless spin group $SO(9)$ only:

$$SO(16) \supset_{ns} SO(9). \tag{2.4}$$

For what concerns the massless spectrum, one considers the maximal symmetric embedding⁷

$$E_{8(8)} \supset_s^{mcs} SO(16) : \mathbf{248} = \mathbf{120} + \mathbf{128}, \tag{2.5}$$

where $\mathbf{128}$ is one of the two chiral spinor irreps. of $SO(16)$. Under (2.4), such two chiral irreps. $\mathbf{128}$ and $\mathbf{128}'$ further decompose as follows:

$$SO(16) \supset_{ns} SO(9) : \begin{cases} \mathbf{128} = \mathbf{84} + \mathbf{44}; \\ \mathbf{128}' = \mathbf{128}, \end{cases} \tag{2.6}$$

where, on the right-hand side, $\mathbf{44}$, $\mathbf{84}$ and $\mathbf{128}$ are the rank-2 symmetric traceless, the rank-3 antisymmetric and the gamma-traceless vector-spinor irreps. of the massless spin group $SO(9)$, respectively. Thus, (2.6) establishes the chiral spinor irrep. $\mathbf{128}$ of the Clifford algebra $SO(16)$ to be irrep. pertaining to the massless *bosonic* spectrum (it branches into the graviton $\mathbf{44}$ and the 3-form $\mathbf{84}$), whereas its conjugate semi-spinor irrep. $\mathbf{128}'$ pertains to the massless *fermionic* spectrum of M -theory (it corresponds to the $D = 11$ gravitino).

2. In $D = 10$ type IIA theory the U -duality is $G_{16}^{10 IIA} = SO(1,1)$ (and thus no continuous \mathcal{R} -symmetry); since this theory is obtained as the Kaluza-Klein S^1 -reduction of M -theory, the relevant chain of maximal embeddings reads

$$E_{8(8)} \supset_{ns} SL(9, \mathbb{R}) \supset_s SO(1,1) \times SL(8, \mathbb{R}); \tag{2.7}$$

note the “*enhancement*” to $SL(9, \mathbb{R})$, consistent with the M -theoretical origin of IIA theory. The corresponding *mcs* level is

$$SO(16) \supset_{ns} SO(9) \supset_s SO(8), \tag{2.8}$$

where $SO(8)$ is the massless spin group. Throughout our analysis, we dub “*next-to-maximal*” (*nm*) those embeddings given by a chain of two maximal embeddings; note that all *nm* embeddings considered in the present investigation are of *maximal* rank, namely they preserve the rank of the original group. For what concerns the IIA

⁷For further subtleties concerning exceptional Lie algebras, see [25] and appendix B further below.

massless spectrum, one considers the branchings of $\mathbf{128}$ (bosons) and $\mathbf{128}'$ (fermions) of the Clifford algebra $SO(16)$ under the nm embedding (2.8):

$$\mathbf{128} = \mathbf{84} + \mathbf{44} = \mathbf{56}_v + \mathbf{28} + \mathbf{35}_v + \mathbf{8}_v + \mathbf{1}; \tag{2.9}$$

$$\mathbf{128}' = \mathbf{128} = \mathbf{56}_s + \mathbf{56}_c + \mathbf{8}_s + \mathbf{8}_c, \tag{2.10}$$

where the subscripts “ v ”, “ s ” and “ c ” respectively stand for *vector*, *spinor*, *conjugate spinor*, and they pertain to the *triality* of $SO(8)$, the little group (spin group) of massless particles in $D = 10$. $\mathbf{56}_i$, $\mathbf{28}$, $\mathbf{35}_i$ and $\mathbf{8}_i$ ($i = v, s, c$) are the rank-3 antisymmetric, adjoint, rank-2 symmetric traceless and vector/spinor irreps. of $SO(8)$, respectively. Thus, the branching (2.9) consistently pertains to the IIA massless *bosonic* spectrum: 3-form $C_{\mu\nu\rho}^{(3)}$ ($\mathbf{56}_v$), B -field $B_{\mu\nu}$ ($\mathbf{28}$), graviton $g_{\mu\nu}$ ($\mathbf{35}_v$), graviphoton $C_\mu^{(1)}$ ($\mathbf{8}_v$) and dilaton scalar field ϕ_{10} ($\mathbf{1}$). On the other hand, the branching (2.10) pertains to the IIA massless *fermionic* spectrum: gravitinos $\mathbf{56}_s$ and $\mathbf{56}_c$ ($s = 3/2$ Majorana-Weyl spinors of opposite chirality), and gauginos $\mathbf{8}_s$ and $\mathbf{8}_c$ ($s = 1/2$ Majorana-Weyl spinors of opposite chirality). This *non-chiral* spectrum can also be deduced by dimensional reduction of the maximal supersymmetric supermultiplet of $D = 11$ supergravity (M -theory).

3. On the other hand, in $D = 10$ type IIB theory the U -duality is $G_{16}^{10}{}^{IIB} = SL(2, \mathbb{R})$, and its mcs is the \mathcal{R} -symmetry $U(1)$, and the relevant nm embedding is given by (1.8), which we report here:

$$E_{8(8)} \supset_s SL(2, \mathbb{R}) \times E_{7(7)} \supset_s SL(2, \mathbb{R}) \times SL(8, \mathbb{R}); \tag{2.11}$$

$$SO(16) \supset_s U(1) \times SU(8) \supset_s U(1) \times SO(8); \tag{2.12}$$

note the “*exceptional enhancement*” to $E_{7(7)}$ in (2.11). For what concerns the IIB massless spectrum, one considers the branching of $\mathbf{128}$ (bosons) and $\mathbf{128}'$ (fermions) of the Clifford algebra $SO(16)$ under the nm embedding (2.12). Under the decomposition

$$\begin{aligned} SU(8) \supset_s SO(8) \\ \mathbf{8} = \mathbf{8}_s, \end{aligned} \tag{2.13}$$

one obtains (disregarding $U(1)$ charges)

$$\mathbf{128} = \mathbf{70}_0 + \mathbf{28} + \overline{\mathbf{28}} + \mathbf{1} + \mathbf{1} = \mathbf{35}_v + \mathbf{35}_c + \mathbf{28} + \mathbf{28} + \mathbf{1} + \mathbf{1}; \tag{2.14}$$

$$\mathbf{128}' = \mathbf{56} + \overline{\mathbf{56}} + \mathbf{8} + \overline{\mathbf{8}} = \mathbf{56}_s + \mathbf{56}_s + \mathbf{8}_s + \mathbf{8}_s. \tag{2.15}$$

Note that, upon (2.13), the rank-4 antisymmetric self-real irrep. $\mathbf{70}$ of $SU(8)$ breaks into $\mathbf{35}_v + \mathbf{35}_c$ of $SO(8)$. Thus, the branching (2.14) consistently pertains to the IIB massless *bosonic* spectrum: graviton $g_{\mu\nu}$ ($\mathbf{35}_v$), 4-form $C^{(4)}$ ($\mathbf{35}_c$), B -field $B_{\mu\nu}$ ($\mathbf{28}$), 2-form $C_{\mu\nu}^{(2)}$ ($\mathbf{28}$), and two scalar fields, namely the dilaton ϕ_{10} and the axion $C^{(0)}$ ($\mathbf{1} + \mathbf{1}$). On the other hand, the branching (2.15) pertains to the IIB massless *fermionic* spectrum: gravitinos $\mathbf{56}_s$ and $\mathbf{56}_s$ ($s = 3/2$ Majorana-Weyl spinors of same chirality) and gauginos $\mathbf{8}_s$ and $\mathbf{8}_s$ ($s = 1/2$ Majorana-Weyl spinors of same chirality). This spectrum is *chiral* and hence cannot be obtained by dimensional reduction of the $D = 11$ M -theory supermultiplet.

4. In $D = 9$ the U -duality is $G_{16}^9 = GL(2, \mathbb{R}) \equiv E_{2(2)}$, and its mcs is the \mathcal{R} -symmetry $U(1)$. There are two possible chains of maximal embeddings, which are equivalent up to redefinitions of $SO(1, 1)$ weights. The first chain, pertinent to a dimensional reduction of M -theory, gives rise to a nm embedding:

$$E_{8(8)} \supset_{ns} SL(9, \mathbb{R}) \supset_s GL(2, \mathbb{R}) \times SL(7, \mathbb{R}); \quad (2.16)$$

$$SO(16) \supset_{ns} SO(9) \supset_s U(1) \times SO(7), \quad (2.17)$$

whereas the second, pertaining to a Kaluza-Klein S^1 -reduction of $D = 10$ IIB theory, determines a “*next-to-next-to-maximal*” (nnm) embedding, because it is 3-stepwise (it is given by a further branching of IIB chain (2.11)):

$$E_{8(8)} \supset_s SL(2, \mathbb{R}) \times E_{7(7)} \supset_s SL(2, \mathbb{R}) \times SL(8, \mathbb{R}) \supset_s GL(2, \mathbb{R}) \times SL(7, \mathbb{R}); \quad (2.18)$$

$$SO(16) \supset_s U(1) \times SU(8) \supset_s U(1) \times SO(8) \supset_s U(1) \times SO(7). \quad (2.19)$$

Besides being equivalent, (2.16)–(2.17) and (2.18)–(2.19) are consistent, because type IIA and IIB theories are equivalent in $D \leq 9$ dimensions (except for half-maximal supergravity in $D = 6$; see further below).

5. In $D = 8$ the U -duality is $G_{16}^8 = SL(2, \mathbb{R}) \times SL(3, \mathbb{R}) \equiv E_{3(3)}$, and its mcs is the \mathcal{R} -symmetry $U(1) \times SU(2) \sim U(2)$. The relevant nm embedding reads⁸ ($SO(6) \sim SU(4)$)

$$E_{8(8)} \supset_{ns} E_{6(6)} \times SL(3, \mathbb{R}) \supset_s SL(2, \mathbb{R}) \times SL(3, \mathbb{R}) \times SL(6, \mathbb{R}); \quad (2.20)$$

$$SO(16) \supset_{ns} USp(8) \times SU(2) \supset_s U(1) \times SU(2) \times SU(4); \quad (2.21)$$

note the “*exceptional enhancement*” to $E_{6(6)}$ in (2.20).

6. In $D = 7$ the U -duality is $G_{16}^7 = SL(5, \mathbb{R}) \equiv E_{4(4)}$, and its mcs is the \mathcal{R} -symmetry $SO(5) \sim USp(4)$. The relevant embedding is maximal non-symmetric:

$$E_{8(8)} \supset_{ns} SL(5, \mathbb{R}) \times SL(5, \mathbb{R}); \quad (2.22)$$

$$SO(16) \supset_{ns} USp(4) \times USp(4). \quad (2.23)$$

Note that in this case there is perfect symmetry between the \mathcal{R} -symmetry and the massless spin sectors.

7. In $D = 6$ (non-chiral $(2, 2)$) maximal theory, the U -duality is $G_{16}^6 = SO(5, 5) \equiv E_{5(5)}$, and its mcs is the \mathcal{R} -symmetry⁹ $SO(5) \times SO(5) \sim USp(4)_L \times USp(4)_R$. The relevant

⁸(2.21) is the $n = 4$ case of the maximal non-symmetric embedding pattern

$$SO(4n) \supset_{ns} SU(2) \times USp(2n);$$

$$\mathbf{Adj}_{SO(4n)} = \mathbf{Adj}_{SU(2)} + \mathbf{Adj}_{USp(2n)} + (\mathbf{3}, \mathbf{A}_{2,0}),$$

where $\mathbf{A}_{2,0}$ is the rank-2 antisymmetric skew-traceless irrep. of $USp(2n)$. For the first appearance of such an embedding in supersymmetry, see [27].

⁹Subscripts “ L ” and “ R ” denote left and right chirality, respectively.

nm embedding reads $(SO(4) \sim SU(2) \times SU(2))$

$$E_{8(8)} \supset_s SO(8,8) \supset_s SO(5,5) \times SO(3,3) \sim SO(5,5) \times SL(4, \mathbb{R}); \quad (2.24)$$

$$SO(16) \supset_s SO(8) \times SO(8) \supset_s SO(5)_L \times SO(3) \times SO(5)_R \times SO(3) \quad (2.25)$$

$$\sim USp(4)_L \times USp(4)_R \times SU(2)_L \times SU(2)_R;$$

note the “*enhancement*” to $SO(8,8)$ in (2.24). Note that in this case both the \mathcal{R} -symmetry and massless spin groups factorize in the direct product of opposite chiralities identical factors. The corresponding Jordan algebra interpretation of (2.24) is as follows:

$$QConf(J_3^{\mathbb{O}_s}) \supset Str_0(J_2^{\mathbb{O}_s}) \times SL(4, \mathbb{R}), \quad (2.26)$$

where $J_3^{\mathbb{O}_s}$ and $J_2^{\mathbb{O}_s} \sim \Gamma_{5,5}$ are the rank-2 and rank-3 Euclidean Jordan algebras over the split octonions \mathbb{O}_s , and $QConf$ and Str_0 respectively denote the *quasi-conformal* and *reduced structure* groups¹⁰ (see e.g. [21] and refs. therein).

8. In $D = 5$ the U -duality undergoes an exceptional enhancement: $G_{16}^5 = E_{6(6)}$, and its mcs is the \mathcal{R} -symmetry $USp(8)$. The relevant embedding is maximal non-symmetric, and it is given by (1.5), which we report here (note that it is the first step of nm embedding (2.20)–(2.21)):

$$E_{8(8)} \supset_{ns} E_{6(6)} \times SL(3, \mathbb{R}); \quad (2.27)$$

$$SO(16) \supset_{ns} USp(8) \times SU(2). \quad (2.28)$$

The corresponding Jordan algebra interpretation of (2.27) is as follows:

$$QConf(J_3^{\mathbb{O}_s}) \supset Str_0(J_3^{\mathbb{O}_s}) \times SL(3, \mathbb{R}), \quad (2.29)$$

and it is a particular non-compact, real version of the *Jordan-pair* embeddings of exceptional Lie algebras recently considered in [22]. Note that the $SU(2)$ in (2.28) is the *principal* $SU(2)$ in $SL(3, \mathbb{R})$ in (2.27).

9. In $D = 4$ the U -duality is $G_{16}^4 = E_{7(7)}$, and its mcs is the \mathcal{R} -symmetry $SU(8)$. The relevant embedding is maximal symmetric (note that it is the first step of chains (2.11)–(2.12) and (2.18)–(2.19)):

$$E_{8(8)} \supset_s E_{7(7)} \times SL(2, \mathbb{R}); \quad (2.30)$$

$$SO(16) \supset_s SU(8) \times U(1). \quad (2.31)$$

The corresponding Jordan algebra interpretation of (2.30) is as follows:

$$QConf(J_3^{\mathbb{O}_s}) \supset Conf(J_3^{\mathbb{O}_s}) \times SL(2, \mathbb{R}), \quad (2.32)$$

where $Conf$ denotes the *conformal* group of $J_3^{\mathbb{O}_s}$ (see e.g. [21] and refs. therein). Similar Jordan-algebraic interpretations can be given for other supergravities in various dimensions.

¹⁰In theories related to Euclidean Jordan algebras J_3 of rank 3, the *quasi-conformal* $QConf(J_3)$, *conformal* $Conf(J_3)$ and *reduced structure* $Str_0(J_3)$ groups are the U -duality groups in $D = 3, 4$ and 5 dimensions, respectively. In particular, $Conf(J_3)$ is nothing but the automorphism group $Aut(\mathfrak{M}(J_3))$ of the corresponding Freudenthal triple system [19–21].

2.2 $N = 12$

In the “pure” theory with $N = 12$, the $D = 3$ U -duality group is $G_{12}^3 = E_{7(-5)}$, with mcs $SO(12) \times SU(2)_{CV}$, where $SO(12)$ is the Clifford algebra for massless particles with $\mathcal{N} = 24$ supersymmetries. The $SU(2)_{CV}$ factor pertains to the so-called *Clifford vacuum (CV)*, which is generally present for *non-maximal* theories ($N < 16$), and it indicates further degeneracy of the Clifford algebra symmetry. In this case, $SU(2)_{CV}$ can be also explained by recalling that this theory shares the very same bosonic sector of a *matter-coupled* supergravity with $N = 4$ [19, 20], in which it is the \mathcal{R} -symmetry of the hypermultiplets’ sector.

This theory can consistently be uplifted only to $D = 4$ and $D = 5$.

1. In $D = 5$ the U -duality is $G_{12}^5 = SU^*(6)$, and its mcs is the \mathcal{R} -symmetry $USp(6)$. The relevant embedding is maximal non-symmetric:

$$E_{7(-5)} \supset_{ns} SU^*(6) \times SL(3, \mathbb{R}); \quad (2.33)$$

$$SO(12) \times SU(2)_{CV} \supset_{ns} USp(6) \times SU(2)_J, \quad (2.34)$$

where we introduced the subscript “ J ” in order to discriminate between the Clifford vacuum $SU(2)_{CV}$ and the $SU(2)_J$ pertaining to the massless spin group in $D = 5$. Note that the embedding (2.33) is maximal non-symmetric, while the embedding (2.34) is non-maximal non-symmetric.

2. In $D = 4$ the U -duality is $G_{12}^4 = SO^*(12)$, and its mcs is the \mathcal{R} -symmetry $U(6)$. The relevant embedding is maximal symmetric:

$$E_{7(-5)} \supset_s SO^*(12) \times SL(2, \mathbb{R}); \quad (2.35)$$

$$SO(12) \times SU(2)_{CV} \supset_s SU(6) \times U(1) \times U(1)_J, \quad (2.36)$$

and it pertains to the so-called c^* -map (see e.g. [29], and refs. therein).

2.3 $N = 10$

In the “pure” theory with $N = 10$, the $D = 3$ U -duality group is $G_{10}^3 = E_{6(-14)}$, with mcs $SO(10) \times SO(2)_{CV}$, where $SO(10)$ is the Clifford algebra for massless particles with $\mathcal{N} = 20$ supersymmetries. In this case, $SO(2)_{CV}$ can be also explained as [add. . .]

This theory can be uplifted only to $D = 4$, in which the U -duality is $G_{10}^4 = SU(5, 1)$, and its mcs is the \mathcal{R} -symmetry $U(5)$. The relevant embedding is maximal symmetric:

$$E_{6(-14)} \supset_s SU(5, 1) \times SL(2, \mathbb{R}); \quad (2.37)$$

$$SO(10) \times SO(2)_{CV} \supset_s SU(5) \times U(1) \times U(1)_J. \quad (2.38)$$

3 $N = 8, 6$ matter coupled theories

3.1 $N = 8$

Half-maximal theories with $N = 8$ exist in $3 \leq D \leq 10$; moreover, for $D = 6$ two inequivalent theories exist, i.e. the non-chiral IIA (1, 1) and the chiral IIB (2, 0).

The $D = 3$ U -duality group is $G_8^3 = SO(8, D - 2 + m)$, where m is the number of matter multiplets in $D = 3$ other than those coming from the reduction of the gravity multiplet in D dimensions. Furthermore, $mcs(G_8^3) = SO(8) \times SO(D - 2 + m)_{CV}$, where $SO(8)$ is the Clifford algebra for massless particles with $\mathcal{N} = 16$ supersymmetries, and $SO(D - 2 + m)_{CV}$ is the *Clifford vacuum* symmetry.

The relevant chain of maximal embeddings leading to the embedding of the D -dimensional Ehlers group $SL(D - 2, \mathbb{R})$ into $SO(8, D - 2 + m)$ depends on the dimension and on the type of theory. We anticipate that embeddings (1.2) and (2.1) are *maximal* in $D = 4$ (symmetric) and *next-to-maximal* in $5 \leq D \leq 10$.

- For $D \geq 5$ (and $D = 6$ type IIA (1,1)), it is given by the following chain of two maximal symmetric steps:

$$\begin{aligned} SO(8, D - 2 + m) \supset_s SO(D - 2, D - 2) \times SO(10 - D, m) \\ \supset_s SL(D - 2, \mathbb{R}) \times SO(1, 1) \times SO(10 - D, m) \end{aligned} \quad (3.1)$$

The group commuting with $SL(D - 2, \mathbb{R})$ inside $SO(8, D - 2 + m)$ is nothing but the D -dimensional U -duality group $G_8^D = SO(1, 1) \times SO(10 - D, m)$. Note the “*enhancement*” to $SO(D - 2, D - 2) \times SO(10 - D, m)$. Furthermore, it is worth remarking that also for $m = 0$ the *Clifford vacuum* degeneracy is still present with an $SO(D - 2)_{CV}$ factor; this is an extra spin quantum number carried by the $SO(8)$ Clifford algebra spinor. In fact, by considering the *mcs* level of the chain (3.1), one obtains

$$\begin{aligned} SO(8)_{\text{Clifford}} \times SO(D - 2 + m)_{CV} \\ \supset_s SO(D - 2) \times SO(D - 2)_{CV} \times SO(10 - D) \times SO(m)_{CV} \\ \supset_s SO(D - 2)_J \times SO(10 - D)_{\mathcal{R}} \times SO(m)_{CV}, \end{aligned} \quad (3.2)$$

where the D -dimensional massless spin group $SO(D - 2)_J = mcs(SL(D - 2, \mathbb{R}))$ is *diagonally* embedded into $SO(D - 2) \times SO(D - 2)_{CV}$, and the \mathcal{R} -symmetry is $SO(10 - D)$. $SO(m)_{CV}$ is the part of *Clifford vacuum* symmetry due to matter coupling.

- For $D = 4$, the maximal symmetric embedding reads:

$$SO(8, 2 + m) \supset_s SO(2, 2) \times SO(6, m) \sim SL(2, \mathbb{R})_{\text{Ehlers}} \times SL(2, \mathbb{R}) \times SO(6, m), \quad (3.3)$$

and it pertains to the so-called *c*-map* (see e.g. [29], and refs. therein). The group commuting with $SL(2, \mathbb{R})_{\text{Ehlers}}$ inside $SO(8, 2 + m)$ is the 4-dimensional U -duality group $G_8^4 = SL(2, \mathbb{R}) \times SO(6, m)$. Also in this case for $m = 0$ the *Clifford vacuum* degeneracy is still present with an $SO(2)_{CV}$ factor. In fact, by considering the *mcs* level of (3.3), one obtains the following maximal symmetric embedding ($SO(6) \sim SU(4)$):

$$\begin{aligned} SO(8)_{\text{Clifford}} \times SO(2 + m)_{CV} \\ \supset_s SO(2)_J \times SO(2)_{CV} \times SO(6) \times SO(m)_{CV} \sim U(1)_J \times U(4)_{\mathcal{R}} \times SO(m)_{CV}, \end{aligned} \quad (3.4)$$

where $SO(2)_J = mcs(SL(2, \mathbb{R})_{\text{Ehlers}})$, and the \mathcal{R} -symmetry is $SO(2)_{CV} \times SO(6) \sim U(4)_{\mathcal{R}}$. Moreover, $SO(m)_{CV}$ is the part of *Clifford vacuum* symmetry due to matter coupling.

- For $D = 6$ type IIB $(2, 0)$, it suffices to start with $SO(8, 3 + m)$, and the maximal symmetric embedding reads as follows:

$$SO(8, 3 + m) \supset_s SO(3, 3) \times SO(5, m) \sim SL(4, \mathbb{R}) \times SO(5, m). \quad (3.5)$$

The group commuting with $SL(4, \mathbb{R})$ inside $SO(8, 4 + m)$ is the 6-dimensional type IIB U -duality group $G_8^{6, IIB} = SO(5, m)$. The corresponding *mcs* level reads

$$\begin{aligned} &SO(8)_{\text{Clifford}} \times SO(3 + m)_{CV} \\ &\supset_s (SO(3) \times SO(3))_J \times SO(5)_{\mathcal{R}} \times SO(m)_{CV} \sim SO(4)_J \times USp(4)_{\mathcal{R}} \times SO(m)_{CV}, \end{aligned} \quad (3.6)$$

where $SO(4)_J = mcs(SL(4, \mathbb{R})_{\text{Ehlers}})$, and the \mathcal{R} -symmetry is $SO(5) \sim USp(4)$. Furthermore, $SO(m)_{CV}$ is the part of *Clifford vacuum* symmetry due to matter coupling.

All cases in $4 \leq D \leq 10$ dimensions are reported in tables 3 and 4.

3.2 $N = 6$

Theories with $N = 6$ exist only in $D = 3, 4$.

The $D = 3$ U -duality group is $G_6^3 = SU(4, m + 1)$, where m is the number of matter multiplets in $D = 3$ other than those coming from the reduction of the gravity multiplet in 4 dimensions. Furthermore, $mcs(G_6^3) = SU(4) \times U(m + 1)_{CV}$, where $SU(4) \sim SO(6)$ is the Clifford algebra for massless particles with $\mathcal{N} = 12$ supersymmetries, and $U(m + 1)_{CV}$ is the *Clifford vacuum* symmetry.

The embedding of the 4-dimensional Ehlers group $SL(2, \mathbb{R})$ into $SU(4, m + 1)$ is maximal and symmetric:

$$SU(4, m + 1) \supset_s SL(2, \mathbb{R}) \times U(3, m), \quad (3.7)$$

and at the *mcs* level:

$$SU(4) \times SU(m + 1) \times U(1) \supset_s U(1)_J \times U(3) \times U(m), \quad (3.8)$$

where $D = 4$ U -duality group is $G_6^4 = SU(3, m)$, and the \mathcal{R} -symmetry is $U(3)$. $U(m)$ is the $D = 4$ *Clifford vacuum* symmetry, which is related to the number of matter multiplets.

4 $N = 4$ matter coupled *symmetric* theories

Quarter-maximal theories with $N = 4$ exist in $3 \leq D \leq 6$; in particular, in $D = 6$ they are chiral $(1, 0)$ theories. The new feature of $N = 4$ theories is the possible existence of two different types of matter multiplets, namely vector and hyper multiplets, transforming in different ways under the \mathcal{R} -symmetry, which is $U(2)$ in $D = 4$ and $USp(2)$ in $D = 5, 6$.

D	$SO(8, D-2+m) \supset G_D^8(m) \times SL(D-2, \mathbb{R})$	type
10	$SO(8, 8+m) \supset SO(1, 1) \times SO(m) \times SL(8, \mathbb{R})$	nm, ns
9	$SO(8, 7+m) \supset SO(1, 1) \times SO(1, m) \times SL(7, \mathbb{R})$	nm, ns
8	$SO(8, 6+m) \supset SO(1, 1) \times SO(2, m) \times SL(6, \mathbb{R})$	nm, ns
7	$SO(8, 5+m) \supset SO(1, 1) \times SO(3, m) \times SL(5, \mathbb{R})$	nm, ns
6, IIA	$SO(8, 4+m) \supset SO(1, 1) \times SO(4, m) \times SL(4, \mathbb{R})$	nm, ns
6, IIB	$SO(8, 3+m) \supset SO(5, m) \times SL(4, \mathbb{R})$	max, s
5	$SO(8, 3+m) \supset SO(1, 1) \times SO(5, m) \times SL(3, \mathbb{R})$	nm, ns
4	$SO(8, 2+m) \supset (SL(2, \mathbb{R}) \times SO(6, m)) \times SL(2, \mathbb{R})$	max, s

Table 3. Embedding $G_8^3 \supset G_8^D \times SL(D-2, \mathbb{R})$ (1.2) for *half-maximal* supergravity theories ($N = 8$) in $10 \geq D \geq 4$ Lorentzian space-time dimensions.

In the following treatment, we will only consider theories based on *symmetric* Abelian-vector multiplets' scalar manifolds, which is a restriction to $D = 4$ (Kähler) and $D = 5$ (real) special geometry; these theories will be denoted as¹¹ *symmetric* $N = 4$ theories.

In $D = 4, 5$, *symmetric* theories are classified by two infinite sequences, as well as by isolated cases given by the so-called “*magical*” models.

We will also not consider the (D -independent) hypermultiplets' quaternionic scalar manifolds.

¹¹We will not consider here the so-called *non-Jordan symmetric sequence* (see e.g. [30] and refs. therein) in $D = 5$, based on vector multiplets' real special symmetric scalar manifolds $\frac{SO(1,n)}{SO(n)}$, which gives rise to *non-symmetric* coset manifolds in $D = 4$ and in $D = 3$.

D	$SO(8) \times SO(D-2+m) \supset H_D^8(m) \times SO(D-2)$	type
10	$SO(8) \times SO(8+m) \supset SO(m) \times SO(8)$	nm, ns
9	$SO(8) \times SO(7+m) \supset SO(m) \times SO(7)$	nm, ns
8	$SO(8) \times SO(6+m) \supset SO(2) \times SO(m) \times SO(6)$	nm, ns
7	$SO(8) \times SO(5+m) \supset SO(3) \times SO(m) \times SO(5)$	nm, ns
6, IIA	$SO(8) \times SO(4+m) \supset SO(4) \times SO(m) \times SO(4)$	nm, ns
6, IIB	$SO(8) \times SO(3+m) \supset SO(5) \times SO(m) \times SO(4)$	max, s
5	$SO(8) \times SO(3+m) \supset SO(5) \times SO(m) \times SO(3)$	nm, ns
4	$SO(8) \times SO(2+m) \supset (SO(2) \times SO(6) \times SO(m)) \times SO(2)$	max, s

Table 4. Embedding $H_8^3 \supset H_8^D \times SO(D-2)$ (2.1) [23] for *half-maximal* supergravity theories ($N = 8$) in $10 \geq D \geq 4$ Lorentzian space-time dimensions. Since *matter coupling* is allowed, H_8^3 and in general H_8^D entail both *half-maximal* \mathcal{R} -symmetry and *Clifford vacuum* symmetry.

For $N = 4$, we recall that the Clifford algebra decomposes as

$$SO(4) \sim SU(2)_v \times SU(2)_h, \tag{4.1}$$

where $SU(2)_v$ pertains to the $D = 3$ reduction of $D = 4$ vector multiplets, while $SU(2)_h$ is related to the hypermultiplet sector, which is insensitive to the number of space-time dimensions in which the *quarter-maximal* theory is defined (namely, $3 \leq D \leq 6$). Since we disregard hypermultiplets, in the treatment below we only consider $SU(2)_v$ (and thus we remove the subscript “ v ”), which will be a commuting factor in the *mcs* of the $D = 3$ U -duality group G_4^4 .

4.1 Minimal coupling infinite sequence and “pure” $D = 4$ supergravity

We start by considering the infinite sequence of $D = 3$ quaternionic Kähler symmetric spaces

$$\frac{\mathrm{SU}(2, 1+n)}{\mathrm{SU}(2) \times \mathrm{SU}(1+n) \times \mathrm{U}(1)}, \quad (4.2)$$

which can be uplifted only to $D = 4$, giving rise to Maxwell-Einstein supergravity models *minimally coupled* to n vector multiplets [33]. The $D = 3$ U -duality group is $G_4^3 = \mathrm{SU}(2, 1+n)$.

The embedding of the 4-dimensional Ehlers group $SL(2, \mathbb{R})$ into $\mathrm{SU}(2, n+1)$ is maximal and symmetric:

$$\mathrm{SU}(2, 1+n) \supset_s \mathrm{SL}(2, \mathbb{R}) \times \mathrm{U}(1, n), \quad (4.3)$$

and at the *mcs* level:

$$\mathrm{SU}(2) \times \mathrm{SU}(1+n) \times \mathrm{U}(1) \supset_s \mathrm{U}(1)_J \times \mathrm{U}(n) \times \mathrm{U}(1)_{\mathcal{R}}, \quad (4.4)$$

where $D = 4$ U -duality group is $G_4^4 = \mathrm{U}(1, n)$. $\mathrm{U}(1)_{\mathcal{R}}$ in (4.4) is the part of $D = 4$ \mathcal{R} -symmetry $\mathrm{U}(2)$ under which the $D = 4$ vector multiplets are charged, whereas the $\mathrm{U}(n)$ factor correspond to $D = 4$ Clifford vacuum symmetry (completely due to *matter coupling*).

By merging (4.3) and (4.4), the following c -map is obtained [31]:

$$\mathbb{CP}^n \equiv \frac{\mathrm{SU}(1, n)}{\mathrm{U}(n)} \xrightarrow{c} \frac{\mathrm{SU}(2, 1+n)}{\mathrm{SU}(2) \times \mathrm{SU}(1+n) \times \mathrm{U}(1)}, \quad (4.5)$$

where \mathbb{CP}^n denotes the *complex projective* (non-compact) spaces.

Note that for $n = 0$ the quaternionic manifold (4.2) is not only Kähler, but also *special Kähler*, and it is an example of Einstein space with self-dual Weyl curvature (see e.g. [34], and refs. therein). It is usually called the *universal hypermultiplet*, and it corresponds to the c -map of “pure” $\mathcal{N} = 2$ supergravity in $D = 4$, obtained as “ $n = 0$ limit” of the \mathbb{CP}^n sequence; namely, by specifying $n = 0$ in (4.5) [31]:

$$\emptyset \xrightarrow{c} \frac{\mathrm{SU}(2, 1)}{\mathrm{SU}(2) \times \mathrm{U}(1)}. \quad (4.6)$$

Correspondingly, for $n = 0$ (4.3) and (4.4) respectively read

$$\mathrm{SU}(2, 1) \supset_s \mathrm{SL}(2, \mathbb{R}) \times \mathrm{U}(1) \sim \mathrm{U}(1, 1); \quad (4.7)$$

$$\mathrm{SU}(2) \times \mathrm{U}(1) \supset_s \mathrm{U}(1)_J \times \mathrm{U}(1)_{\mathcal{R}}, \quad (4.8)$$

and thus the 4 bosonic massless states of $\mathcal{N} = 2$, $D = 4$ “pure” supergravity are in the $\mathbf{2}_{\mathbb{C}}$ of $\mathrm{SU}(2) \times \mathrm{U}(1) \sim \mathrm{U}(2) = \text{mcs}(\mathrm{SU}(2, 1))$.

4.2 “Pure” $D = 5, 6$ supergravity and T^3 and ST^2 models in $D = 4$

4.2.1 $D = 5$

Within the framework under consideration, “pure” $D = 5$ supergravity can be obtained as $D = 5$ uplift of the so-called $\mathcal{N} = 2$, $D = 4$ T^3 model, whose vector multiplet’s scalar

span the symmetric special Kähler manifold $SL(2, \mathbb{R})/U(1)$ (with Ricci scalar curvature $R = -2/3$ [32]), and whose $D = 3$ U -duality group is $G_{4,T^3}^3 = G_{2(2)}$.

The embedding of the 5-dimensional Ehlers group $SL(3, \mathbb{R})$ into G_{4,T^3}^3 is maximal and *non-symmetric* (see e.g. [28] and refs. therein):

$$G_{2(2)} \supset_{ns} SL(3, \mathbb{R}), \tag{4.9}$$

and at the *mcs* level:

$$SU(2) \times SU(2) \supset_s SO(3)_J \sim SU(2)_J, \tag{4.10}$$

where the $D = 5$ massless spin group $SO(3)_J$ is *diagonally* embedded into $SU(2) \times SU(2) = mcs(G_{2(2)})$. The 8 bosonic massless states of $\mathcal{N} = 2$, $D = 5$ “*pure*” supergravity are in the $(4, \mathbf{2})$ of *mcs* ($G_{2(2)}$) itself.

By merging (4.9) and (4.10), the following c -map is obtained¹² [31]:

$$\frac{SL(2, \mathbb{R})}{U(1)} \Big|_{T^3} \xrightarrow{c} \frac{G_{2(2)}}{SU(2) \times SU(2)}. \tag{4.11}$$

The corresponding Jordan algebra interpretation of (4.9) reads

$$QConf(\mathbb{R}) \supset_s SL(3, \mathbb{R}), \tag{4.12}$$

because the T^3 model is related to the (non-generic) simple rank-3 Euclidean Jordan algebra given by the reals \mathbb{R} (see tables 5-8).

4.2.2 $D = 6$

Analogously, “*pure*” $D = 6$ $(1, 0)$ chiral supergravity¹³ can be obtained as $D = 6$ uplift of the so-called $\mathcal{N} = 2$, $D = 4$ ST^2 model, whose vector multiplets’ scalars span the symmetric special Kähler manifold $[SL(2, \mathbb{R})/U(1)]^2$, and whose $D = 3$ U -duality group is $G_{4,ST^2}^3 = SO(4, 3)$.

The embedding of the 6-dimensional Ehlers group $SL(4, \mathbb{R})$ into G_{4,ST^2}^3 is maximal and *symmetric*:

$$SO(4, 3) \supset_s SO(3, 3) \sim SL(4, \mathbb{R}), \tag{4.13}$$

and at the *mcs* level:

$$SO(4) \times SO(3) \supset_s SO(3) \times SO(3) \sim SO(4)_J, \tag{4.14}$$

where the $D = 6$ massless spin group is $SO(4)_J$. The 12 bosonic massless states of “*pure*” $D = 6$ $(1, 0)$ supergravity are in the $(4, \mathbf{3})$ of $SO(4) \times SO(3) = mcs(SO(4, 3))$.

¹²Attention should be paid to distinguish $\frac{SL(2, \mathbb{R})}{U(1)} \Big|_{T^3}$ ($R = -2/3$) from the $n = 1$ element of the \mathbb{CP}^n infinite sequence treated above, namely the \mathbb{CP}^1 space (axio-dilatonic $\mathcal{N} = 2$, $D = 4$ supergravity), which has $R = -2$. Note that $R = -2$ and $R = -2/3$ are the unique two values for which the Kähler manifold $\frac{SL(2, \mathbb{R})}{U(1)}$ is a *special* Kähler manifold [32].

¹³We here disregard the various conditions to be fulfilled for *anomaly-free* chiral supergravity theories in $D = 6$ (see e.g. [35–39]).

By merging (4.13) and (4.14), the following c -map is obtained [31]:

$$\left[\frac{\mathrm{SL}(2, \mathbb{R})}{\mathrm{U}(1)} \right]^2 \xrightarrow{c} \frac{\mathrm{SO}(4, 3)}{\mathrm{SO}(4) \times \mathrm{SO}(3)}. \quad (4.15)$$

The corresponding Jordan algebra interpretation of (4.13) reads

$$QConf(\mathbb{R} \oplus \mathbf{\Gamma}_{1,0}) \supset_s \mathrm{SL}(4, \mathbb{R}), \quad (4.16)$$

because the ST^2 model is related to the (non-generic) semi-simple rank-3 Euclidean Jordan algebra given by $\mathbb{R} \oplus \mathbf{\Gamma}_{1,0} \sim \mathbb{R} \oplus \mathbb{R}$.

4.3 The Jordan symmetric infinite sequence

The aforementioned ST^2 model is actually the first element of the so-called *Jordan symmetric sequence* of *quarter-maximal* theories.

The $D = 3$ U -duality group is $G_4^3 = \mathrm{SO}(4, D - 2 + n)$, where n is the number of matter multiplets in $D = 3$ other than those coming from the reduction of the gravity multiplet in D dimensions. Furthermore, $mcs(G_8^3) = \mathrm{SO}(4) \times \mathrm{SO}(D - 2 + n)_{CV}$; as mentioned, $\mathrm{SO}(4) \sim \mathrm{SU}(2)_v \times \mathrm{SU}(2)_h$ is the Clifford algebra for massless particles with $\mathcal{N} = 8$ supersymmetries, and $\mathrm{SO}(D - 2 + n)_{CV}$ is the *Clifford vacuum* symmetry.

Let us consider the relevant chain of maximal embeddings leading to the embedding of the D -dimensional Ehlers group¹⁴ $SL(D - 2, \mathbb{R})$ into $\mathrm{SO}(4, D - 2 + n)$.

4.3.1 $D = 6$

In $D = 6$, it suffices to start from $G_4^3 = \mathrm{SO}(4, 3 + n)$, and the corresponding maximal symmetric embedding reads

$$\mathrm{SO}(4, 3 + n) \supset_s \mathrm{SO}(3, 3) \times \mathrm{SO}(1, n) \sim \mathrm{SL}(4, \mathbb{R}) \times \mathrm{SO}(1, n), \quad (4.17)$$

and at the mcs level:

$$\mathrm{SO}(4) \times \mathrm{SO}(3 + n) \supset_s \mathrm{SO}(3) \times \mathrm{SO}(3) \times \mathrm{SO}(n) \sim \mathrm{SO}(4) \times \mathrm{SO}(n), \quad (4.18)$$

where n is the number of matter (tensor) multiplets in $D = 6$. The group commuting with $SL(4, \mathbb{R})$ inside $\mathrm{SO}(4, 3 + n)$ is nothing but the 6-dimensional U -duality group of tensor multiplets $G_4^6 = \mathrm{SO}(1, n)$.

4.3.2 $D = 5$

For $D = 5$, one branches once more from (4.17), getting:

$$\mathrm{SO}(4, 3 + n) \supset_s \mathrm{SL}(4, \mathbb{R}) \times \mathrm{SO}(1, n) \supset_s \mathrm{SL}(3, \mathbb{R}) \times \mathrm{SO}(1, 1) \times \mathrm{SO}(1, n), \quad (4.19)$$

and at the mcs level:

$$\mathrm{SO}(4) \times \mathrm{SO}(3 + n) \supset_s \mathrm{SO}(4) \times \mathrm{SO}(n) \supset_s \mathrm{SO}(3) \times \mathrm{SO}(n), \quad (4.20)$$

where $n + 1$ is the number of matter (vector) multiplets in $D = 5$. The group commuting with $SL(3, \mathbb{R})$ inside $\mathrm{SO}(4, 3 + n)$ is nothing but the 5-dimensional U -duality group $G_4^5 = \mathrm{SO}(1, 1) \times \mathrm{SO}(1, n)$. Note the “*enhancement*” to $SL(4, \mathbb{R}) \times \mathrm{SO}(1, n)$ in (4.19).

¹⁴Note that, consistently, for $n = 0$ (in $D = 5$ and $D = 6$) and $n = 1$ (in $D = 4$), one re-obtains the case of the ST^2 model treated above.

4.3.3 $D = 4$

For $D = 4$, the embedding is maximal and symmetric:

$$SO(4, 2+n) \supset_s SO(2, 2) \times SO(2, n) \sim SL(2, \mathbb{R})_{\text{Ehlers}} \times SL(2, \mathbb{R}) \times SO(2, n), \quad (4.21)$$

and at the *mcs* level:

$$SO(4) \times SO(2+n) \supset_s SO(2)_J \times SO(2) \times SO(2) \times SO(n), \quad (4.22)$$

where n is the number of matter (vector) multiplets in $D = 4$. The group commuting with $SL(2, \mathbb{R})_{\text{Ehlers}}$ inside $SO(4, 2+n)$ is nothing but the 4-dimensional U -duality group $G_4^4 = SL(2, \mathbb{R}) \times SO(2, n)$. By merging (4.21) and (4.22), one obtains the following c -map:

$$\frac{SL(2, \mathbb{R})}{U(1)} \times \frac{SO(2, n)}{SO(2) \times SO(n)} \xrightarrow{c} \frac{SO(4, n+2)}{SO(4) \times SO(n+2)}. \quad (4.23)$$

4.4 *Magical models*

Let us now consider the isolated cases of symmetric $N = 8$ *quarter-maximal* theories, the so-called *magical* models [19, 20]. They are associated to rank-2 (in $D = 6$) and rank-3 (in $D = 5$) Euclidean Jordan algebras over the four normed division algebras \mathbb{O} (octonions), \mathbb{H} (quaternions), \mathbb{C} (complex numbers) and \mathbb{R} (real numbers), and to the Freudenthal triple systems over such algebras (in $D = 4$). Consequently, they can be parametrized in terms of the real dimension of the relevant division algebra, namely $q = 8, 4, 2, 1$ for $\mathbb{O}, \mathbb{H}, \mathbb{C}$ and \mathbb{R} , respectively. In this respect, the T^3 model treated above corresponds to $q = -2/3$.

We will now analyze the relevant embeddings in $D = 4, 5$ and 6.

4.4.1 $D = 4$

In $D = 4$, the magic models are related to the Freudenthal triple system $\mathfrak{M}(J_3^{\mathbb{A}})$ over the rank-3 *simple* Euclidean Jordan algebra $J_3^{\mathbb{A}}$ ($\mathbb{A} = \mathbb{O}, \mathbb{H}, \mathbb{C}, \mathbb{R}$). The $D = 3$ and $D = 4$ U -duality groups are nothing but the *quasi-conformal* and *conformal* group of $J_3^{\mathbb{A}}$, respectively, and they are related by the following maximal *symmetric* embedding:

$$G_4^3(q) \supset_s SL(2, \mathbb{R})_{\text{Ehlers}} \times G_4^4(q), \quad (4.24)$$

with *mcs* level involving the $D = 4$ massless spin group:

$$mcs(G_4^3(q)) \supset_s SO(2)_J \times mcs(G_4^4(q)). \quad (4.25)$$

(4.24)–(4.25) correspond to the following c^* -map symmetric embedding of the corresponding scalar manifolds in $D = 3$ (para-quaternionic pseudo-Riemannian) and $D = 4$ (special Kähler):

$$\frac{G_4^4(q)}{mcs(G_4^4(q))} \xrightarrow{c^*} \frac{G_4^3(q)}{SL(2, \mathbb{R}) \times G_4^4(q)}. \quad (4.26)$$

The various cases are listed in tables 5 and 6.

$\mathfrak{M}(J_3^{\mathbb{A}})$	$G_4^3(q) \supset_s G_4^4(q) \times SL(2, \mathbb{R})$	type
$\mathfrak{M}(J_3^{\mathbb{O}}) (q = 8)$	$E_{8(-24)} \supset E_{7(-25)} \times SL(2, \mathbb{R})$	max, s
$\mathfrak{M}(J_3^{\mathbb{H}}) (q = 4)$	$E_{7(-5)} \supset SO^*(12) \times SL(2, \mathbb{R})$	max, s
$\mathfrak{M}(J_3^{\mathbb{C}}) (q = 2)$	$E_{6(2)} \supset SU(3, 3) \times SL(2, \mathbb{R})$	max, s
$\mathfrak{M}(J_3^{\mathbb{R}}) (q = 1)$	$F_{4(4)} \supset Sp(6, \mathbb{R}) \times SL(2, \mathbb{R})$	max, s
$\mathfrak{M}(\mathbb{R}) (q = -2/3)$	$G_{2(2)} \supset SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$	max, s

Table 5. Embedding $G_4^3(q) \supset G_4^4(q) \times_s SL(2, \mathbb{R})_{\text{Ehlers}}$ for *magical* Maxwell-Einstein supergravity theories ($N = 8$) in $D = 4$ Lorentzian space-time dimensions. Also the case of T^3 model ($q = -2/3$) is reported.

4.4.2 $D = 5$

In $D = 5$, the magic models are related to $J_3^{\mathbb{A}}$'s themselves. The $D = 5$ U -duality group is the *reduced structure* group of $J_3^{\mathbb{A}}$, and the embedding of the $D = 5$ Ehlers group $SL(3, \mathbb{R})$ into the $D = 3$ U -duality group is maximal and *non-symmetric*:

$$G_4^3(q) \supset_{ns} SL(3, \mathbb{R}) \times G_4^5(q), \tag{4.27}$$

with *mcs* level involving the $D = 5$ massless spin group:

$$mcs(G_4^3(q)) \supset_s SO(3)_J \times mcs(G_4^5(q)). \tag{4.28}$$

The various cases are listed in tables 7 and 8.

4.4.3 $D = 6$

In $D = 6$, the magic models are related to the rank-2 Jordan algebra $J_2^{\mathbb{A}} \sim \Gamma_{1,q+1}$ (where “ \sim ” here denotes a vector space isomorphism). Namely, the $D = 6$ U -duality group is nothing but the *reduced structure* group of $J_2^{\mathbb{A}}$ itself, with the exception of the cases corresponding to $q = 4$ and $q = 2$, which have a further factor¹⁵ $\mathcal{A}_{q=2} = SO(3)$ resp.

¹⁵We note that the non-triviality of the factor group \mathcal{A}_q in the $D = 6$ U -duality group is related to the reality properties of the spinors within the rank-2 Jordan algebras over the quaternions ($J_2^{\mathbb{H}} \sim \Gamma_{1,5}$) and

$\mathfrak{M}(J_3^{\mathbb{A}})$	$mcs(G_4^3(q)) \supset_s mcs(G_4^4(q)) \times SO(2)_J$	type
$\mathfrak{M}(J_3^{\mathbb{O}}) (q = 8)$	$E_{7(-133)} \times SU(2) \supset E_{6(-78)} \times U(1) \times SO(2)_J$	max, s
$\mathfrak{M}(J_3^{\mathbb{H}}) (q = 4)$	$SO(12) \times SU(2) \supset U(6) \times SO(2)_J$	max, s
$\mathfrak{M}(J_3^{\mathbb{C}}) (q = 2)$	$SU(6) \times SU(2) \supset S(U(3) \times U(3)) \times SO(2)_J$	max, s
$\mathfrak{M}(J_3^{\mathbb{R}}) (q = 1)$	$USp(6) \times SU(2) \supset U(3) \times SO(2)_J$	max, s
$\mathfrak{M}(\mathbb{R}) (q = -2/3)$	$SU(2) \times SU(2) \supset U(1) \times SO(2)_J$	max, s

Table 6. Embedding $mcs(G_4^3(q)) \supset_s mcs(G_4^4(q)) \times SO(2)_J$ for *magical* Maxwell-Einstein supergravity theories ($N = 8$) in $D = 4$ Lorentzian space-time dimensions. Also the case of T^3 model ($q = -2/3$) is reported.

$\mathcal{A}_{q=1} = SO(2)$ in the U -duality group. The embedding of the $D = 6$ Ehlers group $SL(4, \mathbb{R})$ into the $D = 3$ U -duality group is obtained by a two-steps chain of maximal and *symmetric* embeddings ($\mathcal{A}_q = Id, SO(3), SO(2), Id$ respectively for $q = 8, 4, 2, 1$):

$$G_4^3(q) \supset_s SO(4, q+4) \times \mathcal{A}_q \supset_s SL(4, \mathbb{R}) \times SO(1, q+1) \times \mathcal{A}_q, \quad (4.29)$$

with mcs level involving the $D = 6$ massless spin group:

$$mcs(G_4^3(q)) \supset_s SO(4)_J \times SO(q+1) \times mcs(\mathcal{A}_q). \quad (4.30)$$

Note the “*enhancement*” to $SO(4, q+4) \times \mathcal{A}_q$ in (4.29). The various cases are listed in tables 9 and 10.

5 Cosets with $\chi = 0$ and *Poincaré duality*

From the previous treatment, a class of non-compact, pseudo-Riemannian homogeneous spaces can be naturally constructed, with general structure:

$$M_N^D \equiv \frac{G_N^3}{G_N^D \times SL(D-2, \mathbb{R})}, \quad (5.1)$$

over the complex numbers ($J_2^{\mathbb{C}} \sim \Gamma_{1,3}$), which are respectively pseudo-real (quaternionic) and complex (see e.g. table 2 of [40]).

$J_3^{\mathbb{A}}$	$G_4^3(q) \supset_s G_4^5(q) \times SL(3, \mathbb{R})$	type
$J_3^{\mathbb{O}}(q=8)$	$E_{8(-24)} \supset E_{6(-26)} \times SL(3, \mathbb{R})$	max, <i>ns</i>
$J_3^{\mathbb{H}}(q=4)$	$E_{7(-5)} \supset SU^*(6) \times SL(3, \mathbb{R})$	max, <i>ns</i>
$J_3^{\mathbb{C}}(q=2)$	$E_{6(2)} \supset SL(3, \mathbb{C}) \times SL(3, \mathbb{R})$	max, <i>ns</i>
$J_3^{\mathbb{R}}(q=1)$	$F_{4(4)} \supset SL(3, \mathbb{R}) \times SL(3, \mathbb{R})$	max, <i>ns</i>
$\mathbb{R}(q=-2/3)$	$G_{2(2)} \supset SL(3, \mathbb{R})$	max, <i>ns</i>

Table 7. Embedding $G_4^3(q) \supset G_4^5(q) \times_s SL(3, \mathbb{R})_{\text{Ehlers}}$ for *magical* Maxwell-Einstein supergravity theories ($N = 8$) in $D = 5$ Lorentzian space-time dimensions. The $D = 5$ uplift of T^3 model is “*pure*” minimal supergravity

determined by the embedding of the direct product of the D -dimensional Ehlers group $SL(D - 2, \mathbb{R})$ and of the D -dimensional U -duality group G_N^D of a supergravity with $\mathcal{N} = 2N$ supersymmetries into the U -duality group of the same theory reduced to $D = 3$ (Lorentzian) space-time dimensions. From previous sections, such an embedding can be *maximal* or non-maximal (namely, *next-to-maximal*), and *symmetric* or *non-symmetric*, but, as mentioned, it always preserves the rank of the group (1.3), as well as the non-compact rank of the $D = 3$ coset G_N^3/H_N^3 (1.4).

Interestingly, the cosets M_N^D 's (5.1) all share the same feature: they have an equal number of compact and non-compact generators, thus implying the their *coset character* χ [14, 15] to be *vanishing*:

$$\chi(M_N^D) \equiv nc(M_N^D) - c(M_N^D) = 0. \tag{5.2}$$

This property can also be related to the “*mcs* counterpart” of the class of cosets (5.1), given by the compact, Riemannian homogeneous spaces with general structure

$$\widehat{M}_N^D \equiv \frac{mcs(G_N^3)}{mcs(G_N^D) \times SO(D - 2)_J}, \tag{5.3}$$

determined by the embedding of the direct product of the D -dimensional massless spin group $SO(D - 2) = mcs(SL(D - 2, \mathbb{R}))$ and of $H_N^D = mcs(G_N^D)$ into $H_N^3 = mcs(G_N^3)$. As

$J_3^{\mathbb{A}}$	$mcs(G_4^3(q)) \supset_s mcs(G_4^5(q)) \times SO(3)_J$	type
$J_3^{\mathbb{O}} (q = 8)$	$E_{7(-133)} \times SU(2) \supset F_{4(-52)} \times SO(3)_J$	max, ns
$J_3^{\mathbb{H}} (q = 4)$	$SO(12) \times SU(2) \supset USp(6) \times SO(3)_J$	max, ns
$J_3^{\mathbb{C}} (q = 2)$	$SU(6) \times SU(2) \supset SU(3) \times SO(3)_J$	max, ns
$J_3^{\mathbb{R}} (q = 1)$	$USp(6) \times SU(2) \supset SU(2)_P \times SO(3)_J$	max, ns
$\mathbb{R} (q = -2/3)$	$SU(2) \times SU(2) \supset SO(3)_{J,D}$	max, ns

Table 8. Embedding $mcs(G_4^3(q)) \supset mcs(G_4^5(q)) \times_s SO(3)_J$ for *magical* Maxwell-Einstein supergravity theories ($N = 8$) in $D = 5$ Lorentzian space-time dimensions. $SU(2)_P$ denotes the *principal* $SU(2)$, whereas the subscript “ D ” stands for *diagonal embedding*

the M_N^D 's (5.1), also the \widehat{M}_N^D 's (5.3) can be of various types, namely *maximal* or *next-to-maximal*, *symmetric* or *non-symmetric*.

However, \widehat{M}_N^D 's (5.3) all share the same property: the number of compact or non-compact generators of M_N^D 's (5.1) is always *equal* to the (real) dimension of the corresponding \widehat{M}_N^D 's themselves. This is a consequence of (5.2) as well as the general formula on the signature of a pseudo-Riemannian coset G/H (see e.g. [15])

$$\begin{aligned} c(G/H) &= \dim_{\mathbb{R}}(mcs(G)) - \dim_{\mathbb{R}}(mcs(H)); \\ nc(G/H) &= \dim_{\mathbb{R}}(G) - \dim_{\mathbb{R}}(H) - c(G/H), \end{aligned} \tag{5.4}$$

from which thus follows that the compact generators of M_N^D are the very generators of the corresponding \widehat{M}_N^D :

$$nc(M_N^D) = c(M_N^D) = \dim_{\mathbb{R}}(\widehat{M}_N^D) \tag{5.5}$$

Along this line, further elaboration is possible. Indeed, it generally holds that

$$\dim_{\mathbb{R}} \left[\frac{G_N^3}{G_N^D \times SL(D-2, \mathbb{R})} \right] = 2 \dim_{\mathbb{R}} \left[\frac{H_N^3}{H_N^D \times SO(D-2)} \right]. \tag{5.6}$$

A possible interpretation of these results is as follows. In a supergravity theory in D space-time (Lorentzian) dimensions, the number of bosonic massless degrees of freedom other than the scalar and graviton ones is given by the difference between the dimension

$J_2^{\mathbb{A}}$	$G_4^3(q) \supset \text{SO}(1, q+1) \times A_q \times \text{SL}(4, \mathbb{R})$	type
$J_2^{\mathbb{O}} (q=8)$	$E_{8(-24)} \supset \text{SO}(1, 9) \times \text{SL}(4, \mathbb{R})$	nm, ns
$J_2^{\mathbb{H}} (q=4)$	$E_{7(-5)} \supset \text{SO}(1, 5) \times \text{SO}(3) \times \text{SL}(4, \mathbb{R})$	nm, ns
$J_2^{\mathbb{C}} (q=2)$	$E_{6(2)} \supset \text{SO}(1, 3) \times \text{SO}(2) \times \text{SL}(4, \mathbb{R})$	nm, ns
$J_2^{\mathbb{R}} (q=1)$	$F_{4(4)} \supset \text{SO}(1, 2) \times \text{SL}(4, \mathbb{R})$	nm, ns

Table 9. Embedding $G_4^3(q) \supset_{ns} G_4^6(q) \times \text{SL}(4, \mathbb{R})_{\text{Ehlers}}$ ($G_4^6(q) = \text{SO}(1, q+1) \times A_q$) for chiral *magical* Maxwell-Einstein supergravity theories ($N=8$) in $D=6$ Lorentzian space-time dimensions. Recall $\text{SO}(1, 5) \sim \text{SU}^*(4)$, $\text{SO}(1, 3) \sim \text{SL}(3, \mathbb{C})$, $\text{SO}(1, 2) \sim \text{SL}(2, \mathbb{R})$.

$J_2^{\mathbb{A}}$	$mcs(G_4^3(q)) \supset mcs(G_4^6(q)) \times \text{SO}(4)_J$	type
$J_2^{\mathbb{O}} (q=8)$	$E_{7(-133)} \times \text{SU}(2) \supset \text{SO}(9) \times \text{SO}(4)_J$	nm, ns
$J_2^{\mathbb{H}} (q=4)$	$\text{SO}(12) \times \text{SU}(2) \supset \text{SO}(5) \times \text{SO}(3) \times \text{SO}(4)_J$	nm, ns
$J_2^{\mathbb{C}} (q=2)$	$\text{SU}(6) \times \text{SU}(2) \supset \text{SO}(3) \times \text{SO}(2) \times \text{SO}(4)_J$	nm, ns
$J_2^{\mathbb{R}} (q=1)$	$\text{USp}(6) \times \text{SU}(2) \supset \text{SO}(2) \times \text{SO}(4)_J$	nm, ns

Table 10. Embedding $mcs(G_4^3(q)) \supset_{ns} mcs(G_4^6(q)) \times \text{SO}(4)_J$ for chiral *magical* Maxwell-Einstein supergravity theories ($N=8$) in $D=6$ Lorentzian space-time dimensions.

of the Clifford algebra and the sum of the dimensions of the D -dimensional massless spin group and of the D -dimensional “Clifford symmetry” (i.e., \mathcal{R} -symmetry + Clifford vacuum degeneracy due to matter coupling, if any).

Section 5.1 lists the cosets M_N^D 's (5.1) and their “mcs counterparts” \widehat{M}_N^D 's (5.3) for all N 's and D 's treated in the present investigation. Then, in sections 5.2 and 5.3 an interpretation of the vanishing character (5.2) will be given in terms of *Poincaré duality*, or equivalently of *Hodge involution* acting on the cohomology of M_N^D 's.

5.1 The cosets

5.1.1 $N = 16$

The specification of (5.1) and (5.3) to *maximal supergravity* ($N = 16$) give rise the following spaces

$$M_{16}^D \equiv \frac{G_{16}^3}{G_{16}^D \times \text{SL}(D-2, \mathbb{R})} = \frac{E_{8(8)}}{G_{16}^D \times \text{SL}(D-2, \mathbb{R})}; \quad (5.7)$$

$$\widehat{M}_{16}^D \equiv \frac{H_{16}^3}{H_{16}^D \times \text{SO}(D-2)} = \frac{\text{SO}(16)}{H_{16}^D \times \text{SO}(D-2)}; \quad (5.8)$$

they are listed in table 11, along with their number of compact and non-compact generators. Among M_{16}^D 's, the unique maximal and *symmetric* coset is the one pertaining to $D = 4$ (cfr. (2.30)):

$$M_{16}^4 \equiv \frac{G_{16}^3}{G_{16}^4 \times \text{SL}(2, \mathbb{R})} = \frac{E_{8(8)}}{E_{7(7)} \times \text{SL}(2, \mathbb{R})}, \quad (5.9)$$

which is a rank-4 *para-quaternionic* space, as resulting from the classification of [41]. Also the corresponding

$$\widehat{M}_{16}^4 = \frac{\text{SO}(16)}{\text{SU}(8) \times \text{SO}(2)} \quad (5.10)$$

is a maximal and *symmetric* space among \widehat{M}_{16}^D 's.

5.1.2 $N = 12$

The specification of (5.1) and (5.3) to supergravity with $N = 12$ in $D = 5$ and in $D = 4$ respectively reads

$$M_{12}^5 \equiv \frac{G_{12}^3}{G_{12}^5 \times \text{SL}(3, \mathbb{R})} = \frac{E_{7(-5)}}{SU^*(6) \times \text{SL}(3, \mathbb{R})}, \quad c = nc = 45; \quad (5.11)$$

$$\widehat{M}_{12}^5 \equiv \frac{H_{12}^3}{H_{12}^5 \times \text{SO}(3)_J} = \frac{\text{SO}(12) \times \text{SU}(2)}{USp(6) \times \text{SO}(3)_J}; \quad (5.12)$$

$$M_{12}^4 \equiv \frac{G_{12}^3}{G_{12}^4 \times \text{SL}(2, \mathbb{R})} = \frac{E_{7(-5)}}{SO^*(12) \times \text{SL}(2, \mathbb{R})}, \quad c = nc = 32; \quad (5.13)$$

$$\widehat{M}_{12}^4 \equiv \frac{H_{12}^3}{H_{12}^4 \times \text{SO}(2)} = \frac{\text{SO}(12) \times \text{SU}(2)}{\text{SU}(6) \times \text{U}(1) \times \text{SO}(2)_J}. \quad (5.14)$$

They all are maximal cosets, but M_{12}^5 and \widehat{M}_{12}^5 are non-symmetric, whereas M_{12}^4 and \widehat{M}_{12}^4 are symmetric.

The values of $c = nc$ given in (5.11) and (5.13) match the ones of the magical quarter-maximal ($N = 4$) theory for $q = 4$ (see (5.40) and (5.44), respectively); indeed, these theories share the same bosonic sector, and they are both related to J_3^{H} .

D	M_{16}^D	\widehat{M}_{16}^D	$c(M_{16}^D) = nc(M_{16}^D)$
11	$\frac{E_{8(8)}}{\text{SL}(9, \mathbb{R})}$	$\frac{\text{SO}(16)}{\text{SO}(9)}$	84
10, <i>IIA</i>	$\frac{E_{8(8)}}{\text{SO}(1,1) \times \text{SL}(8, \mathbb{R})}$	$\frac{\text{SO}(16)}{\text{SO}(8)}$	92
10, <i>IIB</i>	$\frac{E_{8(8)}}{\text{SL}(2, R) \times \text{SL}(8, \mathbb{R})}$	$\frac{\text{SO}(16)}{\text{SO}(2) \times \text{SO}(8)}$	91
9	$\frac{E_{8(8)}}{\text{GL}(2, R) \times \text{SL}(7, \mathbb{R})}$	$\frac{\text{SO}(16)}{\text{SO}(2) \times \text{SO}(7)}$	98
8	$\frac{E_{8(8)}}{(\text{SL}(2, \mathbb{R}) \times \text{SL}(3, \mathbb{R})) \times \text{SL}(6, \mathbb{R})}$	$\frac{\text{SO}(16)}{(\text{SO}(2) \times \text{SO}(3)) \times \text{SO}(6)}$	101
7	$\frac{E_{8(8)}}{\text{SL}(5, \mathbb{R}) \times \text{SL}(5, \mathbb{R})}$	$\frac{\text{SO}(16)}{\text{SO}(5) \times \text{SO}(5)}$	100
6	$\frac{E_{8(8)}}{\text{SO}(5,5) \times \text{SL}(4, \mathbb{R})}$	$\frac{\text{SO}(16)}{\text{SO}(5) \times \text{SO}(5) \times \text{SO}(4)}$	94
5	$\frac{E_{8(8)}}{E_{6(6)} \times \text{SL}(3, \mathbb{R})}$	$\frac{\text{SO}(16)}{\text{USp}(8) \times \text{SO}(3)}$	81
4	$\frac{E_{8(8)}}{E_{7(7)} \times \text{SL}(2, \mathbb{R})}$	$\frac{\text{SO}(16)}{\text{SU}(8) \times \text{SO}(2)}$	56

Table 11. Pseudo-Riemannian non-compact $E_{8(8)}$ -cosets M_{16}^D (5.7) and Riemannian compact $\text{SO}(16)$ -cosets \widehat{M}_{16}^D (5.8) of *maximal* supergravity theories ($N = 16$) in $11 \geq D \geq 4$ Lorentzian space-time dimensions. The number of compact generators c (equal to the number nc of non-compact generators) of M_{16}^D is also listed. All cosets M_{16}^D have *vanishing character*.

5.1.3 $N = 10$

The specification of (5.1) and (5.3) to supergravity with $N = 10$ in $D = 4$ gives rise to the following symmetric spaces

$$M_{10}^5 \equiv \frac{G_{10}^3}{G_{10}^4 \times SL(2, \mathbb{R})} = \frac{E_{6(-14)}}{SU(5, 1) \times SL(2, \mathbb{R})}, \quad c = nc = 20; \quad (5.15)$$

$$\widehat{M}_{10}^4 \equiv \frac{H_{10}^3}{H_{10}^4 \times SO(2)} = \frac{SO(10) \times U(1)}{SU(5) \times U(1) \times U(1)_J}. \quad (5.16)$$

M_{10}^5 is a rank-4 *para-quaternionic* coset.

5.1.4 $N = 8$

The specification of (5.1) and (5.3) to *half-maximal supergravity* ($N = 8$) gives rise to the following spaces

$$M_8^D \equiv \frac{G_8^3}{G_8^D \times SL(D-2, \mathbb{R})} = \frac{SO(8, D-2+m)}{G_8^D \times SL(D-2, \mathbb{R})}; \quad (5.17)$$

$$\widehat{M}_8^D \equiv \frac{H_8^3}{H_8^D \times SO(D-2)} = \frac{SO(8) \times SO(D-2+m)}{H_8^D \times SO(D-2)}; \quad (5.18)$$

they are listed in table 12, along with their number of compact and non-compact generators.

Among M_8^D 's, the unique maximal and *symmetric* cosets are the ones pertaining to $D = 6$ IIB and $D = 4$ (cfr. (3.3)):

$$M_8^{6, IIB} \equiv \frac{G_8^3}{G_8^{6, IIB} \times SL(2, \mathbb{R})} = \frac{SO(8, 3+m)}{SO(5, m) \times SL(4, \mathbb{R})}; \quad (5.19)$$

$$M_8^4 \equiv \frac{G_8^3}{G_8^4 \times SL(2, \mathbb{R})} = \frac{SO(8, 2+m)}{SL(2, \mathbb{R}) \times SO(6, m) \times SL(2, \mathbb{R})}. \quad (5.20)$$

5.1.5 $N = 6$

The specification of (5.1) and (5.3) to supergravity with $N = 6$ in $D = 4$ gives rise to the following symmetric spaces

$$M_6^4 \equiv \frac{G_6^3}{G_6^4 \times SL(2, \mathbb{R})} = \frac{SU(4, m+1)}{SU(3, m) \times SL(2, \mathbb{R})}, \quad c = nc = 2m + 7; \quad (5.21)$$

$$\widehat{M}_6^4 \equiv \frac{H_6^3}{H_6^4 \times SO(2)} = \frac{SU(4) \times SU(m+1) \times U(1)}{U(3) \times U(m) \times U(1)_J}. \quad (5.22)$$

5.1.6 $N = 4$ *symmetric*

Minimal Coupling. The specification of (5.1) and (5.3) to *minimally coupled* Maxwell-Einstein supergravity with $N = 4$ in $D = 4$ gives rise to the following symmetric spaces

$$M_4^4 \equiv \frac{G_4^3}{G_4^4 \times SL(2, \mathbb{R})} = \frac{SU(2, 1+n)}{U(1, n) \times SL(2, \mathbb{R})}, \quad c = nc = 2n + 2; \quad (5.23)$$

$$\widehat{M}_4^4 \equiv \frac{H_4^3}{H_4^4 \times SO(2)} = \frac{SU(2) \times SU(1+n) \times U(1)}{U(n) \times U(1) \times U(1)}. \quad (5.24)$$

D	M_8^D	\widehat{M}_8^D	$c(M_8^D) = nc(M_8^D)$
10	$\frac{SO(8,8+m)}{(SO(1,1) \times SO(m)) \times SL(8, \mathbb{R})}$	$\frac{SO(8) \times SO(8+m)}{SO(m) \times SO(8)}$	$8m + 28$
9	$\frac{SO(8,7+m)}{(SO(1,1) \times SO(1,m)) \times SL(7, \mathbb{R})}$	$\frac{SO(8) \times SO(7+m)}{SO(m) \times SO(7)}$	$7m + 28$
8	$\frac{SO(8,6+m)}{(SO(1,1) \times SO(2,m)) \times SL(6, \mathbb{R})}$	$\frac{SO(8) \times SO(6+m)}{(SO(2) \times SO(m)) \times SO(6)}$	$6m + 27$
7	$\frac{SO(8,5+m)}{(SO(1,1) \times SO(3,m)) \times SL(5, \mathbb{R})}$	$\frac{SO(8) \times SO(5+m)}{(SO(3) \times SO(m)) \times SO(5)}$	$5m + 25$
6, <i>IIA</i>	$\frac{SO(8,4+m)}{(SO(1,1) \times SO(4,m)) \times SL(4, \mathbb{R})}$	$\frac{SO(8) \times SO(4+m)}{(SO(4) \times SO(m)) \times SO(4)}$	$4m + 22$
6, <i>IIB</i>	$\frac{SO(8,3+m)}{SO(5,m) \times SL(4, \mathbb{R})}$	$\frac{SO(8) \times SO(3+m)}{(SO(5) \times SO(m)) \times SO(4)}$	$3m + 15$
5	$\frac{SO(8,3+m)}{(SO(1,1) \times SO(5,m)) \times SL(3, \mathbb{R})}$	$\frac{SO(8) \times SO(3+m)}{(SO(5) \times SO(m)) \times SO(3)}$	$3m + 18$
4	$\frac{SO(8,2+m)}{(SL(2, \mathbb{R}) \times SO(6,m)) \times SL(2, \mathbb{R})}$	$\frac{SO(8) \times SO(2+m)}{(SO(2) \times SO(6) \times SO(m)) \times SO(2)}$	$2m + 12$

Table 12. Pseudo-Riemannian non-compact M_8^D (5.17) and Riemannian compact cosets \widehat{M}_8^D (5.18) of *half-maximal* supergravity theories ($N = 8$) in $10 \geq D \geq 4$ Lorentzian space-time dimensions. The number of compact generators c (equal to the number nc of non-compact generators) of M_8^D is also listed. All cosets M_8^D have *vanishing character*.

M_4^4 has rank 1 for $n = 0$, and rank 2 for $n \geq 1$, and it is *para-quaternionic*. It is nothing but a suitable pseudo-Riemannian form of the manifold (4.2) itself, namely the c^* -map of the rank-1 symmetric *special Kähler* maximal coset in $D = 4$:

$$\mathbb{CP}^n \equiv \frac{SU(1, n)}{U(n)} \xrightarrow{c^*} \frac{SU(2, 1 + n)}{U(1, n) \times SL(2, \mathbb{R})}. \tag{5.25}$$

T^3 Model. The specification of (5.1) and (5.3) to the so-called T^3 model in $D = 4$ gives rise to the following symmetric spaces

$$M_{4,T^3}^4 \equiv \frac{G_{4,T^3}^3}{G_{4,T^3}^4 \times \text{SL}(2, \mathbb{R})} = \frac{G_{2(2)}}{\text{SL}(2, \mathbb{R}) \times \text{SL}(2, \mathbb{R})_{\text{Ehlers}}}, \quad c = nc = 4; \quad (5.26)$$

$$\widehat{M}_{4,T^3}^4 \equiv \frac{mcs(G_{4,T^3}^3)}{H_{4,T^3}^4 \times \text{SO}(2)} = \frac{\text{SU}(2) \times \text{SU}(2)}{\text{U}(1) \times \text{U}(1)}. \quad (5.27)$$

M_{4,T^3}^4 is rank-2 *para-quaternionic*. It is nothing but a suitable pseudo-Riemannian form of the manifold in the r.h.s. of (4.11), namely the c^* -map of the rank-1 symmetric *special Kähler* maximal coset in $D = 4$:

$$\left. \frac{\text{SL}(2, \mathbb{R})}{\text{U}(1)} \right|_{T^3} \xrightarrow{c^*} \frac{G_{2(2)}}{\text{SL}(2, \mathbb{R}) \times \text{SL}(2, \mathbb{R})}. \quad (5.28)$$

ST^2 Model. The specification of (5.1) and (5.3) to the so-called ST^2 model in $D = 4$ gives rise to the following symmetric spaces

$$M_{4,ST^2}^4 \equiv \frac{G_{4,ST^2}^3}{G_{4,ST^2}^4 \times \text{SL}(2, \mathbb{R})_{\text{Ehlers}}} = \frac{\text{SO}(4, 3)}{\text{SL}(2, \mathbb{R}) \times \text{SL}(2, \mathbb{R}) \times \text{SL}(2, \mathbb{R})_{\text{Ehlers}}}, \quad c = nc = 6; \quad (5.29)$$

$$\widehat{M}_{4,ST^2}^4 \equiv \frac{mcs(G_{4,ST^2}^3)}{H_{4,ST^2}^4 \times \text{SO}(2)} = \frac{\text{SO}(4) \times \text{SO}(3)}{\text{U}(1) \times \text{U}(1) \times \text{U}(1)}. \quad (5.30)$$

Jordan symmetric sequence. As mentioned above, the ST^2 model can be regarded as the first element of the so-called *Jordan symmetric sequence* of *quarter-maximal* theories. The specification of (5.1) and (5.3) to such a sequence in $D = 6$, $D = 5$ and $D = 4$ respectively gives rise to the following spaces:

$D = 6$:

$$M_4^6 \equiv \frac{G_4^3}{G_4^6 \times \text{SL}(4, \mathbb{R})} = \frac{\text{SO}(4, 3+n)}{\text{SO}(1, n) \times \text{SL}(4, \mathbb{R})}, \quad c = nc = 3n + 3; \quad (5.31)$$

$$\widehat{M}_4^6 \equiv \frac{H_4^3}{H_4^6 \times \text{SO}(4)} = \frac{\text{SO}(4) \times \text{SO}(3+n)}{\text{SO}(3) \times \text{SO}(n) \times \text{SO}(3)}; \quad (5.32)$$

M_4^6 and \widehat{M}_4^6 are maximal and *symmetric* spaces.

$D = 5$:

$$M_4^5 \equiv \frac{G_4^3}{G_4^5 \times \text{SL}(3, \mathbb{R})} = \frac{\text{SO}(4, 3+n)}{\text{SO}(1, 1) \times \text{SO}(1, n) \times \text{SL}(3, \mathbb{R})}, \quad c = nc = 3n + 6; \quad (5.33)$$

$$\widehat{M}_4^5 \equiv \frac{H_4^3}{H_4^5 \times \text{SO}(3)} = \frac{\text{SO}(4) \times \text{SO}(3+n)}{\text{SO}(n) \times \text{SO}(3)}; \quad (5.34)$$

M_4^5 and \widehat{M}_4^5 are non-maximal and *non-symmetric* spaces.

$D = 4$:

$$M_4^4 \equiv \frac{G_4^3}{G_4^4 \times \text{SL}(2, \mathbb{R})_{\text{Ehlers}}} = \frac{SO(4, 2+n)}{\text{SL}(2, \mathbb{R}) \times \text{SO}(2, n) \times \text{SL}(2, \mathbb{R})_{\text{Ehlers}}}, \quad c = nc = 2n + 4; \quad (5.35)$$

$$\widehat{M}_4^4 \equiv \frac{mcs(G_4^3)}{H_4^4 \times \text{SO}(2)} = \frac{\text{SO}(4) \times \text{SO}(2+n)}{\text{U}(1) \times \text{U}(1) \times \text{SO}(n) \times \text{U}(1)}. \quad (5.36)$$

M_4^4 and \widehat{M}_4^4 are maximal and *symmetric* spaces. M_4^4 is *para-quaternionic* and it has rank 2 in the case $n = 0$ and rank 3 for $n \geq 1$; it is nothing but a suitable pseudo-Riemannian form of the manifold in the r.h.s. of (4.23), namely the c^* -map of the symmetric *special Kähler* maximal coset in $D = 4$:

$$\frac{\text{SL}(2, \mathbb{R})}{\text{U}(1)} \times \frac{\text{SO}(2, n)}{\text{SO}(2) \times \text{SO}(n)} \xrightarrow{c^*} \frac{SO(4, 2+n)}{\text{SL}(2, \mathbb{R}) \times \text{SO}(2, n) \times \text{SL}(2, \mathbb{R})_{\text{Ehlers}}}. \quad (5.37)$$

Magical models.

$D = 4$: The specification of (5.1) and (5.3) to *magical* models in $D = 4$ gives rise to maximal symmetric spaces. Their general structure reads

$$M_4^4(q) \equiv \frac{G_4^3(q)}{G_4^4(q) \times \text{SL}(2, \mathbb{R})_{\text{Ehlers}}}; \quad (5.38)$$

$$\widehat{M}_4^4(q) \equiv \frac{H_4^3(q)}{H_4^4(q) \times \text{SO}(2)}, \quad (5.39)$$

listed in table 13. The number of compact and non-compact generators of $M_4^4(q)$ can be q -parametrized as follows:

$$c(M_4^4(q)) = nc(M_4^4(q)) = 6q + 8 = \dim_{\mathbb{R}}(\mathbf{R}(G_4^4(q))), \quad (5.40)$$

where \mathbf{R} is the symplectic irrep. of the $D = 4$ U -duality group $G_4^4(q)$ in which the Abelian two-form field strengths sit; see subsection 5.2 for further analysis. Thus, the split of the generators of $M_4^4(q)$ into a signature $(nc, c = nc)$ is consistent with the *Ehlers-doublet* irrep. $(\mathbf{R}, \mathbf{2})$ of $G_4^4(q) \times \text{SL}(2, \mathbb{R})_{\text{Ehlers}}$. Moreover, $M_4^4(q)$ is a rank-4 *pseudo-quaternionic* space, given by the c^* -map of the corresponding symmetric *special Kähler* maximal coset in $D = 4$:

$$\frac{G_4^4(q)}{mcs(G_4^4(q))} \xrightarrow{c^*} \frac{G_4^3(q)}{G_4^4(q) \times \text{SL}(2, \mathbb{R})}. \quad (5.41)$$

$D = 5$: The specification of (5.1) and (5.3) to *magical* models in $D = 5$ gives rise to the maximal, *non-symmetric* spaces listed in table 14. Their general structure reads

$$M_4^5(q) \equiv \frac{G_4^3(q)}{G_4^5(q) \times \text{SL}(3, \mathbb{R})}; \quad (5.42)$$

$$\widehat{M}_4^5(q) \equiv \frac{H_4^3(q)}{H_4^5(q) \times \text{SO}(3)}. \quad (5.43)$$

The number of compact and non-compact generators of $M_4^5(q)$ can be q -parametrized as follows:

$$c(M_4^5(q)) = nc(M_4^5(q)) = 9(q+1) = \dim_{\mathbb{R}}(\mathcal{R}, \mathbf{3}), \quad (5.44)$$

where $(\mathcal{R}, \mathbf{3})$ is the irrep. of $G_4^5(q) \times \text{SL}(3, \mathbb{R})_{\text{Ehlers}}$. Thus, the split of the generators of $M_4^5(q)$ into a signature $(nc, c = nc)$ is consistent with a pair of *Jordan-triplet* irreps. $(\mathcal{R}, \mathbf{3})$ (see subsection 5.2 for further analysis).

$D = 6$: The specification of (5.1) and (5.3) to *magical* models in $D = 6$ respectively gives rise to the non-maximal, *non-symmetric* spaces listed in table 15.¹⁶ Their general structure reads

$$M_4^6(q) \equiv \frac{G_4^3(q)}{G_4^6(q) \times \text{SL}(4, \mathbb{R})}; \quad (5.45)$$

$$\widehat{M}_4^6(q) \equiv \frac{H_4^3(q)}{H_4^6(q) \times \text{SO}(4)}, \quad (5.46)$$

where the U -duality group $G_4^6(q)$ in $D = 6$ reads $\text{SO}(1, q+1) \times \mathcal{A}_q$. The number of compact and non-compact generators of $M_4^6(q)$ can be q -parametrized as follows:

$$c(M_4^6(q)) = nc(M_4^6(q)) = 11q + 6. \quad (5.47)$$

The meaning of $11q + 6$ and the covariant split in terms of irreps. of $\text{SO}(1, q+1) \times mcs(\mathcal{A}_q) \times \text{SO}(4)$ will be discussed in subsection 5.2.

5.2 Poincaré duality

We are now going to analyze the signature split of the manifolds M_N^D (5.1), focussing on the maximal ($N = 32$) and magical quarter-maximal cases ($N = 8$).

Nicely, the split signature of M_N^D covariantly decomposes under $mcs(G_N^D) \times \text{SO}(D-2)_J$ into a pair of sets of irreps., which are related by *Poincaré duality* (alias electric-magnetic duality). In other words, the signature of the pseudo-Riemannian manifolds M_N^D 's naturally arrange the spectrum of $p > 0$ massless forms of the corresponding supergravity theory into a pair of sets of irreps. of $mcs(G_N^D) \times \text{SO}(D-2)_J$, which are interchanged under *Poincaré duality*.

As a consequence, the $\chi = 0$ feature of the manifolds M_N^D (5.1) is actually *Poincaré-duality-invariant* (or, equivalently, *electric-magnetic duality-invariant*).

5.2.1 $N = 16$

1. $D = 11$ (M -theory): the relevant manifold is maximal non-symmetric:

$$M_{16}^{11} = \frac{E_{8(8)}}{SL(9, \mathbb{R})} : \begin{bmatrix} & c & nc \\ E_{8(8)} : & 120 & 128 \\ SL(9, \mathbb{R}) : & 36 & 44 \\ M_{16}^{11} : & 84 & 84 \end{bmatrix}. \quad (5.48)$$

¹⁶Note that the results on $c = nc$ for $q = 8$ (magical *exceptional* supergravity) in $D = 4, 5, 6$ match the results holding for maximal supergravity in the same dimensions. This is not surprising, because maximal ($N = 16$) and exceptional ($N = 4$) theories are respectively related to $J_3^{\text{O}s}$ and J_3^{O} , the unique difference given by the split vs. division form of the octonionic algebra \mathbb{O} .

$\mathfrak{M}(J_3^A)$	$M_4^4(q)$	$\widehat{M}_4^4(q)$	$c(M_4^4(q)) = nc(M_4^4(q))$
$\mathfrak{M}(J_3^{\mathbb{O}}) (q = 8)$	$\frac{E_{8(-24)}}{E_{7(-25)} \times SL(2, \mathbb{R})}$	$\frac{E_{7(-133)} \times SU(2)}{E_{6(-78)} \times U(1) \times SO(2)}$	56
$\mathfrak{M}(J_3^{\mathbb{H}}) (q = 4)$	$\frac{E_{7(-5)}}{SO^*(12) \times SL(2, \mathbb{R})}$	$\frac{SO(12) \times SU(2)}{SU(6) \times U(1) \times SO(2)}$	32
$\mathfrak{M}(J_3^{\mathbb{C}}) (q = 2)$	$\frac{E_{6(2)}}{SU(3,3) \times SL(2, \mathbb{R})}$	$\frac{SU(6) \times SU(2)}{S(U(3) \times U(3)) \times SO(2)}$	20
$\mathfrak{M}(J_3^{\mathbb{R}}) (q = 1)$	$\frac{F_{4(4)}}{Sp(6, \mathbb{R}) \times SL(2, \mathbb{R})}$	$\frac{USp(6) \times SU(2)}{SU(3) \times U(1) \times SO(2)}$	14
$\mathfrak{M}(\mathbb{R}) (q = -2/3)$	$\frac{G_{2(2)}}{SL(2, \mathbb{R}) \times SL(2, \mathbb{R})}$	$\frac{SU(2) \times SU(2)}{U(1) \times SO(2)_J}$	4

Table 13. Pseudo-Riemannian, non-compact, maximal, *para-quaternionic symmetric* cosets $M_4^4(q)$ (5.38) and Riemannian, compact, maximal cosets $\widehat{M}_4^4(q)$ (5.39) of *magic quarter-maximal* supergravity theories ($N = 4$) in $D = 4$ Lorentzian space-time dimensions. Also the T^3 model ($q = -2/3$) is reported. The number of compact generators c (equal to the number nc of non-compact generators) of $M_4^4(q)$ is also listed. All cosets $M_4^4(q)$ have *vanishing character*.

Such a signature splitting is covariant with respect to $SO(9) = mcs(SL(9, \mathbb{R}))$:

$$\begin{aligned} E_{8(8)} \supset_{ns} SL(9, \mathbb{R}); \\ \mathbf{248} = \mathbf{80} + \mathbf{84} + \mathbf{84}'; \end{aligned} \tag{5.49}$$

$$\begin{aligned} SL(9, \mathbb{R}) \xrightarrow{mcs} SO(9); \\ \mathbf{84}^{(')} = \mathbf{84}. \end{aligned} \tag{5.50}$$

Therefore, the split $(c, nc) = (84, 84)$ can be interpreted as the split into two Poincaré-dual $\mathbf{84}$'s of $SO(9)$; namely, the 3-form potential (coupled to $M2$ -brane) and its *Poincaré dual* 6-form potential (coupled to $M5$ -brane):

$$(c, nc) = (84, 84) = \mathbf{84}_{M2} + \mathbf{84}_{M5} \text{ of } SO(9). \tag{5.51}$$

2. $D = 10$ IIA: the relevant manifold is non-maximal and non-symmetric:

$$M_{16}^{10 \text{ IIA}} = \frac{E_{8(8)}}{SO(1, 1) \times SL(8, \mathbb{R})} : \begin{bmatrix} & c & nc \\ E_{8(8)} : & 120 & 128 \\ SO(1, 1) \times SL(8, \mathbb{R}) : & 28 & 36 \\ M_{16}^{10 \text{ IIA}} : & 92 & 92 \end{bmatrix}. \tag{5.52}$$

J_3^A	$M_4^5(q)$	$\widehat{M}_4^5(q)$	$c(M_4^5(q)) = nc(M_4^5(q))$
$J_3^{\mathbb{O}}(q=8)$	$\frac{E_{8(-24)}}{E_{6(-26)} \times SL(3, \mathbb{R})}$	$\frac{E_{7(-133)} \times SU(2)}{F_{4(-52)} \times SU(2) \times SO(3)_J}$	81
$J_3^{\mathbb{H}}(q=4)$	$\frac{E_{7(-5)}}{SU^*(6) \times SL(3, \mathbb{R})}$	$\frac{SO(12) \times SU(2)}{USp(6) \times SO(3)}$	45
$J_3^{\mathbb{C}}(q=2)$	$\frac{E_{6(2)}}{SL(3, \mathbb{C}) \times SL(3, \mathbb{R})}$	$\frac{SU(6) \times SU(2)}{SU(3) \times SO(3)}$	27
$J_3^{\mathbb{R}}(q=1)$	$\frac{F_{4(4)}}{SL(3, \mathbb{R}) \times SL(3, \mathbb{R})}$	$\frac{USp(6) \times SU(2)}{SU(2)_P \times SO(3)_J}$	18
$\mathbb{R}(q=-2/3)$	$G_{2(2)} \supset SL(3, \mathbb{R})$	$\frac{SU(2) \times SU(2)}{SO(3)_{J,D}}$	3

Table 14. Pseudo-Riemannian, non-compact, maximal, *non-symmetric* cosets $M_4^5(q)$ (5.42) and Riemannian, compact, maximal cosets $\widehat{M}_4^5(q)$ (5.43) of *magic quarter-maximal* supergravity theories ($N = 4$) in $D = 5$ Lorentzian space-time dimensions. Also the $D = 5$ uplift of T^3 model ($q = -2/3$), namely *minimal “pure”* supergravity, is reported. The number of compact generators c (equal to the number nc of non-compact generators) of $M_4^5(q)$ is also listed. All cosets $M_4^5(q)$ have *vanishing character*.

Such a signature splitting is covariant with respect to $SO(8) = mcs(SO(1, 1) \times SL(8, \mathbb{R}))$. Indeed, disregarding $SO(1, 1)$ weights, it holds that:

$$E_{8(8)} \supset_{nm} SO(1, 1) \times SL(8, \mathbb{R}); \tag{5.53}$$

$$\mathbf{248} = \mathbf{63} + \mathbf{1} + \mathbf{8} + \mathbf{8}' + \mathbf{28} + \mathbf{28}' + \mathbf{56} + \mathbf{56}';$$

$$SL(8, \mathbb{R}) \xrightarrow{mcs} SO(8); \tag{5.54}$$

$$\mathbf{8}^{(l)}, \mathbf{28}^{(l)}, \mathbf{56}^{(l)} = \mathbf{8}_v, \mathbf{28}, \mathbf{56}_v.$$

Therefore, the split $(c, nc) = (92, 92)$ can be interpreted as the split into two sets of Poincaré-dual irreps. of $SO(8)$; namely, the graviphoton $C_\mu^{(1)} \mathbf{8}_v$, the 2-form $B_{\mu\nu} \mathbf{28}$, the 3-form $C_{\mu\nu\rho}^{(3)} \mathbf{56}_v$ potentials, and their *Poincaré duals*, namely the 7-form $\widetilde{C}_{\mu_1 \dots \mu_7}$, 6-form $\widetilde{B}_{\mu_1 \dots \mu_6}$ and 5-form $\widetilde{C}_{\mu_1 \dots \mu_5}$ potentials:

$$(c, nc) = (92, 92) = \left(\begin{matrix} \mathbf{8}_v \\ C^{(1)} \end{matrix} + \begin{matrix} \mathbf{28} \\ B^{(2)} \end{matrix} + \begin{matrix} \mathbf{56}_v \\ C^{(3)} \end{matrix} \right) + \left(\begin{matrix} \mathbf{8}_v \\ \widetilde{C}^{(7)} \end{matrix} + \begin{matrix} \mathbf{28} \\ \widetilde{B}^{(6)} \end{matrix} + \begin{matrix} \mathbf{56}_v \\ \widetilde{C}^{(5)} \end{matrix} \right) \text{ of } SO(8). \tag{5.55}$$

J_2^A	$M_4^6(q)$	$\widehat{M}_4^6(q)$	$c(M_4^6(q)) = nc(M_4^6(q))$
$J_2^\circ (q = 8)$	$\frac{E_{8(-24)}}{SO(1,9) \times SL(4, \mathbb{R})}$	$\frac{E_{7(-133)} \times SU(2)}{SO(9) \times SO(4)_J}$	94
$J_2^{\mathbb{H}} (q = 4)$	$\frac{E_{7(-5)}}{SO(1,5) \times SO(3) \times SL(4, \mathbb{R})}$	$\frac{SO(12) \times SU(2)}{SO(5) \times SO(3) \times SO(4)_J}$	50
$J_2^{\mathbb{C}} (q = 2)$	$\frac{E_{6(2)}}{SO(1,3) \times SO(2) \times SL(4, \mathbb{R})}$	$\frac{SU(6) \times SU(2)}{SO(3) \times SO(2) \times SO(4)_J}$	28
$J_2^{\mathbb{R}} (q = 1)$	$\frac{F_{4(4)}}{SO(1,2) \times SL(4, \mathbb{R})}$	$\frac{USp(6) \times SU(2)}{SO(2) \times SO(4)_J}$	17

Table 15. Pseudo-Riemannian, non-compact, non-maximal, *non-symmetric* cosets $M_4^6(q)$ (5.45) and Riemannian, compact, non-maximal, *non-symmetric* cosets $\widehat{M}_4^6(q)$ (5.46) of *magic* (1, 0) chiral supergravity theories ($N = 4$) in $D = 6$ Lorentzian space-time dimensions. The number of compact generators c (equal to the number nc of non-compact generators) of $M_4^6(q)$ is also listed. All cosets $M_4^6(q)$ have *vanishing character*.

3. $D = 10$ IIB: the relevant manifold is non-maximal and non-symmetric:

$$M_{16}^{10 \text{ IIB}} = \frac{E_{8(8)}}{SL(2, \mathbb{R}) \times SL(8, \mathbb{R})} : \begin{bmatrix} & c & nc \\ E_{8(8)} : & 120 & 128 \\ SL(2, \mathbb{R}) \times SL(8, \mathbb{R}) : & 29 & 37 \\ M_{16}^{10 \text{ IIB}} : & 91 & 91 \end{bmatrix}. \quad (5.56)$$

Such a signature splitting is covariant with respect to $SO(8) \times SO(2) = mcs(SL(8, \mathbb{R}) \times SL(2, \mathbb{R}))$. Indeed, it holds that:

$$E_{8(8)} \supset_{nm} SL(8, \mathbb{R}) \times SL(2, \mathbb{R}); \quad (5.57)$$

$$\mathbf{248} = (\mathbf{63}, \mathbf{1}) + (\mathbf{1}, \mathbf{3}) + (\mathbf{70}, \mathbf{1}) + (\mathbf{28}, \mathbf{2}) + (\mathbf{28}', \mathbf{2});$$

$$SL(8, \mathbb{R}) \times SL(2, \mathbb{R}) \supset^{mcs} SO(8) \times SO(2);$$

$$(\mathbf{8}, \mathbf{1}) = (\mathbf{8}_v, \mathbf{1})$$

$$(\mathbf{28}^{(\prime)}, \mathbf{2}) = (\mathbf{28}, \mathbf{2}); \quad (5.58)$$

$$(\mathbf{70}, \mathbf{1}) = (\mathbf{35}_s, \mathbf{1}) + (\mathbf{35}_c, \mathbf{1}).$$

Therefore, the split $(c, nc) = (91, 91)$ can be interpreted as the split into two sets of Poincarè-dual irreps. of $SO(8) \times SO(2)$; namely, the 2-form $C_{\mu\nu}^{(2)}$ ($\mathbf{28}, \mathbf{2}$) and 4-form $C_{\mu_1 \dots \mu_4}^{(4)}$ ($\mathbf{35}_s, \mathbf{1}$) potentials, and their *Poincarè duals*, namely the 6-form $\widetilde{C}_{\mu_1 \dots \mu_6}$

$(\mathbf{28}, \mathbf{2})$ and the 4-form $C_{\mu_1 \dots \mu_4}^{(4)} (\mathbf{35}_c, \mathbf{1})$ potentials:

$$(c, nc) = (91, 91) = \left(\begin{matrix} (\mathbf{28}, \mathbf{2}) \\ C^{(2)} \end{matrix} + \begin{matrix} (\mathbf{35}_s, \mathbf{1}) \\ C^{(4)} \end{matrix} \right) + \left(\begin{matrix} (\mathbf{28}, \mathbf{2}) \\ \tilde{C}^{(6)} \end{matrix} + \begin{matrix} (\mathbf{35}_c, \mathbf{1}) \\ C^{(4)} \end{matrix} \right) \text{ of } SO(8) \times SO(2). \quad (5.59)$$

4. $D = 9$: the relevant manifold is non-maximal and non-symmetric:

$$M_{16}^9 = \frac{E_{8(8)}}{GL(2, \mathbb{R}) \times SL(7, \mathbb{R})} : \begin{bmatrix} & c & nc \\ E_{8(8)} : & 120 & 128 \\ GL(2, \mathbb{R}) \times SL(7, \mathbb{R}) : & 22 & 30 \\ M_{16}^9 : & 98 & 98 \end{bmatrix}. \quad (5.60)$$

Such a signature splitting is covariant with respect to $SO(7) \times SO(2) = mcs(SL(7, \mathbb{R}) \times GL(2, \mathbb{R}))$. Indeed, disregarding $SO(1, 1)$ weights, it holds that:

$$\begin{aligned} E_{8(8)} \supset_{nm} SL(7, \mathbb{R}) \times GL(2, \mathbb{R}); \\ \mathbf{248} = (\mathbf{48}, \mathbf{1}) + (\mathbf{1}, \mathbf{1}) + (\mathbf{1}, \mathbf{3}) \\ + (\mathbf{7}, \mathbf{1}) + (\mathbf{7}', \mathbf{1}) + (\mathbf{7}, \mathbf{2}) + (\mathbf{7}', \mathbf{2}) + (\mathbf{21}, \mathbf{2}) + (\mathbf{21}', \mathbf{2}) + (\mathbf{35}, \mathbf{1}) + (\mathbf{35}', \mathbf{1}); \end{aligned} \quad (5.61)$$

$$\begin{aligned} SL(7, \mathbb{R}) \supset_{mcs} SO(7); \\ (\mathbf{7}^{(l)}, \mathbf{21}^{(l)}, \mathbf{35}) = (\mathbf{7}, \mathbf{21}, \mathbf{35}). \end{aligned} \quad (5.62)$$

Therefore, the split $(c, nc) = (98, 98)$ can be interpreted as the split into two sets of Poincarè-dual irreps. of $SO(7) \times SO(2)$; namely, the graviphotons $(\mathbf{7}, \mathbf{1})$ and $(\mathbf{7}, \mathbf{2})$, the 2-form $(\mathbf{21}, \mathbf{2})$ and the 3-form $(\mathbf{35}, \mathbf{1})$ potentials, and their *Poincarè duals*, namely the 6-forms $(\mathbf{7}, \mathbf{1})$ and $(\mathbf{7}, \mathbf{2})$ duals of graviphotons, the 5-form $(\mathbf{21}, \mathbf{2})$ and the 4-form $(\mathbf{35}, \mathbf{1})$ potentials:

$$(c, nc) = (98, 98) = \begin{cases} (\mathbf{7}, \mathbf{1}) + (\mathbf{7}, \mathbf{2}) + (\mathbf{21}, \mathbf{2}) + (\mathbf{35}, \mathbf{1}) \\ + \\ (\mathbf{7}, \mathbf{1}) + (\mathbf{7}, \mathbf{2}) + (\mathbf{21}, \mathbf{2}) + (\mathbf{35}, \mathbf{1}) \end{cases} \text{ of } SO(7) \times SO(2). \quad (5.63)$$

5. $D = 8$: the relevant manifold is non-maximal and non-symmetric:

$$M_{16}^8 = \frac{E_{8(8)}}{SL(3, \mathbb{R}) \times SL(2, \mathbb{R}) \times SL(6, \mathbb{R})} : \begin{bmatrix} & c & nc \\ E_{8(8)} : & 120 & 128 \\ SL(3, \mathbb{R}) \times SL(2, \mathbb{R}) \times SL(6, \mathbb{R}) : & 19 & 27 \\ M_{16}^8 : & 101 & 101 \end{bmatrix}. \quad (5.64)$$

Such a signature splitting is covariant with respect to

$$SO(6) \times SO(2) \times SO(3) = mcs(SL(6, \mathbb{R}) \times SL(2, \mathbb{R}) \times SL(3, \mathbb{R})). \quad (5.65)$$

Therefore, the split $(c, nc) = (81, 81)$, which is related to the so-called *Jordan pairs* (see e.g. [22]), can be interpreted as the split into two sets of Poincarè-dual irreps. of $\text{SO}(3) \times \text{USp}(8)$; namely, the 27 graviphotons A_μ (**3, 27**), and their *Poincarè duals*, namely the 27 2-forms $\tilde{A}_{\mu\nu}$ (**3, 27**):

$$(c, nc) = (81, 81) = (\mathbf{3}, \mathbf{27}) + (\mathbf{3}, \mathbf{27}) \text{ of } \text{SO}(3) \times \text{USp}(8). \quad (5.81)$$

Note that the **3** of the massless spin group $\text{SO}(3) \equiv \text{SO}(3)_J$ corresponds to the three physical polarizations of the graviphotons in $D = 5$.

9. $D = 4$: the relevant manifold is para-quaternionic, maximal and symmetric:

$$M_{16}^4 = \frac{E_{8(8)}}{E_{7(7)} \times \text{SL}(2, \mathbb{R})} : \begin{bmatrix} & c & nc \\ E_{8(8)} : & 120 & 128 \\ E_{7(7)} \times \text{SL}(2, \mathbb{R}) : & 64 & 72 \\ M_{16}^4 : & 56 & 56 \end{bmatrix}. \quad (5.82)$$

Such a signature splitting is covariant with respect to $\text{SU}(8) \times \text{SO}(2)_J = mcs(E_{7(7)} \times \text{SL}(2, \mathbb{R}))$. Indeed, it holds that:

$$\begin{aligned} E_{8(8)} \supset_{ns} \text{SL}(2, \mathbb{R}) \times E_{7(7)}; \\ \mathbf{248} = (\mathbf{3}, \mathbf{1}) + (\mathbf{1}, \mathbf{133}) + (\mathbf{2}, \mathbf{56}); \end{aligned} \quad (5.83)$$

$$\begin{aligned} \text{SL}(2, \mathbb{R}) \times E_{7(7)} \supset_{mcs} \text{SO}(2)_J \times \text{SU}(8); \\ (\mathbf{2}, \mathbf{56}) = (\mathbf{2}, \mathbf{28}) + (\mathbf{2}, \overline{\mathbf{28}}). \end{aligned} \quad (5.84)$$

Therefore, the split $(c, nc) = (56, 56)$, which corresponds to a pair of Freudenthal systems $\mathfrak{M}(J_3^{\text{O}_s})$, can be interpreted as the split into two sets of Poincarè-dual irreps. of $\text{SO}(2)_J \times \text{SU}(8)$; namely, the 28 graviphotons A_μ (**2, 28**), and their *Poincarè-Hodge duals*, namely the 28 graviphotons \tilde{A}_μ (**2, $\overline{28}$**):

$$(c, nc) = (56, 56) = (\mathbf{2}, \mathbf{28}) + (\mathbf{2}, \overline{\mathbf{28}}) \text{ of } \text{SO}(2)_J \times \text{SU}(8). \quad (5.85)$$

Note that the **2** of the massless spin group $\text{SO}(2)_J$ corresponds to the two physical polarizations of the graviphotons in $D = 4$.

5.2.2 $N = 4$ *Magical* models

$D = 4$: the relevant manifold is para-quaternionic, maximal and symmetric (recall (5.38) and (5.40)):

$$M_4^4(q) \equiv \frac{G_4^3(q)}{G_4^4(q) \times \text{SL}(2, \mathbb{R})_{\text{Ehlers}}} : (c, nc) = (6q + 8, 6q + 8). \quad (5.86)$$

Such a signature splitting is covariant with respect to $mcs(G_4^4(q)) \times \text{SO}(2)_J$. Indeed, it holds that:

$$\begin{aligned} G_4^3(q) \supset_s \text{SL}(2, \mathbb{R}) \times G_4^4(q); \\ \text{Adj}_{G_4^3} = (\mathbf{3}, \mathbf{1}) + (\mathbf{1}, \text{Adj}_{G_4^4}) + (\mathbf{2}, \mathbf{R}); \end{aligned} \quad (5.87)$$

$$\begin{aligned} \text{SL}(2, \mathbb{R}) \times G_4^4 \supset_{mcs} \text{SO}(2)_J \times mcs(G_4^4); \\ (\mathbf{2}, \mathbf{R}) = (\mathbf{2}, \mathbf{1}) + (\mathbf{2}, \mathcal{R}) + (\mathbf{2}, \mathbf{1}) + (\mathbf{2}, \overline{\mathcal{R}}), \end{aligned} \quad (5.88)$$

where the bar here denotes the conjugate irrep. \mathbf{R} (dim= $6q+8$) denotes the relevant symplectic irrep. of G_4^4 into which the vectors sit, and \mathcal{R} ($\overline{\mathcal{R}}$) is its electric (magnetic) $D=5$ counterpart, of dimension $3q+3$. The irrep. \mathbf{R} is given by:¹⁷

$$\begin{array}{l} q : \quad 8 \quad 4 \quad 2 \quad 1 \quad -2/3 \\ G_4^4 : E_{7(-25)} \quad SO^*(12) \quad SU(3,3) \quad Sp(6, \mathbb{R}) \quad SL(2, \mathbb{R}) \\ \mathbf{R} : \quad \mathbf{56} \quad \mathbf{32}' \quad \mathbf{20} \quad \mathbf{14}' \quad \mathbf{4} \end{array} \quad (5.89)$$

On the other hand, the irrep. \mathcal{R} is given by:

$$\begin{array}{l} q : \quad 8 \quad 4 \quad 2 \quad 1 \quad -2/3 \\ G_4^5 : E_{6(-26)} \quad SU^*(6) \quad SL(3, \mathbb{C}) \quad SL(3, \mathbb{R}) \quad SL(2, \mathbb{R}) \\ \mathcal{R} : \quad \mathbf{27} \quad \mathbf{15} \quad \mathbf{9} = (\mathbf{3}, \overline{\mathbf{3}}) \quad \mathbf{6}' \quad \mathbf{1} \end{array} \quad (5.90)$$

Therefore, the split of signature of $M_4^4(q)$, which corresponds to a pair of Freudenthal systems $\mathfrak{M}(J_3^A)$, can be interpreted as the split into two sets of Poincarè-dual irreps. of $SO(2)_J \times mcs(G_4^4)$; namely, the $D=4$ graviphoton $A_\mu(\mathbf{2}, \mathbf{1})$ and the $3q+3$ matter vectors $(\mathbf{2}, \mathcal{R})$, and their *Poincarè duals*, namely the graviphoton $A_\mu(\mathbf{2}, \mathbf{1})$ and the $3q+3$ matter vectors $(\mathbf{2}, \overline{\mathcal{R}})$:

$$(c, nc) = (6q+8, 6q+8) = ((\mathbf{2}, \mathbf{1}) + (\mathbf{2}, \mathcal{R})) + ((\mathbf{2}, \mathbf{1}) + (\mathbf{2}, \overline{\mathcal{R}})) \text{ of } SO(2)_J \times mcs(G_4^4). \quad (5.91)$$

Note that the $\mathbf{2}$ of the massless spin group $SO(2)_J$ corresponds to the two physical polarizations of the graviphotons.

$D=5$: the relevant manifold is para-quaternionic, maximal and non-symmetric (recall (5.42) and (5.44)):

$$M_4^5(q) \equiv \frac{G_4^3(q)}{G_4^5(q) \times SL(3, \mathbb{R})} : (c, nc) = (9(q+1), 9(q+1)). \quad (5.92)$$

Such a signature splitting is covariant with respect to $mcs(G_4^5) \times SO(3)$. Indeed, it holds that:

$$\begin{aligned} G_4^3(q) \supset_s SL(3, \mathbb{R}) \times G_4^5(q); \\ \mathbf{Adj}_{G_4^3} = (\mathbf{8}, \mathbf{1}) + (\mathbf{1}, \mathbf{Adj}_{G_4^5}) + (\mathbf{3}, \mathcal{R}) + (\mathbf{3}', \mathcal{R}'); \end{aligned} \quad (5.93)$$

$$\begin{aligned} SL(3, \mathbb{R}) \times G_4^5(q) \supset^{mcs} SO(3) \times mcs(G_4^5); \\ (\mathbf{3}^{(\prime)}, \mathcal{R}^{(\prime)}) = (\mathbf{3}, \mathbf{1}) + (\mathbf{3}, \mathfrak{A}), \end{aligned} \quad (5.94)$$

where the prime here denotes the non-compact analogue of conjugation. \mathfrak{A} (dim= $3q+2$) denotes the relevant irrep. of $mcs(G_4^5)$ into which the $D=5$ matter vectors sit. Therefore, the split of signature of $M_4^5(q)$, which corresponds to a *Jordan pair* (see e.g. [22]), can be interpreted as the split into two sets of Poincarè-dual irreps. of $SO(3)_J \times mcs(G_4^5)$; namely, the $D=5$ graviphoton $A_\mu(\mathbf{3}, \mathbf{1})$ and the $3q+2$ matter

¹⁷Actually, in the case $q=4$, $\mathbf{32}'$ is the *conjugate* of the irreps. $\mathbf{32}$ in which the vectors sit; see appendix B for further detail.

vectors $(\mathbf{3}, \mathfrak{A})$, and their *Poincaré duals*, namely the graviphoton A_μ $(\mathbf{3}, \mathbf{1})$ and the $3q + 2$ matter vectors $(\mathbf{2}, \mathfrak{A})$:

$$\begin{aligned} (c, nc) &= (9(q+1), 9(q+1)) \\ &= ((\mathbf{3}, \mathbf{1}) + (\mathbf{3}, \mathfrak{A})) + ((\mathbf{3}, \mathbf{1}) + (\mathbf{3}, \mathfrak{A})) \text{ of } \text{SO}(3) \times mcs(G_4^5). \end{aligned} \quad (5.95)$$

Note that the $\mathbf{3}$ of the massless spin group $\text{SO}(3) \equiv \text{SO}(3)_J$ corresponds to the three physical polarizations of the vectors in $D = 5$.

$D = 6$: the relevant manifold is maximal and non-symmetric (recall (5.45) and (5.47)):

$$M_4^6(q) \equiv \frac{G_4^3(q)}{\text{SO}(1, q+1) \times \mathcal{A}_q \times \text{SL}(4, \mathbb{R})} : (c, nc) = (11q+6, 11q+6). \quad (5.96)$$

Such a signature splitting is covariant with respect to $\text{SO}(q+1) \times mcs(\mathcal{A}_q) \times \text{SO}(4)$. Indeed, it holds that:

$$\begin{aligned} G_4^3(q) &\supset_s \text{SL}(4, \mathbb{R}) \times \text{SO}(1, q+1) \times \mathcal{A}_q; \\ \text{Adj}_{G_4^3} &= (\mathbf{15}, \mathbf{1}, \mathbf{1}) + (\mathbf{1}, \text{Adj}_{\text{SO}(1, q+1)}, \mathbf{1}) + (\mathbf{1}, \mathbf{1}, \text{Adj}_{\mathcal{A}_q}) \\ &+ (\mathbf{4}, \mathbf{Spin}, \mathbf{2}) + (\mathbf{4}', \mathbf{Spin}', \mathbf{2}) + (\mathbf{6}, \mathbf{q} + \mathbf{2}, \mathbf{1}); \end{aligned} \quad (5.97)$$

$$\begin{aligned} \text{SL}(4, \mathbb{R}) \times \text{SO}(1, q+1) \times \mathcal{A}_q &\stackrel{mcs}{\supset} \text{SU}(2) \times \text{SU}(2) \times \text{SO}(q+1) \times mcs(\mathcal{A}_q); \\ (\mathbf{4}^{(l)}, \mathbf{Spin}^{(l)}, \mathbf{2}) &= (\mathbf{2}, \mathbf{2}, \mathbf{Spin}, \mathbf{2}), \\ (\mathbf{6}, \mathbf{q} + \mathbf{2}, \mathbf{1}) &= (\mathbf{3}, \mathbf{1}, \mathbf{q} + \mathbf{1}, \mathbf{1}) + (\mathbf{1}, \mathbf{3}, \mathbf{q} + \mathbf{1}, \mathbf{1}) + (\mathbf{3}, \mathbf{1}, \mathbf{1}, \mathbf{1}) + (\mathbf{1}, \mathbf{3}, \mathbf{1}, \mathbf{1}). \end{aligned} \quad (5.98)$$

In (5.97), \mathbf{Spin} , \mathbf{Spin}' and $\mathbf{q} + \mathbf{2}$ respectively denote the two conjugate chiral (semi)spinors and the vector irreps. of $\text{SO}(1, q+1) \sim \text{SL}(2, \mathbb{A})$ (with $\mathbb{A} = \mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$ for $q = 1, 2, 4, 8$, respectively), whereas in the right-hand side of (5.98) \mathbf{Spin} and $\mathbf{q} + \mathbf{1}$ respectively denote the spinor and vector irreps. of $\text{SO}(q+1)$. The irrep. \mathbf{Spin} of $\text{SO}(q+1)$ is given by:

$$\begin{array}{rcccc} q : & 8 & 4 & 2 & 1 \\ \text{SO}(q+1) : & \text{SO}(9) & \text{SO}(5) & \text{SO}(3) & \text{SO}(2) \\ \mathbf{Spin} : & \mathbf{16} & \mathbf{4} & \mathbf{2} & \mathbf{2} \end{array} \quad (5.99)$$

Thus, through these branchings, the resulting pair of Poincaré-dual irreps. of $\text{SU}(2) \times \text{SU}(2) \times \text{SO}(q+1) \times mcs(\mathcal{A}_q)$ irreps. is composed by: *i*) the physical polarizations $(\mathbf{2}, \mathbf{2}, \mathbf{Spin}, \mathbf{2})$ of massless 1-forms and the physical polarizations of 2-forms, which (under the assumption of gravity sector to be anti-self-dual) split into $(\mathbf{1}, \mathbf{3}, \mathbf{1}, \mathbf{1})$ (anti-self-dual gravity sector) and $(\mathbf{3}, \mathbf{1}, \mathbf{q} + \mathbf{1}, \mathbf{1})$ (self-dual matter sector); *ii*) the physical polarizations $(\mathbf{2}, \mathbf{2}, \mathbf{Spin}, \mathbf{2})$ of massless 3-forms and the physical polarizations of Poincaré-dual 2-forms, which split into $(\mathbf{3}, \mathbf{1}, \mathbf{1}, \mathbf{1})$ (self-dual Poincaré-dual gravity sector) and $(\mathbf{1}, \mathbf{3}, \mathbf{q} + \mathbf{1}, \mathbf{1})$ (anti-self-dual Poincaré-dual matter sector). The real dimension of each set of such irreps. can be computed as (here square brackets denote the integer part)

$$2^{\lfloor q/2 \rfloor + 2 + (1 - \delta_{q,8})} + 3(q+2) = 11q + 6, \quad (5.100)$$

and thus corresponds to the signature split of $M_4^6(q)$ in terms of irreps. of $SU(2) \times SU(2) \times SO(q+1) \times mcs(\mathcal{A}_q)$:

$$(c, nc) = (11q + 6, 11q + 6) = \begin{cases} (\mathbf{2}, \mathbf{2}, \mathbf{Spin}, \mathbf{2}) + (\mathbf{1}, \mathbf{3}, \mathbf{1}, \mathbf{1}) + (\mathbf{3}, \mathbf{1}, \mathbf{q} + \mathbf{1}, \mathbf{1}) \\ + \\ (\mathbf{2}, \mathbf{2}, \mathbf{Spin}, \mathbf{2}) + (\mathbf{3}, \mathbf{1}, \mathbf{1}, \mathbf{1}) + (\mathbf{1}, \mathbf{3}, \mathbf{q} + \mathbf{1}, \mathbf{1}). \end{cases} \quad (5.101)$$

5.3 Hodge involution and coset cohomology

A general property of the cosets M_N^D 's (5.1) resides in the fact that the *Hodge involution*¹⁸

$$* : \Lambda^d \mapsto *\Lambda^d = \Lambda^{D-2-d} \quad (5.102)$$

acts as a symmetry of the coset cohomology, where the differential forms of order d are associated to d -fold antisymmetric irreps. Λ^d of $SO(D-2) = mcs(SL(D-2, \mathbb{R}))$.

Note that, out of the possible forms belonging to the cohomology of $SO(D-2) = mcs(SL(D-2, \mathbb{R}))$, the coset M_N^D (5.1) precisely singles out the physical massless $p > 0$ forms of the corresponding supergravity theory with $\mathcal{N} = 2N$ supersymmetries in D (Lorentzian) space-time dimensions. Indeed, by casting the Cartan decomposition of the cosets M_N^D 's (5.1) in manifestly $SO(D-2)$ -covariant way, the Lie algebra of M_N^D itself branches as

$$\mathfrak{g}_3^N \ominus (\mathfrak{g}_D^N \oplus \mathfrak{sl}(D-2, \mathbb{R})) \sim \sum_d n_d \Lambda^d + \sum_d n_d *\Lambda^d, \quad (5.103)$$

manifestly $SO(D-2)$ -cov.

where \mathfrak{g}_3^N and \mathfrak{g}_D^N respectively are the Lie algebras of G_3^N and G_D^N , and n_d is the (real) dimension of the relevant irreps. of the U -duality group G_D^N in D dimensions. Note that the r.h.s. of (5.103) is manifestly invariant under the Hodge involution $*$ (5.102). Thus, the vanishing character (5.2) of cosets M_N^D 's (5.1) trivially follows from

$$c(M_N^D) = \sum_d n_d \binom{D-2}{d} = \sum_d n_d \binom{D-2}{D-2-d} = nc(M_N^D). \quad (5.104)$$

By recalling formula (5.5), $c(M_N^D) = nc(M_N^D)$ can also be computed as the real dimension of the “*mcs* counterparts” of cosets M_N^D 's (5.1), namely of the cosets \widehat{M}_N^D (5.3).

In maximal theories ($N = 16$), by recalling the embedding (2.2) and table 2, one can trace back the fact that only d -fold antisymmetric irreps. Λ^d 's occur in (5.103) to the embedding

$$\begin{aligned} SO(16) \supset \mathcal{R}_D^{16} \times SO(D-2)_J \\ \mathbf{Adj}_{SO(16)} \equiv \mathbf{16} \times_a \mathbf{16} = \mathbf{Adj}_{SO(D-2)} + \sum_d n_d \Lambda^d. \end{aligned} \quad (5.105)$$

¹⁸The involutive or anti-involutive property $*^2\Lambda^d = \pm\Lambda^d$ generally depends on the signature and the dimension of the group manifold whose cohomology is under consideration. In this case, the relevant group is $SO(D-2) = mcs(SL(D-2, \mathbb{R}))$, and thus $*^2\Lambda^d = \Lambda^d$ for $D-2 = 4n$, while $*^2\Lambda^d = -\Lambda^d$ for $D-2 = 4n+2$ ($n \in \mathbb{N}$).

Namely, in $SO(D - 2)_J$ the antisymmetric rank-2 tensor product of spinor irreps. only contain antisymmetric d -fold irreps. (see e.g. [26]). We will consider here three examples, namely $D = 11$ and $D = 10$ (type IIA and IIB).

(I) In maximal supergravity ($N = 16$) in $D = 11$, $d = 3$ and $n_d = 1$, thus (5.103) and (5.104) specifies as follows:

$$\mathfrak{e}_{8(8)} \ominus \mathfrak{sl}(9, \mathbb{R}) \sim \Lambda^3 + * \Lambda^3 = \Lambda^3 + \Lambda^6 = \mathbf{84} + \mathbf{84}; \tag{5.106}$$

manifestly $SO(9)$ -cov.

$$c \left(\frac{E_{8(8)}}{SL(9, \mathbb{R})} \right) = \binom{9}{3} = \binom{9}{6} = nc \left(\frac{E_{8(8)}}{SL(9, \mathbb{R})} \right) = 84 = \dim_{\mathbb{R}} \left(\frac{SO(16)}{SO(9)} \right). \tag{5.107}$$

In terms of the Cartan decomposition of the maximal non-symmetric Riemannian compact coset $\widehat{M}_{16}^{11} = SO(16)/SO(9)$, the result (5.107) can be obtained as a consequence of the maximal non-symmetric embedding¹⁹ (cfr. (2.2) and table 2)

$$\begin{aligned} \mathfrak{so}(16) \supset_{ns} \mathfrak{so}(9) \\ \mathbf{16} = \mathbf{16} \\ \mathbf{Adj}_{SO(16)} \equiv (\mathbf{16} \times \mathbf{16})_a = \mathbf{Adj}_{SO(9)} + \Lambda_{\mathbf{84}}^3. \end{aligned} \tag{5.108}$$

$\mathbf{120}$ $\mathbf{36}$

(II) In maximal $D = 10$ IIA supergravity, the relevant values are $d = 1, 2, 3$ with $n_1 = n_2 = n_3 = 1$, and thus (5.103) and (5.104) specifies as follows:

$$\begin{aligned} \mathfrak{e}_{8(8)} \ominus (\mathfrak{sl}(8, \mathbb{R}) \oplus \mathfrak{so}(1, 1)) \\ \sim \Lambda^1 + \Lambda^2 + \Lambda^3 + * \Lambda^1 + * \Lambda^2 + * \Lambda^3 = \Lambda^1 + \Lambda^2 + \Lambda^3 + \Lambda^7 + \Lambda^6 + \Lambda^5 \tag{5.109} \\ \text{manifestly } SO(8)\text{-cov.} \\ = (\mathbf{8}_v + \mathbf{28} + \mathbf{56}_v) + (\mathbf{8}_v + \mathbf{28} + \mathbf{56}_v); \\ c \left(\frac{E_{8(8)}}{SO(1, 1) \times SL(8, \mathbb{R})} \right) = 8 + \binom{8}{2} + \binom{8}{3} = \binom{8}{7} + \binom{8}{6} + \binom{8}{5} \\ = nc \left(\frac{E_{8(8)}}{SO(1, 1) \times SL(8, \mathbb{R})} \right) = 92 = \dim_{\mathbb{R}} \left(\frac{SO(16)}{SO(8)} \right). \end{aligned} \tag{5.110}$$

In terms of the Cartan decomposition of the non-maximal non-symmetric Riemannian compact coset $\widehat{M}_{16}^{10 IIA} = SO(16)/SO(8)$, the result (5.110) can be obtained as a

¹⁹The embedding (5.108) actually follow from a Theorem due to Dynkin [42, 43], applied to the *self-conjugate* spinor irrep. $\mathbf{16}$ of $SO(9)$:

$$SO(9) : \mathbf{16} \times_s \mathbf{16} = \Lambda^0 + \Lambda^1 + \Lambda^4 = \mathbf{1} + \mathbf{9} + \mathbf{126}.$$

consequence of the next-to-maximal non-symmetric embedding ($\mathbf{Adj} = \Lambda^2$; (cfr. (2.2) and table 2)

$$\begin{aligned}
 \mathfrak{so}(16) &\supset_{ns} \mathfrak{so}(8) \\
 \mathbf{16} &= \mathbf{8}_s + \mathbf{8}_c \\
 \mathbf{Adj}_{\mathbf{SO}(16)} &\equiv (\mathbf{16} \times \mathbf{16})_a = (\mathbf{8}_s + \mathbf{8}_c) \times_a (\mathbf{8}_s + \mathbf{8}_c) \\
 &\quad \mathbf{120} \\
 &= \mathbf{8}_s \times_a \mathbf{8}_s + \mathbf{8}_c \times_a \mathbf{8}_c + \mathbf{8}_s \times \mathbf{8}_c = \mathbf{Adj}_{\mathbf{SO}(8)} + \Lambda_{\mathbf{8}_v}^1 + \Lambda_{\mathbf{28}}^2 + \Lambda_{\mathbf{56}_v}^3.
 \end{aligned} \tag{5.111}$$

(III) In maximal $D = 10$ IIB supergravity, the relevant values are $d = 2, 4$ with $n_2 = 2n_4 = 2$, and thus (5.103) and (5.104) specifies as follows:

$$\begin{aligned}
 \mathfrak{e}_{8(8)} \ominus (\mathfrak{sl}(8, \mathbb{R}) \oplus \mathfrak{sl}(2, \mathbb{R})) &\sim 2\Lambda^2 + \Lambda^4 + 2 * \Lambda^1 + * \Lambda^4 = 2\Lambda^2 + \Lambda^4 + 2\Lambda^6 + \Lambda^4 \\
 &\quad \text{manifestly SO(8)-cov.} \\
 &= ((\mathbf{28}, \mathbf{2}) + (\mathbf{35}_s, \mathbf{1})) + ((\mathbf{28}, \mathbf{2}) + (\mathbf{35}_c, \mathbf{1})); \tag{5.112}
 \end{aligned}$$

$$\begin{aligned}
 c \left(\frac{E_{8(8)}}{\mathbf{SO}(1, 1) \times \mathbf{SL}(8, \mathbb{R})} \right) &= 2 \binom{8}{2} + \binom{8}{4} = 2 \binom{8}{6} + \binom{8}{4} \\
 &= nc \left(\frac{E_{8(8)}}{\mathbf{SO}(1, 1) \times \mathbf{SL}(8, \mathbb{R})} \right) = 91 \\
 &= \dim_{\mathbb{R}} \left(\frac{\mathbf{SO}(16)}{\mathbf{SO}(8) \times \mathbf{SO}(2)} \right). \tag{5.113}
 \end{aligned}$$

In terms of the Cartan decomposition of the non-maximal non-symmetric Riemannian compact coset $\widehat{M}_{16}^{10 \text{ IIB}} = \mathbf{SO}(16)/(\mathbf{SO}(8) \times \mathbf{SO}(2))$, the result (5.113) can be obtained as a consequence of the next-to-maximal non-symmetric embedding (cfr. (2.2) and table 2):

$$\begin{aligned}
 \mathfrak{so}(16) &\supset_{ns} \mathfrak{so}(8) \oplus \mathfrak{so}(2) \\
 \mathbf{16} &= (\mathbf{8}_s, \mathbf{2}) \\
 \mathbf{Adj}_{\mathbf{SO}(16)} &\equiv (\mathbf{16} \times \mathbf{16})_a = (\mathbf{8}_s, \mathbf{2}) \times_a (\mathbf{8}_s, \mathbf{2}) = (\mathbf{8}_s \times_a \mathbf{8}_s, \mathbf{2} \times_s \mathbf{2}) + (\mathbf{8}_s \times_s \mathbf{8}_s, \mathbf{2} \times_a \mathbf{2}) \\
 &\quad \mathbf{120} \\
 &= (\mathbf{28}, \mathbf{3}) + (\mathbf{1}, \mathbf{1}) + (\mathbf{35}_s, \mathbf{1}) = \mathbf{Adj}_{\mathbf{SO}(8), 0} + \mathbf{Adj}_{\mathbf{SO}(2), 0} + \Lambda_{\mathbf{28}_2}^2 + \Lambda_{\mathbf{28}_{-2}}^2 + \Lambda_{\mathbf{35}_{s, 0}}^4, \tag{5.114}
 \end{aligned}$$

where in the last step the subscripts denote the charges of the $D = 10$ IIB \mathcal{R} -symmetry $\mathfrak{so}(2)$.

Finally, we present below the same analysis for other two “pure” supergravities:

(IV) In $N = 12$ supergravity (which shares the same bosonic sector of the quaternionic magical theory with $N = 4$ [19, 20]) in $D = 5$, $d = 1$ and $n_d = 15$, thus (5.103)

and (5.104) specifies as follows:

$$\begin{aligned} \mathfrak{e}_{7(-5)} \ominus (\mathfrak{su}^*(6) \oplus \mathfrak{sl}(3, \mathbb{R})) &\sim (14+1)\Lambda^1 + (14+1)*\Lambda^1 = (14+1)\Lambda^1 + (14+1)\Lambda^2 \\ &\quad \text{manifestly SO(3)-cov.} \\ &= (14+1)\mathbf{3} + (14+1)\mathbf{3} \end{aligned} \quad (5.115)$$

$$\begin{aligned} c\left(\frac{E_{7(-5)}}{SU^*(6) \times SL(3, \mathbb{R})}\right) &= (14+1)\mathbf{3} = (14+1)\binom{3}{2} = nc\left(\frac{E_{7(-5)}}{SU^*(6) \times SL(3, \mathbb{R})}\right) \\ &= 45 = \dim_{\mathbb{R}}\left(\frac{SO(12)}{USp(6)}\right). \end{aligned} \quad (5.116)$$

In terms of the Cartan decomposition of the maximal non-symmetric Riemannian compact coset $\widehat{M}_{12}^5 = SO(12)/USp(6)$, the result (5.116) can be obtained as a consequence of the maximal non-symmetric embedding

$$\begin{aligned} \mathfrak{so}(12) \supset_{ns} \mathfrak{usp}(6) \oplus \mathfrak{su}(2) \\ \mathbf{12} &= (\mathbf{6}, \mathbf{2}) \\ \mathbf{Adj}_{SO(12)} &\equiv (\mathbf{12} \times \mathbf{12})_a = (\mathbf{6}, \mathbf{2}) \times_a (\mathbf{6}, \mathbf{2}) = (\mathbf{6} \times_a \mathbf{6}, \mathbf{2} \times_s \mathbf{2}) + (\mathbf{6} \times_s \mathbf{6}, \mathbf{2} \times_a \mathbf{2}) \\ &\quad \mathbf{66} \\ &= (\mathbf{14}, \mathbf{3}) + \underset{\mathbf{Adj}_{SU(2)}}{(\mathbf{1}, \mathbf{3})} + \underset{\mathbf{Adj}_{USp(6)}}{(\mathbf{21}, \mathbf{1})}, \end{aligned} \quad (5.117)$$

where the $D = 5$ massless spin algebra $\mathfrak{su}(2)$ is not modded out in order to determine \widehat{M}_{12}^5 , and it corresponds to the “extra” $USp(6)$ (\mathcal{R} -symmetry-)singlet, a peculiar feature of this extended supergravity theory (which makes it amenable to an $N = 4$ interpretation).

(V) In $N = 10$ supergravity in $D = 4$, $d = 1$ and $n_d = 10$, thus (5.103) and (5.104) specifies as follows:

$$\begin{aligned} \mathfrak{e}_{6(-14)} \ominus (\mathfrak{su}(5, 1) \oplus \mathfrak{sl}(2, \mathbb{R})) &\sim 10\Lambda^1 + 10*\Lambda^1 = 10\Lambda^1 + 10\Lambda^1 = (10)\mathbf{2} + (10)\mathbf{2} \\ &\quad \text{manifestly SO(2)-cov.} \\ c\left(\frac{E_{6(-14)}}{SU(5, 1) \times SL(2, \mathbb{R})}\right) &= 10 \cdot 2 = nc\left(\frac{E_{6(-14)}}{SU(5, 1) \times SL(2, \mathbb{R})}\right) \\ &= 40 = \dim_{\mathbb{R}}\left(\frac{SO(10)}{U(5)}\right). \end{aligned} \quad (5.118)$$

In terms of the Cartan decomposition of the maximal non-symmetric Riemannian compact coset $\widehat{M}_{10}^4 = SO(10)/U(5)$, the result (5.118) can be obtained as a consequence of the maximal symmetric embedding

$$\begin{aligned} \mathfrak{so}(10) \supset_s \mathfrak{su}(5) \oplus \mathfrak{u}(1) \\ \mathbf{10} &= \mathbf{5}_1 + \bar{\mathbf{5}}_{-1} \\ \mathbf{Adj}_{SO(10)} &\equiv (\mathbf{10} \times \mathbf{10})_a = (\mathbf{5}_1 + \bar{\mathbf{5}}_{-1}) \times_a (\mathbf{5}_1 + \bar{\mathbf{5}}_{-1}) \\ &\quad \mathbf{45} \\ &= \mathbf{5}_1 \times_a \mathbf{5}_1 + \bar{\mathbf{5}}_{-1} \times_a \bar{\mathbf{5}}_{-1} + \mathbf{5}_1 \times \bar{\mathbf{5}}_{-1} = \mathbf{10}_2 + \bar{\mathbf{10}}_{-2} + \mathbf{24}_0 + \mathbf{1}_0, \end{aligned} \quad (5.119)$$

where the subscripts denote the charges with respect to the $D = 4$ massless spin algebra $\mathfrak{u}(1)$.

6 Conclusion

In this paper we have analyzed some consequences of the super-Ehlers structure of N -extended supergravity theories in $D \geq 4$ space-time dimensions. As the Ehlers $\text{SL}(D-2, \mathbb{R})$ is an off-shell symmetry of the Lagrangian [10–12], so there should exist an Hamiltonian formulation of light-cone supergravity in which U -duality G_N^D is an off-shell symmetry. Moreover, *at least* for any amount of supersymmetry $N \geq 4$, the Ehlers group can be regarded as the commutant of G_N^D itself inside the U -duality group G_N^3 in $D = 3$.

The pseudo-Riemannian manifolds pertaining to the embedding of the super-Ehlers group $G_N^D \times \text{SL}(D-2, \mathbb{R})$ into G_N^3 , namely the cosets M_N^D 's (5.1), have been found to exhibit an interesting invariance under the Hodge involution (5.102), acting on the cohomology of M_N^D , which in turn singles out only the physical massless $p > 0$ forms of the corresponding supergravity theory, regarded as p -fold antisymmetric irreps. Λ^p of the massless spin group $\text{SO}(D-2)_J = \text{mcs}(\text{SL}(D-2, \mathbb{R})_{\text{Ehlers}})$ in D (Lorentzian) space-time dimensions.

The symmetry under the Hodge involution (5.102) implies all the cosets M_N^D 's (5.1) to have a vanishing *character*, namely to have the same number of compact and non-compact generators: $c(M_N^D) = nc(M_N^D)$. Such a number, along with its manifestly $\text{SO}(D-2)_J$ -covariant decomposition in terms of physical massless $p > 0$ forms, can be computed by considering the Cartan decomposition of the cosets \widehat{M}_N^D 's (5.3), which can be regarded as the “*mcs* counterpart” of M_N^D 's (5.1). Indeed, the embedding of $\text{SO}(D-2)_J$ inside $H_N^3 \equiv \text{mcs}(G_N^3)$ is such that the generators of \widehat{M}_N^D split only into antisymmetric tensor irreps. of $\text{SO}(D-2)_J$ itself, with multiplicities given by irreps. of $H_N^D \equiv \text{mcs}(G_N^D)$.

The approach of this paper may be relevant for the analysis of the issue of ultraviolet divergences in supergravity theories with maximal or non-maximal supersymmetry, by exploiting the light-cone formulation, along the lines e.g. of [1, 10–12].

Acknowledgments

We would like to thank V. S. Varadarajan for an enlightening discussion on the Hodge involution.

A.M. would like to thank the Department of Physics and Astronomy, University of California at Los Angeles, where this project was completed, for kind hospitality and stimulating environment.

The work of S.F. has been supported by the ERC Advanced Grant no. 226455, Supersymmetry, Quantum Gravity and Gauge Fields (SUPERFIELDS), and in part by DOE Grant DE-FG03-91ER40662.

A Embeddings

Let us start by recalling some useful definitions.

Given two semisimple Lie groups G' and G , generated by the Lie algebras \mathfrak{g}' , \mathfrak{g} , respectively, if $G' \subset G$ (proper inclusion), we say that G' is *maximal* in G iff there is no proper subalgebra \mathfrak{g}'' of \mathfrak{g} containing \mathfrak{g}' . If G' and G are complex semisimple Lie groups such that




Figure 1. $\mathfrak{e}_{8(8)}$ Dynkin diagram.

$G' \subset G$, the embedding of G' into G is *regular* iff one can find a basis of \mathfrak{g}' consisting of elements of a Cartan subalgebra \mathfrak{h} of \mathfrak{g} and shift-generators E_α corresponding to roots α of \mathfrak{g} relative to \mathfrak{h} [42]. Regular subalgebras \mathfrak{g}' of a semisimple Lie algebra \mathfrak{g} can be constructed using the simple procedure defined by Dynkin in [42]: the Dynkin diagram of \mathfrak{g}' can be obtained as a truncation of the extended diagram of \mathfrak{g} . When considering *real forms* G' , G of complex semisimple Lie groups $G'_\mathbb{C}$, $G_\mathbb{C}$, we say that $G' \subset G$ is regularly embedded in G iff the complexification $\mathfrak{g}'_\mathbb{C}$ of \mathfrak{g}' is regularly embedded in the complexification $\mathfrak{g}_\mathbb{C}$ of \mathfrak{g} . The embedding of G' into G is *symmetric* iff we can write $\mathfrak{g} = \mathfrak{g}' \oplus \mathfrak{p}$, such that $[\mathfrak{g}', \mathfrak{p}] \subset \mathfrak{p}$ and $[\mathfrak{p}, \mathfrak{p}] \subset \mathfrak{g}'$. Finally the embedding is *rank-preserving* iff $\text{rank}(\mathfrak{g}') = \text{rank}(\mathfrak{g})$.

A.1 The embeddings $\text{SL}(D - 2, \mathbb{R}) \times \text{E}_{11-D(11-D)} \subset \text{E}_{8(8)}$

The $D = 5$ case $\text{SL}(3, \mathbb{R}) \times \text{E}_{6(6)} \subset \text{E}_{8(8)}$. The embedding of $\mathfrak{sl}(3, \mathbb{R}) \oplus \mathfrak{e}_{6(6)} \subset \mathfrak{e}_{8(8)}$ is regular and can be described using Dynkin's construction [42]. Let us number the simple roots of $\mathfrak{e}_{8(8)}$ so that the D_7 sub-Dynkin diagram consists of the roots $\alpha_2, \dots, \alpha_8$, with α_2 and α_3 on the two symmetric legs, and α_1 is the D_7 -spinor weight attached to α_3 , see figure 1. The $\mathfrak{e}_{8(8)}$ Cartan matrix reads:

$$\langle \alpha_i, \alpha_j \rangle = \begin{pmatrix} 2 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 2 & -1 & 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}. \tag{A.1}$$

In an orthonormal basis the simple roots α_i read:

$$\begin{aligned} \alpha_1 &= -\frac{1}{2}(\epsilon_1 + \epsilon_2 + \epsilon_3 + \epsilon_4 + \epsilon_5 + \epsilon_6 - \epsilon_7 - \epsilon_8), \\ \alpha_2 &= \epsilon_6 + \epsilon_7; \quad \alpha_3 = \epsilon_6 - \epsilon_7; \quad \alpha_4 = \epsilon_5 - \epsilon_6; \quad \alpha_5 = \epsilon_4 - \epsilon_5; \quad \alpha_6 = \epsilon_3 - \epsilon_4; \quad \alpha_7 = \epsilon_2 - \epsilon_3; \\ \alpha_8 &= \epsilon_1 - \epsilon_2 \end{aligned} \tag{A.2}$$

Let us denote by $\Delta_+[\mathfrak{e}_{8(8)}] = \{\alpha = \sum_{i=1}^8 n^i \alpha_i\}$ the set of positive roots of $\mathfrak{e}_{8(8)}$. The $\mathfrak{e}_{6(6)}$ subalgebra is defined by the sub-Dynkin diagram consisting of the simple roots α_a ,

$a = 1, \dots, 6$. The 36 positive $\mathfrak{e}_{6(6)}$ -roots be denoted by γ_A , so that:

$$\Delta_+[\mathfrak{e}_{6(6)}] = \left\{ \gamma_A = \sum_{a=1}^6 n_A^a \alpha_a \right\}. \quad (\text{A.3})$$

Furthermore let us consider the following positive roots β_x , $x = 1, 2, 3$:

$$\begin{aligned} \beta_1 &= \alpha_8 = \epsilon_1 - \epsilon_2 ; \quad \beta_2 = 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 6\alpha_4 + 5\alpha_5 + 4\alpha_6 + 3\alpha_7 + \alpha_8 = \epsilon_2 + \epsilon_8 ; \\ \beta_3 &= 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 6\alpha_4 + 5\alpha_5 + 4\alpha_6 + 3\alpha_7 + 2\alpha_8 = \epsilon_1 + \epsilon_8 . \end{aligned} \quad (\text{A.4})$$

One can easily verify that β_x generate an $\mathfrak{sl}(3, \mathbb{R})$ -root space which is orthogonal to $\Delta_+[\mathfrak{e}_{6(6)}]$: $\beta_x \cdot \gamma_A = 0$. We have then constructed an $\mathfrak{sl}(3, \mathbb{R}) \oplus \mathfrak{e}_{6(6)}$ subalgebra of $\mathfrak{e}_{8(8)}$:

$$\begin{aligned} \mathfrak{sl}(3, \mathbb{R}) &= \text{Span}(H_{\beta_1}, H_{\beta_2}, E_{\pm\beta_1}, E_{\pm\beta_2}, E_{\pm\beta_3}), \\ \mathfrak{e}_{6(6)} &= \text{Span}(H_{\alpha_a}, E_{\pm\gamma_A})_{\substack{a=1, \dots, 6 \\ A=1, \dots, 36}} \end{aligned} \quad (\text{A.5})$$

Within $\mathfrak{sl}(3, \mathbb{R}) \oplus \mathfrak{e}_{6(6)}$ we can identify its maximal compact subalgebra $\mathfrak{so}(3) \oplus \mathfrak{usp}(8)$, which is a maximal subalgebra of $\mathfrak{so}(16)$:

$$\begin{aligned} \mathfrak{so}(16) &= \text{Span}(E_\alpha - E_{-\alpha})_{\alpha \in \Delta_+[\mathfrak{e}_{8(8)}]}, \\ \mathfrak{so}(3) &= \text{Span}(E_{\beta_x} - E_{-\beta_x})_{x=1,2,3}, \\ \mathfrak{usp}(8) &= \text{Span}(E_{\gamma_A} - E_{-\gamma_A})_{A=1, \dots, 36}. \end{aligned} \quad (\text{A.6})$$

With respect to this $\text{SO}(3) \times \text{USp}(8)$ subgroup of $\text{SO}(16)$ the coset space

$$\mathfrak{K} = \mathfrak{e}_{8(8)} \ominus \mathfrak{so}(16) = \text{Span}(H_{\alpha_i}, E_\alpha + E_{-\alpha})_{\alpha \in \Delta_+[\mathfrak{e}_{8(8)}]; i=1, \dots, 8}, \quad (\text{A.7})$$

should decompose as follows:

$$\begin{aligned} \mathfrak{K} &= \mathfrak{K}[\mathfrak{sl}(3, \mathbb{R})] \oplus \mathfrak{K}[\mathfrak{e}_{6(6)}] \oplus (\mathbf{3}, \mathbf{27}), \\ \mathfrak{K}[\mathfrak{sl}(3, \mathbb{R})] &= \mathfrak{sl}(3, \mathbb{R}) \ominus \mathfrak{so}(3) = \text{Span}(H_{\beta_1}, H_{\beta_2}, E_{\beta_x} + E_{-\beta_x})_{x=1,2,3} = (\mathbf{5}, \mathbf{1}), \\ \mathfrak{K}[\mathfrak{e}_{6(6)}] &= \mathfrak{e}_{6(6)} \ominus \mathfrak{usp}(8) = \text{Span}(H_{\alpha_a}, E_{\gamma_A} + E_{-\gamma_A})_{\substack{a=1, \dots, 6 \\ A=1, \dots, 36}} = (\mathbf{1}, \mathbf{42}), \end{aligned} \quad (\text{A.8})$$

Generalizing to $\text{SL}(D-2, \mathbb{R}) \times \mathbf{E}_{11-D(11-D)} \subset \mathbf{E}_8(8)$. The above construction is extended to define the embedding of $\mathfrak{sl}(D-2, \mathbb{R}) \oplus \mathfrak{e}_{11-D(11-D)} \subset \mathfrak{e}_{8(8)}$, $D \geq 4$, following the same recipe by Dynkin. The embedding of $\mathfrak{e}_{11-D(11-D)}$ is defined by deleting in the $\mathfrak{e}_{8(8)}$ Dynkin diagram the last $D-3$ simple roots to the right, namely $\alpha_{12-D}, \dots, \alpha_8$, see figure 2. The set of positive roots of $\mathfrak{e}_{11-D(11-D)}$ reads:

$$\Delta_+[\mathfrak{e}_{11-D(11-D)}] = \{\gamma_A\} = \left\{ \epsilon_a \pm \epsilon_b, \left[\frac{\epsilon_8}{2} - \sum_{\alpha=1}^{D-3} \frac{\epsilon_\alpha}{2} + \left(\sum_{a=D-2}^7 \pm \frac{\epsilon_a}{2} \right)_{\text{odd}+} \right] \right\}, \quad (\text{A.9})$$

where those in square brackets are the weights of a chiral spinorial representation of the $\mathfrak{so}(10-D, 10-D)$ subalgebra of $\mathfrak{e}_{11-D(11-D)}$. The set of positive roots of $\mathfrak{sl}(D-2, \mathbb{R})$ reads:

$$\Delta_+[\mathfrak{sl}(D-2, \mathbb{R})] = \{\beta_x\} = \{\epsilon_\alpha - \epsilon_\beta, \epsilon_\alpha + \epsilon_8\}, \quad (\text{A.10})$$




Figure 2. The filled circles define the $\mathfrak{e}_{11-D(11-D)}$ sub-Dynkin diagram, while the thick circle represents the exceptional root $-\psi$, $\psi = \epsilon_1 + \epsilon_8$ being the highest root of \mathfrak{e}_8 , which, together with the other roots in the rectangles, defines the Dynkin diagram of $\mathfrak{sl}(D-2, \mathbb{R})$.

where $\alpha, \beta = 1, \dots, D-3$, $\beta > \alpha$ and $x = 1, \dots, (D-3)(D-2)/2$. One can easily verify that the two root systems are orthogonal, namely: $\beta_x \cdot \gamma_A = 0$.

This defines the $\mathfrak{sl}(D-2, \mathbb{R}) \oplus \mathfrak{e}_{11-D(11-D)}$ subalgebra of $\mathfrak{e}_{8(8)}$:

$$\begin{aligned} \mathfrak{sl}(D-2, \mathbb{R}) &= \text{Span}(H_{\alpha_{13-D}}, \dots, H_{\alpha_8}, H_{\epsilon_1 + \epsilon_8}, E_{\pm\beta_x})_{\beta_x \in \Delta_+[\mathfrak{sl}(D-2, \mathbb{R})]}, \\ \mathfrak{e}_{11-D(11-D)} &= \text{Span}(H_{\alpha_a}, E_{\pm\gamma_A})_{\substack{a=1, \dots, 11-D \\ \gamma_A \in \Delta_+[\mathfrak{e}_{11-D(11-D)]}}, D=4, \dots, 8, \\ \mathfrak{e}_{2(2)} &= \text{Span}(H_{\alpha_1}, H_\lambda, E_{\pm\alpha_1}), D=9, \end{aligned} \tag{A.11}$$

where the generators $H_{\alpha_{13-D}}, \dots, H_{\alpha_8}$, in the first line, are not counted for $D=4$, for which the only Cartan generator of $\mathfrak{sl}(2, \mathbb{R})$ is $H_{\epsilon_1 + \epsilon_8}$. In the $D=9$ case, the vector λ in the last line is: $\lambda = \epsilon_7 - \alpha_1/4$ and is orthogonal to the β_x and to α_1 .

As far as the corresponding maximal compact subalgebra $\mathfrak{so}(D-2) \oplus \mathfrak{H}_D$ is concerned, it can be constructed as follows:

$$\begin{aligned} \mathfrak{so}(D-2) &= \text{Span}(E_{\beta_x} - E_{-\beta_x})_{\beta_x \in \Delta_+[\mathfrak{sl}(D-2, \mathbb{R})]}, \\ \mathfrak{H}_D &= \text{Span}(E_{\gamma_A} - E_{-\gamma_A})_{\gamma_A \in \Delta_+[\mathfrak{e}_{11-D(11-D)}]}, \end{aligned} \tag{A.12}$$

where \mathfrak{H}_D is the maximal compact subalgebra of $\mathfrak{e}_{11-D(11-D)}$.

In the $D=10$ case we need to consider the type IIA and type IIB descriptions in which the relevant subgroups of $E_{8(8)}$ are $\text{SL}(8, \mathbb{R}) \times \text{SO}(1, 1)$ and $\text{SL}(8, \mathbb{R})' \times \text{SL}(2, \mathbb{R})$, respectively.²⁰ Their embeddings are illustrated in figure 6. In the former case the U -duality group is $\text{SO}(1, 1)$ and is generated by the Cartan generator $\sum_{i=1}^8 H_{\epsilon_i} - 2H_{\epsilon_8}$.

²⁰We shall omit the prime in the following.




Figure 3. From top to bottom: Satake diagram of $\mathfrak{e}_{8(-24)}$ and embeddings of the $\mathfrak{e}_{6(-26)} \oplus \mathfrak{sl}(3, \mathbb{R})$ and $\mathfrak{so}(1, 9) \oplus \mathfrak{sl}(4, \mathbb{R})$ diagrams inside the extension of the $\mathfrak{e}_{8(-24)}$ one.

A.2 Other embeddings

Embeddings considered here were also dealt with in [17]. Here we provide a detailed and explicit construction of a number of embeddings in terms of the generators of the corresponding Lie algebras, using the notation of [14]. Let us start discussing in detail the embeddings of $E_{6(-26)} \times \text{SL}(3, \mathbb{R})$ and $\text{SO}(1, 9) \times \text{SL}(4, \mathbb{R})$ inside $E_{8(-24)}$. At the level of the corresponding Lie algebras, these embeddings are illustrated in figure 3, where the *Satake diagrams* of $\mathfrak{e}_{6(-26)} \oplus \mathfrak{sl}(3, \mathbb{R})$ and $\mathfrak{so}(1, 9) \oplus \mathfrak{sl}(4, \mathbb{R})$ are obtained from the $\mathfrak{e}_{8(-24)}$ one once again using Dynkin’s procedure of extending the latter and canceling a suitable simple root. Let us briefly review the definition of Satake diagrams for non-split (i.e. non-maximally-non-compact) Lie algebras and the construction of the $\mathfrak{e}_{6(-26)} \oplus \mathfrak{sl}(3, \mathbb{R})$ and $\mathfrak{so}(1, 9) \oplus \mathfrak{sl}(4, \mathbb{R})$ generators in terms of a canonical basis of the complex \mathfrak{e}_8 . The latter consists of a basis $\{H_{\epsilon_i}\}$, $i = 1 \dots, 8$, of Cartan generators, with respect to which the \mathfrak{e}_8 roots are defined, and shift operators $E_\alpha, E_{-\alpha}$, α being the 120 positive roots. The real form $\mathfrak{e}_{8(-24)}$ is characterized by a Cartan subalgebra \mathfrak{h} which splits into the direct sum of a subspace \mathfrak{h}^{nc} of non-compact generators (i.e. generators which are odd with respect to the Cartan involution τ ²¹) and a subspace \mathfrak{h}^c of compact generators, defined in terms of the $\{H_{\epsilon_i}\}$ as follows:

$$\mathfrak{h} = \mathfrak{h}^{nc} \oplus \mathfrak{h}^c ; \quad \mathfrak{h}^c = \text{Span}(i H_{\alpha_2}, i H_{\alpha_3}, i H_{\alpha_4}, i H_{\alpha_5}) ; \quad \mathfrak{h}^{nc} = \text{Span}(H_{\epsilon_1}, H_{\epsilon_2}, H_{\epsilon_3}, H_{\epsilon_8}), \tag{A.13}$$

Note that \mathfrak{h}^c is the Cartan subalgebra of an $\mathfrak{so}(8)$ subalgebra of $\mathfrak{e}_{8(-24)}$ whose Dynkin diagram is defined by the black roots in figure 3. The \mathfrak{e}_8 positive roots split into a 12-dimensional sub-space $\Delta_+^{(0)}[\mathfrak{e}_8]$ of roots having null restriction to \mathfrak{h}^{nc} and a 108-dimensional

²¹We can always find a suitable basis for the matrix representation of the generators so that $\tau(M) = -M^\dagger$. This means that we shall regard compactness and non-compactness of a generator to be synonyms, in any matrix representation, of being anti-hermitian and hermitian, respectively. Moreover, in our conventions, $E_{-\alpha} = -\tau(E_\alpha) = E_\alpha^\dagger$.

space $\bar{\Delta}_+[\mathfrak{e}_8]$ of roots with a non-trivial restriction to \mathfrak{h}^{nc} :

$$\Delta_+[\mathfrak{e}_8] = \Delta_+^{(0)}[\mathfrak{e}_8] \oplus \bar{\Delta}_+[\mathfrak{e}_8]. \quad (\text{A.14})$$

The conjugation σ with respect to $\mathfrak{e}_{8(-24)}$ is the conjugation on the complex \mathfrak{e}_8 which leaves the elements of the subalgebra $\mathfrak{e}_{8(-24)}$ invariant. It defines a correspondence between \mathfrak{e}_8 -roots $\alpha \leftrightarrow \alpha^\sigma$ such that $\sigma(E_\alpha) \propto E_{\alpha^\sigma}$. The couple of roots α, α^σ satisfies the property:

$$\alpha|_{\mathfrak{h}^{nc}} = \alpha^\sigma|_{\mathfrak{h}^{nc}}; \quad \alpha|_{\mathfrak{h}^c} = -\alpha^\sigma|_{\mathfrak{h}^c}. \quad (\text{A.15})$$

Clearly if $\alpha \in \Delta_+^{(0)}[\mathfrak{e}_8]$, $\alpha^\sigma = -\alpha$, while if $\alpha \in \bar{\Delta}_+[\mathfrak{e}_8]$ and $\alpha|_{\mathfrak{h}^c} = 0$, we have $\alpha^\sigma = \alpha$. Thus if $\alpha \in \bar{\Delta}_+[\mathfrak{e}_8]$, to each couple of nilpotent generators E_α and $\sigma(E_\alpha)$ in \mathfrak{e}_8 , there corresponds a couple of nilpotent generators in $\mathfrak{e}_{8(-24)}$ given by the σ -invariant combinations $i(E_\alpha - \sigma(E_\alpha))$, $E_\alpha + \sigma(E_\alpha)$, which can be both brought to an upper-triangular form, for all α . If, on the other hand, $\alpha \in \Delta_+^{(0)}[\mathfrak{e}_8]$, the same combinations define *compact* $\mathfrak{so}(8)$ generators $i(E_\alpha + E_{-\alpha})$, $E_\alpha - E_{-\alpha}$.

To summarize, the $\mathfrak{e}_{8(-24)}$ generators can be expressed in terms of the \mathfrak{e}_8 canonical basis as follows:

$$\begin{aligned} \mathfrak{e}_{8(-24)} &= \mathfrak{h} \oplus \mathfrak{l}_+ \oplus \mathfrak{l}_- \oplus \mathfrak{m}_0, \\ \mathfrak{l}_+ &= \text{Span} [i(E_\alpha - \sigma(E_\alpha)), E_\alpha + \sigma(E_\alpha)]_{(\alpha, \alpha^\sigma) \in \bar{\Delta}_+[\mathfrak{e}_8]}, \\ \mathfrak{l}_- &= \text{Span} [i(E_{-\alpha} - \sigma(E_{-\alpha})), E_{-\alpha} + \sigma(E_{-\alpha})]_{(\alpha, \alpha^\sigma) \in \bar{\Delta}_+[\mathfrak{e}_8]}, \\ \mathfrak{m}_0 &= \text{Span} [i(E_\alpha + E_{-\alpha}), E_\alpha - E_{-\alpha}]_{\alpha \in \Delta_+^{(0)}[\mathfrak{e}_8]}, \end{aligned} \quad (\text{A.16})$$

The 112-dimensional solvable Lie algebra $\mathfrak{s}_0 = \mathfrak{h}^{nc} \oplus \mathfrak{l}_+$ is the one defined by the Iwasawa decomposition of $\mathfrak{e}_{8(-24)}$ with respect to $\mathfrak{e}_{7(-133)} \oplus \mathfrak{su}(2)$, and its generators, in a suitable basis, can all be represented by upper-triangular matrices. The centralizer of \mathfrak{h}^{nc} is the $\mathfrak{so}(8)$ subalgebra given by $\mathfrak{h}^c \oplus \mathfrak{m}_0$ and is also contained inside the subalgebras $\mathfrak{e}_{6(-26)}$ and $\mathfrak{so}(1, 9)$, as it is apparent from figure 3.

The $\mathfrak{e}_{6(-26)}$ generators in terms of the above $\mathfrak{e}_{8(-24)}$ ones are easily written:

$$\begin{aligned} \mathfrak{e}_{6(-26)} &= \mathfrak{h}' \oplus \mathfrak{l}'_+ \oplus \mathfrak{l}'_- \oplus \mathfrak{m}_0, \\ \mathfrak{l}'_+ &= \text{Span} [i(E_\alpha - \sigma(E_\alpha)), E_\alpha + \sigma(E_\alpha)]_{(\alpha, \alpha^\sigma) \in \bar{\Delta}_+[\mathfrak{e}_8] \cap \Delta_+[\mathfrak{e}_6]}, \\ \mathfrak{l}'_- &= \text{Span} [i(E_{-\alpha} - \sigma(E_{-\alpha})), E_{-\alpha} + \sigma(E_{-\alpha})]_{(\alpha, \alpha^\sigma) \in \bar{\Delta}_+[\mathfrak{e}_8] \cap \Delta_+[\mathfrak{e}_6]}, \\ \mathfrak{m}_0 &= \text{Span} [i(E_\alpha + E_{-\alpha}), E_\alpha - E_{-\alpha}]_{\alpha \in \Delta_+^{(0)}[\mathfrak{e}_8]}, \end{aligned} \quad (\text{A.17})$$

where $\Delta_+[\mathfrak{e}_6]$ are the \mathfrak{e}_6 -positive roots in the \mathfrak{e}_8 -root system, while

$$\mathfrak{h}' = \mathfrak{h}^{mc} \oplus \mathfrak{h}^c; \quad \mathfrak{h}^{mc} = \text{Span}(H_{\epsilon_1 + \epsilon_2 - \epsilon_8}, H_{\epsilon_3}). \quad (\text{A.18})$$

The $\mathfrak{sl}(3, \mathbb{R})$ subalgebra commuting with $\mathfrak{e}_{6(-26)}$ has the following form:

$$\mathfrak{sl}(3, \mathbb{R}) = \text{Span} [H_{\epsilon_1 - \epsilon_2}, H_{-\epsilon_1 - \epsilon_8}, E_{\pm\beta_x}], \quad \{\beta_x\} = \{\epsilon_1 - \epsilon_2, \epsilon_8 + \epsilon_1, \epsilon_8 + \epsilon_2\},$$

note that $\beta_x^\sigma = \beta_x$.




Figure 4. Embeddings $SL(3, \mathbb{R}) \times SU^*(6) \subset E_{7(-5)}$ and $SL(3, \mathbb{C}) \times SL(3, \mathbb{R}) \subset E_{6(2)}$. The thick circle is, as usual, the opposite of the highest root of the corresponding algebra.

Finally the $\mathfrak{so}(1, 9) \subset \mathfrak{e}_{6(-26)}$ generators read:

$$\begin{aligned}
 \mathfrak{so}(1, 9) &= \mathfrak{h}'' \oplus \mathfrak{l}''_+ \oplus \mathfrak{l}''_- \oplus \mathfrak{m}_0, \\
 \mathfrak{l}''_+ &= \text{Span} [i (E_\alpha - \sigma(E_\alpha)), E_\alpha + \sigma(E_\alpha)]_{(\alpha, \alpha\sigma) \in \bar{\Delta}_+[\mathfrak{e}_8] \cap \Delta_+[\mathfrak{so}(10)]}, \\
 \mathfrak{l}''_- &= \text{Span} [i (E_{-\alpha} - \sigma(E_{-\alpha})), E_{-\alpha} + \sigma(E_{-\alpha})]_{(\alpha, \alpha\sigma) \in \bar{\Delta}_+[\mathfrak{e}_8] \cap \Delta_+[\mathfrak{so}(10)]}, \\
 \mathfrak{m}_0 &= \text{Span} [i (E_\alpha + E_{-\alpha}), E_\alpha - E_{-\alpha}]_{\alpha \in \Delta_+^{(0)}[\mathfrak{e}_8]}, \tag{A.19}
 \end{aligned}$$

where $\Delta_+[\mathfrak{so}(10)]$ are the roots of the complex $\mathfrak{so}(10)$ algebra within \mathfrak{e}_8 -root system, and

$$\mathfrak{h}'' = \mathfrak{h}''^{nc} \oplus \mathfrak{h}^c; \quad \mathfrak{h}''^{nc} = \text{Span}(H_{\epsilon_1 + \epsilon_2 + \epsilon_3 - \epsilon_8}). \tag{A.20}$$

The $\mathfrak{sl}(4, \mathbb{R})$ subalgebra commuting with $\mathfrak{so}(1, 9)$ is described by the following generators:

$$\mathfrak{sl}(4, \mathbb{R}) = \text{Span} [H_{\epsilon_1 - \epsilon_2}, H_{\epsilon_2 - \epsilon_3}, H_{-\epsilon_1 - \epsilon_8}, E_{\pm\beta_x}], \quad \{\beta_x\} = \{\epsilon_\alpha - \epsilon_\beta, \epsilon_8 + \epsilon_\alpha\}_{\alpha < \beta, \alpha, \beta = 1, 2, 3},$$

By the same token we can prove other embeddings, like $SL(3, \mathbb{R}) \times SU^*(6) \subset E_{7(-5)}$ and $SL(3, \mathbb{C}) \times SL(3, \mathbb{R}) \subset E_{6(2)}$, see figure 4. In the latter case there is a subtlety which is not apparent from the truncation of the extended Satake diagram: The bottom-right diagram in figure 4 would naively suggest that the roots $\alpha_1, \alpha_3, \alpha_5, \alpha_6$ define two commuting $\mathfrak{sl}(3, \mathbb{R})$ subalgebras. This is however not the case since, as represented by the lower arrows, the conjugation σ corresponding to the real form $\mathfrak{e}_{6(2)}$ inside the complex \mathfrak{e}_6 , maps α_1 and α_3 into $\alpha_1^\sigma = \alpha_6$ and $\alpha_3^\sigma = \alpha_5$, respectively. As a consequence of this the \mathfrak{e}_6 shift generators corresponding to the two orthogonal $\mathfrak{sl}(3, \mathbb{R})$ root spaces are mixed together in σ -invariant combinations inside $\mathfrak{e}_{6(2)}$, which make the shift generators of a $\mathfrak{sl}(3, \mathbb{C})$ subalgebra. This subalgebra also contains the two non-compact combinations $H_{\alpha_1} + H_{\alpha_6}, H_{\alpha_2} + H_{\alpha_5}$ and the two compact combinations $i(H_{\alpha_1} - H_{\alpha_6}), i(H_{\alpha_2} - H_{\alpha_5})$ of the \mathfrak{e}_6 Cartan generators.

In figure 5 the embeddings $SL(4, \mathbb{R}) \times SO(3) \times SO(1, 5) \subset E_{7(-5)}$ and $SL(2, \mathbb{C}) \times SL(4, \mathbb{R}) \times SO(2) \subset E_{6(2)}$ are illustrated.




Figure 5. Embeddings $SL(4, \mathbb{R}) \times SO(3) \times SO(1, 5) \subset E_{7(-5)}$ and $SL(2, \mathbb{C}) \times SL(4, \mathbb{R}) \times SO(2) \subset E_{6(2)}$.

A.3 General features

One can generalize the above discussion and show that, as a general feature of the embeddings considered in this work, the \mathfrak{g}_N^3 algebra, and its super-Ehlers subalgebra $\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)$ can be written in the forms:

$$\mathfrak{g}_N^3 = \mathfrak{h} \oplus \mathfrak{l}_+ \oplus \mathfrak{l}_- \oplus \mathfrak{m}_0 ; \quad \mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2) = \mathfrak{h} \oplus \hat{\mathfrak{l}}_+ \oplus \hat{\mathfrak{l}}_- \oplus \mathfrak{m}_0 . \tag{A.21}$$

Note that, as a consequence of the regularity of the embedding and properties (1.3), (1.4), their Cartan subalgebras

$$\mathfrak{h} = \mathfrak{h}^{nc} \oplus \mathfrak{h}^c , \tag{A.22}$$

can be chosen to coincide, where $\dim(\mathfrak{h}^{nc})$ is the non-compact rank of the two groups. This is implicit in Dynkin’s construction of the $\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)$ algebra by truncating the extended diagram of \mathfrak{g}_N^3 . Moreover the centralizer of \mathfrak{h}^{nc} , which is the compact algebra $\mathfrak{h}^c \oplus \mathfrak{m}_0$, is common to the two algebras:

$$\mathfrak{h}^c \oplus \mathfrak{m}_0 \subset \mathfrak{g}_N^3 \cap [\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)] . \tag{A.23}$$

For a split (maximally non-compact) \mathfrak{g}_N^3 , $\mathfrak{h}^c = \mathfrak{m}_0 = \emptyset$ and $\alpha^\sigma = \alpha$.

The nilpotent spaces $\mathfrak{l}_\pm, \hat{\mathfrak{l}}_\pm$ have the form:

$$\begin{aligned} \mathfrak{l}_\pm &= \text{Span} [i(E_{\pm\alpha} - \sigma(E_{\pm\alpha})), E_{\pm\alpha} + \sigma(E_{\pm\alpha})]_{(\alpha, \alpha^\sigma) \in \bar{\Delta}_+[\mathfrak{g}_N^3]} , \\ \hat{\mathfrak{l}}_\pm &= \text{Span} [i(E_{\pm\alpha} - \sigma(E_{\pm\alpha})), E_{\pm\alpha} + \sigma(E_{\pm\alpha})]_{(\alpha, \alpha^\sigma) \in \bar{\Delta}_+[\mathfrak{g}_N^3] \cap \Delta_+[\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)]} , \end{aligned} \tag{A.24}$$

where, as usual, $\bar{\Delta}_+[\mathfrak{g}_N^3]$ denotes the set of positive roots of the (complexification of) \mathfrak{g}_N^3 with non-trivial restriction to \mathfrak{h}^{nc} , and $\Delta_+[\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)]$ the set of positive roots of the (complexification of) $\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)$, which is a subset of $\Delta_+[\mathfrak{g}_N^3]$. Thus in general we have:

$$\hat{\mathfrak{l}}_{\pm} \subset \mathfrak{l}_{\pm}. \quad (\text{A.25})$$

We can then write the coset space as follows:

$$\mathfrak{g}_N^3 \ominus [\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)] = \mathfrak{N}^+ \oplus \mathfrak{N}^-, \quad (\text{A.26})$$

where $\mathfrak{N}^{\pm} = \mathfrak{l}_{\pm} \ominus \hat{\mathfrak{l}}_{\pm}$. Semisimplicity of \mathfrak{g}_N^3 and \mathfrak{g}_N^D implies that $\dim(\mathfrak{l}_+) = \dim(\mathfrak{l}_-)$ and $\dim(\hat{\mathfrak{l}}_+) = \dim(\hat{\mathfrak{l}}_-)$, so that $\dim(\mathfrak{N}^+) = \dim(\mathfrak{N}^-)$. More precisely, in a suitable basis, for each strictly-upper-triangular matrix M_+ representing an element in \mathfrak{N}^+ , its (strictly-lower-triangular) hermitian-conjugate $M_- = M_+^{\dagger} = -\tau(M_+)$ represents an element in \mathfrak{N}^- : The former is given by a generator either of the form $i(E_{\alpha} - \sigma(E_{\alpha}))$ or $E_{\alpha} + \sigma(E_{\alpha})$, for some $\alpha \in \bar{\Delta}_+[\mathfrak{g}_N^3] \ominus \bar{\Delta}_+[\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)]$, the latter will either be $-i(E_{-\alpha} - \sigma(E_{-\alpha}))$ or $E_{-\alpha} + \sigma(E_{-\alpha})$, corresponding to the same α . Thus if $\{L_{\ell}^+\}$, $\ell = 1, \dots, \dim(\mathfrak{N}^{\pm})$, is a basis of \mathfrak{N}^+ , $\{L_{\ell}^-\} = \{-\tau(L_{\ell}^+)\}$ is a basis of \mathfrak{N}^- and we can also write the coset space in the form:

$$\mathfrak{g}_N^3 \ominus [\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)] = \mathfrak{N}^c \oplus \mathfrak{N}^{nc}, \quad (\text{A.27})$$

where

$$\mathfrak{N}^{nc} = \text{Span}(L_{\ell}^+ + L_{\ell}^-); \quad \mathfrak{N}^c = \text{Span}(L_{\ell}^+ - L_{\ell}^-), \quad (\text{A.28})$$

which are the eigenspaces of τ on $\mathfrak{N}^+ \oplus \mathfrak{N}^-$ corresponding to the eigenvalues -1 and $+1$, respectively. These subspaces define representations with respect to the compact group $\text{SO}(D-2) \times mcs(G_N^D)$. With respect to the G_N^3 -invariant scalar product on \mathfrak{g}_N^3 , \mathfrak{N}^c and \mathfrak{N}^{nc} have negative and positive signatures, respectively. Since

$$\dim(\mathfrak{N}^c) = \dim(\mathfrak{N}^{nc}), \quad (\text{A.29})$$

the manifold M_N^D in (5.1) has vanishing character, being

$$c(M_N^D) = \dim(\mathfrak{N}^c) = \dim(\mathfrak{N}^{nc}) = nc(M_N^D),$$

as also proven in Sect 5.3. We shall come back on this issue in appendix C.

B $\mathfrak{so}(8,8)$ outer automorphisms and dual subalgebras of $\mathfrak{e}_{8(8)}$

Consider in the maximal $D = 3$ theory the effect of an $\text{O}(8,8)$ ‘‘reflection’’ of the form:

$$\mathcal{O}_k = \begin{pmatrix} \mathbf{1}_8 - \mathbf{D}_k & \mathbf{D}_k \\ \mathbf{D}_k & \mathbf{1}_8 - \mathbf{D}_k \end{pmatrix}, \quad (\text{B.1})$$

where each block is an 8×8 matrix and \mathbf{D}_k is the zero-matrix except for only an *odd number* k of 1s along the diagonal. Such transformation, which belongs to the $\text{O}(8)$ subgroup of $\text{O}(8,8)$, is an outer automorphism of the D_8 algebra whose effect, modulo Weyl




Figure 6. Outer automorphism of the D_8 subalgebra of \mathfrak{e}_8 and two inequivalent $\mathfrak{sl}(8, \mathbb{R})$ subalgebras of $\mathfrak{e}_{8(8)}$.

transformations of the same algebra, is to interchange α_2 with α_3 in figure 1. While it is a symmetry of the D_8 Dynkin diagram, it is not a symmetry of the $\mathfrak{e}_{8(8)}$ one, as it changes the $SO(8, 8)$ -chirality of the α_1 root, which is a D_8 -spinorial weight [25]. In particular this outer automorphism may map inequivalent subalgebras \mathfrak{g} , \mathfrak{g}' of $\mathfrak{so}(8, 8)$ into one another. This is the case of subalgebras \mathfrak{g} (and thus \mathfrak{g}') which are the direct sum of commuting A_k -algebras with odd rank k . In mathematical language such *dual* subalgebras are said to be *linearly equivalent*, i.e. in any matrix representation they are equivalent through conjugation by means of a matrix, which is however not necessarily a representation of an $SO(8, 8)$ element, as it is the case for the outer automorphisms. Equivalence therefore implies linear equivalence though the reverse implication is not true. With respect to \mathfrak{g} and \mathfrak{g}' , a same spinorial representation of $\mathfrak{so}(8, 8)$, and thus the adjoint representation of the whole $\mathfrak{e}_{8(8)}$, will branch differently. They are clearly inequivalent $\mathfrak{e}_{8(8)}$ -subalgebras. Examples are given in [44]: $\mathfrak{g} = \mathfrak{sl}(8)$, $\mathfrak{sl}(6) \oplus \mathfrak{sl}(2)$, $\mathfrak{sl}(4) \oplus \mathfrak{sl}(4)$, etc., see Fig 6.

What has been said for $\mathfrak{so}(8, 8)$ also holds for the $\mathfrak{so}^*(16)$ and $\mathfrak{so}(16)$ subalgebras of $\mathfrak{e}_{8(8)}$. For instance there are two inequivalent $\mathfrak{u}(8)$, $\mathfrak{u}'(8)$ in either $\mathfrak{so}^*(16)$ or $\mathfrak{so}(16)$. One contains the R -symmetry algebras $\mathfrak{su}(8)$, $\mathfrak{usp}(8)$, etc. of the higher dimensional parent maximal supergravities, the other dual subalgebras $\mathfrak{su}'(8)$, $\mathfrak{usp}'(8)$, etc. which are not contained in the chain of exceptional duality algebras $\mathfrak{e}_{7(7)}$, $\mathfrak{e}_{6(6)}$ etc.

Let us briefly recall the relation between outer automorphisms of $\mathfrak{so}(8, 8)$ and dualities. Consider the toroidal reduction of the $D = 11$ theory down to $D = 3$ (in the Einstein frame). The Kaluza-Klein ansatz for the metric reads:

$$G_{\hat{\mu}\hat{\nu}}^{(11)} = \begin{pmatrix} e^{2\xi} g_{\mu\nu}^{(3)} + G_{pq} G_\mu^p G_\nu^q & G_{np} G_\mu^p \\ G_{mp} G_\nu^p & G_{mn} \end{pmatrix}; \quad \xi = -\frac{1}{2} \log(\det(G_{mn})), \quad (\text{B.2})$$

where $\hat{\mu}, \hat{\nu} = 0, \dots, 10$, $\mu, \nu = 0, 1, 2$, $m, n = 3, \dots, 10$ and the internal metric of T^8 is

conveniently written as follows:

$$\mathbf{G} = (G_{mp}) = \mathbf{E}\mathbf{E}^T = \hat{\mathbf{E}}\mathbf{D}^2\hat{\mathbf{E}}^T, \quad (\text{B.3})$$

where $\mathbf{E} = (E_m^a)$ is the vielbein of the coset $\text{GL}(8, \mathbb{R})/\text{SO}(8)$, $a = 3, \dots, 10$, written as the product of a matrix $\hat{\mathbf{E}}$ which only depends on the axionic moduli associated with the off-diagonal components of the metric times the diagonal matrix $\mathbf{D} = (D_m^a) = (e^{\sigma_a} \delta_m^a)$. The exponentials e^{σ_a} can be viewed as the internal radii R_a . The bosonic section of the $D = 3$ Lagrangian reads:²²

$$\begin{aligned} e^{-1} \mathcal{L}_3 = & \frac{R}{2} - \frac{1}{2} \partial_\mu \vec{h} \cdot \partial^\mu \vec{h} - \frac{1}{2} \sum_{a < b} e^{2(\sigma_b - \sigma_a)} P_{\mu a}{}^b P^\mu{}_a{}^b - \frac{1}{2} \sum_a e^{-2(\sigma_a - \xi)} F_{\mu a} F^\mu{}_a - \\ & - \frac{1}{4} \sum_{a, b} e^{2(\sigma_a + \sigma_b + \xi)} F_\mu{}^{ab} F^{\mu ab} - \frac{1}{12} \sum_{a, b, c} e^{-2(\sigma_a + \sigma_b + \sigma_c)} F_{\mu abc} F^\mu{}_{abc}, \end{aligned} \quad (\text{B.4})$$

where $P_{\mu a}{}^b \equiv (\hat{\mathbf{E}}^{-1} \partial_\mu \hat{\mathbf{E}})_a{}^b$ and the dialtonic vector \vec{h} has the following form in the (ϵ_i) orthonormal basis:

$$\vec{h} = \sum_{a=3}^9 \left(\sigma_a + \frac{\sigma_{10}}{2} \right) \epsilon_{a-2} + \left(\frac{\sigma_{10}}{2} + \sum_{a=3}^9 \sigma_a \right) \epsilon_8. \quad (\text{B.5})$$

The field strengths $F_{\mu a}$ and $F_\mu{}^{ab}$ are associated with the scalars χ_n and χ^{mn} dual in $D = 3$ to the vectors G_μ^m and $A_{\mu mn}$ respectively, while $F_{\mu abc}$ is the one pertaining to the scalars A_{mnp} . In these conventions, the lower (or upper) internal $\text{SO}(8)$ -indices a, b, c of these field strengths are related to the $\text{SL}(8, \mathbb{R})$ indices m, n, p by means of $\hat{\mathbf{E}}$ (or $\hat{\mathbf{E}}^{-1}$). For instance:

$$F_{\mu abc} = \hat{E}_a{}^m \hat{E}_b{}^n \hat{E}_c{}^p F_{\mu mnp}; \quad F_{\mu mnp} = \partial_\mu A_{mnp}. \quad (\text{B.6})$$

The above Lagrangian can also be written in the more compact form:

$$e^{-1} \mathcal{L}_3 = \frac{R}{2} - \frac{1}{2} \partial_\mu \vec{h} \cdot \partial^\mu \vec{h} - \frac{1}{2} \sum_{\alpha \in \Delta_+[\mathfrak{e}_{8(8)}} e^{-2\alpha \cdot \vec{h}} \Phi_\mu^{(\alpha)} \Phi^{\mu(\alpha)}, \quad (\text{B.7})$$

where the one-forms $\Phi_\mu^{(\alpha)}$ are associated with each of the $\mathfrak{e}_{8(8)}$ -positive roots α [46–48].²³ It is useful to express the various radial moduli σ_a in terms of the corresponding fields $\hat{\sigma}_a$ in the $D = 10$ string frame:

$$\sigma_a = \hat{\sigma}_a - \frac{\phi}{3}, \quad a = 3, \dots, 9; \quad \sigma_{10} = \frac{2}{3} \phi, \quad (\text{B.8})$$

we find:

$$\vec{h} = \sum_{i=1}^8 h_i \epsilon_i = \sum_{a=3}^9 \hat{\sigma}_a \epsilon_{a-2} + \left(-2\phi + \sum_{a=3}^9 \hat{\sigma}_a \right) \epsilon_8. \quad (\text{B.9})$$

²²We adopt the mostly plus signature for the metric.

²³The representation (B.7) of the $D = 3$ Lagrangian applies to all $D = 3$ supergravities. In the general (non necessarily maximal) case, \vec{h} is a suitable dilaton-dependent vector in the \mathfrak{h}^{nc} subspace of the Cartan subalgebra of \mathfrak{g}_N^3 , while α are the restrictions to \mathfrak{h}^{nc} of the \mathfrak{g}_N^3 positive roots (*restricted roots*, see [14]).

The outer automorphism \mathcal{O}_k in (B.1) has the effect of changing the sign to an odd number of ϵ_a , or, equivalently, to their coefficients in \vec{h} :

$$\epsilon_{i_\ell} \rightarrow -\epsilon_{i_\ell} \ ; \ \ell = 1, \dots, k. \tag{B.10}$$

To see this let us consider the effect of \mathcal{O}_k on the dilatonic part of the coset representative of $O(8,8)/[O(8) \times O(8)]$, which has the following form:

$$\mathbf{D}(\vec{h}) = \begin{pmatrix} (e^{h_i} \delta_i^j) & \mathbf{0} \\ \mathbf{0} & (e^{-h_i} \delta_i^j) \end{pmatrix}. \tag{B.11}$$

We see that:

$$\mathcal{O}_k^{-1} \mathbf{D}(\vec{h}) \mathcal{O}_k = \mathbf{D}(\vec{h}'), \tag{B.12}$$

where $h'_{i_\ell} = -h_{i_\ell}$, $h'_{i \neq i_\ell} = h_{i \neq i_\ell}$, $\ell = 1, \dots, k$. If i_ℓ run between 1 and 7, this transformation amounts to a T -duality along the internal directions $y^{i_\ell+2}$ [25, 45]:

$$R'_{i_\ell+2} = e^{\hat{\sigma}'_{i_\ell+2}} = e^{-\hat{\sigma}_{i_\ell+2}} = \frac{1}{R_{i_\ell+2}} \ ; \ \phi' = \phi - \sum_{\ell=1}^k \hat{\sigma}_{i_\ell+2}. \tag{B.13}$$

These transformations map type IIA into type IIB theory. If $k = 1$ and $i_\ell = 8$ then there is an S -duality involved: $\hat{\sigma}'_i = \hat{\sigma}_i$ and $\phi' = -\phi + \sum_{a=3}^9 \hat{\sigma}_a$.

Instead of considering inequivalent T-dual subalgebras $\mathfrak{g}, \mathfrak{g}' \subset \mathfrak{so}(8,8)$ within a same $\mathfrak{e}_{8(8)}$ algebra, we may adopt an equivalent point of view and consider a same subalgebra $\mathfrak{g} \subset \mathfrak{so}(8,8)$ within two $\mathfrak{e}_{8(8)}$ algebras, called in [25] $\mathfrak{e}_{8(8)}^+$ and $\mathfrak{e}_{8(8)}^-$,²⁴ defined respectively by completing the $\mathfrak{so}(8,8)$ Dynkin diagram with spinorial weights of different chiralities, namely attaching the weight α_1 to α_3 , as in figure 1, or a weight α'_1 to α_2 , defined as follows:

$$\alpha_1 = -\frac{1}{2} \left(\sum_{i=1}^8 \epsilon_i \right) + \epsilon_7 + \epsilon_8 \xrightarrow{\text{T-duality along } y^9} \alpha'_1 = -\frac{1}{2} \left(\sum_{i=1}^8 \epsilon_i \right) + \epsilon_8. \tag{B.14}$$

This is useful if, for instance, we fix the $\mathfrak{g} = \mathfrak{gl}(8, \mathbb{R}) \subset \mathfrak{so}(8,8)$ group to be the same in the type IIA and type IIB settings. Then the different $\mathfrak{gl}(8, \mathbb{R})$ -weights defining the dimensionally reduced type IIA and type IIB forms are obtained by branching the adjoint representations of $\mathfrak{e}_{8(8)}^+$ and $\mathfrak{e}_{8(8)}^-$, respectively, with respect to the common $\mathfrak{gl}(8, \mathbb{R})$, [25].

The doubling of the equivalence classes inside a D_n algebra into dual pairs, discussed above, does not occur if the subalgebra is the sum of commuting algebras in the case in which either all of them are of type A_k with *even* rank k , or at least one of them is of type D [42]. This is consistent with the fact observed in subsection 2.1, that the $SL(7, \mathbb{R})$ $D = 9$ Ehlers subgroups of $SO(8,8)$ which pertain to the type IIA and IIB descriptions are equivalent. The same rule guarantees that, in $D = 6$, the $SO(5,5) \times SL(4, \mathbb{R})$ subgroups of $SO(8,8)$ in the type IIA and IIB settings, are equivalent.

²⁴Actually in [25] only the $D = 4$ theory was considered, the T -duality group being $O(6,6)$ in this case, and the algebras $\mathfrak{e}_{7(7)}^\pm$ defined.

C Poincaré duality and level decomposition

Consider now the branching of the adjoint representation of \mathfrak{g}_N^3 with respect to $\mathrm{SL}(D-2) \times G_N^D$:

$$\begin{aligned} \mathrm{Adj}_{G_N^3} &\rightarrow (\mathrm{Adj}_{\mathrm{SL}(D-2)}, \mathbf{1}) \oplus (\mathbf{1}, \mathrm{Adj}_{G_N^D}) \bigoplus_d \mathfrak{N}_d, \\ \mathfrak{N}_d &= \left[(\Lambda^d, \mathcal{R}_d) \oplus (*\Lambda^d, \mathcal{R}'_d) \right], \end{aligned} \tag{C.1}$$

where it is understood that if $(\Lambda^d, \mathcal{R}_d) = (*\Lambda^d, \mathcal{R}'_d)$, they are counted just once in \mathfrak{N}_d . In light of our discussion in appendix A.3, we can write the coset space as the carrier of a representation $\bigoplus_d \mathfrak{N}_d$, namely rewrite eq. (A.26) as follows:

$$\mathfrak{g}_N^3 \ominus (\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)) = \mathfrak{N}^+ \oplus \mathfrak{N}^- = \bigoplus_d \mathfrak{N}_d. \tag{C.2}$$

In fact each subspace \mathfrak{N}_d splits into conjugate nilpotent subalgebras as follows:

$$\mathfrak{N}_d = \mathfrak{N}_d^+ \oplus \mathfrak{N}_d^-, \quad \mathfrak{N}_d^+ = \mathfrak{N}_d \cap \mathfrak{N}^+, \quad \mathfrak{N}_d^- = \mathfrak{N}_d \cap \mathfrak{N}^- = \tau(\mathfrak{N}_d^+), \tag{C.3}$$

this being a consequence of the property: $\tau(\mathfrak{N}_d) = \mathfrak{N}_d$. Each nilpotent subalgebra \mathfrak{N}_d^+ or \mathfrak{N}_d^- separately defines a representation with respect to (the adjoint action of) G_N^D and the subgroup $\mathrm{GL}(D-3) \subset \mathrm{SL}(D-2)$, though not with respect to $\mathrm{SL}(D-2)$ itself. We can decompose each space \mathfrak{N}_d into eigenspaces of the Cartan involution τ , consisting of compact and non-compact generators:

$$\mathfrak{N}_d = \mathfrak{N}_d^c \oplus \mathfrak{N}_d^{nc}, \quad \mathfrak{N}_d^c = \mathfrak{N}^c \cap \mathfrak{N}_d, \quad \mathfrak{N}_d^{nc} = \mathfrak{N}^{nc} \cap \mathfrak{N}_d. \tag{C.4}$$

These subspaces define representations with respect to the compact group $\mathrm{SO}(D-2) \times \mathrm{mcs}(G_N^D)$ and, moreover

$$\dim(\mathfrak{N}_d^c) = \dim(\mathfrak{N}_d^{nc}). \tag{C.5}$$

For the sake of simplicity, let us consider a split (maximally non-compact) \mathfrak{g}_N^3 . Then each \mathfrak{N}_d will be generated by shift operators corresponding to a certain set of positive roots $\alpha^{(d)}$ and their negatives:

$$\mathfrak{N}_d = \mathrm{Span}(E_{\alpha^{(d)}}, E_{-\alpha^{(d)}})_{\alpha^{(d)} \in \Delta_+[\mathfrak{g}_N^3]}, \tag{C.6}$$

and the conjugate nilpotent subalgebras are $\mathfrak{N}_d^+ = \mathrm{Span}(E_{\alpha^{(d)}})$ and $\mathfrak{N}_d^- = \mathrm{Span}(E_{-\alpha^{(d)}})$. The eigenspaces \mathfrak{N}_d^{nc} , \mathfrak{N}_d^c of the Cartan involution, consisting of compact and non-compact generators read:

$$\mathfrak{N}_d^c = \mathrm{Span}(E_{\alpha^{(d)}} - E_{-\alpha^{(d)}})_{\alpha^{(d)} \in \Delta_+[\mathfrak{g}_N^3]}; \quad \mathfrak{N}_d^{nc} = \mathrm{Span}(E_{\alpha^{(d)}} + E_{-\alpha^{(d)}})_{\alpha^{(d)} \in \Delta_+[\mathfrak{g}_N^3]}. \tag{C.7}$$

Each positive root $\alpha^{(d)}$ corresponds to a $D=3$ scalar field in the Lagrangian (B.7). For a given d the roots $\alpha^{(d)}$ are defined by the *level decomposition* of the \mathfrak{g}_N^3 -roots with respect to the root which is truncated out of its extended diagram in order to define the $\mathfrak{g}_N^D \oplus \mathfrak{sl}(D-2)$ -subdiagram.²⁵

²⁵In the non-split case, one should consider the level decomposition of the restricted roots. Level decompositions are a common procedure in the E_{10} and E_{11} approaches to maximal supergravity [49, 50].

Let us illustrate this procedure in the maximal theory. As shown in appendix A, the $\mathfrak{e}_{11-D(11-D)} \times \mathfrak{sl}(D-2)$ diagram is obtained by deleting from the $\mathfrak{e}_{8(8)}$ -extended Dynkin diagram the root α_{12-D} . The $\mathfrak{sl}(D-2)$ subalgebra is defined by the simple roots $\alpha_{13-D}, \dots, \alpha_8, -\psi$, $\psi = \epsilon_1 + \epsilon_8$ being the $\mathfrak{e}_{8(8)}$ highest root, while its $\mathfrak{gl}(D-3)$ subalgebra only by the roots $\alpha_{13-D}, \dots, \alpha_8$. Writing a generic $\mathfrak{e}_{8(8)}$ positive root in the simple root basis:

$$\alpha = \sum_{i=1}^8 n_i \alpha_i, \tag{C.8}$$

the positive integer n_i defines the *level* of α with respect to α_i . Let us consider the level-decomposition with respect to the root α_{12-D} for dimensions $D < 9$, namely the values of n_{12-D} defining the roots $\alpha^{(d)}$.²⁶

$D = 4$. In the case of $D = 4$ we have 63 roots with $n_8 = 0$, corresponding to the $\mathfrak{e}_{7(7)}$ -positive roots. The level $n_8 = 1$ roots are 56 and are the $\alpha^{(1)}$ -roots whose shift generators $E_{\pm\alpha^{(1)}}$ define the carrier space of the $\mathfrak{N}_{d=1} = (\mathbf{1}, \mathbf{56})$ representation. The level $n_8 = 2$ root defines, with its negative, the shift generators in the quotient $\mathfrak{sl}(D-2) \ominus \mathfrak{gl}(D-3) = \mathfrak{sl}(2) \ominus \mathfrak{gl}(1)$, which are the two shift generators of the Ehlers group.

$D = 5$. Consider now the $D = 5$ case. There are 37 level- $n_7 = 0$ roots corresponding to the positive roots of $\mathfrak{e}_{6(6)} \oplus \mathfrak{gl}(2)$. The level- $n_7 = 1$ roots are 54 and define in $\mathfrak{N}_{d=1}^+$ a subspace in the $(\mathbf{2}, \mathbf{27})$ -representation of $\mathrm{SL}(D-3) \times E_{6(6)} = \mathrm{SL}(2) \times E_{6(6)}$, while the 27 level- $n_7 = 2$ roots define a subspace in the $(\mathbf{1}, \mathbf{27}')$ with respect to the same group. The space $\mathfrak{N}_{d=1}^-$ will be the carrier of the conjugate representations. Together, the level $n_7 = 1, 2$ roots and their negatives define the space $\mathfrak{N}_{d=1} = \mathfrak{N}_{d=1}^+ \oplus \mathfrak{N}_{d=1}^- = (\mathbf{3}, \mathbf{27}) \oplus (\mathbf{3}', \mathbf{27}')$, and are collectively denoted by $\alpha^{(1)}$. Finally the 2 level- $n_7 = 3$ roots, with their negative, define the generators of the coset $\mathfrak{sl}(D-2) \ominus \mathfrak{gl}(D-3) = \mathfrak{sl}(3) \ominus \mathfrak{gl}(2)$.

$D = 6$. As far as the $D = 6$ case is concerned, the 23 level- $n_6 = 0$ roots are positive roots of $\mathfrak{gl}(3) \oplus \mathfrak{so}(5, 5)$, while the 48 level- $n_6 = 1$ and the 16 level- $n_6 = 3$ roots define generators in $\mathfrak{N}_{d=1}^+$ transforming in the $(\mathbf{3}, \mathbf{16})$ and $(\mathbf{1}, \mathbf{16}')$ of $\mathrm{SL}(D-3) \times \mathrm{SO}(5, 5) = \mathrm{SL}(3) \times \mathrm{SO}(5, 5)$, respectively. These are the $\alpha^{(1)}$ roots, which, together with their negatives, define the $\mathfrak{N}_{d=1} = (\mathbf{4}, \mathbf{16}) \oplus (\mathbf{4}', \mathbf{16}')$ space. The roots $\alpha^{(2)}$ ($d = 2$) are 30 and have $n_6 = 2$. The corresponding space $\mathfrak{N}_{d=2}^+$ is the carrier of a $(\mathbf{3}, \mathbf{10})$ representation of $\mathrm{SL}(3) \times \mathrm{SO}(5, 5)$, while $E_{\pm\alpha^{(2)}}$ generate the $\mathfrak{N}_{d=2} = (\mathbf{6}, \mathbf{10})$. Finally the 3 level- $n_6 = 4$ roots, with their negative, define the generators of the coset $\mathfrak{sl}(D-2) \ominus \mathfrak{gl}(D-3) = \mathfrak{sl}(4) \ominus \mathfrak{gl}(3)$.

A similar pattern occurs in the higher- D cases.

²⁶More precisely, the level n^i is the grading of the generator E_α with respect to the $\mathrm{SO}(1, 1)$ generator H_{λ^i} (i.e. $[H_{\lambda^i}, E_\alpha] = n^i E_\alpha$), λ^i being the \mathfrak{g}_N^3 simple weights. The level decomposition is defined by the Cartan generator which is orthogonal to the Cartan subalgebra of $\mathfrak{g}_N^D \oplus \mathfrak{gl}(D-3)$ (and therefore commutes with $\mathfrak{g}_N^D \oplus \mathfrak{gl}(D-3)$). In the maximal theory, for $D < 9$, the relevant Cartan generator is $H_{\lambda_{12-D}}$ and thus the level to consider is n_{12-D} . For $D = 9$ the generator is $H_{\lambda_2} + H_{\lambda_3}$ and so we shall consider the decomposition with respect to the integer $n = n_2 + n_3$. In the type IIA $D = 10$ description, the generator is $H_{\lambda_1} + 2H_{\lambda_2}$ and the decomposition will be effected with respect to $n = n_1 + 2n_2$.

$D = 7$. For $D = 7$, we have 16 roots with $n_5 = 0$ which are the roots of $\mathfrak{gl}(4) \oplus \mathfrak{sl}(5)$. The 40 $n_5 = 1$ and the 10 $n_5 = 4$ roots define subspaces of $\mathfrak{N}_{d=1}^+$ in the $(4, \mathbf{10}')$ and $(\mathbf{1}, \mathbf{10})$ of $\mathrm{SL}(D-3) \times \mathrm{SL}(5) = \mathrm{SL}(4) \times \mathrm{SL}(5)$, respectively. These are the $\alpha^{(1)}$ -roots and the space $\mathfrak{N}_{d=1} = \mathfrak{N}_{d=1}^+ \oplus \mathfrak{N}_{d=1}^-$ is the carrier of the representation $(\mathbf{5}, \mathbf{10}') \oplus (\mathbf{5}', \mathbf{10})$ of $\mathrm{SL}(5) \times \mathrm{SL}(5)$. The $\alpha^{(2)}$ -roots consist in the 30 level- $n_5 = 2$ and the 20 level- $n_5 = 3$ roots defining the representations $(\mathbf{6}, \mathbf{5})$ and $(\mathbf{4}', \mathbf{5}')$ of $\mathrm{SL}(4) \times \mathrm{SL}(5)$ in $\mathfrak{N}_{d=2}^+$, respectively. The space $\mathfrak{N}_{d=2} = \mathfrak{N}_{d=2}^+ \oplus \mathfrak{N}_{d=2}^-$ then defines the representation $(\mathbf{10}, \mathbf{5}) \oplus (\mathbf{10}', \mathbf{5}')$ of $\mathrm{SL}(5) \times \mathrm{SL}(5)$. Finally the 4 level- $n_5 = 5$ roots, with their negative, define the generators of the coset $\mathfrak{sl}(D-2) \ominus \mathfrak{gl}(D-3) = \mathfrak{sl}(5) \ominus \mathfrak{gl}(4)$.

$D = 8$. In the $D = 8$ case the 14 $n_4 = 0$ roots are the positive roots of $\mathfrak{gl}(5) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(3)$. The $\alpha^{(1)}$ s consist of the 30 $n_4 = 1$ and the 6 $n_4 = 5$ roots defining the $\mathrm{SL}(5) \times \mathrm{SL}(2) \times \mathrm{SL}(3)$ -representations $(\mathbf{5}, \mathbf{2}, \mathbf{3}') \oplus (\mathbf{1}, \mathbf{2}, \mathbf{3})$ in $\mathfrak{N}_{d=1}^+$ which, together with the conjugate representations in $\mathfrak{N}_{d=1}^-$, complete the $\mathfrak{N}_{d=1} = (\mathbf{6}, \mathbf{2}, \mathbf{3}') \oplus (\mathbf{6}', \mathbf{2}, \mathbf{3})$ of $\mathrm{SL}(6) \times \mathrm{SL}(2) \times \mathrm{SL}(3)$. The $\alpha^{(2)}$ s are defined by the 30 $n_4 = 2$ and 15 $n_4 = 4$ roots, corresponding to the representation $(\mathbf{10}, \mathbf{1}, \mathbf{3}) \oplus (\mathbf{5}', \mathbf{1}, \mathbf{3}')$ in $\mathfrak{N}_{d=2}^+$, so that $\mathfrak{N}_{d=2} = \mathfrak{N}_{d=2}^+ \oplus \mathfrak{N}_{d=2}^- = (\mathbf{15}, \mathbf{1}, \mathbf{3}) \oplus (\mathbf{15}', \mathbf{1}, \mathbf{3}')$ of $\mathrm{SL}(6) \times \mathrm{SL}(2) \times \mathrm{SL}(3)$. The $\alpha^{(3)}$ roots are the 20 ones with $n_4 = 3$. They define the $\mathfrak{N}_{d=3}^+$ space in the $(\mathbf{10}, \mathbf{2}, \mathbf{1})$ of $\mathrm{SL}(5) \times \mathrm{SL}(2) \times \mathrm{SL}(3)$ which, together with $\mathfrak{N}_{d=3}^-$, complete the $\mathfrak{N}_{d=3} = (\mathbf{20}, \mathbf{2}, \mathbf{1})$ of $\mathrm{SL}(6) \times \mathrm{SL}(2) \times \mathrm{SL}(3)$. The remaining 5 roots with $n_4 = 6$, with their negative, define the generators of the coset $\mathfrak{sl}(D-2) \ominus \mathfrak{gl}(D-3) = \mathfrak{sl}(6) \ominus \mathfrak{gl}(5)$.

$D = 9$. The same analysis applies to $D = 9$, although in this case we shall consider the sum $n = n_2 + n_3$. There are 16 roots with $n = 0$, which are the positive roots of the algebra $\mathfrak{gl}(6) \oplus \mathfrak{gl}(2)$. The $\alpha^{(1)}$ roots consist of the 18 with $n = 1$ and the 3 with $n = 6$, defining the $\mathrm{SL}(6) \times \mathrm{GL}(2)$ -representations $(\mathbf{6}, \mathbf{2}_{+3} + \mathbf{1}_{-4}) \oplus (\mathbf{1}, \mathbf{2}_{-3} + \mathbf{1}_{+4})$ in $\mathfrak{N}_{d=1}^+$ which, together with the conjugate representations in $\mathfrak{N}_{d=1}^-$, complete the $\mathfrak{N}_{d=1} = (\mathbf{7}, \mathbf{2}_{+3} + \mathbf{1}_{-4}) \oplus (\mathbf{7}', \mathbf{2}_{-3} + \mathbf{1}_{+4})$ of $\mathrm{SL}(7) \times \mathrm{GL}(2)$. The $\alpha^{(2)}$ s are the 30 roots with $n = 2$ and the 12 with $n = 5$ defining the $\mathrm{SL}(6) \times \mathrm{GL}(2)$ -representations $(\mathbf{15}, \mathbf{2}_{-1}) \oplus (\mathbf{6}', \mathbf{2}_{+1})$ in $\mathfrak{N}_{d=2}^+$, so that $\mathfrak{N}_{d=2} = \mathfrak{N}_{d=2}^+ \oplus \mathfrak{N}_{d=2}^- = (\mathbf{21}, \mathbf{2}_{-1}) \oplus (\mathbf{21}', \mathbf{2}_{+1})$. Next we have to consider the 20 $n = 3$ and the 15 $n = 4$ roots which make the $\alpha^{(3)}$ and define the $\mathrm{SL}(6) \times \mathrm{GL}(2)$ -representations $(\mathbf{20}, \mathbf{1}_{+2}) \oplus (\mathbf{15}', \mathbf{1}_{-2})$ in $\mathfrak{N}_{d=3}^+$, so that $\mathfrak{N}_{d=3} = \mathfrak{N}_{d=3}^+ \oplus \mathfrak{N}_{d=3}^- = (\mathbf{35}, \mathbf{1}_{+2}) \oplus (\mathbf{35}', \mathbf{1}_{-2})$. Finally the 6 $n = 7$ roots, with their negative, define the generators of the coset $\mathfrak{sl}(D-2) \ominus \mathfrak{gl}(D-3) = \mathfrak{sl}(7) \ominus \mathfrak{gl}(6)$.

$D = 10$, IIB. In the $D = 10$ case we have to distinguish between the type IIA and IIB theories. In the type IIB setting we need consider the level n_3 with respect to α_3 . The 22 roots with $n_3 = 0$ are the positive roots of $\mathfrak{gl}(7) \oplus \mathfrak{sl}(2)$, $\mathfrak{sl}(2)$ being the U -duality group. In this case we only have $d = 2, 4$. The $\alpha^{(2)}$ s consist of the 42 $n_3 = 1$ and the 14 $n_3 = 3$ defining the $\mathrm{SL}(7) \times \mathrm{SL}(2)$ -representations $(\mathbf{21}, \mathbf{2}) \oplus (\mathbf{7}', \mathbf{2})$ in $\mathfrak{N}_{d=2}^+$, so that $\mathfrak{N}_{d=2} = \mathfrak{N}_{d=2}^+ \oplus \mathfrak{N}_{d=2}^- = (\mathbf{28}, \mathbf{2}) \oplus (\mathbf{28}', \mathbf{2})$ of $\mathrm{SL}(8) \times \mathrm{SL}(2)$. Next we have the 35 roots with $n_3 = 2$, which are the $\alpha^{(4)}$ s and define the $(\mathbf{35}, \mathbf{1})$ of $\mathrm{SL}(7) \times \mathrm{SL}(2)$ in $\mathfrak{N}_{d=4}^+$, so that $\mathfrak{N}_{d=4} = \mathfrak{N}_{d=4}^+ \oplus \mathfrak{N}_{d=4}^- = (\mathbf{70}, \mathbf{1})$. There are 7 $n_3 = 4$ roots which, with their negative, define the generators of the coset $\mathfrak{sl}(D-2) \ominus \mathfrak{gl}(D-3) = \mathfrak{sl}(8) \ominus \mathfrak{gl}(7)$.

$D = 10$, IIA. As far as the type IIA description is concerned, the level to consider for the decomposition is the sum $n = n_1 + 2n_2$. In this case we only have $d = 1, 2, 3$. There are

21 $n = 0$ roots which are the positive roots of $\mathfrak{gl}(7) \oplus \mathfrak{so}(1, 1)$, $\mathfrak{so}(1, 1)$ being the U -duality algebra. The $\alpha^{(1)}$ roots consist of the 7 roots with $n = 1$ and the single $n = 7$ root defining the $\mathrm{SL}(7) \times \mathrm{SO}(1, 1)$ -representation $\mathbf{7}_{+3} \oplus \mathbf{1}_{-3}$ in $\mathfrak{N}_{d=1}^+$ which, together with the conjugate representations in $\mathfrak{N}_{d=1}^-$, complete the $\mathfrak{N}_{d=1} = \mathbf{8}_{+3} \oplus \mathbf{8}_{-3}$ of $\mathrm{SL}(8) \times \mathrm{SO}(1, 1)$. Next we consider the 21 roots with $n = 2$ and the 7 with $n = 6$, whose shift operators generating $\mathfrak{N}_{d=2}^+$ transform in the $\mathbf{21}_{-2} \oplus \mathbf{7}'_{+2}$ with respect to $\mathrm{SL}(7) \times \mathrm{SO}(1, 1)$. These roots define then the $\alpha^{(2)}$ and $\mathfrak{N}_{d=2} = \mathfrak{N}_{d=2}^+ \oplus \mathfrak{N}_{d=2}^- = \mathbf{28}_{-2} \oplus \mathbf{28}'_{+2}$ of $\mathrm{SL}(8) \times \mathrm{SO}(1, 1)$. The $\alpha^{(3)}$ s consist of the 35 $n = 3$ and the 21 $n = 6$ roots defining the $\mathrm{SL}(7) \times \mathrm{SO}(1, 1)$ -representation $\mathbf{35}_{+1} \oplus \mathbf{21}'_{-1}$ in $\mathfrak{N}_{d=3}^+$ which, together with the conjugate representations in $\mathfrak{N}_{d=3}^-$, complete the $\mathfrak{N}_{d=3} = \mathbf{56}_{+1} \oplus \mathbf{56}'_{-1}$. There are no roots with $n = 5$ while those with $n = 8$ are 7 and, with their negative, define the generators of the coset $\mathfrak{sl}(D - 2) \ominus \mathfrak{gl}(D - 3) = \mathfrak{sl}(8) \ominus \mathfrak{gl}(7)$.

$D = 11$. We end this analysis with the $D = 11$ case discussed in the previous section. The relevant level decomposition is with respect to the root α_2 . With $n_2 = 0$ we have the positive roots of $\mathfrak{gl}(8)$. In this case we only have $d = 3$ and the $\alpha^{(3)}$ -roots consist of the 56 $n_2 = 1$ and the 28 $n_2 = 2$ ones defining the $\mathrm{SL}(8) \times \mathrm{SO}(1, 1)$ -representation $\mathbf{56}_{+1} \oplus \mathbf{28}'_{+2}$ in $\mathfrak{N}_{d=3}^+$ which, together with the conjugate representations in $\mathfrak{N}_{d=3}^-$, completes the $\mathfrak{N}_{d=3} = \mathbf{84} \oplus \mathbf{84}'$. Finally the 8 $n_2 = 3$ roots, with their negative, define the generators of the coset $\mathfrak{sl}(D - 2) \ominus \mathfrak{gl}(D - 3) = \mathfrak{sl}(9) \ominus \mathfrak{gl}(8)$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] M. Goroff and J.H. Schwarz, *D-dimensional gravity in the light cone gauge*, *Phys. Lett. B* **127** (1983) 61 [INSPIRE].
- [2] E. Cremmer and B. Julia, *The $N = 8$ Supergravity Theory. 1. The Lagrangian*, *Phys. Lett. B* **80** (1978) 48 [INSPIRE].
- [3] E. Cremmer and B. Julia, *The $\mathrm{SO}(8)$ Supergravity*, *Nucl. Phys. B* **159** (1979) 141 [INSPIRE].
- [4] C. Hull and P. Townsend, *Unity of superstring dualities*, *Nucl. Phys. B* **438** (1995) 109 [[hep-th/9410167](#)] [INSPIRE].
- [5] E. Cremmer, B. Julia, H. Lü and C. Pope, *Higher dimensional origin of $D = 3$ coset symmetries*, [hep-th/9909099](#) [INSPIRE].
- [6] H. Lü and C. Pope, *P-brane solitons in maximal supergravities*, *Nucl. Phys. B* **465** (1996) 127 [[hep-th/9512012](#)] [INSPIRE].
- [7] N. Marcus and J.H. Schwarz, *Three-Dimensional Supergravity Theories*, *Nucl. Phys. B* **228** (1983) 145 [INSPIRE].
- [8] E. Cremmer, *Supergravities in 5 Dimensions*, in *Superspace and Supergravity*, S.W. Hawking and M. Rocek eds., Cambridge University Press, Cambridge, U.K. (1981).
- [9] B. Julia, *Group Disintegrations*, in *Superspace and Supergravity*, S.W. Hawking and M. Rocek eds., Cambridge University Press, Cambridge, U.K. (1981).

- [10] S. Ananth, L. Brink and P. Ramond, *Eleven-dimensional supergravity in light-cone superspace*, *JHEP* **05** (2005) 003 [[hep-th/0501079](#)] [[INSPIRE](#)].
- [11] L. Brink, S.-S. Kim and P. Ramond, $E_{8(8)}$ in Light Cone Superspace, *JHEP* **07** (2008) 113 [[arXiv:0804.4300](#)] [[INSPIRE](#)].
- [12] L. Brink, S.-S. Kim and P. Ramond, $E_{7(7)}$ on the Light Cone, *JHEP* **06** (2008) 034 [[arXiv:0801.2993](#)] [[INSPIRE](#)].
- [13] B. de Wit, A. Tollsten and H. Nicolai, *Locally supersymmetric $D = 3$ nonlinear σ -models*, *Nucl. Phys. B* **392** (1993) 3 [[hep-th/9208074](#)] [[INSPIRE](#)].
- [14] S. Helgason, *Differential Geometry, Lie Groups and Symmetric Spaces*, Academic Press, New York, U.S.A. (1978).
- [15] R. Gilmore, *Lie Groups, Lie Algebras, and Some of Their Applications*, Dover Publications (2006).
- [16] A. Keurentjes, *The Group theory of oxidation*, *Nucl. Phys. B* **658** (2003) 303 [[hep-th/0210178](#)] [[INSPIRE](#)].
- [17] A. Keurentjes, *The Group theory of oxidation 2: Cosets of nonsplit groups*, *Nucl. Phys. B* **658** (2003) 348 [[hep-th/0212024](#)] [[INSPIRE](#)].
- [18] N. Marcus, A. Sagnotti and J.H. Schwarz, *Infinite symmetry algebras of extended supergravity theories*, *Nucl. Phys. B* **243** (1984) 335 [[INSPIRE](#)].
- [19] M. Günaydin, G. Sierra and P. Townsend, *Exceptional Supergravity Theories and the MAGIC Square*, *Phys. Lett. B* **133** (1983) 72 [[INSPIRE](#)].
- [20] M. Günaydin, G. Sierra and P. Townsend, *The Geometry of $N = 2$ Maxwell-Einstein Supergravity and Jordan Algebras*, *Nucl. Phys. B* **242** (1984) 244 [[INSPIRE](#)].
- [21] M. Günaydin, *Lectures on Spectrum Generating Symmetries and U-duality in Supergravity, Extremal Black Holes, Quantum Attractors and Harmonic Superspace*, [arXiv:0908.0374](#) [[INSPIRE](#)].
- [22] P. Truini, *Exceptional Lie Algebras, $SU(3)$ and Jordan Pairs*, [arXiv:1112.1258](#) [[INSPIRE](#)].
- [23] W. Nahm, *Supersymmetries and their Representations*, *Nucl. Phys. B* **135** (1978) 149 [[INSPIRE](#)].
- [24] E. Cremmer, B. Julia and J. Scherk, *Supergravity Theory in Eleven-Dimensions*, *Phys. Lett. B* **76** (1978) 409 [[INSPIRE](#)].
- [25] M. Bertolini and M. Trigiante, *Regular RR and NS-NS BPS black holes*, *Int. J. Mod. Phys. A* **15** (2000) 5017 [[hep-th/9910237](#)] [[INSPIRE](#)].
- [26] R. D'Auria, S. Ferrara, M. Lledó and V. Varadarajan, *Spinor algebras*, *J. Geom. Phys.* **40** (2001) 101 [[hep-th/0010124](#)] [[INSPIRE](#)].
- [27] S. Ferrara, C.A. Savoy and B. Zumino, *General massive multiplets in extended supersymmetry*, *Phys. Lett. B* **100** (1981) 393 [[INSPIRE](#)].
- [28] G. Compere, S. de Buyl, E. Jamsin and A. Virmani, G_2 Dualities in $D = 5$ Supergravity and Black Strings, *Class. Quant. Grav.* **26** (2009) 125016 [[arXiv:0903.1645](#)] [[INSPIRE](#)].
- [29] E. Bergshoeff, W. Chemissany, A. Ploegh, M. Trigiante and T. Van Riet, *Generating Geodesic Flows and Supergravity Solutions*, *Nucl. Phys. B* **812** (2009) 343 [[arXiv:0806.2310](#)] [[INSPIRE](#)].
- [30] L. Borsten, M.J. Duff, S. Ferrara, A. Marrani and W. Rubens, *Small Orbits*, *Phys. Rev. D* **85** (2012) 086002 [[arXiv:1108.0424](#)] [[INSPIRE](#)].

- [31] S. Cecotti, S. Ferrara and L. Girardello, *Geometry of Type II Superstrings and the Moduli of Superconformal Field Theories*, *Int. J. Mod. Phys. A* **4** (1989) 2475 [INSPIRE].
- [32] E. Cremmer and A. Van Proeyen, *Classification of Kähler manifolds in $N = 2$ vector multiplet supergravity couplings*, *Class. Quant. Grav.* **2** (1985) 445 [INSPIRE].
- [33] J. Luciani, *Coupling of $O(2)$ Supergravity with Several Vector Multiplets*, *Nucl. Phys. B* **132** (1978) 325 [INSPIRE].
- [34] J. Bagger and E. Witten, *Matter Couplings in $N = 2$ Supergravity*, *Nucl. Phys. B* **222** (1983) 1 [INSPIRE].
- [35] A. Salam and E. Sezgin, *Anomaly freedom in chiral supergravities*, *Phys. Scripta* **32** (1985) 283.
- [36] S. Randjbar-Daemi, A. Salam, E. Sezgin and J. Strathdee, *An Anomaly Free Model in Six-Dimensions*, *Phys. Lett. B* **151** (1985) 351 [INSPIRE].
- [37] S. Ferrara, F. Riccioni and A. Sagnotti, *Tensor and vector multiplets in six-dimensional supergravity*, *Nucl. Phys. B* **519** (1998) 115 [hep-th/9711059] [INSPIRE].
- [38] F. Riccioni and A. Sagnotti, *Consistent and covariant anomalies in six-dimensional supergravity*, *Phys. Lett. B* **436** (1998) 298 [hep-th/9806129] [INSPIRE].
- [39] H. Nishino and E. Sezgin, *New couplings of six-dimensional supergravity*, *Nucl. Phys. B* **505** (1997) 497 [hep-th/9703075] [INSPIRE].
- [40] M. Günaydin, H. Samtleben and E. Sezgin, *On the Magical Supergravities in Six Dimensions*, *Nucl. Phys. B* **848** (2011) 62 [arXiv:1012.1818] [INSPIRE].
- [41] D.V. Alekseevskii, *Classification of quaternionic spaces with a transitive solvable group of motions*, *Math. USSR Izvestija* **9** (1975) 297.
- [42] E.B. Dynkin, *Semisimple Subalgebras of Semisimple Lie Algebras*, *American Mathematical Society Translations, Series 2* **6** (1957) 111.
- [43] E.B. Dynkin, *The Maximal Subgroups of the Classical Groups*, *American Mathematical Society Translations, Series 2* **6** (1957) 245.
- [44] A. Minchenko, *The Semisimple Subalgebras of Exceptional Lie Algebras*, *Trans. Moscow Math. Soc.* **67** (2006) 225.
- [45] H. Lü, C. Pope and K. Stelle, *Weyl group invariance and p -brane multiplets*, *Nucl. Phys. B* **476** (1996) 89 [hep-th/9602140] [INSPIRE].
- [46] L. Andrianopoli, R. D’Auria, S. Ferrara, P. Fré and M. Trigiante, *RR scalars, U duality and solvable Lie algebras*, *Nucl. Phys. B* **496** (1997) 617 [hep-th/9611014] [INSPIRE].
- [47] L. Andrianopoli, R. D’Auria, S. Ferrara, P. Fré, R. Minasian and M. Trigiante, *Solvable Lie algebras in type IIA, type IIB and M theories*, *Nucl. Phys. B* **493** (1997) 249 [hep-th/9612202] [INSPIRE].
- [48] E. Cremmer, B. Julia, H. Lü and C. Pope, *Dualization of dualities. 1.*, *Nucl. Phys. B* **523** (1998) 73 [hep-th/9710119] [INSPIRE].
- [49] A. Kleinschmidt and H. Nicolai, *E_{10} and $SO(9,9)$ invariant supergravity*, *JHEP* **07** (2004) 041 [hep-th/0407101] [INSPIRE].
- [50] E.A. Bergshoeff, M. de Roo, S.F. Kerstan, T. Ortín and F. Riccioni, *IIA ten-forms and the gauge algebras of maximal supergravity theories*, *JHEP* **07** (2006) 018 [hep-th/0602280] [INSPIRE].