

Progress on the symmetric Strassen conjecture.

Original

Progress on the symmetric Strassen conjecture / Carlini, E., Catalisano, M.V., Chiantini, L.. - In: JOURNAL OF PURE AND APPLIED ALGEBRA. - ISSN 0022-4049. - STAMPA. - 219:8(2015), pp. 3149-3157. [10.1016/j.jpaa.2014.10.006]

Availability:

This version is available at: 11583/2617538 since: 2015-09-30T15:04:59Z

Publisher:

Elsevier

Published

DOI:10.1016/j.jpaa.2014.10.006

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

PROGRESS ON THE SYMMETRIC STRASSEN CONJECTURE

ENRICO CARLINI, MARIA VIRGINIA CATALISANO, AND LUCA CHIANTINI

ABSTRACT. Let F and G be homogeneous polynomials in disjoint sets of variables. We prove that the Waring rank is additive, thus proving the symmetric Strassen conjecture, when either F or G is a power, or F and G have two variables, or either F or G has small rank.

1. INTRODUCTION

In his famous result of 1969 Strassen showed that it is possible to multiply two 2×2 matrices using seven basic operations rather than eight, see [Lan08] for more on this. Using this fact, a better algorithm was produced to multiply matrices of any size and this was proved to have the best possible computational complexity by Winograd in [Win71].

After Strassen's result, it was clear that even straightforward procedures can require fewer operations than expected. In [Str73] Strassen formulated his well known *additive conjecture* for bilinear maps: *Given bilinear maps ϕ, ψ and two pairs of matrices A, B , and C, D the computational complexity of simultaneously computing $\phi(A, B)$ and $\psi(C, D)$ is the sum of the complexities of ϕ and ψ .* The conjecture stands open since its formulation in 1973, for some partial results see [FW84].

Strassen conjecture can be naturally stated in terms of tensors and the notion of tensor rank, see [Lan12]. Note that an analogue of the additive conjecture for *approximate* complexity (border rank, in more recent terminology) does not hold (see [Sch81]).

We will focus on the relevant case of symmetric tensors, that is, the case of homogeneous polynomials, also known as forms. The *rank*, or *Waring rank*, of a degree d form F is

$$\text{rk}(F) = \min\{r : F = L_1^d + \dots + L_r^d \text{ for linear forms } L_i\}.$$

We can now state the symmetric version of the Strassen Additivity Conjecture (SAC)

Conjecture. (SAC). Let $d > 1$. If $F \in \mathbb{C}[x_0, \dots, x_n]$ and $G \in \mathbb{C}[y_0, \dots, y_m]$ are non-zero forms of degree d , then

$$\text{rk}(F + G) = \text{rk}(F) + \text{rk}(G).$$

Also the symmetric version of the conjecture stands open. A relevant contribution to its study is the 2012 paper [CCG12] where SAC is proved for the sum of (several) monomials.

In this paper we contribute to the study of SAC by proving that the conjecture holds for one and two variables and for forms of small rank. More precisely, we prove the following theorem.

Theorem 1.1. *Let $d > 1$ and let $F \in \mathbb{C}[x_0, \dots, x_n]$, $G \in \mathbb{C}[y_0, \dots, y_m]$ be non-zero degree d forms. If $n = 0$ or $m = 0$ or $n, m = 1$, then $\text{rk}(F + G) = \text{rk}(F) + \text{rk}(G)$.*

In Corollary 3.2 we show that SAC holds even with no assumptions on n and m if one of the forms satisfies a condition on the rank (see the statement of the corollary).

We obtain Theorem 1.1 using algebraic geometry. In particular, we exploit a deep knowledge of the Hilbert functions of finite sets of points. A similar approach was considered also in [BB12]. We will use the strong structural result contained in [BGM94]. Our techniques can be extended to study SAC in more than two variables, but we need some genericity assumptions on the forms F and G ; we are currently investigating this approach.

2. PRELIMINARIES

In this section we state our notation and we recall the main results that we will use.

Notation 2.1. For a zero-dimensional scheme $Z \subset \mathbb{P}^N$, we will denote with h_Z the Hilbert function of Z , that is the function which associates to any integer n the dimension of the image of the map $H^0 \mathcal{O}_{\mathbb{P}^N}(n) \rightarrow H^0 \mathcal{O}_Z(n)$. We will denote with Dh_Z the first difference of the Hilbert function, that is $Dh_Z(n+1) = h_Z(n+1) - h_Z(n)$.

Notation 2.2. In this paper, F and G will denote forms of degree d , with $d > 1$, in disjoint sets of variables. Namely, $F \in \mathbb{C}[x_0, \dots, x_n]$ and $G \in \mathbb{C}[y_0, \dots, y_m]$.

We will use the following formulation of SAC. Let $\text{rk}(F) = r$, $\text{rk}(G) = s$, and write

$$F = F_1^d + \dots + F_r^d \text{ and } G = G_1^d + \dots + G_s^d,$$

for linear forms $F_i \in \mathbb{C}[x_0, \dots, x_n]$ and $G_i \in \mathbb{C}[y_0, \dots, y_m]$. With these notation SAC is equivalent to the fact that it is not possible to have

$$(1) \quad F_1^d + \dots + F_r^d + G_1^d + \dots + G_s^d = H_1^d + \dots + H_t^d$$

where $t < r + s$ and the H_i are linear forms in $\mathbb{C}[x_0, \dots, x_n, y_0, \dots, y_m]$.

We begin with a reduction step.

Lemma 2.3. *To prove SAC as stated in (1), it is enough to consider*

$$H_i \notin \mathbb{C}[x_0, \dots, x_n] \cup \mathbb{C}[y_0, \dots, y_m].$$

Proof. Assume that we have F, G as in (1), with a of the H_i 's (say the first a of them) belonging to $\mathbb{C}[x_0, \dots, x_n]$, and b of the H_i 's (say the last b) belonging to $\mathbb{C}[y_0, \dots, y_m]$.

Notice that $F' = F_1^d + \dots + F_r^d - H_1^d - \dots - H_a^d$ is a form of degree d in $\mathbb{C}[x_0, \dots, x_n]$, whose rank is at least $r - a$. Otherwise we could write F as a form of rank less than $r - a$ plus the sum of a powers of linear forms, contradicting the fact that $\text{rk}(F) = r$.

Similarly, $G' = G_1^d + \dots + G_s^d - H_{t-b+1}^d - \dots - H_t^d$ is a form of degree d in $\mathbb{C}[y_0, \dots, y_m]$, whose rank is at least $s - b$.

Then $F' + G' = H_{a+1}^d + \cdots + H_{t-b+1}^d$, with $\text{rk}(F') \geq r - a$, $\text{rk}(G') \geq s - b$. The remaining forms H_i are $t - a - b < r - a + s - b$ and none of them lies either in $\mathbb{C}[x_0, \dots, x_n]$ or in $\mathbb{C}[y_0, \dots, y_m]$. Assuming that SAC holds in this situation yields a contradiction, thus proving the conjecture for any choice of the forms H_i . \square

We recall some basic properties of binary forms, that is forms in two variables.

Remark 2.4. The linear span of a rational normal curve of degree d contains all binary forms in a given set of variables. Hence, any set of $i \geq d + 1$ degree d binary forms is linearly dependent. Moreover, any degree d binary form has rank at most d , see [CS11] for more details.

3. ONE VARIABLE

Assume that either F or G is a polynomial in one variable. In this case, a direct computation proves that SAC holds.

Proposition 3.1. *If $n = 0$ or $m = 0$, then SAC holds.*

Proof. We will prove the statement for $m = 0$. Let $F(x_0, \dots, x_n) \neq 0$ be a degree $d > 1$ form in $k[x_0, \dots, x_n]$ and let $G(y_0) = y_0^d$. We set

$$P(x_0, \dots, x_n, y_0) = F(x_0, \dots, x_n) + G(y_0),$$

and we will show that $\text{rk}(P) = \text{rk}(F) + \text{rk}(G) = \text{rk}(F) + 1$.

Let $\text{rk}(F) = r$. For $r = 1$, the conclusion immediately follows, so assume $r > 1$. Clearly $\text{rk}(P) \leq r + 1$. If $\text{rk}(P) < r$, since $F(x_0, \dots, x_n) = P(x_0, \dots, x_n, 0)$, we get a contradiction.

We now prove that it is not possible to have $\text{rk}(P) = r$. Let

$$P = (\alpha_1 y_0 + L_1)^d + \cdots + (\alpha_r y_0 + L_r)^d,$$

where the L_i are linear forms in x_0, \dots, x_n . By setting $y_0 = 0$ we have that

$$F = L_1^d + \cdots + L_r^d,$$

and thus

$$L_1^d + \cdots + L_r^d + y_0^d = (\alpha_1 y_0 + L_1)^d + (\alpha_2 y_0 + L_2)^d + \cdots + (\alpha_r y_0 + L_r)^d.$$

Since $G \neq 0$, the α_i 's cannot all be equal to zero. Assume that $\alpha_1 \neq 0$. Let γ be a d -th root of the identity. Since $d > 1$, we may assume that $\gamma \neq \alpha_1$.

Now set

$$y_0 = \frac{L_1}{\gamma - \alpha_1},$$

so that

$$(\alpha_1 y_0 + L_1)^d = \left(\frac{\gamma L_1}{\gamma - \alpha_1} \right)^d = \left(\frac{L_1}{\gamma - \alpha_1} \right)^d = y_0^d.$$

Hence we get

$$F = \left(\alpha_2 \left(\frac{L_1}{\gamma - \alpha_1} \right) + L_2 \right)^d + \cdots + \left(\alpha_r \left(\frac{L_1}{\gamma - \alpha_1} \right) + L_r \right)^d.$$

It follows that $\text{rk}(F) < r$, a contradiction. Hence $\text{rk}(P) = r + 1$. \square

To extend this result, we use apolarity theory, see [Ger96, IK99]. Recall that $(F^\perp)_1$ is the vector space of linear differential operator vanishing on a form F . Following [Car06] and [BBS08] we say that $F \in \mathbb{C}[x_0, \dots, x_n]$ *essentially involves* N variables if

$$\dim(F^\perp)_1 = n + 1 - N.$$

In other words, F essentially involves N variables if there exists a linear change of variables such that $F \in \mathbb{C}[z_0, \dots, z_{N-1}] \subset \mathbb{C}[x_0, \dots, x_n]$.

Corollary 3.2. *If F essentially involves $\text{rk}(F)$ variables, or G essentially involves $\text{rk}(G)$ variables, then SAC holds.*

Proof. We let $r = \text{rk}(F)$, $s = \text{rk}(G)$ and we prove the statement in the case that G essentially involves s variables. After a change of variables we may assume

$$G = y_0^d + \dots + y_{s-1}^d.$$

Arguing by induction on s , we see that $F + y_0^d + \dots + y_{s-2}^d$ has rank $r + s - 1$, thus by Proposition 3.1, $F + G = (F + y_0^d + \dots + y_{s-2}^d) + y_{s-1}^d$ has rank $(r + s - 1) + 1 = r + s$. \square

4. SAC AND THE HILBERT FUNCTION

Here we introduce our algebro-geometric approach to SAC with the relative notation.

Notation 4.1. As above let $F = F_1^d + \dots + F_r^d$, $r = \text{rk}(F)$ and $G = G_1^d + \dots + G_s^d$, $s = \text{rk}(G)$, where the forms F_i and the G_i have degree one. We introduce the sets

$$\begin{aligned} Z(F) &= \{[F_1], \dots, [F_r]\} \subset \mathbb{P}^n = \mathbb{P}(\mathbb{C}[x_0, \dots, x_n]), \\ Z(G) &= \{[G_1], \dots, [G_s]\} \subset \mathbb{P}^m = \mathbb{P}(\mathbb{C}[y_0, \dots, y_m]). \end{aligned}$$

We will argue by contradiction assuming that

$$(2) \quad F + G = H_1^d + \dots + H_t^d,$$

for $t < r + s$ and linear forms H_i . Finally, we set

$$\begin{aligned} Z(H) &= \{[H_1], \dots, [H_t]\} \subset \mathbb{P}^{n+m+1} = \mathbb{P}(\mathbb{C}[x_0, \dots, x_n, y_0, \dots, y_m]), \\ Z &= Z(H) \cup Z(F) \cup Z(G) \subset \mathbb{P}^{n+m+1}. \end{aligned}$$

We will use the following geometric remarks.

Remark 4.2. The standard exact sequence of sheaves

$$0 \rightarrow \mathcal{I}_Z(d) \rightarrow \mathcal{O}_{\mathbb{P}^N}(d) \rightarrow \mathcal{O}_Z \rightarrow 0$$

yields

$$0 \rightarrow H^0 \mathcal{I}_Z(d) \rightarrow H^0 \mathcal{O}_{\mathbb{P}^N}(d) \rightarrow H^0 \mathcal{O}_Z \rightarrow H^1 \mathcal{I}_Z(d) \rightarrow 0.$$

By the definitions, we have that $h_Z(d) = |Z| (= \dim H^0 \mathcal{O}_Z)$ if and only if $H^1 \mathcal{I}_Z(d) = 0$. In this case, we will say that Z is *separated in degree d* or that Z *imposes independent conditions to degree d forms*.

Remark 4.3. The equality in (2) implies that the following are linearly dependent forms:

$$F_1^d, \dots, F_r^d, G_1^d, \dots, G_m^d, H_1^d, \dots, H_t^d.$$

Thus, by Lemma 1 in [BB12], Z does not impose independent conditions to forms of degree d . If we denote with ν the d -th Veronese map, one of the points of $\nu(Z)$, say F_i^d , depends linearly on the others. Thus, any linear form which vanishes on

$\nu(Z) \setminus \{F_i^d\}$ also vanishes at F_i^d . Hence, any form of degree d vanishing on $Z \setminus \{F_i\}$ also vanishes at F_i . In other words, Z is not separated in degree d , and the ideal sheaf of Z is such that

$$h^1 \mathcal{I}_Z(d) > 0.$$

Remark 4.4. Let $L_F \simeq \mathbb{P}^n$ and $L_G \simeq \mathbb{P}^m$, be two skew linear subspaces of \mathbb{P}^{n+m+1} , containing $Z(F)$ and $Z(G)$, respectively. The following holds: if for some $i > 0$

$$h^1 \mathcal{I}_{Z(F)}(i) = h^1 \mathcal{I}_{Z(G)}(i) = 0,$$

then

$$h^1 \mathcal{I}_{Z(F) \cup Z(G)}(i) = 0.$$

To see this, we argue as follows. If P is a point of $Z(F)$, consider a degree i hypersurface $X \subset L_F$ that contains all the points of $Z(F) \setminus \{P\}$. The cone on X with vertex L_G is a hypersurface of degree i in \mathbb{P}^{n+m+1} . This cone separates P from the other points of $Z(F) \cup Z(G)$. A similar argument works if $P \in Z(G)$. Thus $Z(F) \cup Z(G)$ is separated in degree i .

The next two lemmas will be useful in the following section.

Lemma 4.5. *Let W be a set of w distinct points spanning a linear space of dimension at least three. If $w \leq 2u - 2$ and $h^1 \mathcal{I}_W(u - 2) > 0$, then there exists a line that contains at least u points of W .*

Proof. Consider the first difference Dh_W of the Hilbert function of W . Since

$$h^1 \mathcal{I}_W(u - 2) > 0,$$

then $Dh_W(u - 1) > 0$, so that $Dh_W(i) > 0$ for all $i = 0, \dots, u - 1$. Moreover $Dh_W(1) \geq 3$, since the points span at least a space of dimension 3. Since $w \leq 2u - 2$, we cannot have $Dh_W(i) \geq 2$ for $i = 2, \dots, u - 2$. It follows that there are at least two values $2 \leq i_1, i_2 \leq u - 1$ for which $Dh_W(i_1) = Dh_W(i_2) = 1$. By Theorem 3.6 in [BGM94] it follows that we may assume the two steps are $i_1 = u - 2$ and $i_2 = u - 1$ and moreover u points of W are aligned. \square

Lemma 4.6. *Let $W_1, W_2 \subset \mathbb{P}^n$ be finite sets of points such that W_1 is contained in a proper linear subspace R , and $W_2 \cap R = \emptyset$. If for some integer u one has:*

- (1) $h^1 \mathcal{I}_{W_2}(u - 1) = 0$ and
- (2) the cardinality of W_1 is at most $u + 1$,

then $h^1 \mathcal{I}_W(u) = 0$, where $W = W_1 \cup W_2$.

Proof. If we let w_i be the cardinality of W_i , then it is easy to see that

$$h^0 \mathcal{O}_{\mathbb{P}^n}(u) - w_1 - w_2 \leq h^0 \mathcal{I}_W(u).$$

We now use Castelnuovo's inequality, see [CCG11] Section 3 for more details and [AH95] Section 2 for a proof. This inequality yields

$$h^0 \mathcal{I}_W(u) \leq h^0 \mathcal{I}_{Res}(u - 1) + h^0 \mathcal{I}_{Tr}(u),$$

where Tr is the trace of W on a generic hyperplane $H \supseteq R \supset W_1$ and Res is the residue of W with respect to H . More precisely $\mathcal{I}_{Tr} = \mathcal{I}_{W,H} = \mathcal{I}_{W_1}$ and $\mathcal{I}_{Res} = \mathcal{I}_W : \mathcal{I}_H = \mathcal{I}_{W_2}$, that is $Res = W_2$ and $Tr = W_1$. From (2) we get that $h^0 \mathcal{I}_{Tr}(u) = h^0 \mathcal{O}_{\mathbb{P}^{n-1}}(u) - w_1$. From (1) we get that $h^0 \mathcal{I}_{Res}(u - 1) = h^0 \mathcal{O}_{\mathbb{P}^n}(u - 1) - w_2$. Hence, the result follows. \square

5. FORMS IN TWO VARIABLES

In this section, we assume that F, G are forms in two variables, that is we assume $n = m = 1$, and we prove that SAC holds in this setting. We let $r = \text{rk}(F)$, $s = \text{rk}(G)$ and we assume that

$$F_1^d + \cdots + F_r^d + G_1^d + \cdots + G_s^d = H_1^d + \cdots + H_t^d$$

for $t < r + s$ and for linear forms $F_i \in \mathbb{C}[x_0, x_1]$ and $G_i \in \mathbb{C}[y_0, y_1]$, while the H_i are linear forms involving all the four variables y_0, y_1, x_0 , and x_1 (see Lemma 2.3).

Remark 5.1. By Proposition 3.1 and its Corollary 3.2, we may assume $r, s \geq 3$. Moreover, as noted in Remark 2.4 the rank of a form of degree d in two variables is at most d . So we have $r, s \leq d$.

We treat the cases of large r, s using the Apolarity Lemma, see [IK99].

Lemma 5.2. *If $r = s = d$, then SAC holds.*

Proof. If F is a binary degree d form of rank d , then the Hilbert-Burch Theorem (see Theorem 20.15 in [Eis95]) yields that $F^\perp = (A, B)$ where $\deg(A) = 2$, $\deg(B) = d$ and A is a square. From this we see that $F = LM^{d-1}$ for some linear forms L and M , thus F is a monomial. The conclusion follows as the rank is additive on coprime monomials, see [CCG12]. \square

From now on we assume that:

$$s \leq r \text{ and } s \leq d - 1.$$

We now show that $Z(H)$ spans \mathbb{P}^3 and that at most d points of $Z(H)$ lie on a line.

Lemma 5.3. *There is no plane containing $Z(H)$.*

Proof. Since $Z(H) \subset \mathbb{P}(\mathbb{C}[x_0, x_1, y_0, y_1]) = \mathbb{P}^3$, we use apolarity theory to describe the dual space \mathbb{P}^{3*} . Thus, \mathbb{P}^{3*} can be identified with the space of linear differential operators on the polynomial ring $\mathbb{C}[x_0, x_1, y_0, y_1]$. We now argue by contradiction. If a plane containing $Z(H)$ exists, then the plane is the vanishing locus of some differential operator ∂ . Thus ∂ vanishes on $Z(H)$, and thus $\partial(H) = \partial(F + G) = 0$, that is $\partial(F) = -\partial(G)$. Note that $\partial(F) \in \mathbb{C}[x_0, x_1]$ and $\partial(G) \in \mathbb{C}[y_0, y_1]$ are forms of degree $d - 1 > 0$, hence $\partial(F) = \partial(G) = 0$. The latter is impossible, since F, G are binary forms of rank at least three. \square

Remark 5.4. If t is the minimal integer such that $F + G = H_1^d + \cdots + H_t^d$, then we may assume that at most d points of $Z(H)$ lie on a line.

Indeed, if H_1, \dots, H_{d+1} lie on a line, then by Remark 2.4 there is a linear relation between their d -th powers, say $H_1^d = a_2 H_2^d + \cdots + a_{d+1} H_{d+1}^d$, so that

$$F + G = (1 + a_2)H_2^d + \cdots + (1 + a_{d+1})H_{d+1}^d + H_{d+2}^d + \cdots + H_t^d$$

against the minimality of t .

In our notation $Z \subset \mathbb{P}^3 = \mathbb{P}(\mathbb{C}[x_0, x_1, y_0, y_1])$ and the subsets $Z(F)$ and $Z(G)$ span two skew lines, namely the lines corresponding to $\mathbb{P}(\mathbb{C}[x_0, x_1])$ and $\mathbb{P}(\mathbb{C}[y_0, y_1])$. We will denote these lines with $R_F \supset Z(F)$ and $R_G \supset Z(G)$.

Lemma 5.5. *There is no line R containing d points of $Z(H)$ and intersecting R_G .*

Proof. Assume by contradiction that such a line R exists, say $R \supset \{H_1, \dots, H_d\}$. The projection π from R_G to R_F is obtained by setting $y_0 = y_1 = 0$. Since π contracts R to a point, it follows that F is generated by the d -th powers of the images of H_1 and H_{d+1}, \dots, H_t . So the rank of F is at most $t - d + 1$. On the other hand $t - d + 1 < r + s - d + 1 \leq r$, since we are assuming $s \leq d - 1$. Hence we have a contradiction. \square

We now prove SAC for two variables.

Theorem 5.6. *Let $d > 1$. If $F \in \mathbb{C}[x_0, x_1]$ and $G \in \mathbb{C}[y_0, y_1]$ are non-zero degree d forms, then*

$$\text{rk}(F + G) = \text{rk}(F) + \text{rk}(G).$$

Proof. We will use all the previous notation and we may assume what follows:

- $3 \leq r \leq d$, $3 \leq s \leq d - 1$, and thus $d \geq 4$ (see Remark 5.1 and Proposition 5.2).
- $Z(H)$ does not intersect $R_F \cup R_G$ (see Lemma 2.3).
- $Z(H)$ spans \mathbb{P}^3 (see Lemma 5.3).
- At most d points of $Z(H)$ lie on a line (see Remark 5.4).

By contradiction assume that $t < r + s$ and thus $t \leq 2d - 2$. We let

$$Z = Z(F) \cup Z(G) \cup Z(H)$$

and we will find the contradiction $h^1\mathcal{I}_Z(d) = 0$ (see Remark 4.3).

Case 1: there is no line containing d points of $Z(H)$. Lemma 4.5, with $u = d$, yields that $h^1\mathcal{I}_{Z(H)}(d - 2) = 0$. Since $Z(G)$ has cardinality at most $d - 1$, we apply Lemma 4.6 with $u = d - 1$, and obtain $h^1\mathcal{I}_{Z(H) \cup Z(G)}(d - 1) = 0$. Finally, since $Z(F)$ has cardinality at most d , we may apply Lemma 4.6 with $u = d$, and obtain $h^1\mathcal{I}_Z(d) = 0$, a contradiction.

Case 2: there are d points of $Z(H)$ on a line R . By Lemma 5.5, R cannot intersect R_G . R may or may not intersect R_F and, in the first case, the point $R \cap R_F$ may or may not lie in $Z(F)$. In any case, set $\Phi = R \cap R_F$ and set $Z' = Z(H) \setminus R$, $Z'' = (Z(H) \cap R) \cup \Phi$. Notice that Z' contains at most $d - 2$ points and Z'' contains at most $d + 1$ points. Notice also that Z is contained in $Z(F) \cup Z(G) \cup Z'' \cup Z'$, with equality if and only if either $\Phi = \emptyset$ or Φ is not a point of $Z(F)$.

Since the cardinality of Z' is at most $d - 2$, then $h^1\mathcal{I}_{Z'}(d - 3) = 0$. Because $Z(G)$ has cardinality at most $d - 1$, we apply Lemma 4.6 with $u = d - 2$, and obtain $h^1\mathcal{I}_{Z' \cup Z(G)}(d - 2) = 0$. Since $Z(F) \setminus \Phi$ has cardinality at most d , we may apply Lemma 4.6 with $u = d - 1$, and obtain $h^1\mathcal{I}_{Z' \cup Z(G) \cup (Z(F) \setminus \Phi)}(d - 1) = 0$. Finally, notice that $Z' \cup Z(G) \cup (Z(F) \setminus \Phi)$ cannot intersect R . Thus, since Z'' has cardinality at most $d + 1$, we apply Lemma 4.6 with $u = d$, and we obtain

$$h^1\mathcal{I}_{Z' \cup Z(G) \cup (Z(F) \setminus \Phi) \cup Z''}(d) = 0.$$

Since $Z \subseteq Z' \cup Z(G) \cup (Z(F) \setminus \Phi) \cup Z''$ we get the contradiction

$$h^1\mathcal{I}_Z(d) \leq h^1\mathcal{I}_{Z' \cup Z(G) \cup (Z(F) \setminus \Phi) \cup Z''}(d) = 0.$$

\square

Theorem 1.1 is now proved by collecting the previous results.

Proof of Theorem 1.1. The cases $n = 0$ and $m = 0$ follow from Proposition 3.1 and Theorem 5.6 yields the rest of the statement. \square

REFERENCES

- [AH95] J. Alexander and A. Hirschowitz. Polynomial interpolation in several variables. *J. Algebraic Geom.*, 4(2):201–222, 1995.
- [BB12] E. Ballico and A. Bernardi. Decomposition of homogeneous polynomials with low rank. *Math. Z.*, 271(3-4):1141–1149, 2012.
- [BBS08] A. Białynicki-Birula and A. Schinzel. Representations of multivariate polynomials by sums of univariate polynomials in linear forms. *Colloq. Math.*, 112(2):201–233, 2008.
- [BGM94] A. Bigatti, A. V. Geramita, and J. C. Migliore. Geometric consequences of extremal behavior in a theorem of Macaulay. *Trans. Amer. Math. Soc.*, 346(1):203–235, 1994.
- [Car06] E. Carlini. Reducing the number of variables of a polynomial. In *Algebraic geometry and geometric modeling*, Math. Vis., pages 237–247. Springer, Berlin, 2006.
- [CCG11] E. Carlini, M. V. Catalisano, and A. V. Geramita. 3-dimensional sundials. *Cent. Eur. J. Math.*, 9(5):949–971, 2011.
- [CCG12] E. Carlini, M.V. Catalisano, and A. V. Geramita. The solution to the Waring problem for monomials and the sum of coprime monomials. *J. Algebra*, 370:5–14, 2012.
- [CS11] Gonzalo Comas and Malena Seiguer. On the rank of a binary form. *Found. Comput. Math.*, 11(1):65–78, 2011.
- [Eis95] D. Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [FW84] E. Feig and S. Winograd. On the direct sum conjecture. *Linear Algebra Appl.*, 63:193–219, 1984.
- [Ger96] A.V. Geramita. Inverse systems of fat points: Waring’s problem, secant varieties of Veronese varieties and parameter spaces for Gorenstein ideals. In *The Curves Seminar at Queen’s, Vol. X (Kingston, ON, 1995)*, volume 102 of *Queen’s Papers in Pure and Appl. Math.*, pages 2–114. Queen’s Univ., Kingston, ON, 1996.
- [IK99] A. Iarrobino and V. Kanev. *Power sums, Gorenstein algebras, and determinantal loci*, volume 1721 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1999.
- [Lan08] J. M. Landsberg. Geometry and the complexity of matrix multiplication. *Bull. Amer. Math. Soc. (N.S.)*, 45(2):247–284, 2008.
- [Lan12] J. M. Landsberg. *Tensors: geometry and applications*, volume 128 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2012.
- [Sch81] A. Schönhage. Partial and total matrix multiplication. *SIAM J. Comput.*, 10(3):434–455, 1981.
- [Str73] V. Strassen. Vermeidung von Divisionen. *J. Reine Angew. Math.*, 264:184–202, 1973.
- [Win71] S. Winograd. On multiplication of 2×2 matrices. *Linear Algebra and Appl.*, 4:381–388, 1971.

(E. Carlini) DEPARTMENT OF MATHEMATICAL SCIENCES, POLITECNICO DI TORINO, TURIN, ITALY

E-mail address: `enrico.carlini@polito.it`

(E.Carlini) SCHOOL OF MATHEMATICAL SCIENCES, MONASH UNIVERSITY, CLAYTON, AUSTRALIA.

E-mail address: `enrico.carlini@monash.edu`

(M.V. Catalisano) DIME - DIPARTIMENTO DI INGEGNERIA MECCANICA, ENERGETICA, GESTIONALE E DEI TRASPORTI, UNIVERSITY OF GENOA, ITALY.

E-mail address: `catalisano@diptem.unige.it`

(L. Chiantini) DEPARTMENT OF INFORMATION ENGINEERING AND MATHEMATICS, UNIVERSITY OF SIENA, ITALY.

E-mail address: `luca.chiantini@unisi.it`