Breathomics can discriminate between anti IgE-treated and non-treated severe asthma adults

Original

Breathomics can discriminate between anti IgE-treated and non-treated severe asthma adults / Santini, Giuseppe; DI CARLO, Stefano; Benso, Alfredo; Mores, Nadia; Brinkmann, Paul; Valente, Salvatore; Montuschi, Paolo; Macagno, Francesco; Politano, GIANFRANCO MICHELE MARIA; Wagener, Ariane H.; Bansal, Aruna T.; Knobel, Hugo H.; Vink, Anton J.; Rattray, Nicholas; Santonico, Marco; Pennazza, Giorgio; Wang, Yuanyue; Horvath, Ildiko; Djukanovic, Ratko; Polosa, Riccardo; Fowler, Stephen J.; Chanez, Pascal; Chung, Kian F.; Sterk, Peter J.; Montuschi, Paolo. - In: EUROPEAN RESPIRATORY JOURNAL. - ISSN 1399-3003. - ELETTRONICO. - 46:Suppl. 59, abstract number OA1463(2015), pp. 1-1

This version is available at: 11583/2622355 since: 2017-05-02T13:07:29Z

Publisher:

European Respiratory Society

Published

DOI:

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

VOLUME 46 / SUPPLEMENT 59 / SEPTEMBER 2015

EUROPEAN RESPIRATORY *journal*

OFFICIAL SCIENTIFIC JOURNAL OF THE ERS

Abstracts / *25th International Congress*
Amsterdam, Netherlands 26 –30 September 2015

Online ISSN: 1399-3003

ERS

EUROPEAN
RESPIRATORY
SOCIETY

every breath counts

Copyright for individual abstracts remains with the authors.

This abstract supplement has been produced electronically by the European Respiratory Society. The European Respiratory Society is not responsible for errors or omissions in content. The ideas and opinions expressed in this publication do not necessarily reflect those of Coe-Truman and the European Respiratory Society. Products mentioned in this publication should not be construed as an endorsement of the product or the manufacturer's claims. Readers are encouraged to contact the manufacturer with any questions about the features or limitations of the products mentioned. The European Respiratory Society assumes no responsibility for any injury and/or damage to persons or property arising out of or related to any use of the material contained in these abstracts. The reader is advised to check the appropriate medical literature and the product information currently provided by the manufacturer of each drug to be administered to verify the dosage, the method and duration of administration, or contraindications. It is the responsibility of the treating physician or other health care professional, relying on independent experience and knowledge of the patient, to determine drug dosages and the best treatment for the patient. An effort has been made to check generic and trade names, and to verify drug doses. The ultimate responsibility, however, lies with the prescribing physician. Please convey any errors to scientific@ersnet.org.

Citations should be made in the following way: **Authors. Title. Eur Respir J 20015; 46: Suppl. 59, abstract number.**

Table Of Content

160. Phenotyping asthma with biomarkers	2
OA1463: Breathomics can discriminate between anti IgE-treated and non-treated severe asthma adults	2

160. Phenotyping asthma with biomarkers

OA1463

Breathomics can discriminate between anti IgE-treated and non-treated severe asthma adults

Giuseppe Santini¹, Stefano Di Carlo², Alfredo Benso², Nadia Mores¹, Paul Brinkman³, Salvatore Valente⁴, Paolo Montuschi², Francesco Macagno⁴, Gianfranco Politano², Ariane H. Wagener³, Aruna T. Bansal⁵, Hugo H. Knobel⁶, Anton J. Vink⁶, Nicholas Rattray⁷, Marco Santonico⁸, Giorgio Pennazza⁸, Yuanyue Wang⁶, Ildiko Horvath⁹, Ratko Djukanovic¹⁰, Riccardo Polosa¹¹, Stephen J. Fowler⁷, Pascal Chanez¹², Kian F. Chung¹³, Peter J. Sterk³, Paolo Montuschi¹, U-BIOPRED Study Group

¹Pharmacology, Catholic University of the Sacred Heart, Rome, Italy; ²Computer Engineering, Polytechnic of Turin, Turin, Italy; ³Respiratory Medicine, Academic Medical Centre, University of Amsterdam, Amsterdam, Netherlands; ⁴Internal Medicine and Geriatrics, Catholic University of the Sacred Heart, Rome, Italy; ⁵Biostatistics, Acclarogen Ltd., Cambridge, United Kingdom; ⁶Physical Chemistry, Philips Research Laboratories, Eindhoven, Netherlands; ⁷Respiratory Medicine, University of Manchester, Manchester, United Kingdom; ⁸Electronic Engineering, Università Campus Bio-Medico, Rome, Italy; ⁹Respiratory Medicine, Semmelweis University, Budapest, Hungary; ¹⁰Respiratory Medicine, University of Southampton, Southampton, United Kingdom; ¹¹Internal Medicine, University of Catania, Catania, Italy; ¹²Respiratory Medicine, University of Marseille, Marseille, France; ¹³Respiratory Medicine, Imperial College, London, United Kingdom

Rationale: Omalizumab, an anti-IgE monoclonal antibody, is indicated in adults with severe persistent allergic asthma. Exhaled molecular markers can provide phenotypic information in asthma. **Objectives:** Determine whether adults with severe asthma on omalizumab (anti-IgE⁺) have a different breathprint compared with those who were not on anti-IgE therapy (anti-IgE⁻) as assessed by eNoses and gas chromatography/mass spectrometry (GC/MS) (breathomics). **Methods:** This was a cross-sectional analysis of the U-BIOPRED adult cohort. Severe asthma was defined by IMI-criteria [Bel, Thorax 2011]. Anti-IgE⁺ patients were on a regular treatment with s.c. omalizumab (150-375 mg) every 2-4 weeks. Exhaled volatile compounds trapped on adsorption tubes were analysed by a centralized eNose platform (Owlstone Lonestar, two Cyranose 320, Comon Invent, Tor Vergata TEN), including a total of 190 sensors, and GC/MS. Recursive feature elimination (<http://topepo.github.io/caret/rfe.html>) was used for feature selection and random forests, more robust to overfitting, for classification. **Results:** 9 anti-IgE⁺ (females/males 2/7, age 52.6±16.3 years, mean±SD, 1/2/6 current/ex/nonsmokers, pre-bronchodilator FEV₁ 70.6±21.1% predicted value) and 30 anti-IgE⁻ patients (18/12 females/males, age 53.2±14.2 years, 0/16/14 current/ex/nonsmokers, pre-bronchodilator FEV₁ 59.6±30.7% predicted value) were studied. Accuracy of classification is shown in Table 1.

Table 1

eNose/Technique	Variables	Accuracy
TOR	4	0.87
LONE	71	0.83
CYRA2	12	0.82
CI	2	0.87
CYRA1	14	0.75
All eNoses	110	0.85
GC/MS	96	0.83
eNoses+GC/MS	16	0.83

Conclusions: Preliminary results suggest that breathomics can distinguish between anti-IgE⁺ and anti-IgE⁻ severe asthma patients.