

Efficient spectral domain technique for the frequency locking analysis of nonlinear oscillators

Original

Efficient spectral domain technique for the frequency locking analysis of nonlinear oscillators / Bonnin, M.; Traversa, F. L.; Bonani, F.. - In: THE EUROPEAN PHYSICAL JOURNAL PLUS. - ISSN 2190-5444. - ELETTRONICO. - 133:7(2018), p. 246. [10.1140/epjp/i2018-12076-0]

Availability:

This version is available at: 11583/2710432 since: 2018-07-19T12:10:17Z

Publisher:

Springer

Published

DOI:10.1140/epjp/i2018-12076-0

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Springer postprint/Author's Accepted Manuscript

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: <http://dx.doi.org/10.1140/epjp/i2018-12076-0>

(Article begins on next page)

HADAMARD PRODUCTS OF LINEAR SPACES

CRISTIANO BOCCI, ENRICO CARLINI AND JOE KILEEL

ABSTRACT. We describe properties of Hadamard products of algebraic varieties. We show any Hadamard power of a line is a linear space, and we construct star configurations from products of collinear points. Tropical geometry is used to find the degree of Hadamard products of other linear spaces.

1. INTRODUCTION

The concept of Hadamard product, as matrix entry-wise multiplication, is well known in linear algebra: it has nice properties in matrix analysis ([15, 16, 21]) and has applications in both statistics and physics ([22, 23, 24, 25]). Recently, in the papers [8, 9], the authors use this entry-wise multiplication to define a Hadamard product between projective varieties: given varieties $X, Y \subset \mathbb{P}^n$, their Hadamard product $X \star Y$ is the closure of the image of the rational map

$$X \times Y \dashrightarrow \mathbb{P}^n, \quad ([a_0 : \cdots : a_n], [b_0 : \cdots : b_n]) \mapsto [a_0 b_0 : a_1 b_1 : \cdots : a_n b_n].$$

For any projective variety X , we may consider its Hadamard square $X^{\star 2} = X \star X$ and its higher Hadamard powers $X^{\star r} = X \star X^{\star(r-1)}$. In [8], the authors use this definition to describe the algebraic variety associated to the restricted Boltzmann machine, which is the undirected graphical model for binary random variables specified by the bipartite graph $K_{r,n}$. This variety is the r -th Hadamard power of the first secant variety of $(\mathbb{P}^1)^n$. Note that [9] concerns the case $r = 2, n = 4$.

Hadamard products and powers are in fact well-connected to other operations of varieties. They are the multiplicative analogs of joins and secant varieties, and in tropical geometry, tropicalized Hadamard products equal Minkowski sums. It is natural to study properties of this new operation, and see its effect on various varieties. This paper is a first step in that direction. Here is how it is organized.

In Section 2, we start by giving a different definition of the Hadamard product of varieties in terms of projections of Segre products. As a first important result we give a Hadamard version of Terracini's Lemma (Lemma 2.12) which describes the tangent space of $X \star Y$ at $p \star q$ as the span $\langle p \star T_q(Y), q \star T_p(X) \rangle$. We also point out varieties parametrized by monomials are closed under Hadamard product.

In Section 3, our attention is fixed on the Hadamard powers of a line. This case is special and admits direct analysis by projective geometry. Theorem 3.4 is that the powers of a line are linear spaces, and our proof uses the Hadamard version of Terracini's Lemma. Proposition 3.6 then offers explicit equations.

In Section 4, we study the r -th square-free Hadamard power $Z^{\star r}$ of a finite set Z of projective points. We obtain a classification when Z is collinear, using star

2010 *Mathematics Subject Classification.* 14T05, 14N05, 68W30, 14M99.

Key words and phrases. Hadamard products, linear spaces, star configurations, tropical geometry, bracket polynomials.

configurations. In contrast with the standard approach to construct star configurations as intersections of a set of randomly chosen linear spaces, which can give points with complicated coordinates, Theorem 4.7 permits a cheaper construction, easily implementable in computer algebra software.

In Section 5, we recall basic definitions in tropical geometry, and give the precise connection to Hadamard products. Motivated by tropical considerations, we define a refined expected dimension formula for Hadamard products. Interestingly, Hadamard products can have deficient dimension (Example 5.5).

In Section 6, we analyze Hadamard products of linear spaces in general. This case requires the use of tropical machinery, and we derive a result analogous to identifiability in the theory of secant varieties along the way. The paper culminates with a general degree formula for Hadamard products of linear spaces (Theorem 6.8), an extension to include *reciprocal linear spaces* (Corollary 6.10), and then two non-trivial bracket polynomial formulas.

2. PRELIMINARIES

We work over the field of complex numbers \mathbb{C} .

Definition 2.1. Let $H_i \subset \mathbb{P}^n, i = 0, \dots, n$, be the hyperplane $x_i = 0$ and set

$$\Delta_i = \bigcup_{0 \leq j_1 < \dots < j_{n-i} \leq n} H_{j_1} \cap \dots \cap H_{j_{n-i}}.$$

In other words, Δ_i is the i -dimensional variety of points having *at most* $i + 1$ non-zero coordinates. Thus Δ_0 is the set of coordinate points and Δ_{n-1} is the union of the coordinate hyperplanes. Note that elements of Δ_i have *at least* $n - i$ zero coordinates. We have the following chain of inclusions:

$$(2.1) \quad \Delta_0 = \{[1 : 0 : \dots : 0], \dots, [0 : \dots : 0 : 1]\} \subset \Delta_1 \subset \dots \subset \Delta_{n-1} \subset \Delta_n = \mathbb{P}^n.$$

Definition 2.2. Given varieties $X, Y \subset \mathbb{P}^n$ we consider the usual Segre product

$$X \times Y \subset \mathbb{P}^N$$

$$([a_0 : \dots : a_n], [b_0 : \dots : b_n]) \mapsto [a_0 b_0 : a_0 b_1 : \dots : a_n b_n]$$

and we denote with z_{ij} the coordinates in \mathbb{P}^N . Let $\pi : \mathbb{P}^N \dashrightarrow \mathbb{P}^n$ be the projection map from the linear space Λ defined by equations $z_{ii} = 0, i = 0, \dots, n$. The *Hadamard product* of X and Y is

$$X \star Y = \overline{\pi(X \times Y)},$$

where the closure is taken in the Zariski topology.

Definition 2.3. Let $p, q \in \mathbb{P}^n$ be two points of coordinates respectively $[a_0 : a_1 : \dots : a_n]$ and $[b_0 : b_1 : \dots : b_n]$. If $a_i b_i \neq 0$ for some i , their Hadamard product $p \star q$ of p and q , is defined as

$$p \star q = [a_0 b_0 : a_1 b_1 : \dots : a_n b_n].$$

If $a_i b_i = 0$ for all $i = 0, \dots, n$ then we say $p \star q$ is not defined.

Remark 2.4. We have that

$$X \star Y = \overline{\{p \star q : p \in X, q \in Y, p \star q \text{ is defined}\}}.$$

Hence, our definition of Hadamard product corresponds with the one in [8] and [9].

Remark 2.5. To understand the role of the closure operation in Definition 2.2 of Hadamard product we can proceed as follows. If π is defined on all of $X \times Y$, then no closure is needed, by [11, Corollary 14.2]. Otherwise, we can blow up along $\Lambda \cap (X \times Y)$. The points of $X \star Y$ are hence limits of $p(t) \star q(s)$ for curves $p(t) \in X$ and $q(s) \in Y$, in the Euclidean topology by [29, §I.10, Corollary 1]. Thus the closure operation only adds points obtained when $p(t) \star q(s)$ goes to a point of $\Lambda \cap (X \times Y)$.

Remark 2.6. In general, it is difficult to compute $X \star Y$. The situation seems as bad as computing joins. Suppose we have for input the defining homogeneous ideals I and J of X and Y in \mathbb{P}^n , and we wish to output the defining ideal K of $X \star Y$. It may be achieved with an elimination as follows:

- Work in the ring $\mathbb{C}[x_0, \dots, x_n, y_0, \dots, y_n, z_0, \dots, z_n]$ with $3n + 3$ variables.
- Form the ideal $I(x) + J(y) + \langle z_0 - x_0 y_0, \dots, z_n - x_n y_n \rangle$.
- Eliminate the $2n + 2$ variables x_0, \dots, y_n .

However, this elimination is a heavy computation. We know of no efficient algorithm to compute Hadamard products in general.

Note that $X \star Y$ is a variety such that $\dim(X \star Y) \leq \dim(X) + \dim(Y)$ and that $X \star Y$ can be empty even if neither X nor Y is empty. In Section 5, we will give a refined upper bound on $\dim(X \star Y)$.

Note that if X, Y are irreducible, then $X \times Y$ is irreducible, so $X \star Y = \overline{\pi(X \times Y)}$ is irreducible. Also, associativity of Hadamard product of points extends to associativity of Hadamard product of varieties:

Lemma 2.7. *If $X, Y, Z \subset \mathbb{P}^n$ are varieties, then $(X \star Y) \star Z = X \star (Y \star Z)$.*

Proof. In fact, both sides equal

$$\Pi = \overline{\{p \star q \star z : p \in X, q \in Y, z \in Z, p \star q \star z \text{ is defined}\}}.$$

To see $(X \star Y) \star Z \subset \Pi$, let $s \in (X \star Y) \star Z, z \in Z$ and f be a homogeneous polynomial in the ideal $I(\Pi)$. Consider $f(z_0 \cdot -, z_1 \cdot -, \dots, z_n \cdot -)$ a polynomial defined up to scale. It vanishes on $\{p \star q : p \in X, q \in Y, p \star q \text{ is defined}\}$, and hence at s in the closure. The other inclusions then are clear. \square

In what follows we will often work with Hadamard products of a variety with itself, thus we give the following:

Definition 2.8. Given a positive integer r and a variety $X \subset \mathbb{P}^n$, the r -th Hadamard power of X is

$$X^{\star r} = X \star X^{\star(r-1)},$$

where $X^{\star 0} = [1 : \dots : 1]$.

Note that $\dim(X^{\star r}) \leq r \dim(X)$ and the r -th Hadamard power cannot be empty if X is not empty.

As the definition of Hadamard product involves a closure operation, it is not trivial to describe all points of $X^{\star r}$. However, we can say something about the maximal number of zero coordinates of any point in the Hadamard power.

Lemma 2.9. *If $X \cap \Delta_{n-i} = \emptyset$, then $X^{\star r} \cap \Delta_{n-ri+r-1} = \emptyset$.*

Proof. The hypothesis means that all points of X have at most $i-1$ zero coordinates. By Remark 2.5 any point in $X^{\star r}$ is the limit of the Hadamard product of r points in X . Thus, any point in $X^{\star r}$ has at most $ri-r$ zero coordinates. Hence, $X^{\star r} \cap \Delta_{n-ri+r-1} = \emptyset$. \square

In the case of lines we have a stronger result.

Lemma 2.10. *Let $n \geq 2$ and $n \geq r$. If $L \subset \mathbb{P}^n$ is a line such that $L \cap \Delta_{n-2} = \emptyset$, then*

$$L^{\star r} = \bigcup_{p_i \in L} p_1 \star \dots \star p_r,$$

that is, the closure operation is not necessary.

Proof. We use Remark 2.5 and we note that the map π is everywhere defined. \square

Example 2.11. For contrast, consider the curve $X \subset \mathbb{P}^2$ defined by $x_0x_1 - x_2^2 = 0$ and the point $p = [1 : 0 : 2]$. The Hadamard product $p \star X$ is the line $x_1 = 0$. However the point $[0 : 0 : 1]$ cannot be obtained as $p \star x$, with $x \in X$.

We now prove the analogue of Terracini's Lemma for Hadamard products using tangent spaces; recall that for a variety X and a point $p \in X$, $T_p(X)$ denotes the tangent space to X at p .

Lemma 2.12. *Consider varieties $X, Y \subset \mathbb{P}^n$. If $p \in X$ and $q \in Y$ are general points, then*

$$T_{p \star q}(X \star Y) = \langle p \star T_q(Y), q \star T_p(X) \rangle.$$

Moreover, if $p_1, \dots, p_r \in X$ are general points and $p_1 \star \dots \star p_r \in X^{\star r}$ is a general point, then

$$T_{p_1 \star \dots \star p_r}(X^{\star r}) = \langle p_2 \star \dots \star p_r \star T_{p_1}(X), \dots, p_1 \star \dots \star p_{r-1} \star T_{p_r}(X) \rangle.$$

Proof. It is enough to prove the result for X and Y . Since p and q are generic, by generic smoothness, we can find a parametrization of X , respectively of Y , in an analytic neighbourhood of p , respectively of q . Let $p(x)$ with $p(0) = p$, respectively $q(y)$ with $q(0) = q$, be such a parametrization. Thus, a parameterization of $X \star Y$ around $p \star q$, using our base field is \mathbb{C} and Proposition 10.6 in [14], is given by

$$p(x) \star q(y).$$

Thus the tangent space $T_{p \star q}(X \star Y)$ is obtained by picking curves $x(t)$ and $y(t)$, taking the derivative with respect to t at $t = 0$ in the expression

$$p(x(t)) \star q(y(t))$$

and making $x'(0)$ and $y'(0)$ vary. The result follows. \square

We note in passing that toric varieties, in the sense of [33], are closed under Hadamard products.

Remark 2.13. Let $\mathcal{A} = (\mathbf{a}_0, \dots, \mathbf{a}_n)$ be a $d \times (n+1)$ integer matrix in which all column sums are the same. This defines $X \subset \mathbb{P}^n$ as the closure of the image of the monomial map:

$$\begin{aligned} (\mathbb{C}^\times)^d &\longrightarrow \mathbb{P}^n \\ \mathbf{t} &\longmapsto [\mathbf{t}^{\mathbf{a}_0} : \dots : \mathbf{t}^{\mathbf{a}_n}]. \end{aligned}$$

Let \mathcal{B} be an $e \times (n+1)$ integer matrix similarly defining $Y \subset \mathbb{P}^n$. Then $X \star Y$ is defined by the $(d+e) \times (n+1)$ integer matrix which is the vertical concatenation of \mathcal{A} and \mathcal{B} .

It would be interesting to study Hadamard products of *T-varieties with low complexity* ([18]) in future work.

3. POWERS OF A LINE

In this section, our aim is to give a direct and elementary proof by projective geometry that Hadamard powers of a line are linear spaces. This is achieved in Theorem 3.4, and made explicit in Proposition 3.6.

3.1. Multiplying by a point. We begin with useful results about multiplying linear spaces by a point, which amounts to projecting into a coordinate subspace, then acting by an element of that subspace's torus.

Lemma 3.1. *Let $L \subset \mathbb{P}^n$ be a linear space of dimension m . If $p \in \mathbb{P}^n$, then $p \star L$ is either empty or it is a linear space of dimension at most m . If $p \notin \Delta_{n-1}$, then $\dim(p \star L) = m$.*

Proof. We first consider the case $p \notin \Delta_{n-1}$, that is, p has no coordinate equal to zero. We can describe L as the solution set of a linear system of equations of the form $M\mathbf{x} = 0$. Let D_p be the diagonal matrix with entries a choice of coordinates of p and consider the matrix $M' = MD_p^{-1}$. If $q \in L$, then $p \star q$ solves $M'\mathbf{x} = 0$. Conversely, if z solves $M'\mathbf{x} = 0$, then $z = p \star q$ for some $q \in L$. Thus, we have

$$\bigcup_{q \in L} p \star q = \{\mathbf{x} : M'\mathbf{x} = 0\}.$$

Since linear spaces are closed in the Zariski topology, it follows from Remark 2.4 that $p \star L = \{\mathbf{x} : M'\mathbf{x} = 0\}$. This completes the proof since M and M' have the same rank.

Now we consider the case $p \in \Delta_i \setminus \Delta_{i-1}$ for $i < n$, that is, p has exactly $n-i$ coordinates equal to zero. This means that we can find $i+1$ coordinate points spanning a linear space Σ such that $p \in \Sigma$; let Λ be the linear span of the remaining $n-i$ coordinate points. Now consider the projection map from Λ

$$\pi : \mathbb{P}^n \dashrightarrow \Sigma.$$

Choose $q \in \pi^{-1}(p) \setminus \Delta_{n-1}$. The conclusion follows since $p \star L = \pi(q \star L)$. \square

Lemma 3.2. *Let $L \subset \mathbb{P}^n$ be a linear space of dimension $m < n$ and consider points $p, q \in \mathbb{P}^n \setminus \Delta_{n-1}$. If $p \neq q$, $L \cap \Delta_{n-m-1} = \emptyset$, and $\langle p, q \rangle \cap \Delta_{n-m-2} = \emptyset$, then $p \star L \neq q \star L$.*

Proof. Let $p = [a_0 : \cdots : a_n]$ and $q = [b_0 : \cdots : b_n]$. Let $t = n - m$. We prove the result by contradiction dealing with the case $t = 1$ and then using a projection argument.

Case $t = 1$. In this situation, L is a hyperplane. We consider its equation

$$L : \sum_{i=0}^n \alpha_i x_i = 0.$$

Direct computations show that

$$p \star L : \sum_{i=0}^n \frac{\alpha_i}{a_i} x_i = 0 \quad \text{and} \quad q \star L : \sum_{i=0}^n \frac{\alpha_i}{b_i} x_i = 0.$$

Since $L \cap \Delta_0 = \emptyset$ all the coefficients α_i are non-zero. The assumption $p \star L = q \star L$ yield that $\frac{\alpha_i}{a_i} = \frac{\alpha_j}{b_j}$ for all i, j . This is a contradiction as $p \neq q$.

Case $t > 1$. Choose $t - 1$ coordinate points in Δ_0 and denote with Σ their linear span. Note that $\Sigma \subset \Delta_{t-2}$. Let π be the projection from Σ

$$\pi : \mathbb{P}^n \dashrightarrow \mathbb{P}^{n-t+1},$$

that is, π forgets $t - 1$ coordinates. We set

$$p' = \pi(p), q' = \pi(q) \text{ and } L' = \pi(L).$$

Since $L \cap \Sigma \subset L \cap \Delta_{t-1} \subset L \cap \Delta_{n-m-1} = \emptyset$, L' is a linear space of dimension m . Similarly, $p' \neq q'$ since $\langle p, q \rangle \cap \Delta_{t-2} = \emptyset$. Since π simply forgets $t - 1$ coordinates, we have

$$p' \star L' = \pi(p \star L) \text{ and } q' \star L' = \pi(q \star L).$$

Now, we show $p' \star L' \neq q' \star L'$, and hence $p \star L \neq q \star L$. To conclude using the case $t = 1$, we have to check the following in \mathbb{P}^{m+1} :

- (1) $L' \cap \Delta_0 = \emptyset$.
- (2) $p', q' \notin \Delta_m = \emptyset$.

Since $p \notin \Delta_{n-1}$, clearly $p' \notin \Delta_m$, and similarly $q' \notin \Delta_m$. Thus (2) is proved. If $z' \in L' \cap \Delta_0$, then we can choose $z \in L$ such that $\pi(z) = z'$. Thus, $z \in \Delta_{t-1} = \Delta_{n-m-1}$, a contradiction. Hence (1) is proved and the proof is now completed. \square

3.2. Powers of a line. The case of the powers of a line is special. The main reason is that two dimension counts agree in the proof of 3.4.

Lemma 3.3. *Let $n \geq 2$, $n \geq r$ and $L \subset \mathbb{P}^n$ be a line such that $L \cap \Delta_{n-2} = \emptyset$. There exist $p \neq q$ in $L \setminus \Delta_{n-1}$, and for any such: $z \in L$ implies $z \star L^{*r} \subset \langle p \star L^{*r}, q \star L^{*r} \rangle$.*

Proof. Choose a vector space basis of L made of points with no zero coordinates. One exists, since $L \cap \Delta_{n-2} = \emptyset$. Hence z is a linear combination of p and q and:

$$z \star \bigcup_{z_i \in L} z_1 \star \dots \star z_r \subset \langle p \star L^{*r}, q \star L^{*r} \rangle.$$

We can note the LHS equals L^{*r} by Lemma 2.10, or take closure of both sides. \square

Theorem 3.4. *Let $L \subset \mathbb{P}^n$, $n > 1$, be a line. If $L \cap \Delta_{n-2} = \emptyset$, then $L^{*r} \subset \mathbb{P}^n$ is a linear space of dimension $\min\{r, n\}$.*

Proof. It is enough to consider the case $r \leq n$. Indeed, if $r > n$, $\mathbb{P}^n \supseteq L^{*r} \supseteq p \star L^{*n}$ for some $p \in L^{*(r-n)}$ without zero coordinates. The result follows from Lemma 3.1.

We will prove the statement by proving that (i) $\dim(L^{*r}) = r$ and that (ii) $\dim \langle L^{*r} \rangle = r$. We proceed by induction on r . The base case is $r = 1$.

Now we assume the statement to hold for r and we prove it for $r + 1$. To prove (i) we use Lemma 2.12. Thus, we consider general points $p_1, \dots, p_{r+1} \in L \setminus \Delta_{n-1}$ and we have

$$\begin{aligned} T_{p_1 \star \dots \star p_{r+1}}(L^{*(r+1)}) &= \langle p_2 \star \dots \star p_{r+1} \star L, \dots, p_1 \star \dots \star p_r \star L \rangle \\ &= \langle p_{r+1} \star T_{p_1 \star \dots \star p_r}(L^{*r}), p_1 \star T_{p_2 \star \dots \star p_{r+1}}(L^{*r}) \rangle. \end{aligned}$$

Using the inductive hypothesis we conclude that

$$T_{p_1 \star \dots \star p_{r+1}}(L^{\star(r+1)}) = \langle p_{r+1} \star L^{\star r}, p_1 \star L^{\star r} \rangle.$$

Thus (i) follows, since $p_{r+1} \star L^{\star r}$ and $p_1 \star L^{\star r}$ are r dimensional, distinct (by Lemma 3.2 which we can apply because of Lemma 2.9) linear spaces intersecting in an $r-1$ dimensional linear space, namely $p_1 \star p_{r+1} \star L^{\star(r-1)}$, so we conclude

$$\dim(L^{\star(r+1)}) = r + 1.$$

To prove (ii) we use Lemma 3.3 which yields

$$\langle L^{\star(r+1)} \rangle = \left\langle \bigcup_{z \in L} z \star L^{\star r} \right\rangle \subset \langle p \star L^{\star r}, q \star L^{\star r} \rangle$$

for $p \neq q$ and $p, q \in L \setminus \Delta_{n-1}$. Again by Lemma 2.9, Lemma 3.1, Lemma 3.2, and the inductive hypothesis we get that $p \star L^{\star r}$, $q \star L^{\star r}$ are distinct linear spaces of dimension r containing a common linear space of dimension $r-1$, namely $p \star q \star L^{\star(r-1)}$. Hence, $\dim(\langle L^{\star(r+1)} \rangle) = r + 1$ and the proof is now completed. \square

Remark 3.5. If $L \cap \Delta_{n-2} = \emptyset$ fails in Theorem 3.4, $L^{\star r}$ is still linear, but possibly of lower dimension. For example, consider the line $L \subset \mathbb{P}^5$

$$L : \begin{cases} 2x_0 - x_1 = 0 \\ x_1 + 3x_2 - x_4 = 0 \\ 3x_2 - x_3 = 0 \\ 16x_3 - 12x_4 - 3x_5 = 0. \end{cases}$$

In Macaulay2 ([13]), we compute

$$L^{\star 2} : \begin{cases} 9x_2 - x_3 = 0 \\ 192x_1 + 64x_3 - 48x_4 - 9x_5 = 0 \\ 768x_0 + 64x_3 - 48x_4 - 9x_5 = 0. \end{cases}$$

$$L^{\star 3} : \begin{cases} 27x_2 - x_3 = 0 \\ 8x_0 - x_1 = 0 \end{cases}$$

$$L^{\star 4} : \begin{cases} 81x_2 - x_3 = 0 \\ 16x_0 - x_1 = 0 \end{cases}$$

and $\dim(L^{\star r}) = 3$ for all $r \geq 3$.

3.3. Equations. We now make Theorem 3.4 explicit, and express the Plücker coordinates ([28, §III.14]) of $L^{\star r}$ in terms of the Plücker coordinates of the given line L in \mathbb{P}^n . From this, we get canonical equations for $L^{\star r}$.

Proposition 3.6. *Let L in \mathbb{P}^n , $n > 1$, be a line with $L \cap \Delta_{n-2} = \emptyset$, and let $r < n$.*

(1) *If $L = \text{rowspan} \begin{pmatrix} a_{00} & a_{01} & \dots & a_{0n} \\ a_{10} & a_{11} & \dots & a_{1n} \end{pmatrix}$, then*

$$L^{\star r} = \text{rowspan} \begin{pmatrix} a_{00}^r & a_{01}^r & \dots & a_{0n}^r \\ a_{00}^{r-1}a_{10} & a_{01}^{r-1}a_{11} & \dots & a_{0n}^{r-1}a_{1n} \\ \dots & \dots & \dots & \dots \\ a_{10}^r & a_{11}^r & \dots & a_{1n}^r \end{pmatrix}.$$

(2) In terms of Plücker coordinates, if $0 \leq i_0 < i_1 < \dots < i_r \leq n$, then

$$[i_0, i_1, \dots, i_r]_{L^{\star r}} = \prod_{0 \leq j < k \leq r} [i_j, i_k]_L.$$

Proof. By Lemma 3.3, if L is spanned by points p and q , then $L^{\star r}$ is spanned by points $p^{\star r}$, $p^{\star(r-1)} \star q$, \dots , $q^{\star r}$. Part (1) follows. For (2):

$$[i_0, i_1, \dots, i_r]_{L^{\star r}} = \det \begin{pmatrix} a_{0i_0}^r & a_{0i_1}^r & \dots & a_{0i_r}^r \\ a_{0i_0}^{r-1} a_{1i_0} & a_{0i_1}^{r-1} a_{1i_1} & \dots & a_{0i_r}^{r-1} a_{1i_r} \\ \dots & \dots & \dots & \dots \\ a_{1i_0}^r & a_{1i_1}^r & \dots & a_{1i_r}^r \end{pmatrix}.$$

This is a homogenized Vandermonde determinant. It equals:

$$\prod_{0 \leq j < k \leq r} (a_{0i_j} a_{1i_k} - a_{0i_k} a_{1i_j}) = \prod_{0 \leq j < k \leq r} [i_j, i_k]_L. \quad \square$$

The virtue of expressing a formula in terms of Plücker coordinates as above, and also as later on in this paper, is that Plücker coordinates form the most canonical coordinate system on a Grassmannian of linear spaces.

Corollary 3.7. *In Proposition 3.6, if x_0, x_1, \dots, x_n are the homogeneous coordinates of \mathbb{P}^n , then $L^{\star r}$ is cut out by the maximal minors of*

$$\begin{pmatrix} a_{00}^r & a_{01}^r & \dots & a_{0n}^r \\ a_{00}^{r-1} a_{10} & a_{01}^{r-1} a_{11} & \dots & a_{0n}^{r-1} a_{1n} \\ \dots & \dots & \dots & \dots \\ a_{10}^r & a_{11}^r & \dots & a_{1n}^r \\ x_0 & x_1 & \dots & x_n \end{pmatrix}.$$

These are $\binom{n+1}{r+2}$ linear equations with coefficients the Plücker coordinates of $L^{\star r}$ (up to sign). For $r < n-1$, they are not a complete intersection. But for $r = n-1$, there is just one equation, for the hyperplane $L^{\star(n-1)}$:

$$L^{\star(n-1)} : \sum_{i=0}^n \left((-1)^{n+i} \prod_{\substack{0 \leq j < k \leq n \\ j, k \neq i}} [j, k]_L \right) x_i = 0.$$

4. STAR CONFIGURATIONS

In this section, we give an application of the Hadamard powers of a line: Theorem 4.7, which can be viewed as a simple construction of star configurations ([4, 5, 12]). Interest in star configurations has increased in recent years because of their extremal behaviour with respect to many open problems such as, for example, symbolic powers of ideals [3].

Given a finite set of points $Z \subset \mathbb{P}^n$, we can consider the Hadamard powers $Z^{\star r}$. However, it is more interesting to only consider Hadamard products of distinct points in Z .

Definition 4.1. Let $Z \subset \mathbb{P}^n$ be a finite set of points. The r -th *square-free* Hadamard power of Z is

$$Z^{\star r} = \{p_1 \star \dots \star p_r : p_i \in Z \text{ and } p_i \neq p_j \text{ for } i \neq j\}.$$

We want to describe the square-free Hadamard powers of a finite set of points. Thus, we begin with a useful technical result.

We will use the following notion.

Definition 4.2. Let $H_1, \dots, H_m \subset M \subset \mathbb{P}^n$ be linear spaces such that $r = \dim M = \dim H_i + 1$. The linear spaces H_i are said to be in *linear general position in M* if, whenever i_1, \dots, i_j are distinct:

- $\dim(H_{i_1} \cap \dots \cap H_{i_j}) = r - j$, for $j \leq r$.
- $H_{i_1} \cap \dots \cap H_{i_j} = \emptyset$, for $r < j$.

Lemma 4.3. Let $L \subset \mathbb{P}^n$ be a line, let $p_i \in L \setminus \Delta_{n-1}$, $1 \leq i \leq m$, be m distinct points, and $r \leq n$. Set $M = L^{\star r}$ and $H_i = p_i \star L^{\star(r-1)}$, $1 \leq i \leq m$. If $L \cap \Delta_{n-2} = \emptyset$, then whenever i_1, \dots, i_j are distinct:

- $H_{i_1} \cap \dots \cap H_{i_j} = p_{i_1} \star \dots \star p_{i_j} \star L^{\star(r-j)}$, for $j \leq r$.
- $H_{i_1} \cap \dots \cap H_{i_j} = \emptyset$, for $r < j$.

In particular, the linear spaces H_i are in linear general position in M .

Proof. We will prove this for $j \leq r + 1$ by induction, and that suffices.

The base case is $j = 1$. Now we assume the statement for $j \leq r$ and prove it for $j + 1$. Without loss of generality it is enough to show that:

$$H_1 \cap \dots \cap H_{j+1} = p_1 \star \dots \star p_{j+1} \star L^{\star(r-j-1)}.$$

By the inductive hypothesis:

- $H_1 \cap \dots \cap H_j = p_1 \star \dots \star p_j \star L^{\star(r-j)}$, and
- $H_2 \cap \dots \cap H_{j+1} = p_2 \star \dots \star p_{j+1} \star L^{\star(r-j)}$.

Intersecting these, we get:

$$H_1 \cap \dots \cap H_{j+1} = p_1 \star V \cap p_{j+1} \star V,$$

where $V = p_2 \star \dots \star p_j \star L^{\star(r-j)}$. By Theorem 3.4 and Lemma 3.1, V is a linear space of dimension $r - j$. By Lemma 2.9, $V \cap \Delta_{n-(r-j)-1} = \emptyset$, which means Lemma 3.2 applies to give $p_1 \star V \neq p_{j+1} \star V$. Combining with

$$p_1 \star V \cap p_{j+1} \star V \supset p_1 \star \dots \star p_{j+1} \star L^{\star(r-j-1)},$$

and Lemma 3.1 and Theorem 3.4, we complete the induction. \square

Corollary 4.4. Let $L \subset \mathbb{P}^n$ be a line, $r \leq n$, and consider points $p_i, q_j \in L \setminus \Delta_{n-1}$ for $1 \leq i, j \leq r$. If $L \cap \Delta_{n-2} = \emptyset$, and

$$\{p_1, \dots, p_r\} \neq \{q_1, \dots, q_r\},$$

then $p_1 \star \dots \star p_r \neq q_1 \star \dots \star q_r$.

Proof. Set $x = p_1 \star \dots \star p_r$ and $y = q_1 \star \dots \star q_r$. Without loss of generality we may assume $q_1 \neq p_i$ for $1 \leq i \leq r$. Consider the linear spaces

$$p_1 \star L^{\star(r-1)}, \dots, p_r \star L^{\star(r-1)}$$

and

$$q_1 \star L^{\star(r-1)}.$$

They are in linear general position in $L^{\star r}$ by Lemma 4.3. So:

- $x = p_1 \star L^{\star(r-1)} \cap \dots \cap p_r \star L^{\star(r-1)}$, and
- $\emptyset = q_1 \star L^{\star(r-1)} \cap p_1 \star L^{\star(r-1)} \cap \dots \cap p_r \star L^{\star(r-1)}$.

From $y \in q_1 \star L^{\star(r-1)}$, the result follows. \square

Corollary 4.5. *Let $L \subset \mathbb{P}^n$ be a line, $Z \subset L$ be a set of m points and $r \leq n$. If $L \cap \Delta_{n-2} = \emptyset$ and $Z \cap \Delta_{n-1} = \emptyset$, then $Z^{\star r}$ is a set of $\binom{m}{r}$ points.*

We now see that $Z^{\star r}$ for collinear points is a star configuration.

Definition 4.6. A set of $\binom{m}{r}$ points $\mathbb{X} \subset \mathbb{P}^n$ is a *star configuration* if there exist linear spaces $H_1, \dots, H_m \subset M \subset \mathbb{P}^n$ such that:

- $r = \dim M = \dim H_i + 1$.
- H_i are in linear general position in M .
- $\mathbb{X} = \bigcup_{1 \leq i_1 < \dots < i_r \leq m} H_{i_1} \cap \dots \cap H_{i_r}$.

Theorem 4.7. *Let $L \subset \mathbb{P}^n$ be a line, $Z \subset L$ be a set of m points and $r \leq \min\{m, n\}$. If $L \cap \Delta_{n-2} = \emptyset$ and $Z \cap \Delta_{n-1} = \emptyset$, then $Z^{\star r}$ is a star configuration in $M = L^{\star r}$.*

Proof. Let $Z = \{p_1, \dots, p_m\}$ and notice that $Z^{\star r}$ is a set of $\binom{m}{r}$ points by Corollary 4.5. Using Lemma 4.3 we see that the linear spaces

$$H_1 = p_1 \star L^{\star(r-1)}, \dots, H_m = p_m \star L^{\star(r-1)}$$

are in linear general position in $M = L^{\star r}$, and

$$H_{i_1} \cap \dots \cap H_{i_r} = p_{i_1} \star \dots \star p_{i_r}$$

for all $1 \leq i_1 < \dots < i_r \leq m$. The conclusion follows from the definitions of star configuration and square-free Hadamard product. \square

5. TROPICAL CONNECTION

To study Hadamard products of other linear spaces in Section 6, we will need some machinery from tropical geometry. We now review the basics.

Given an irreducible variety $X \subset \mathbb{P}^n$ not contained in Δ_{n-1} , let $I \subset \mathbb{C}[x_0^\pm, \dots, x_n^\pm]$ be the defining ideal of $X - \Delta_{n-1} \subset (\mathbb{C}^*)^{n+1}/\mathbb{C}^*$. The set

$$\text{trop}(X) = \overline{\{\mathbf{w} \in \mathbb{R}^{n+1}/\mathbb{R}\mathbf{1} : \text{in}_{\mathbf{w}}(I) \neq \langle 1 \rangle\}}$$

is the *tropicalization* of X ([26, Theorem 3.2.3]). It is the support of a pure rational polyhedral subfan of the Gröbner fan ([7, §8.4]) of I , inside $\mathbb{R}^{n+1}/\mathbb{R}\mathbf{1}$ ([26, Theorem 3.3.5]). That subfan has positive integer multiplicities attached to its facets:

$$\text{mult}(\sigma) = \sum_{P \in \text{Ass}^{\text{min}}(\text{in}_{\mathbf{w}}(I))} \text{mult}(P, \text{in}_{\mathbf{w}}(I)).$$

Here \mathbf{w} is any point in the relative interior of σ and the sum is over minimal associated primes P of $\text{in}_{\mathbf{w}}(I)$. These multiplicities balance along ridges ([26, Theorem 3.4.14]). A lot of the geometry of X can be recovered from $\text{trop}(X)$.

Tropical geometry provides powerful tools to study Hadamard products, because of the following connection.

Proposition 5.1. *Let $X, Y \subset \mathbb{P}^n$ be irreducible varieties. The tropicalization of the Hadamard product is the Minkowski sum of the tropicalizations:*

$$\text{trop}(X \star Y) = \text{trop}(X) + \text{trop}(Y)$$

as sets. If also $X \times Y \dashrightarrow X \star Y$ is generically δ to 1, then:

$$\text{trop}(X \star Y) = \frac{1}{\delta} (\text{trop}(X) + \text{trop}(Y))$$

as weighted balanced fans.

Proof. The set theoretic statement is [26, Proposition 5.5.11]. The multiplicities statement follows from [26, Remark 5.5.2]. \square

Remark 5.2. We explain the RHS of Proposition 5.1. As a set, it is

$$\text{trop}(X) + \text{trop}(Y) = \{ \mathbf{w}_1 + \mathbf{w}_2 : \mathbf{w}_1 \in \text{trop}(X), \mathbf{w}_2 \in \text{trop}(Y) \} \subset \mathbb{R}^{n+1}/\mathbb{R}\mathbf{1}.$$

Now choose refinements of $\text{trop}(X)$ and $\text{trop}(Y)$ so

$$\{ \sigma_1 + \sigma_2 : \sigma_1 \text{ is a facet of } \text{trop}(X), \sigma_2 \text{ is a facet of } \text{trop}(Y) \}$$

is a polyhedral fan. Put weights on facets σ by

$$\text{mult}(\sigma) = \sum_{\sigma_1, \sigma_2} \text{mult}(\sigma_1) \text{mult}(\sigma_2) [N_\sigma : N_{\sigma_1} + N_{\sigma_2}].$$

Here the sum is over all pairs of facets σ_1, σ_2 in $\text{trop}(X), \text{trop}(Y)$ respectively such that $\sigma = \sigma_1 + \sigma_2$. Also N_σ is the maximal sublattice of $\mathbb{Z}^{n+1}/\mathbb{Z}\mathbf{1}$ parallel to the affine span of σ , similarly for N_{σ_1} and N_{σ_2} , and $[N_\sigma : N_{\sigma_1} + N_{\sigma_2}]$ is the lattice index. Now multiply these weights by $\frac{1}{\delta}$. Finally note the equality of weighted balanced fans is up to common refinement.

Remark 5.3. Proposition 5.1 and Remark 5.2 generalize to several varieties. If $X_1, \dots, X_r \subset \mathbb{P}^n$ are irreducible and $X_1 \times \dots \times X_r \dashrightarrow X_1 \star \dots \star X_r$ is generically δ to 1, then:

$$\text{trop}(X_1 \star \dots \star X_r) = \frac{1}{\delta} (\text{trop}(X_1) + \dots + \text{trop}(X_r))$$

as weighted balanced fans. Multiplicities in $\text{trop}(X_1) + \dots + \text{trop}(X_r)$ are:

$$\text{mult}(\sigma) = \sum_{\sigma_1, \dots, \sigma_r} \text{mult}(\sigma_1) \dots \text{mult}(\sigma_r) [N_\sigma : N_{\sigma_1} + \dots + N_{\sigma_r}].$$

As a first application of tropical geometry, we can say what dimension we expect Hadamard products to have in general.

Proposition 5.4. *Let $X, Y \subset \mathbb{P}^n$ be irreducible varieties. Denote by $H \subset (\mathbb{C}^*)^{n+1}/\mathbb{C}^*$ the largest subtorus acting on both X and Y , and $G \subset (\mathbb{C}^*)^{n+1}/\mathbb{C}^*$ the smallest subtorus having a coset containing X and a coset containing Y . Then*

$$\min\{\dim(X) + \dim(Y) - \dim(H), \dim(G)\}$$

upper bounds $\dim(X \star Y)$. We call it the expected dimension of $X \star Y$.

Proof. Let $V \subset \mathbb{R}^{n+1}/\mathbb{R}\mathbf{1}$ be the common lineality space of $\text{trop}(X)$ and $\text{trop}(Y)$, i.e., the largest linear subspace contained in all cones of the two fans. Let $W \subset \mathbb{R}^{n+1}/\mathbb{R}\mathbf{1}$ be the linear span $\langle \text{trop}(X), \text{trop}(Y) \rangle$. We need three facts:

1. Tropicalization applied to irreducible varieties preserves dimension.
2. Proposition 5.1
3. Here $\text{trop}(H) = V$ and $\text{trop}(G) = W$ (see [9, §3]).

Now we have:

$$\begin{aligned}
\dim(X \star Y) &\stackrel{1.}{=} \dim(\text{trop}(X \star Y)) \\
&\stackrel{2.}{=} \dim(\text{trop}(X) + \text{trop}(Y)) \\
&= \max\{\dim(C + D) : C \text{ and } D \text{ are facets in } \text{trop}(X) \text{ and } \text{trop}(Y)\} \\
&\leq \min\{\dim(\text{trop}(X)) + \dim(\text{trop}(Y)) - \dim(V), \dim(W)\} \\
&\stackrel{3.}{=} \min\{\dim(X) + \dim(Y) - \dim(H), \dim(G)\}. \quad \square
\end{aligned}$$

It is a tricky matter as to when Hadamard products have deficient dimension. Here is an example related to *tropical A-discriminants* ([10]). We verified it in `Macaulay2` and Felipe Rincón's `TropLi` ([31]).

Example 5.5. Let $X \subset \mathbb{P}^{11}$ be the Segre product of \mathbb{P}^2 and \mathbb{P}^3 , i.e., the variety of 3×4 matrices with rank 1. Let $Y \subset \mathbb{P}^{11}$ be the linear subspace of 3×4 matrices with all row and column sums zero. Then $H = 1$, $G = (\mathbb{C}^*)^{12}/\mathbb{C}^*$, so the expected dimension is $\min\{5 + 5 - 0, 11\} = 10$. But $X \star Y = \{3 \times 4 \text{ matrices with rank } \leq 2\}$ has dimension 9. Here $\text{trop}(X)$ is a 5-dimensional classical linear space and $\text{trop}(Y)$ is a 5-dimensional tropicalized linear space with 55 rays and 1656 maximal cones in its *cyclic Bergman fan* (a refinement of the Gröbner subfan). There is no common lineality space, the linear span is $\mathbb{R}^{12}/\mathbb{R}\mathbf{1}$, but the Minkowski sum is 9-dimensional.

6. PRODUCTS OF LINEAR SPACES

Besides powers of a line, the Hadamard product of linear spaces is not linear. We start with two numerical examples to illustrate this.

6.1. Numerical examples.

Example 6.1. Let L be the line in \mathbb{P}^3 through points $[2 : 3 : 5 : 7]$ and $[11 : 13 : 17 : 19]$, and M the line through $[23 : 29 : 31 : 37]$ and $[41 : 43 : 47 : 53]$. Let x_0, x_1, x_2, x_3 be the homogenous coordinates of \mathbb{P}^3 . Using ideal elimination in `Macaulay2`, we compute $L \star M$ to be the quadric surface defined by:

$$\begin{aligned}
&88128x_0^2 - 89280x_0x_1 - 5299632x_1^2 - 817938x_0x_2 + 8896641x_1x_2 - 1481805x_2^2 \\
&\quad - 321510x_0x_3 - 1777545x_1x_3 - 54250x_2x_3 + 116375x_3^2.
\end{aligned}$$

Note the ruling on $L \star M$ is $\{p \star M : p \in L\} \cup \{q \star L : q \in M\}$.

Remark 6.2. In general, for linear spaces $L, M \subset \mathbb{P}^n$, $L \star M$ has big Fano schemes, because $L \star M$ contains the linear spaces $\{p \star M : p \in L\} \cup \{q \star L : q \in M\}$.

Example 6.3. Let P be the 2-plane in \mathbb{P}^5 spanned by $[3 : 1 : 4 : 1 : 5 : 9]$, $[2 : 6 : 5 : 3 : 5 : 8]$ and $[9 : 7 : 9 : 3 : 2 : 3]$. In `Macaulay2`, we compute the Hadamard square $P^{\star 2}$ to be a cubic hypersurface in \mathbb{P}^5 , whose defining polynomial is too big to exhibit here. We also discover that $P^{\star 2}$ is singular in codimension 2, with singular locus $\{p \star p : p \in P\}$. Future work could study singular loci of Hadamard products. A precedent for secant varieties is [27, §5].

6.2. Linear span. One way to quantify nonlinearity of an embedded variety is by the dimension of its linear span.

Proposition 6.4. *Let $L \subset \mathbb{P}^n$ be a generic linear space of dimension m . Then the linear span $\langle L^{\star r} \rangle$ has dimension $\min\{\binom{m+r}{r} - 1, n\}$.*

Proof. Let $L = \langle p_0, p_1, \dots, p_m \rangle$ and correspondingly write:

$$L = \text{rowspan} \begin{pmatrix} a_{00} & a_{01} & \dots & a_{0n} \\ a_{10} & a_{11} & \dots & a_{1n} \\ \dots & \dots & \dots & \dots \\ a_{m0} & a_{m1} & \dots & a_{mn} \end{pmatrix}.$$

By the analogue of Lemma 3.3 for higher dimensional L , we have:

$$\langle L^{\star r} \rangle = \langle p_0^{\star r_0} \star p_1^{\star r_1} \star \dots \star p_m^{\star r_m} : r_i \in \mathbb{Z}_{\geq 0} \text{ and } r_0 + r_1 + \dots + r_m = r \rangle,$$

and correspondingly:

$$L^{\star r} = \text{rowspan} \begin{pmatrix} a_{00}^r & a_{01}^r & \dots & a_{0n}^r \\ \dots & \dots & \dots & \dots \\ \prod a_{i0}^{r_i} & \prod a_{i1}^{r_i} & \dots & \prod a_{in}^{r_i} \\ \dots & \dots & \dots & \dots \\ a_{m0}^r & a_{m1}^r & \dots & a_{mn}^r \end{pmatrix}.$$

The maximal minors of this $\binom{m+r}{r} \times (n+1)$ matrix are the *generalized Vandermonde determinants* from [1]. To finish the proof, we note they are non-zero polynomials in $\mathbb{C}[a_{ij} : 0 \leq i \leq m, 0 \leq j \leq n]$, because each of their coefficients is ± 1 . Indeed, they have no like terms as sums over the symmetric group. \square

Corollary 6.5. *Let $L \subset \mathbb{P}^n$ be a generic linear space of dimension m . Assume $n \geq \binom{m+r}{r} - 1$. Consider points $p_i, q_j \in L$ for $1 \leq i, j \leq r$ such that $p_1 \star \dots \star p_r = q_1 \star \dots \star q_r$. Then $\{p_1, \dots, p_r\} = \{q_1, \dots, q_r\}$.*

Proof. Choose a basis $L = \langle t_0, \dots, t_m \rangle$. Choose lifts $\tilde{p}_i, \tilde{q}_i, \tilde{t}_i$ of p_i, q_i, t_i to the affine cone $\tilde{L} \subset \mathbb{A}^{n+1}$ over L so that $\tilde{p}_1 \star \dots \star \tilde{p}_r = \tilde{q}_1 \star \dots \star \tilde{q}_r$. Expand:

- $\tilde{p}_i = \sum_{j=0}^m \alpha_{ij} \tilde{t}_j$
- $\tilde{q}_i = \sum_{j=0}^m \beta_{ij} \tilde{t}_j$.

Define a ring homomorphism

$$\begin{aligned} \Phi : \mathbb{C}[T_0, \dots, T_m] &\longrightarrow \mathbb{C}^{\times(n+1)} \\ T_i &\longmapsto t_i. \end{aligned}$$

Here the domain is the polynomial ring, and the codomain is the direct product of $n+1$ copies of \mathbb{C} . Then:

$$\Phi \left(\left(\sum_{j=0}^m \alpha_{1j} T_j \right) \dots \left(\sum_{j=0}^m \alpha_{rj} T_j \right) \right) = \Phi \left(\left(\sum_{j=0}^m \beta_{1j} T_j \right) \dots \left(\sum_{j=0}^m \beta_{rj} T_j \right) \right)$$

By Proposition 6.4, $\ker(\Phi)$ contains no homogenous polynomials of degree r . Hence:

$$\left(\sum_{j=0}^m \alpha_{1j} T_j \right) \dots \left(\sum_{j=0}^m \alpha_{rj} T_j \right) = \left(\sum_{j=0}^m \beta_{1j} T_j \right) \dots \left(\sum_{j=0}^m \beta_{rj} T_j \right).$$

Since $\mathbb{C}[T_0, \dots, T_m]$ is a unique factorization domain, the corollary follows. \square

Remark 6.6. Corollaries 4.7 and 6.5 correspond to *identifiability* ([2, 6, 32]) in the theory of secant varieties. A direction for future research is to look for $(\dim(L), r, n)$ such that $L^{*r} \subset \mathbb{P}^n$ is generically *not* identifiable. For secant varieties of symmetric tensors, a classification is [20, Theorem 3.3.3.1].

Remark 6.7. Proposition 6.4 and Corollary 6.5 generalize to several linear spaces. Let $L_1, \dots, L_k \subset \mathbb{P}^n$ be generic linear spaces of dimensions m_1, \dots, m_k . The linear span $\langle L_1^{r_1} \star \dots \star L_k^{r_k} \rangle$ has dimension $\min\left\{\binom{m_1+r_1}{r_1} \dots \binom{m_k+r_k}{r_k} - 1, n\right\}$. If $n \geq \binom{m_1+r_1}{r_1} \dots \binom{m_k+r_k}{r_k} - 1$, then $L_1^{r_1} \star \dots \star L_k^{r_k}$ is identifiable. The proofs are similar.

6.3. Degree formula. As a second application of tropical geometry, we now prove a general degree formula for Hadamard products of linear spaces, when the ambient projective space is of sufficiently high dimension. This is one of our main results.

Theorem 6.8. *Let $L_1, L_2, \dots, L_r \subset \mathbb{P}^n$ be generic linear spaces of dimensions m_1, m_2, \dots, m_r , respectively. Set $m = m_1 + m_2 + \dots + m_r$ and $d = \binom{m_1+m_2+\dots+m_r}{m_1, m_2, \dots, m_r}$. Assume $n \gg 0$. Then $L_1 \star L_2 \star \dots \star L_r$ has dimension m and degree*

$$\begin{cases} d & \text{if } L_i \text{ are pairwise distinct} \\ \frac{d}{r!} & \text{if } L_i \text{ are the same.} \end{cases}$$

In full generality, when L_i form a multiset with multiplicities r_1, \dots, r_k (where $r_1 + \dots + r_k = r$), the dimension is m and the degree is $\frac{d}{(r_1!) \dots (r_k!)}$.

Proof (in full generality). Generically L_i has Plücker coordinates all nonzero, so $\text{trop}(L_i)$ equals the *standard tropical linear space* Λ_{m_i} of dimension m_i ([26, Example 4.2.13]):

$$\text{trop}(L_i) = \Lambda_{m_i} = \bigcup_{0 \leq j_1 < \dots < j_{m_i} \leq n} \text{pos}(\mathbf{e}_{j_1}, \dots, \mathbf{e}_{j_{m_i}}).$$

Here $\mathbf{e}_0, \dots, \mathbf{e}_n$ are the images in $\mathbb{R}^{n+1}/\mathbb{R}\mathbf{1}$ of the standard basis vectors, pos denotes positive span, and all multiplicities are 1.

By Remark 5.3, $\text{trop}(L_1) + \dots + \text{trop}(L_r)$ has support the same as Λ_m , and for each facet:

$$\text{mult}(\sigma) = \sum_{\sigma_1, \dots, \sigma_r} \text{mult}(\sigma_1) \dots \text{mult}(\sigma_r) [N_\sigma : N_{\sigma_1} + \dots + N_{\sigma_r}] = \sum_{\sigma_1, \dots, \sigma_r} 1 = d.$$

By Remark 6.7, $L_1 \times \dots \times L_r \dashrightarrow L_1 \star \dots \star L_r$ is generically $(r_1!) \dots (r_k!)$ to 1. So by Remark 5.3:

$$\text{trop}(L_1 \star \dots \star L_r) = \frac{d}{(r_1!) \dots (r_k!)} \Lambda_m.$$

It follows that $L_1 \star \dots \star L_r$ has dimension m and degree ([26, Corollary 3.6.16]):

$$\begin{aligned} \deg(L_1 \star \dots \star L_r) &= \text{mult}_{\mathbf{0}} \left(\text{trop}(L_1 \star \dots \star L_r) \cap_{\text{st}} \Lambda_{n-m} \right) \\ &= \text{mult}_{\mathbf{0}} \left(\frac{d}{(r_1!) \dots (r_k!)} \Lambda_m \cap_{\text{st}} \Lambda_{n-m} \right) \\ &= \frac{d}{(r_1!) \dots (r_k!)}. \end{aligned}$$

Here we take *stable intersection* ([19]) with the standard tropical linear space of complementary dimension, and then measure the multiplicity of the origin. \square

Remark 6.9. We can be explicit about $n \gg 0$ in Theorem 6.8. From the application of Remark 6.7, the hypothesis is $n \geq \binom{m_1+r_1}{r_1} \dots \binom{m_k+r_k}{r_k} - 1$.

We can extend Theorem 6.8 to include the case of negative exponents. Given a linear space $L \subset \mathbb{P}^n$, its *reciprocal linear space* is:

$$L^{-1} = \overline{\{[a_0^{-1} : \dots : a_n^{-1}] \in \mathbb{P}^n : [a_0 : \dots : a_n] \in L \setminus \Delta_{n-1}\}}.$$

These varieties have good combinatorial properties. In [17], they were used to prove log-concavity of the coefficients of the characteristic polynomial of every realizable matroid. From [30], the defining ideal of L^{-1} has all initial ideals squarefree, and a universal Gröbner basis in bijection with the set of circuits in the matroid realized by L . For familiar examples, note rational normal curves are reciprocal lines.

Corollary 6.10. *Use the notation of Theorem 6.8. Additionally, let $\tilde{L}_1, \dots, \tilde{L}_s$ be generic linear spaces of dimensions $\tilde{m}_1, \dots, \tilde{m}_s$, and set $\tilde{m} = \tilde{m}_1 + \dots + \tilde{m}_s$ and $\tilde{d} = \binom{\tilde{m}_1 + \dots + \tilde{m}_s}{\tilde{m}_1, \dots, \tilde{m}_s}$. Assume $n \gg 0$ and \tilde{L}_i form a multiset with multiplicities s_1, \dots, s_ℓ . Then $L_1 \star \dots \star L_r \star \tilde{L}_1^{-1} \star \dots \star \tilde{L}_s^{-1}$ has dimension $m + \tilde{m}$ and degree:*

$$\binom{n-m}{\tilde{m}} \cdot \frac{d}{(r_1!) \dots (r_k!)} \cdot \frac{\tilde{d}}{(s_1!) \dots (s_\ell!)}.$$

Proof. The tropicalization of the reciprocal linear space \tilde{L}_i^{-1} equals:

$$\text{trop}(\tilde{L}_i^{-1}) = -\Lambda_{\tilde{m}_i} = \bigcup_{0 \leq j_1 < \dots < j_{\tilde{m}_i} \leq n} \text{neg}(\mathbf{e}_{j_1}, \dots, \mathbf{e}_{j_{\tilde{m}_i}}).$$

Here neg denotes negative span, and all multiplicities are 1. It follows that in

$$\text{trop}(L_1) + \dots + \text{trop}(L_r) + \text{trop}(\tilde{L}_1^{-1}) + \dots + \text{trop}(\tilde{L}_s^{-1}),$$

the cones are:

$$(*) \quad \text{pos}(\mathbf{e}_i : i \in I) + \text{neg}(\mathbf{e}_j : j \in J)$$

for $I, J \subset \{0, \dots, n\}$ with $|I| = m$, $|J| = \tilde{m}$ and $I \cap J = \emptyset$. Similar to in the proof of Theorem 6.8, the multiplicities of these cones are all $d\tilde{d}$. Using Remark 6.7, we can clear denominators to deduce that:

$$L_1 \times \dots \times L_r \times \tilde{L}_1^{-1} \times \dots \times \tilde{L}_s^{-1} \dashrightarrow L_1 \star \dots \star L_r \star \tilde{L}_1^{-1} \star \dots \star \tilde{L}_s^{-1}$$

is generically $(r_1!) \dots (r_k!)(s_1!) \dots (s_\ell!)$ to 1. So by Remark 5.3:

$$\text{trop}(L_1 \star \dots \star L_r \star \tilde{L}_1^{-1} \star \dots \star \tilde{L}_s^{-1})$$

consists of the cones (*), each with multiplicity $\frac{d}{(r_1!) \dots (r_k!)} \cdot \frac{\tilde{d}}{(s_1!) \dots (s_\ell!)}$. It follows the dimension of the Hadamard product and its tropicalization are $m + \tilde{m}$.

For the degree, take stable intersection with $\Lambda_{n-m-\tilde{m}}$. To compute the multiplicity of $\mathbf{0}$ using formula ([2]) of [19], pick $v \in \mathbb{R}^{n+1}/\mathbb{R}\mathbf{1}$ to be generic inside the octant $\text{pos}(\mathbf{e}_1, \dots, \mathbf{e}_m) + \text{neg}(\mathbf{e}_{m+1}, \dots, \mathbf{e}_n)$. Then there are $\binom{n-m}{\tilde{m}}$ summands in that formula: indeed for each $I \subset \{m+1, \dots, n\}$, the pair $\sigma_1 = \text{pos}(\mathbf{e}_1, \dots, \mathbf{e}_m) + \text{neg}(\mathbf{e}_i : i \in I)$ and $\sigma_2 = \text{neg}(\mathbf{e}_i : i \in \{m+1, \dots, n\} \setminus I)$ contributes a term, namely:

$$\frac{d}{(r_1!) \dots (r_k!)} \cdot \frac{\tilde{d}}{(s_1!) \dots (s_\ell!)} \cdot 1 \cdot 1.$$

This gives a total of $\binom{n-m}{\tilde{m}} \cdot \frac{d}{(r_1!) \dots (r_k!)} \cdot \frac{\tilde{d}}{(s_1!) \dots (s_\ell!)}$, and the corollary follows. \square

We conclude with two Plücker formulas for products in the regime of Theorem 6.8. They consist of bracket polynomials, like in Proposition 3.6, and correspond to numerical Examples 6.1 and 6.3, respectively.

Example 6.11. Let L and M be generic distinct lines in \mathbb{P}^3 , with Plücker coordinates $[ij] := [i, j]_L$ and $\{ij\} := [i, j]_M$ respectively. Then $L \star M$ is the quadric surface in \mathbb{P}^3 cut out by:

$$\begin{aligned} & [12][13][23]\{12\}\{13\}\{23\} x_0^2 + [02][03][23]\{02\}\{03\}\{23\} x_1^2 \\ & + [01][03][13]\{01\}\{03\}\{13\} x_2^2 + [01][02][12]\{01\}\{02\}\{12\} x_3^2 \\ & - [23]\{23\}([02][13]\{03\}\{12\} + [03][12]\{02\}\{13\}) x_0 x_1 \\ & + [13]\{13\}([01][23]\{03\}\{12\} + [03][12]\{01\}\{23\}) x_0 x_2 \\ & - [12]\{12\}([01][23]\{02\}\{13\} + [02][13]\{01\}\{23\}) x_0 x_3 \\ & - [03]\{03\}([01][23]\{02\}\{13\} + [02][13]\{01\}\{23\}) x_1 x_2 \\ & + [02]\{02\}([01][23]\{03\}\{12\} + [03][12]\{01\}\{23\}) x_1 x_3 \\ & - [01]\{01\}([02][13]\{03\}\{12\} + [03][12]\{02\}\{13\}) x_2 x_3. \end{aligned}$$

To prove this formula, substitute $[ij] = a_{0i}a_{1j} - a_{0j}a_{1i}$, $\{ij\} = b_{0i}b_{1j} - b_{0j}b_{1i}$, and $x_i = (\lambda_0 a_{0i} + \lambda_1 a_{1i})(\mu_0 b_{0i} + \mu_1 b_{1i})$ in `Macaulay2`, and get identically zero.

To find this formula, note the incidence variety

$$\mathbb{X} = \overline{\{(p, L, M) \in \mathbb{P}^3 \times \text{Gr}(2, 4) \times \text{Gr}(2, 4) : p \in L \star M\}}$$

has three group actions:

- \mathfrak{S}_2 acts by switching L and M
- \mathfrak{S}_4 acts by permuting the homogeneous coordinates of \mathbb{P}^3
- $(\mathbb{C}^*)^4/\mathbb{C}^*$ acts by scaling the homogeneous coordinates of \mathbb{P}^3 .

So, the defining equation of \mathbb{X} is $(\mathfrak{S}_2 \times \mathfrak{S}_4)$ -symmetric and \mathbb{Z}^3 -multihomogeneous. Also, specializing $L = M$ should give the square of the linear equation in Corollary 3.7. Lastly, guess the formula has integer coefficients, on the basis of Example 6.1. Putting these clues together, it is straightforward to find the formula.

Example 6.12. Let P be a generic 2-plane in \mathbb{P}^5 , with Plücker coordinates $[ijk] := [i, j, k]_P$. Then $P^{\star 2}$ is the cubic hypersurface in \mathbb{P}^5 . In a defining equation, the coefficient of x_0^3 is $-(-1)^{0+0+0}$ times:

$$[123][124][125][134][135][145][234][235][245][345].$$

The coefficient of $x_0^2 x_1$ is $-(-1)^{0+0+1}$ times:

$$([023][045][124][125][134][135] + [024][035][123][125][134][145] + [025][034][123][124][135][145])[234][235][245][345].$$

The coefficient of $x_0 x_1 x_2$ is $-(-1)^{0+1+2}$ times:

$$\begin{aligned} & [013][024][134][234][125][035][235][045][145][345] + [013][124][034][234][025][135][235][045][145][345] + \\ & [023][014][134][234][125][035][135][045][245][345] + [023][124][034][134][015][135][235][045][245][345] + \\ & [123][014][034][234][025][035][135][145][245][345]. \end{aligned}$$

To get the other coefficients, act on the indices by \mathfrak{S}_6 . We find and prove this as in Example 6.11, here specializing $P = L^{\star 2}$ where L is a line.

Acknowledgements. The authors are grateful to Bernd Sturmfels for suggesting that we investigate multiplicative geometry and reading an earlier draft. The third author also thanks Justin Chen and Steven Sam for helpful conversations.

REFERENCES

- [1] L.F.T. Alonso and C. D’Andrea, *Tropicalization and irreducibility of generalized Vandermonde determinants*, Proceedings of the American Mathematical Society **137** (2009), 3647–3656.
- [2] C. Bocci and L. Chiantini, *On the identifiability of binary Segre products*, J. Alg. Geom. **22** (2013) 1–11.
- [3] C. Bocci and B. Harbourne, *Comparing powers and symbolic powers of ideals*, J. Alg. Geom. **19** (2010) 399–417.
- [4] E. Carlini, E. Guardo and A. Van Tuyl, *Star configurations on generic hypersurfaces*, J. Algebra **407** (2014), 1–20.
- [5] E. Carlini and A. Van Tuyl, *Star configuration points and generic plane curves*, Proc. Amer. Math. Soc. **139** (2011), no. 12, 4181–4192.
- [6] L. Chiantini and C. Ciliberto, *On the concept of k -secant order of a variety*, J. London Math. Soc. **73** (2006) 436–454.
- [7] D.A. Cox, J. Little and D. O’Shea, *Using Algebraic Geometry*, second ed., Springer-Verlag (2005).
- [8] M.A. Cueto, J. Morton and B. Sturmfels, *Geometry of the restricted Boltzmann machine*, Algebraic Methods in Statistics and Probability (eds. M. Viana and H. Wynn), American Mathematical Society, Contemporary Mathematics **516** (2010) 135–153.
- [9] M.A. Cueto, E.A. Tobis and J. Yu, *An implicitization challenge for binary factor analysis*, J. Symbolic Comput. **45** (2010), no. 12, 1296–1315.
- [10] A. Dickenstein, E.M. Feichtner and B. Sturmfels, *Tropical discriminants*, J. Amer. Math. Soc. **20** (2007) 1111–1133.
- [11] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Springer (1999).
- [12] A.V. Geramita, B. Harbourne and J. Migliore, *Star configurations in \mathbb{P}^n* , J. Algebra **376** (2013), 279–299.
- [13] D.R. Grayson and M.E. Stillman: *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [14] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics vol. 52, Springer-Verlag (1977).
- [15] R.A. Horn, *Topics in Matrix Analysis*, Cambridge University Press, 1994.
- [16] R.A. Horn and R. Mathias, *Block-matrix generalizations of Schur’s basic theorems on Hadamard products*, Linear Algebra Appl. **172** (1992) 337–346.
- [17] J. Huh and E. Katz, *Log-concavity of characteristic polynomials and the Bergman fan of matroids*, Mathematische Annalen **354** (2012), 1103–1116.
- [18] N.O. Ilten and H. Süß, *Polarized complexity-1 T -varieties*, Michigan Math. J. **60** (2011), no. 3, 561–578.
- [19] A. Jensen and J. Yu, *Stable intersections of tropical varieties*, arXiv:1309.7064v3.
- [20] J.M. Landsberg, *Tensors: geometry and applications*, Graduate Studies in Mathematics, vol. 128, AMS, Providence, RI (2012).
- [21] S. Liu, *Inequalities involving Hadamard products of positive semidefinite matrices*, J. Math. Ana. Appl. **243** (2000b) 458–463.
- [22] S. Liu, *On the Hadamard Product of Square Roots of Correlation Matrices*, Econometric Theory **18/19** (2002b/2003) 1007/703–704.
- [23] S. Liu and H. Neudecker, W. Polasek, *The Hadamard product and some of its applications in statistics*, Statistics **26** (1995a) no. 4, 365–373.
- [24] S. Liu and H. Neudecker, W. Polasek, *The heteroskedastic linear regression model and the Hadamard product – a note*, J. Econometrics **68** (1995b) 361–366.
- [25] S. Liu and G. Trenkler, *Hadamard, Khatri-Rao, Kronecker and other matrix products*, Int. Jour. of Information and Systems Sciences **4** (1), (2008), 160–177.
- [26] D. Maclagan and B. Sturmfels, *Introduction to tropical geometry*, Graduate Studies in Mathematics vol. 161, AMS, Providence, RI (2015).

- [27] M. Michalek, L. Oeding and P. Zwiernik, *Secant cumulants and toric geometry*, International Mathematics Research Notices, rnu056 (2014).
- [28] E. Miller and B. Sturmfels, *Combinatorial Commutative Algebra*, Graduate Texts in Mathematics vol. 227, Springer (2005).
- [29] D. Mumford, *The Red Book of Varieties and Schemes*, Springer-Verlag (1999).
- [30] N. Proudfoot and D. Speyer, *A broken circuit ring*, Beiträge Algebra Geom. **47** (2006), no. 1, 161–166.
- [31] F. Rincón, *Computing Tropical Linear Spaces*, J. Symbolic Comput. **51** (2013), 86–98.
- [32] V. Strassen, *Rank and optimal computation of generic tensors*, Linear Algebra Appl. **52** (1983) 645–685.
- [33] B. Sturmfels, *Equations defining toric varieties*, Algebraic Geometry - Santa Cruz 1995, Proc. Sympos. Pure Math. **62**, Part 2, AMS, Providence, RI, (1997), 437–449.

CRISTIANO BOCCI, DEPARTMENT OF INFORMATION ENGINEERING AND MATHEMATICS, UNIVERSITY OF SIENA, VIA ROMA, 56 SIENA, ITALY
E-mail address: `cristiano.bocci@unisi.it`

ENRICO CARLINI, SCHOOL OF MATHEMATICAL SCIENCES, MONASH UNIVERSITY, WELLINGTON ROAD, CLAYTON, AUSTRALIA

DEPARTMENT OF MATHEMATICAL SCIENCES, POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI 24, TURIN, ITALY
E-mail address: `enrico.carlini@monash.edu`, `enrico.carlini@polito.it`

JOE KILEEL, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720, USA
E-mail address: `jkileel@math.berkeley.edu`