

Increased tissue oxygenation explains the attenuation of hyperemia upon repetitive pneumatic compression of the lower leg

Original

Increased tissue oxygenation explains the attenuation of hyperemia upon repetitive pneumatic compression of the lower leg / Messere, Alessandro; Ceravolo, Gianluca; Franco, Walter; Maffiodo, Daniela; Ferraresi, Carlo; Roatta, Silvestro. - In: JOURNAL OF APPLIED PHYSIOLOGY. - ISSN 8750-7587. - 123:6(2017), pp. 1451-1460. [10.1152/jappphysiol.00511.2017]

Availability:

This version is available at: 11583/2696430 since: 2018-01-09T10:41:51Z

Publisher:

American Physiological Society

Published

DOI:10.1152/jappphysiol.00511.2017

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

**Increased tissue oxygenation explains the attenuation of hyperaemia upon repetitive
pneumatic compression of the lower leg**

Alessandro Messere¹, Gianluca Ceravolo², Walter Franco², Daniela Maffiodo², Carlo Ferraresi², Silvestro
Roatta¹

Author contribution

AM: conception and design of the experiment, collection, analysis and interpretation of the data, drafting the
manuscript

GC: collection, analysis and interpretation of the data, drafting of the manuscript

WF: design of the experimental set-up, collection, analysis and interpretation of the data

DM: design of the experimental set-up, collection, analysis and interpretation of the data

CF: design of the experiment, critical revision of the manuscript

SR: conception and design of the experiment and critical revision of the manuscript

All authors approved the final version of the manuscript.

¹ Dept. Neuroscience, University of Torino, Torino, Italy

² Dept. of Mechanical and Aerospace Engineering, Politecnico di Torino, Torino, Italy

Running Head Tissue oxygenation modulates compression-induced hyperaemia

Corresponding author

Silvestro Roatta

Dip. Neuroscienze, Università di Torino

C.so Raffaello 30, 10125 Torino, Italy

Email: silvestro.roatta@uni.to.it

26 **ABSTRACT**

27 **Aim**

28 The rapid hyperaemia evoked by muscle compression is short-lived and was recently shown to undergo a
 29 rapid decrease even in spite of continuing mechanical stimulation. The present study aims at investigating
 30 the mechanisms underlying this attenuation which include local metabolic mechanisms, desensitization of
 31 mechano-sensitive pathways, and reduced efficacy of the muscle pump.

32 **Methods**

33 In 10 healthy subjects short sequences of mechanical compressions (n=3-6; 150 mmHg) of the lower leg
 34 were delivered at different inter-stimulus intervals (ranging from 20 to 160 s) through a customized
 35 pneumatic device. Hemodynamic monitoring included near infrared spectroscopy, detecting tissue
 36 oxygenation and blood volume in calf muscles, as well as simultaneous echo-Doppler measurement of
 37 arterial (superficial femoral artery) and venous (femoral vein) blood flow.

38 **Results**

39 The results indicate that: i) a long lasting (>100 s) increase in local tissue oxygenation follows the
 40 compression-induced hyperaemia ; ii) the compression-induced hyperaemia exhibits different patterns of
 41 attenuation depending on the inter-stimulus interval; iii) the amplitude of the hyperaemia is not correlated
 42 with the amount of blood volume displaced by the compression; iv) the extent of attenuation negatively
 43 correlates with tissue oxygenation ($r=-0.78$, $P<0.05$).

44 **Conclusion**

45 Increased tissue oxygenation appears to be the key factor for the attenuation of hyperaemia upon repetitive
 46 compressive stimulation. Tissue oxygenation monitoring is suggested as a useful integration in medical
 47 treatments aimed at improving local circulation by repetitive tissue compression.

49 **NEW AND NOTEWORTHY**

50 This study shows that i) the hyperaemia induced by muscle compression produces a long-lasting increase in
 51 tissue oxygenation; ii) the hyperaemia produced by subsequent muscle compressions exhibits different
 52 pattern of attenuation at different inter-stimulus intervals; iii) the extent of attenuation of the compression-
 53 induced hyperaemia is proportional to the level of oxygenation achieved in the tissue. The results support the
 54 concept that tissue oxygenation is a key variable in blood flow regulation.

56 **Keywords:** muscle blood flow, hyperaemia, muscle compression, tissue oxygenation.

58 **Glossary**

59 IPC intermittent pneumatic compression

60 ISI inter-stimulus interval

61 NIRS near-infrared spectroscopy

62 SRS spatially-resolved spectroscopy

63 THI total haemoglobin index

64 TOI tissue oxygenation index

65

66

67 **INTRODUCTION**

68 Since the seminal work of Mohrman and Sparks (39) several studies have demonstrated that a rapid and
69 transient hyperaemic response can be elicited by a short-lasting muscle compression (10, 30, 38, 56-60).

70 Although the underlying mechanisms have not been fully identified, this phenomenon has been well
71 documented in different experimental models, such as the isolated muscle (39), awake and anesthetized
72 animals (57, 58, 60) and humans (10, 30, 38, 56). In addition a rapid dilatory response to compressive
73 stimuli has also been observed in isolated feed arteries (7). More controversial is the hemodynamic response
74 to repeated compressive stimuli. Kirby et al (30) observed that the response to 5 consecutive compressions
75 was non-significantly attenuated with respect to the response to a single compression. Conversely, Clifford et
76 al (7) using the same pattern of 5 consecutive compressive stimuli on an isolated muscle feed arteries
77 observed a significant increase of the dilatory response as compared to the single compression.

78 In a recent work Turturici and colleagues investigated the blood flow response to a longer lasting sequence
79 of mechanical stimulations (20 compressions, 1 s ON /1 s OFF) reporting that the initial hyperaemic
80 response progressively fades away in spite of continuing stimulation and hypothesized that the
81 mechanosensitive mechanism underlying the response could undergo some kind of transient inactivation
82 (60). In fact, the attenuation of the compression-induced hyperaemia was observed to increase at increasing
83 stimulation frequencies (60). A similar behavior was recently observed also in humans (38).

84 Surprisingly this phenomenon has been poorly described in the several investigations concerning the
85 hyperaemic effect of intermittent pneumatic compressions (IPC) (14, 15, 32-34), and in experimental studies
86 investigating the mechanisms underlying compression and contraction-induced hyperaemia (9, 24, 31, 40,
87 44), with the exception of a short report by Tschakowsky et al (56). In this pioneering investigation the
88 authors observed that repetitive compression of the forearm below heart level exhibited a transient
89 hyperaemia settling to a lower level after 10-20 s from the beginning of the treatment (56). More recently
90 Sheldon et al (47) also reported attenuation of the hyperaemia during IPC treatment, although on a larger

time scale (45 vs. 5 min from the beginning) and observed that the effect was dependent on the frequency of stimulation.

The issue is relevant because improving limb perfusion is a major aim in the treatment of disorders such as the peripheral arterial disease and is pursued in sport medicine for accelerated recovering from fatigue (1, 35). Understanding of the underlying mechanisms is essential for implementing optimal treatments (46). Potential mechanisms underlying attenuation of the hyperaemia during repetitive mechanical stimulation include: 1) inactivation of the mechano-sensitive vasodilatory pathways (60), 2) diminished efficacy of the muscle pump (56), and 3) local regulatory mechanisms that may be activated in response to hyper-perfusion (30, 56). Unfortunately, none of these possibilities is supported by a solid experimental evidence. In particular, 1) mechano-sensitive channels exhibiting inactivation properties have been identified (17, 26), but their actual involvement in the rapid compression-induced dilatation was not ascertained, 2) at high stimulation frequencies incomplete vascular refilling may reduce the contribution of the pump, however, a role for the muscle pump was excluded in a previous animal study (60), and 3) local vasoconstrictory mechanisms are known to act in response to hyper-perfusion but little is known about the actual regulatory variable (O_2 , CO_2 , pH, etc.) and about the strength and timing of this vascular reaction (6, 45). However, in a recent reformulation of the metabolic control of blood flow, a primary role for tissue pO_2 has been postulated (23). According to their model, an excessive rise in O_2 concentration within the tissue would trigger a vasoconstrictory response, mediated by the inhibition of a tonically released vasodilator (23). Along this line, a rise in tissue O_2 occurring during a compression-induced hyperaemia could then trigger a constrictor response and limit further hyperaemic events in response to subsequent mechanical stimuli.

On this basis the present study was aimed to test the following hypotheses: 1) the compression-induced hyperaemia elicits a rise in tissue oxygenation, 2) the attenuation of the hyperaemic response to subsequent compressive stimuli is related to the extent of hyper-oxygenation achieved in the tissue, and 3) the other mechanisms, namely, the intrinsic inactivation of mechano-sensitive pathways and the muscle pump would have a minor role in the attenuation of the hyperaemic response upon repetitive compressive stimulation. In order to assess changes in tissue oxygenation, the near infrared spectroscopy (NIRS) was adopted. By locating the NIRS probe under the compressive cuff, continuous monitoring of local oxygenation and blood volume changes from the relevant muscles was achieved. Moreover, in addition to arterial inflow, venous outflow was also monitored as its response to the compression is an indicator of the extent of filling of the venous compartments and thus, of the efficacy of the muscle pump exerted by compressive stimuli.

MATERIALS AND METHODS

Ethical approval

Ten healthy subjects (8 men and 2 women; age: 27.1 ± 3.0 years; weight: 67.9 ± 11.7 kg; height: 176.7 ± 9.7 cm) were recruited for the present study. All subjects were normotensive and non-obese.

The study conformed to the standards set by the Declaration of Helsinki and was approved by the Local

127 Ethical Committee (Prot. # 60195) and all subjects gave their written informed consent after they were
 128 instructed about purpose and procedures of the experiment.

129

130 *Mechanical leg compressions*

131 A previously tested prototype of IPC device was employed in the present study to deliver controlled and
 132 repeatable compressions to the leg of the subject (19, 20). Briefly the device exerts a compressive action by
 133 inflating five different bladders wrapped around the foot and the calf of the subject, with programmable
 134 pressure levels and timing. In the present study all bladders were inflated simultaneously to a supra-systolic
 135 pressure of 150 mmHg, with inflation and deflation times of about 3 s each. Two digital pulses are generated
 136 by the device to signal the starting time of both inflation and deflation.

137

138 *Near-infrared spectroscopy*

139 Local hemodynamic changes induced by leg compression were measured using a continuous wave NIRS
 140 device (NIRO-200NX, Hamamatsu Photonics, Hamamatsu City, Japan), which, besides the classical
 141 modified-Lambert-Beer method, supports spatially-resolved spectroscopy (SRS) (16, 52). Since mechano-
 142 sensitive vascular reactivity appears to be more prominently expressed by muscular than cutaneous tissues
 143 (57) we focused our attention on SRS parameters which, being less affected by cutaneous circulation,
 144 provide a more specific monitoring of muscle tissue (2, 36, 37). Since NIRS cannot discriminate between
 145 haemoglobin (Hb) and myoglobin (Mb), all measurements always refer to Hb+Mb in the sample volume
 146 (51). In particular, TOI (tissue oxygenation index) indicates the ratio $(MbO_2 + HbO_2) / (Mbtot + Hbtot)$
 147 expressed in percentage, and THI (tissue haemoglobin index) indicates the concentration of (Hb+Mb) in
 148 arbitrary units and is therefore an indicator of blood volume changes. Classical Lambert-Beer Parameters
 149 (O_2Hb and HHb detecting changes in the concentration of oxygenated and deoxygenated (Hb+Mb),
 150 respectively) are only displayed in Fig. 1 and not further considered in the study.

151

152 *Hemodynamic measurements*

153 Measurements of blood velocity in femoral artery and vein were performed simultaneously using two
 154 ultrasound systems (MyLab 25 XVision and MyLab 25 Gold, Esaote, Genoa, Italy) equipped with linear
 155 arrays (LA 523, Esaote, Genoa, Italy). Superficial femoral artery and femoral vein were insonated distally to
 156 the inguinal ligament. Since these instruments could not measure blood velocity and vessel diameter
 157 simultaneously, the latter was measured at the beginning and at the end of every stimulation protocol.
 158 Doppler measurements were performed by extending the sample volume over the whole vessel size,
 159 echographically displayed (transversal approach) in real time. All blood velocity measurements in femoral
 160 artery were obtained with insonation angle of about 60° (operating frequency of 6.6 MHz) instead, a higher
 161 angle of about 70° (operating frequency of 5 MHz) was used in order to avoid saturation of the recording
 162 when assessing the high-speed venous outflow propelled by leg compression. The two probes were placed

few centimeters apart with the ultrasound beam of the proximal probe oriented proximally and the one of the distal probe oriented distally, in order to avoid interference between the measurements.

Experimental set-up

A schematic representation of the experimental setup is reported in Fig 1 A. All experiments were performed in a quiet room with a constant ambient temperature of about 22-23 °C. The subject sat upright on an adjustable chair with the back supported by a back rest.

The NIRS probe was located on the lateral head of gastrocnemius muscle of the right leg (inter-optode distance = 4 cm). The IPC device was wrapped around the lower leg, over the NIRS probe. The two echographic probes were maintained in place by dedicated holders for the whole duration of the protocol.

Experimental protocol

After 15 min of rest, an initial series of 3 compressive stimuli with inter-stimulus interval (ISI) of 160 s was delivered to the subject. After other 4 min of rest four series of 6 compressive stimuli were delivered at different frequency (ISI= 20, 40, 60 and 80 s) in randomized order and separated by 4-min resting intervals. Femoral artery and femoral vein diameters were collected at the beginning and at the end of every stimulation protocol. Diameters were measured along a single direction, since both vessels present a circular cross-section in these experimental conditions. Average diameter of the artery was calculated as $(D_s + 2 \cdot D_d)/3$, D_s being the systolic and D_d the diastolic diameter.

Data acquisition and processing

The NIRS signals were digitally acquired along with both Doppler audio signals and the digital synchronism signal from the IPC device by a single acquisition system (CED Micro 1041, Cambridge Electronic Design, Cambridge, UK) and stored on the computer for later analysis with Spike2 software (version 6.10, Cambridge Electronic Design, UK).

A specific algorithm was implemented in the Spike2 script language to calculate blood velocity from Doppler audio signals (11, 25). Briefly, power spectra of the audio signals were computed by the Fast Fourier Transform over non-overlapping epochs lasting 25.6 ms. From each spectrum the maximum frequency of the signal (corresponding to maximum blood velocity) was estimated according to D'Alessio (11), then the mean frequency was calculated as the average of all frequencies below the maximum, weighted according to spectral amplitude (25). The mean frequency was then time-averaged over each cardiac cycle and converted into blood velocity, $BV = (MF \cdot C)/(2F \cdot \cos\theta)$, where MF is the mean frequency calculated from Doppler shift, C the averaging speed of ultrasound in soft tissue (1540 m/s), F the operating frequency of the Doppler, and θ the insonation angle). Blood flow, in ml/min, was then calculated

197 as mean blood velocity times cross-sectional area of the vessel ($BF = BV * \pi r^2 * 60$, where BV is the blood
198 velocity expressed in cm/s, and πr^2 the cross sectional area of the vessel in cm^2).

199 The response to each compressive stimulus was characterized by: pre-compression arterial blood flow,
200 calculated as the average over the 4 s preceding the compression; pre-compression TOI; pre-compression
201 THI; peak arterial blood flow, as the hyperaemic peak reached after the compression; Δ TOI, calculated as
202 the difference between the peak TOI reached after the compression and pre-compression TOI; displaced
203 blood volume, calculated as the product of the area under the curve of the venous blood velocity response
204 and the cross-sectional area of vein.

205 In addition, the amplitude of the hyperaemic response was also calculated as the difference between peak
206 arterial flow and pre-compression flow.

207 In order to assess the extent of attenuation of the response throughout the experimental protocol, changes in
208 blood flow and blood volume were normalized to the changes observed in response to the first delivered
209 compressive stimulus.

210

211 **Statistics**

212 To examine the effect of repetitive leg compression performed at different ISI on peak blood flow, displaced
213 blood volume, pre-compression THI and pre-compression TOI, a two-way repeated-measures ANOVA was
214 used with factors ISI and repetition (GraphPad Prism v 6.0, GraphPad Software, La Jolla, CA). When
215 significance was found, a Dunnett's post hoc test was performed to assess significant changes within each
216 series with respect to the response to the first stimulus. Pearson's coefficient was used to assess the
217 correlation between different variables. All data are expressed as means \pm standard deviation in the text and
218 means \pm standard error in diagrams. The level of statistical significance was set at $P < 0.05$.

219

220

221 **RESULTS**

222

223 *Single leg compression*

224 A typical response to a single compressive stimulus is reported in figure 1B. Venous blood velocity exhibits
225 a prompt and short-lasting increase, peaking 1.7 ± 0.2 s after the beginning and terminating before the end of
226 the compression. The blood volume displaced by compression was on average 28.3 ± 14.8 ml. The increase
227 in arterial blood flow starts immediately after deflation and peaks in 4.9 ± 1.4 s passing from a basal value of
228 74.5 ± 22.7 ml min⁻¹ to 260.2 ± 83.3 ml min⁻¹ during the peak (peak flow is 3.6 ± 1.0 of baseline). Blood
229 flow generally returns within 15-25 s The response in tissue oxygenation is further delayed. TOI slowly
230 increases (from 66.4 ± 5.1 to 78.0 ± 4.0 %) and peaks after 20.6 ± 5.1 s from deflation. Local changes in
231 blood volume are detected by THI exhibiting a rapid decrease during compression followed by a slower
232 return to the basal level, in agreement with the changes in venous and arterial blood flow, respectively.

233

234 *Repeated leg compressions*

235 The hemodynamic response to repetitive leg compression at different ISI is summarized in Fig 2, each
 236 column representing the response to a single stimulus. The upper two rows show the response in terms of
 237 peak arterial blood flow and displaced venous blood volume, both variables exhibiting a significant
 238 dependence on ISI ($p < 0.01$) and repetition ($p < 0.01$). It can be observed that when ISI = 160 s the response to
 239 subsequent stimuli is unchanged. Unchanged response in terms of peak arterial flow and displaced blood
 240 volume is also observed in response to the first compression in each series. Instead, both parameters exhibit a
 241 progressive attenuation although with different time course at ISI ranging from 20 to 80 s. In particular, the
 242 hyperaemia is consistently reduced starting from the second stimulus in the series, at ISI ranging from 20 to
 243 60 s, while displaced blood volume is consistently reduced at ISI = 20 and 40 s, starting from the third
 244 stimulus. A peculiar pattern is observed at ISI = 80 s where hyperaemia is only attenuated in response to
 245 even and not to odd stimuli, while, at the same time displaced blood volume remains unaffected.

246 NIRS parameters, shown in the lower rows of fig 2, exhibited a significant dependence on repetition
 247 ($p < 0.01$) but not on ISI, along with a significant interaction between the two factors. It can be observed that
 248 pre-compression THI, which can be considered an indicator of vascular filling, qualitatively parallels the
 249 changes in displaced blood volume, remaining unchanged at large ISI and exhibiting the most marked
 250 reduction at ISI = 20 s. Pre-compression TOI exhibits instead marked increases at all ISIs lower than 160 s
 251 starting from the second stimulus in the sequence. It is interesting to observe that its pattern of change is
 252 opposite to peak blood flow: i.e., hyperaemic peak is higher if the pre-compression TOI is lower. Note also
 253 that the oscillating pattern previously observed in peak blood flow at ISI = 80 s is also exhibited by pre-
 254 compression TOI in an opposite way.

255 In order to provide a better understanding of the interplay between the different parameters in the peculiar
 256 response to repetitive compression at ISI = 80 s, original tracings are reported from a representative subject
 257 in Fig. 3. As described in Fig. 1, the first stimulus elicits a marked hyperaemia which results in a marked
 258 increase in oxygenation. The following compression, which occurs when the tissue oxygenation is still high,
 259 now elicits a much smaller hyperaemia, resulting in a proportionally smaller increase in TOI and attenuated
 260 vascular refilling in THI. The third compression occurs when the TOI is almost returned to basal levels and
 261 the elicited hyperaemia resumes its original size. Although it cannot be fully appreciated with this time scale,
 262 the venous blood flow response is comparable in all instances as well as the pre-compression level reached
 263 by THI.

264 Another representative recording illustrating the pattern at ISI = 20 s is reported in fig 4. Note the
 265 disappearance of the hyperaemic response to the second and subsequent stimuli in spite of the fact that
 266 arterial blood flow is returned to basal level. A weak hyperaemia reappears only in response to the last
 267 stimulus, when also TOI is almost returned to basal level. Note that THI indicates that blood volume is

almost fully returned to basal level after the first stimulus (thanks to the marked hyperaemia) but not afterwards. Accordingly, the venous response is markedly reduced after the third and subsequent stimuli.

In general a good correlation was found between the peak blood flow during hyperaemia and the ensuing increase in oxygenation as shown in fig 5 A, in which all subjects have been pooled and each dot represents the response to a single compression. The overall r is 0.76 ($p < 0.05$). When individually computed for the different subjects r ranged between 0.72 and 0.95 ($p < 0.05$) (average 0.78 ± 0.1).

On the contrary the hyperaemic response was not correlated with the amount of displaced blood volume as shown in Fig. 5B ($r = 0.34$, individual r ranging between -0.4 and $+0.3$).

Fig 5C shows the correlation between pre-compression TOI and the peak of the hyperaemic response which is exhibiting an overall $r = -0.434$ ($p < 0.05$), however a much higher within- subject correlation is observed: -0.78 ± 0.06 , individual r ranging between 0.7 and 0.9 ($p < 0.05$).

In Fig. 5D the amplitude of the hyperaemic response (= peak flow-basal flow) instead of peak flow is plot vs. pre-compression TOI. While the general pictures resembles that of Fig. 5C, it is here better evidenced that the hyperaemia can be almost abolished at high TOI levels. Moreover, the slope of the regression lines, m , allows to quantify the dependence of the hyperaemic response on tissue oxygenation. On average, $m = -0.082 \pm 0.0026$ meaning that the compression-induced hyperaemia is attenuated by 8% per unitary increase of TOI, with respect to its full amplitude (the one that is evoked in resting conditions).

Changes in vessel size

A slight increase in vessel diameter was detected from the comparison of measurements performed at the beginning and at completion of the experimental protocol in both femoral artery (from 6.0 ± 0.8 to 6.2 ± 0.8 mm, $p < 0.05$) and vein (from 8.3 ± 0.9 to 8.6 ± 1.3 mm, $p < 0.05$)

DISCUSSION

For the first time a comprehensive approach has been employed for the investigation of the rapid compression-induced hyperaemia and its adaptation upon repetitive stimulation, which includes continuous assessment of NIRS indicators of changes in local tissue oxygenation and blood volume as well as simultaneous monitoring of arterial inflow and venous outflow. This allowed us to describe the early hyperaemic changes taking place at the beginning of IPC treatments at different frequencies, and to confirm our initial hypotheses: i) the compression-induced hyperaemia elicits proportional increases in local tissue oxygenation; ii) the extent of attenuation of the hyperaemic response to subsequent stimuli is related to the current level of tissue oxygenation; iii) the extent of attenuation is not strictly dependent on the extent of vascular filling and on the ISI, therefore the attenuation cannot be attributed to the reduced efficacy of the muscle pump or to a simple, time-dependent, inactivation mechanism of mechano-sensitive pathways.

302

303 *Compression-induced hyperaemia increases tissue oxygenation*

304 A novel observation of the present study is that muscle compression elicits a prominent increase in local
 305 tissue oxygenation. This increase is consequent to the induced hyperaemia but is much longer lasting. This
 306 aspect is important because it reveals that the return to “control conditions” is not achieved at the end of the
 307 hyperaemia, which normally occurs within 15-25 s and may instead require up to 100 - 200 s. This pattern
 308 has never been reported for compression-induced hyperaemia but it is in agreement with what occurs in the
 309 rapid-onset hyperaemia induced by short contractions (53).

310 It is generally accepted that an increase in perfusion, with unchanged metabolism, increases tissue
 311 oxygenation (3, 12). In the present condition, different factors could contribute to the observed TOI increase
 312 in response to compression-induced hyperaemia: 1) depletion of the venous-compartment, which alters the
 313 proportion of arterial/venous blood in the sample volume; 2) increased Hb saturation in venous blood due to
 314 decreased oxygen extraction, given that the hyperaemia occurs in a condition of constant metabolism; 3)
 315 increased saturation of myoglobin. The voiding of venous compartment does not seem to affect the TOI
 316 signal considerably, as no relevant changes are observed immediately after the compression, including those
 317 associated with large blood volume changes (see original tracings in Figs. 1, 3 and 4). Unfortunately, NIRS
 318 cannot discriminate between Mb and Hb saturation nor between arterial and venous compartments, thus no
 319 univocal explanation can be provided. Irrespective of the underlying reason, the increase in tissue
 320 oxygenation was a very consistent feature of the hemodynamic response to the compression of the resting
 321 muscle and exhibited a good correlation with the amplitude of hyperaemia (Fig. 5A).

322

323 *Is compression-induced hyperaemia attenuated by increased tissue oxygenation?*

324 Several lines of evidence from the present study support the finding that elevated tissue oxygenation is the
 325 factor responsible for the attenuation of the hyperaemia and for the reduced responsiveness to the mechanical
 326 stimulus. By looking at the original tracings of Fig. 3 it can be observed that the response to the second
 327 compression is smaller as compared to the first and the third responses, while TOI is higher than baseline.
 328 The same is visible in Fig.4: the hyperaemic response almost disappears during the initial high oxygenation
 329 phase and only later exhibits a tendency to recover, concomitantly with a decrease in TOI. This dependence
 330 of peak hyperaemia on pre-compression TOI is also supported by the histograms of Fig. 2 (see opposite
 331 patterns of peak blood flow and pre-compression TOI) and is quantitatively assessed by the correlations in
 332 Fig. 5 C and D. Moreover, it appears to be rather linear and rather similar between different subjects.
 333 According to these indications, the amplitude of the hyperaemic response is attenuated by 8 ± 2 % per
 334 unitary increase of TOI meaning that an increase in TOI by 12.5 points virtually abolishes the response.

Notably, the dependence of the active vessel dilatation on tissue oxygenation may explain why the same short sequence of compressive stimuli elicited opposite effects in vitro (7), where tissue hyperoxia does not take place, and in vivo (30).

In the several studies investigating hemodynamic effects during IPC treatments this pattern of adaptation of the hyperaemia has not been described, possibly because the attention was focused on medium-long term rather than on early effects. Although different devices and patterns of stimulation have been used in previous investigations, an increase in limb perfusion is generally reported, ranging between 20 and 240 %, and being assessed at 5-60 min from the beginning of the treatment (9, 15, 24, 33, 40, 44, 47), which also appear to be little dependent on the stimulation frequency (47). These results are not readily comparable with the present ones because no steady state was reached in our study. It is reasonable to expected that a certain stable increase in perfusion is obtained with prolonged stimulation, once steady tissue oxygenation is achieved.

Underlying mechanisms and implications

As discussed above, the attenuation of the mechano-sensitive dilatory response to multiple compressions could result as a reaction of the tissue to the hyper-perfusion (generated in response to the first stimulus), which entails the washout of metabolites and alteration of the local milieu in which PO_2 is a most relevant variable (4, 23, 27). It is well known that low oxygenation stimulates vasodilatation and, conversely, that increased oxygenation leads to vasoconstriction, although the effects generally observed in humans exposed to increased levels of inspired PO_2 are rather small (5, 62). In the latter study, increasing arterial PO_2 from 100 to 2100 mmHg increased resting vascular conductance only by 20-25% and reduced functional hyperaemia by 20% (5). However it must be observed that tissue PO_2 is differently affected by increased arterial PO_2 and hyper-perfusion. In fact while the hyperbaric hypoxia at 2100 mmHg increases the amount of oxygen carried to the tissue by about 30% (5) a 2-fold increase in perfusion results in a 200% increase in oxygen flow. In early studies reactions to hyper-perfusion were investigated on isolated preparations with externally-controlled blood supply (21). However these studies could not provide a clear indication of the time course of the local tissue response, nor could they discriminate between “metabolic” and myogenic response, given that hyper-perfusion was produced by increased perfusion pressure which also resulted in increased transmural pressure (48). In this respect, the compression-induced hyperaemia offers a peculiar model of (transient) tissue hyper-perfusion, characterized by unchanged tissue metabolism, unchanged arterial PO_2 and most likely unchanged neuro-hormonal drive.

The prompt counter-reaction to the compression-induced hyperaemia and the concomitant inactivation of the mechano-sensitive dilatation upon increased tissue oxygenation fits with the “bang-bang” model of blood flow control, recently proposed by Golub & Pittman (23) according to which the feedback signal (O_2^- , whose concentration increases in response to increased O_2 availability) carries the information of excessive

370 perfusion and operates a vasoconstriction by inactivating the tonically released vasodilators (namely, nitric
371 oxide), aim of this regulation being to protect the tissue from hyperoxia and prevent excessive perfusion.

372 Accordingly, the vascular mechano-sensitivity, which is considered to mediate the rapid dilatation and the
373 anticipatory (feed-forward) hyperaemia at the beginning of exercise (8, 30, 43, 60) is promptly abolished if
374 the exercise does not take place, due to the hyper-oxygenation produced by the hyper-perfusion. Instead, in
375 the case of exercise the hyper-oxygenation is quickly reduced even below control levels (18) by increased
376 metabolism and no limitation to vasodilation takes place, which results in the "functional hyperaemia". The
377 same mechanism is likely to explain why both passive movement hyperaemia is attenuated upon repeated
378 stimulation (54, 55) and contraction-induced hyperaemia is attenuated after a sequence of muscle
379 compressions (38).

380 Surprisingly, with one exception (34) no study has ever included NIRS in the characterization of the
381 hyperaemic response to compression and IPC. Although tissue oxygenation can be considered a major
382 outcome of perfusion, in the short term it does not strictly follow arterial blood flow, e.g., in Fig. 2 TOI is
383 maintained at high levels for some time, after the end of hyperaemia. On this basis, it might be more
384 appropriate to monitor TOI rather than blood flow in order to better appreciate the actual effects of the
385 treatment. In addition, adopting NIRS as the monitoring technique gives the possibility to assess the effects
386 specifically on the tissue of interest, as compared to the more global information provided by blood flow in
387 an large supplying artery.

388

389 *Alternative hypothesis 1: Vascular refilling and the muscle pump*

390 The parallelism observed between changes in pre-compression THI and in displaced blood volume (Fig.2),
391 suggests that pre-compression THI is a good indicator of current vascular filling and that its changes mostly
392 reflect volume changes of the venous compartment. By observing its time course after the compressive
393 stimulus we can detect a fast refilling phase, associated to the possible concomitant hyperaemia, and a
394 subsequent slow phase, associated to "resting" blood flow. At high ISI, i.e., 80 and 160 s, a complete
395 vascular refill is granted by both a consistent hyperaemia and a large time interval. Accordingly, the
396 compressive action of the device displaces comparable amount of blood volume at every stimulus. At lower
397 ISI, the lack of hyperaemia and/or insufficient time for the slow phase to yield a significant contribution may
398 result in incomplete vascular refilling and in a reduction of the blood volume displaced by the subsequent
399 compression. This observation is in agreement with the study by Delis and colleagues who reported 3 to 4
400 compressions per minute (i.e., ISI = 20 or 15 s) as the optimum stimulation frequency to maintain low
401 venous pressure in the treated limb (13). Valic et al (61), in the anesthetized dog estimated a refilling time of
402 less than 1 s due to the large contraction-induced hyperaemia. Based on direct foot venous pressure
403 estimation, two human studies reported refilling times of 16 - 40 s after 10 tip-toe movements (42) and
404 pneumatic compression (22). In the present conditions the refill could take place in 10-15 s through the rapid

405 phase in the presence of large hyperaemia but could otherwise require more than one minute when
 406 hyperaemia was blunted (Fig. 3).

407 According to the “muscle pump” effect, an increase in intramuscular pressure empties the venous
 408 compartments producing a decrease in venous pressure, which in turn increases the artero-venous pressure
 409 gradient thus contributing to the ensuing hyperaemia. This mechanism is activated both with active muscle
 410 contraction as well as with the compression of the passive muscle and has been often considered to explain
 411 the larger hyperaemic responses observed when compressing (10, 56) or contracting (41, 50) limbs muscles
 412 below as compared to above heart level. However the issue is still debated (7, 29, 49) due to the conflicting
 413 evidence provided by other studies (24, 28, 61). In particular, Jasperse et al. investigated the effect of
 414 positional differences on reactive hyperaemia, as a model of hyperaemia dissociated from the muscle pump.
 415 They showed that also reactive hyperaemia is larger when evoked below, with respect to above heart level,
 416 suggesting that positional effects may be secondary to differences in driving pressure rather than to the
 417 muscle pump. The present results support this view through a complementary model, i.e., the muscle pump
 418 action dissociated from the hyperaemia. This particular condition was observed in several instances such as
 419 the responses to the second compression at ISI ranging from 20 to 80 s (in Fig 2 and in Fig 4), in which
 420 maintained vascular filling and compression-displaced blood volume, i.e., an effective muscle pump, was
 421 associated with a considerably reduced hyperaemia, as compared to the first compression in the series. This
 422 proves that the muscle pump mechanism is not involved in the attenuation of the hyperaemia in response to
 423 multiple compressions. Whether the muscle pump plays a role in the hyperaemic response to the first
 424 compressive stimulus cannot be ruled out based on the present data. In fact, from scatter plot in Fig 5B we
 425 can observe that the largest hyperaemic responses were never associated with low displaced blood volume,
 426 which suggests that adequate vascular filling may be a necessary condition to express the full response.
 427 Investigating the mechanisms behind compression-induced hyperaemia was not an aim of this study; further
 428 investigations will be necessary to elucidate this issue.

429

430 *Alternative hypothesis 2: Desensitization of mechano-sensitive pathways*

431 It was previously observed that the hyperaemic response to the compressive stimulus progressively reduced
 432 to 26% of its original amplitude, with decreasing ISI from 4 min to 2 s (60). On this basis the hypothesis was
 433 put forward that the attenuation could be due to some transient inactivation (desensitization) of mechano-
 434 sensitive dilatory mechanisms. This hypothesis was supported by the observation that desensitization upon
 435 repeated activation is a characteristic of certain vascular mechano-sensitive channels (17, 26). A subsequent
 436 human study in which similar stimulation protocols were applied to the forearm, qualitatively confirmed the
 437 attenuation pattern, although with a less gradual dependence on the ISI (38)

438 The up-and-down pattern exhibited by compression-induced hyperaemia at ISI = 80 s (Fig. 2 and Fig. 3)
 439 seems to exclude a simple, frequency-dependent, desensitization mechanism of mechano-sensitive pathways,
 440 as previously hypothesized (38, 60). More complex desensitization patterns possibly affecting multiple

mechanosensitive pathways cannot be excluded based on the present data. However, in order to explain the peculiar hyperaemic responses observed at ISI=80 s, such desensitization pattern should exhibit an up-and-down time course, as exhibited by TOI, which would appear a quite unlikely coincidence.

Limitations

Manual assessment of insonation angles, as required with the transversal approach, is not very accurate and may introduce systematic errors in the calculation of absolute flow values. This is particularly true for assessment of venous blood flow since a wide angle between the vessel axis and the ultrasound beam had to be adopted in order to avoid saturation of the velocity signal (aliasing). However, the analysis was here focused on relative changes, thereby eliminating errors associated with measurement of the insonation angle.

Diameter of the femoral vein was not continuously monitored. Possible enlargement of the vessel during the passage of the blood volume displaced by the compression may have resulted in underestimation of venous flow.

Diameter of both femoral artery and vein exhibited a small increase throughout the experimental protocol, which was not accounted for. This may also have led to increasing underestimation of blood flow with time. Since the sequence of the series was randomized this aspect should not have affected the results.

Conclusions

This study allowed to prove that the attenuation of hyperaemia upon repetitive limb compression is not dependent on vascular filling and the muscle pump nor on a simple ISI-dependent desensitization of mechano-sensitive pathways. In addition, strong evidence is provided, supporting the concept that tissue hyper-oxygenation is the key signal underlying the inactivation of the rapid dilatory response to muscle compression. This evidence is however indirect and other studies are necessary to conclusively prove this assertion.

Irrespective of the underlying mechanisms, inactivation of mechano-sensitive pathways may almost completely abolish the compression-induced hyperemia, suggesting a role for this phenomenon in protecting the tissue from hyperperfusion and oxidative stress.

The hyperaemic response to muscle compression is proposed as a peculiar model for the investigation of the response to hyper-perfusion characterized by constant arterial pO_2 , constant tissue metabolism as well as modest or absent systemic reactions.

Finally, tissue oxygenation monitoring is recommended to assess the efficacy of IPC treatments, oriented to improve blood perfusion in limbs.

474 **Acknowledgements**

475 We are grateful to the Laboratory of Engineering of Neuromuscular System and Motor Rehabilitation
476 (LISiN, Politecnico di Torino) for lending us the NIRS device. We also thank all the volunteers enrolled in
477 the study.

478 **Grants**

479 This research was supported by the University of Torino.

480 **Disclosures**

481 No competing interest to declare.

REFERENCES

1. **Barnett A.** Using recovery modalities between training sessions in elite athletes: does it help? *Sports Med* 36: 781-796, 2006.
2. **Canova D, Roatta S, Bosone D, and Micieli G.** Inconsistent detection of changes in cerebral blood volume by near infrared spectroscopy in standard clinical tests. *J Appl Physiol* 110: 1646-1655, 2011.
3. **Canova D, Roatta S, Micieli G, and Bosone D.** Cerebral oxygenation and haemodynamic effects induced by nimodipine in healthy subjects. *Funct Neurol* 27: 169-176, 2013.
4. **Casey DP, and Joyner MJ.** Local control of skeletal muscle blood flow during exercise: influence of available oxygen. *J Appl Physiol* (1985) 111: 1527-1538, 2011.
5. **Casey DP, Joyner MJ, Claus PL, and Curry TB.** Hyperbaric hyperoxia reduces exercising forearm blood flow in humans. *Am J Physiol Heart Circ Physiol* 300: H1892-1897, 2011.
6. **Clifford PS, and Hellsten Y.** Vasodilatory mechanisms in contracting skeletal muscle. *J Appl Physiol* (1985) 97: 393-403, 2004.
7. **Clifford PS, Kluess HA, Hamann JJ, Buckwalter JB, and Jasperse JL.** Mechanical compression elicits vasodilatation in rat skeletal muscle feed arteries. *J Physiol* 572: 561-567, 2006.
8. **Clifford PS, and Tschakovsky ME.** Rapid vascular responses to muscle contraction. *Exerc Sport Sci Rev* 36: 25-29, 2008.
9. **Creelius AR, Kirby BS, Luckasen GJ, Larson DG, and Dinunno FA.** Mechanisms of rapid vasodilation following a brief contraction in human skeletal muscle. *Am J Physiol Heart Circ Physiol* 305: H29-40, 2013.
10. **Credeur DP, Holwerda SW, Restaino RM, King PM, Crutcher KL, Laughlin MH, Padilla J, and Fadel PJ.** Characterizing rapid-onset vasodilation to single muscle contractions in the human leg. *J Appl Physiol* (1985) 118: 455-464, 2015.
11. **D'Alessio T.** 'Objective' algorithm for maximum frequency estimation in Doppler spectral analysers. *Med Biol Eng Comput* 23: 63-68, 1985.
12. **Davis SL, Fadel PJ, Cui J, Thomas GD, and Crandall CG.** Skin blood flow influences near-infrared spectroscopy-derived measurements of tissue oxygenation during heat stress. *J Appl Physiol* 100: 221-224, 2006.
13. **Delis KT, Azizi ZA, Stevens RJ, Wolfe JH, and Nicolaides AN.** Optimum intermittent pneumatic compression stimulus for lower-limb venous emptying. *Eur J Vasc Endovasc Surg* 19: 261-269, 2000.
14. **Delis KT, Husmann MJ, Cheshire NJ, and Nicolaides AN.** Effects of intermittent pneumatic compression of the calf and thigh on arterial calf inflow: a study of normals, claudicants, and grafted arteriopathies. *Surgery* 129: 188-195, 2001.
15. **Delis KT, Labropoulos N, Nicolaides AN, Glenville B, and Stansby G.** Effect of intermittent pneumatic foot compression on popliteal artery haemodynamics. *Eur J Vasc Endovasc Surg* 19: 270-277, 2000.
16. **Delpy DT, Cope M, van der Zee P, Arridge S, Wray S, and Wyatt J.** Estimation of optical pathlength through tissue from direct time of flight measurement. *Phys Med Biol* 33: 1433-1442, 1988.
17. **Earley S, and Brayden JE.** Transient receptor potential channels in the vasculature. *Physiol Rev* 95: 645-690, 2015.
18. **Felici F, Quaresima V, Fattorini L, Sbriccoli P, Filligoi GC, and Ferrari M.** Biceps brachii myoelectric and oxygenation changes during static and sinusoidal isometric exercises. *J Electromyogr Kinesiol* 19: e1-11, 2009.
19. **Ferraresi C, Maffiodo D, and Hajimirzaalian H.** A model-based method for the design of intermittent pneumatic compression systems acting on humans. *Proc Inst Mech Eng H* 228: 118-126, 2014.
20. **Ferraresi C, Maffiodo D, and Hajimirzaalian H.** Simulation and Control of a Robotic Device for Cardio-Circulatory Rehabilitation. In: *Advances in Robot Design and Intelligent Control*, edited by Borangiu TSpringer, 2016, p. 357-365.
21. **Folkow B.** Intravascular pressure as a factor regulating the tone of the small vessels. *Acta Physiol Scand* 17: 289-310, 1949.
22. **Gaskell P, and Parrott JC.** The effect of a mechanical venous pump on the circulation of the feet in the presence of arterial obstruction. *Surg Gynecol Obstet* 146: 583-592, 1978.
23. **Golub AS, and Pittman RN.** Bang-bang model for regulation of local blood flow. *Microcirculation* 20: 455-483, 2013.
24. **Gonzalez-Alonso J, Mortensen SP, Jeppesen TD, Ali L, Barker H, Damsgaard R, Secher NH, Dawson EA, and Dufour SP.** Haemodynamic responses to exercise, ATP infusion and thigh compression in humans: insight into the role of muscle mechanisms on cardiovascular function. *J Physiol* 586: 2405-2417, 2008.
25. **Guiot C, Roatta S, Piccoli E, Saccomandi F, and Todros T.** Quantitative Doppler measures in coiled vessels: investigation on excised umbilical veins. *Ultrasound Med Biol* 25: 1465-1473, 1999.
26. **Honore E, Patel AJ, Chemin J, Suchyna T, and Sachs F.** Desensitization of mechano-gated K2P channels. *Proc Natl Acad Sci U S A* 103: 6859-6864, 2006.
27. **Jackson WF.** Arteriolar oxygen reactivity: where is the sensor and what is the mechanism of action? *J Physiol* 594: 5055-5077, 2016.

- 541 28. **Jasperse JL, Shoemaker JK, Gray EJ, and Clifford PS.** Positional differences in reactive hyperemia
542 provide insight into initial phase of exercise hyperemia. *J Appl Physiol* (1985) 119: 569-575, 2015.
- 543 29. **Joyner MJ, and Casey DP.** Regulation of increased blood flow (hyperemia) to muscles during exercise: a
544 hierarchy of competing physiological needs. *Physiol Rev* 95: 549-601, 2015.
- 545 30. **Kirby BS, Carlson RE, Markwald RR, Voyles WF, and Dinunno FA.** Mechanical influences on skeletal
546 muscle vascular tone in humans: insight into contraction-induced rapid vasodilatation. *J Physiol* 583: 861-874, 2007.
- 547 31. **Kirby BS, Markwald RR, Smith EG, and Dinunno FA.** Mechanical effects of muscle contraction do not
548 blunt sympathetic vasoconstriction in humans. *Am J Physiol Heart Circ Physiol* 289: H1610-1617, 2005.
- 549 32. **Labropoulos N, Leon LR, Jr., Bhatti A, Melton S, Kang SS, Mansour AM, and Borge M.** Hemodynamic
550 effects of intermittent pneumatic compression in patients with critical limb ischemia. *J Vasc Surg* 42: 710-716, 2005.
- 551 33. **Labropoulos N, Watson WC, Mansour MA, Kang SS, Littooy FN, and Baker WH.** Acute effects of
552 intermittent pneumatic compression on popliteal artery blood flow. *Arch Surg* 133: 1072-1075, 1998.
- 553 34. **Manfredini F, Lamberti N, Malagoni AM, Zamboni C, Basaglia N, Mascoli F, Manfredini R, and**
554 **Zamboni P.** Reliability of the vascular claudication reporting in diabetic patients with peripheral arterial disease: a
555 study with near-infrared spectroscopy. *Angiology* 66: 365-374, 2014.
- 556 35. **Martin JS, Friedenreich ZD, Borges AR, and Roberts MD.** Acute Effects of Peristaltic Pneumatic
557 Compression on Repeated Anaerobic Exercise Performance and Blood Lactate Clearance. *J Strength Cond Res* 29:
558 2900-2906, 2015.
- 559 36. **Messere A, and Roatta S.** Influence of cutaneous and muscular circulation on spatially resolved versus
560 standard Beer-Lambert near-infrared spectroscopy. *Physiol Rep* 1: e00179, 2013.
- 561 37. **Messere A, and Roatta S.** Local and remote thermoregulatory changes affect NIRS measurement in forearm
562 muscles. *Eur J Appl Physiol* 115: 2281-2291, 2015.
- 563 38. **Messere A, Turturici M, Mollo G, and Roatta S.** Repetitive muscle compression reduces vascular mechano-
564 sensitivity and the hyperemic response to muscle contraction. *J Physiol Pharmacol* In Press.
- 565 39. **Mohrman DE, and Sparks HV.** Myogenic hyperemia following brief tetanus of canine skeletal muscle. *Am J*
566 *Physiol* 227: 531-535, 1974.
- 567 40. **Mortensen SP, Thaning P, Nyberg M, Saltin B, and Hellsten Y.** Local release of ATP into the arterial
568 inflow and venous drainage of human skeletal muscle: insight from ATP determination with the intravascular
569 microdialysis technique. *J Physiol* 589: 2011.
- 570 41. **Nadland IH, Walloe L, and Toska K.** Effect of the leg muscle pump on the rise in muscle perfusion during
571 muscle work in humans. *Eur J Appl Physiol* 105: 829-841, 2009.
- 572 42. **Nicolaides AN, and Zukowski AJ.** The value of dynamic venous pressure measurements. *World J Surg* 10:
573 919-924, 1986.
- 574 43. **Poole DC, Ferreira LF, Behnke BJ, Barstow TJ, and Jones AM.** The final frontier: oxygen flux into muscle
575 at exercise onset. *Exerc Sport Sci Rev* 35: 166-173, 2007.
- 576 44. **Roseguini BT, Arce-Esquivel AA, Newcomer SC, Yang HT, Terjung R, and Laughlin MH.** Intermittent
577 pneumatic leg compressions enhance muscle performance and blood flow in a model of peripheral arterial
578 insufficiency. *J Appl Physiol* (1985) 112: 1556-1563, 2012.
- 579 45. **Secomb TW.** Theoretical models for regulation of blood flow. *Microcirculation* 15: 765-775, 2008.
- 580 46. **Sheldon RD, Roseguini BT, Laughlin MH, and Newcomer SC.** New insights into the physiologic basis for
581 intermittent pneumatic limb compression as a therapeutic strategy for peripheral artery disease. *J Vasc Surg* 58: 1688-
582 1696, 2013.
- 583 47. **Sheldon RD, Roseguini BT, Thyfault JP, Crist BD, Laughlin MH, and Newcomer SC.** Acute impact of
584 intermittent pneumatic leg compression frequency on limb hemodynamics, vascular function, and skeletal muscle gene
585 expression in humans. *J Appl Physiol* (1985) 112: 2099-2109, 2012.
- 586 48. **Shepherd JT** editor. *Circulation to skeletal muscle*. Bethesda: American Physiological Society, 1983, p. 319-
587 370.
- 588 49. **Sheriff D.** Point: The muscle pump raises muscle blood flow during locomotion. *J Appl Physiol* (1985) 99:
589 371-372; discussion 374-375, 2005.
- 590 50. **Shoemaker JK, Tschakovsky ME, and Hughson RL.** Vasodilation contributes to the rapid hyperemia with
591 rhythmic contractions in humans. *Can J Physiol Pharmacol* 76: 418-427, 1998.
- 592 51. **Spires J, Lai N, Zhou H, and Saidel GM.** Hemoglobin and myoglobin contributions to skeletal muscle
593 oxygenation in response to exercise. *Adv Exp Med Biol* 701: 347-352, 2011.
- 594 52. **Suzuki S, Takasaki S, Ozaki T, and Kobayashi Y.** Tissue oxygenation monitor using NIR spatially resolved
595 spectroscopy. In: *Optical Tomography and Spectroscopy of Tissue III*. San Jose, CA: 1999, p. 582-592.
- 596 53. **Towse TF, Slade JM, Ambrose JA, DeLano MC, and Meyer RA.** Quantitative analysis of the
597 postcontractile blood-oxygenation-level-dependent (BOLD) effect in skeletal muscle. *J Appl Physiol* (1985) 111: 27-39,
598 2011.
- 599 54. **Trinity JD, Groot HJ, Layec G, Rossman MJ, Ives SJ, Runnels S, Gmelch B, Bledsoe A, and Richardson**
600 **RS.** Nitric oxide and passive limb movement: a new approach to assess vascular function. *J Physiol* 590: 1413-1425,
601 2012.

55. Trinity JD, McDaniel J, Venturelli M, Fjeldstad AS, Ives SJ, Witman MA, Barrett-O'Keefe Z, Amann M, Wray DW, and Richardson RS. Impact of body position on central and peripheral hemodynamic contributions to movement-induced hyperemia: implications for rehabilitative medicine. *Am J Physiol Heart Circ Physiol* 300: H1885-1891, 2011.
56. Tschakovsky ME, Shoemaker JK, and Hughson RL. Vasodilation and muscle pump contribution to immediate exercise hyperemia. *Am J Physiol* 271: H1697-1701, 1996.
57. Turturici M, Mohammed M, and Roatta S. Evidence that the contraction-induced rapid hyperemia in rabbit masseter muscle is based on a mechanosensitive mechanism, not shared by cutaneous vascular beds. *J Appl Physiol* 113: 524-531, 2012.
58. Turturici M, and Roatta S. Compression-induced hyperaemia in the rabbit masseter muscle: a model to investigate vascular mechano-sensitivity of skeletal muscle. *Physiol Meas* 34: 307-314, 2013.
59. Turturici M, and Roatta S. Effects of gadolinium chloride on basal flow and compression-induced rapid hyperemia in the rabbit masseter muscle. *J Physiol Pharmacol* 65: 409-415, 2014.
60. Turturici M, and Roatta S. Inactivation of mechano-sensitive dilatation upon repetitive mechanical stimulation of the musculo-vascular network in the rabbit. *J Physiol Pharmacol* 64: 299-308, 2013.
61. Valic Z, Buckwalter JB, and Clifford PS. Muscle blood flow response to contraction: influence of venous pressure. *J Appl Physiol* 98: 72-76, 2005.
62. Welch HG, Bonde-Petersen F, Graham T, Klausen K, and Secher N. Effects of hyperoxia on leg blood flow and metabolism during exercise. *J Appl Physiol Respir Environ Exerc Physiol* 42: 385-390, 1977.

LEGENDS TO FIGURES

Fig 1

Experimental setup and typical hemodynamic response to a compressive stimulus.

A) The experimental setup includes: the IPC system for the compression of the lower limb, eco-Doppler monitoring of blood flow from femoral vein and femoral artery, and NIRS monitoring at lateral head of gastrocnemius muscle. B) Typical response to leg compression in a representative subject. From top to bottom: blood velocity in femoral vein (BVFV), blood velocity in femoral artery (BVFA), tissue oxygenation index (TOI), total hemoglobin index (THI), changes in oxygenated hemoglobin (O₂Hb) and in deoxygenated hemoglobin (HHb) and the synchronism signal (Sync.), the thick and thin bars indicating start of inflation and deflation of the cuff, respectively.

Fig 2

Hemodynamic responses to repetitive compression at different inter-stimulus intervals (ISI).

The ISI is indicated at the bottom of each column of bar- diagrams; each bar refers to the response to a single compressive stimulus. From top to bottom: Peak (arterial) blood flow, displaced (venous) blood volume, Pre-compression THI (indicating local vascular filling reached before the delivery of the compressive stimulus); Pre-compression TOI (indicating local tissue oxygenation before the stimulus). For the first three variables

642 and for each subject, responses have been normalized to the response to the first stimulus in the 160-s series
 643 (white bar). * significantly different from the first response in the series ($p < 0.05$)

644 **Fig 3**

645 **Original recordings of the response to repetitive leg compression at inter-stimulus interval = 80 s, from**
 646 **a representative subject.**

647 Notations as in Fig.1. Note the pattern of response of arterial blood velocity in relation to tissue oxygenation.
 648 The dotted line represents the initial TOI baseline.

649 **Fig 4**

650 **Original recordings of the response to repetitive leg compression at inter-stimulus interval = 20 s, from**
 651 **a representative subject.**


652 Notations as in Fig.1. Note the complete disappearance of the hyperaemic response (BVFA) after the first
 653 compressive stimulation, as long as tissue oxygenation (TOI) remains elevated, and the agreement between
 654 the displaced blood volume (area under BVFV) and the current vascular filling (THI).

655 **Fig 5**


656 **Scatter plots for assessing the correlation between different variables.** Each dot indicates the response to
 657 a single compressive stimulus in a single subject. Notations as in Fig. 2. ($n=10$). Straight lines indicate linear
 658 regressions for individual subjects. Note that: the increase in tissue oxygenation is related to the peak blood
 659 flow (A); Peak blood flow is not related to the displaced blood volume (B) but is inversely related to pre-
 660 compression oxygenation level. In D the amplitude of the hyperaemic response (peak-baseline) is plot vs
 661 pre-compression TOI to indicate that at high oxygenation levels the hyperaemic response may be almost
 662 completely abolished.


663


A)




B)












A)



B)



C)



D)

