POLITECNICO DI TORINO
Repository ISTITUZIONALE

On the periodic writing of cubic irrationals and a generalization of Rédei functions

Original
On the periodic writing of cubic irrationals and a generalization of Rédei functions / Murru, N.. - In: INTERNATIONAL JOURNAL OF NUMBER THEORY. - ISSN 1793-0421. - 11:3(2015), pp. 779-799.

Availability:
This version is available at: 11583/2723276 since: 2019-01-20T18:50:01Z

Publisher:
World Scientific Publishing

Published
DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

18 November 2026

On the periodic writing of cubic irrationals and a generalization of Rédei functions

Nadir Murru
Department of Mathematics, University of Turin
Via Carlo Alberto 10, Turin, 10123, ITALY
nadir.murru@unito.it

Abstract

In this paper, we provide a periodic representation (by means of periodic rational or integer sequences) for any cubic irrationality. In particular, for a root α of a cubic polynomal with rational coefficients, we study the Cerruti polynomials $\mu_n^{(0)}(z)$, $\mu_n^{(1)}(z)$, and $\mu_n^{(2)}(z)$, which are defined via

$$(z+\alpha)^2 = \mu_n^{(0)}(z) + \alpha\mu_n^{(1)}(z) + \alpha^2\mu_n^{(2)}(z).$$

Using these polynomials, we show how any cubic irrational can be written periodically as a ternary continued fraction. A periodic multi-dimensioanl continued fraction (with pre–period of length 2 and period of length 3) is proved convergent to a given cubic irrationality, by using the algebraic properties of cubic irrationalities and linear recurrent sequences.

1 Introduction

Continued fractions yields to a representation of any real number α by means of a sequence of integers as follows:

$$\alpha = a_0 + \cfrac{1}{a_1 + \cfrac{1}{a_2 + \cfrac{1}{a_3 + \cdots}}},$$

where $a_0, a_1, ...$ are evaluated by

$$\begin{cases} a_k = [\alpha_k] \\ \alpha_{k+1} = \dfrac{1}{\alpha_k - a_k} \quad \text{if } \alpha_k \text{ is not integer} \end{cases} \qquad k = 0, 1, 2...$$

Continued fractions are generalized by *ternary* continued fractions, which represent a pair of real numbers (α, β) by means of two sequences of integers as follows:

$$\alpha = a_0 + \cfrac{1}{b_1 + \cfrac{1}{b_3 + \cfrac{1}{a_2 + \cfrac{1}{\ddots}}}} \Bigg/ \cfrac{1}{a_3 + \cfrac{1}{\ddots}} \Bigg/ \cfrac{1}{b_2 + \cfrac{1}{a_3 + \cfrac{1}{\ddots}}} \Bigg/ \cfrac{1}{a_1 + \cfrac{1}{b_3 + \cfrac{1}{a_2 + \cfrac{1}{a_3 + \cfrac{1}{\ddots}}}}}$$

$$\text{and} \quad \beta = b_0 + \cfrac{1}{b_2 + \cfrac{1}{a_3 + \cfrac{1}{\ddots}}} \Bigg/ \cfrac{1}{a_1 + \cfrac{1}{b_3 + \cfrac{1}{a_2 + \cfrac{1}{a_3 + \cfrac{1}{\ddots}}}}} \tag{1}$$

The a_i's and b_i's can be determined by the Jacobi algorithm

$$\begin{cases} a_n = [\alpha_n] \\ b_n = [\beta_n] \\ \alpha_{n+1} = \dfrac{1}{\beta_n - [\beta_n]} \\ \beta_{n+1} = \dfrac{\alpha_n - [\alpha_n]}{\beta_n - [\beta_n]} \end{cases}, \tag{2}$$

see the VI volume of the book Ges. Werke published in 1891, [19]. In 1907, Perron [27] developed a generalization of this algorithm for higher dimensions (for a complete survey about the Jacobi–Perron algorithm see [10] and [30]). Further results on the Jacobi–Perron algorithm can be found in [28] and [36]. Many variants of the Jacobi algorithm arise by modifying the function used in Eqs. (2) for determining the a_i's and b_i's (in the classical Jacobi algorithm this function is the floor function). In this sense, many different algorithms are summarized and compared in [11] and [12]. Moreover, Tamura and Yasutomi [33], [34] recently presented a modified Jacobi–Perron algorithm. Similarly, the Jacobi–Perron algorithm has been modified using different functions, e.g., in [17], [37], [25], [32]. See the beautiful book of Schweiger [31] for a guide about multidimensional continued fractions and [21] for a new geometric vision of multidimensional continued fractions.

A very interesting approach can be found in the works [15], [3], [13],

where a multidimensional continued fraction related to triangle sequences is studied. Moreover, in [6] a generalization of the Minkowski question–mark function is developed. Finally, a completely different approach to multidimensional continued fractions can be found in [20].

In this paper, we study ternary continued fractions by means of a new family of polynomials, named Cerruti polynomials. Indeed, we construct ternary continued fractions such that their convergents are related to this kind of polynomials. In particular, we develop ternary continued fractions only for cubic irrationalities starting from the knowledge of their minimal polynomial. In this way, we will provide a periodic representation via ternary continued fractions for all cubic irrationals.

The development of multidimensional continued fractions is related to the Hermite problem. Indeed, in 1839 Hermite (see the letters published in 1850, [18]) posed to Jacobi the problem of generalizing the construction of continued fractions to higher dimensions. In particular, he asked for a method of representing algebraic irrationalities by means of periodic sequences that can highlight algebraic properties and possibly provide rational approximations. Hermite especially focused the attention on cubic irrationalities.

Continued fractions completely solve this problem for quadratic irrationalities, but the problem for algebraic numbers of degree ≥ 3 is still open. Indeed, the Jacobi–Perron algorithm, when periodic, provides sequences that converge to cubic irrationalities. However, the viceversa has never been proved. In [7], [8], [9], Bernstein studied and proved the periodicity of the Jacobi–Perron algorithm for a vast class of algebraic irrationals.

The previous mentioned modifications of the Jacobi–Perron algorithm have the same problem. None algorithm has been proved to become periodic when the input is a cubic irrational. Thus, not exists any algorithm that provides a periodic representation for a given cubic irrationality.

In this paper, the Hermite problem has been approached finding a periodic representation for an irrational number satisfying the cubic equation $x^3 - px^2 - qx - r = 0$, with p, q, r rational numbers. In other words, the problem of writing cubic irrationals as a periodic sequence of rational or integer numbers has been solved. The Hermite problem can not be claimed completely solved, since the periodic representation does not derive from an algorithm defined over all real numbers and it is derived only starting from the knowledge of the minimal polynomial of the cubic irrational. However, the results obtained are important, since for the first time we have found a periodic representation for any cubic irrational.

The periodic representation has been directly found by means of elementary techniques that only involve the algebraic properties of cubic irrationalities and the properties of linear recurrent sequences. Moreover, an algorithm, which provides a periodic sequence when a fixed cubic irrational is given in input, is derived.

In section 2, the fundamental properties of the multidimensional continued fraction (named ternary continued fraction), derived from the Jacobi algorithm, are presented. In section 3, the main case (α root largest in modulus of $x^3 - px^2 - qx - r$) is treated. A periodic expansion that converges to the pair $(\frac{r}{\alpha}, \alpha)$ is shown. In section 4, all the remaining cases of cubic irrationalities satisfying a cubic equation $x^3 - px^2 - qx - r = 0$ are treated. In this way, we can say that a real number α is a cubic irrational $\Leftrightarrow$ α can be represented by means of a periodic ternary continued fraction (that is convergent).

The iteration on cubic irrationalities of the map in section 5 provides (according to the Jacobi algorithm) the periodic expansion found in the previous sections. Section 6 is devoted to the conclusions.

2 Ternary continued fractions

The Jacobi algorithm associates a pair of integer sequences to a pair of real numbers by Eqs (2). It follows that

$$\begin{cases} \alpha_n = a_n + \dfrac{\beta_{n+1}}{\alpha_{n+1}} \\ \beta_n = b_n + \dfrac{1}{\alpha_{n+1}} \end{cases}$$

Therefore, the real numbers α and β are represented by the sequences $(a_n)_{n=0}^{\infty}$, $(b_n)_{n=0}^{\infty}$ as in (3) and we call *ternary* continued fraction (as named, e.g., in [14] and [23], where these objects have been studied independently from the generating algorithm) such a pair of objects representing the numbers α and β and we write

$$(\alpha, \beta) = [\{a_0, a_1, a_2, ...\}, \{b_0, b_1, b_2, ...\}], \tag{3}$$

where a_i's and b_i's are called *partial quotients*.

Remark 2.1. Ternary continued fraction are also called bifurcating continued fraction as in [2] and [16].

Similarly to classical continued fractions, the notion of convergent is introduced as follows (for a complete survey of the Jacobi–Perron algorithm see [10]):

$$[\{a_0, a_1, ..., a_n\}, \{b_0, b_1, ..., b_n\}] = (\frac{A_n}{C_n}, \frac{B_n}{C_n}), \quad \forall n \geq 0$$

$$\lim_{n\to\infty} \frac{A_n}{C_n} = \alpha, \quad \lim_{n\to\infty} \frac{B_n}{C_n} = \beta$$

are the *n–th convergents* of the ternary continued fraction (3), where A_n, B_n, C_n satisfy the following recurrent relations

$$\begin{cases} A_n = a_n A_{n-1} + b_n A_{n-2} + A_{n-3} \\ B_n = a_n B_{n-1} + b_n B_{n-2} + B_{n-3} \\ C_n = a_n C_{n-1} + b_n C_{n-2} + C_{n-3} \end{cases} , \quad \forall n \geq 1 \tag{4}$$

with initial conditions

$$\begin{cases} A_{-2} = 1, & A_{-1} = 0, & A_0 = a_0 \\ B_{-2} = 0, & B_{-1} = 1, & B_0 = b_0 \\ C_{-2} = 0, & C_{-1} = 0, & C_0 = 1 \end{cases}$$

Furthermore, a matricial approach is known. Indeed, it is easy to prove by induction that

$$\begin{pmatrix} a_0 & 1 & 0 \\ b_0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} ... \begin{pmatrix} a_n & 1 & 0 \\ b_n & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} = \begin{pmatrix} A_n & A_{n-1} & A_{n-2} \\ B_n & B_{n-1} & B_{n-2} \\ C_n & C_{n-1} & C_{n-2} \end{pmatrix} \tag{5}$$

$$\Updownarrow$$

$$(\frac{A_n}{C_n}, \frac{B_n}{C_n}) = [\{a_0, ..., a_n\}, \{b_0, ..., b_n\}],$$

for $n = 0, 1, 2, ...$

Remark 2.2. A ternary continued fraction (1) can converge to a pair of real numbers although the partial quotients are not obtained by the Jacobi algorithm. Thus, it is possible to study convergence of ternary continued fractions independently from the origin of the partial quotients. In sections 3 and 4, we find the partial quotients a_i's and b_i's such that the ternary continued fraction (1) converges to a given cubic irrational. In Section 5, these partial quotients are obtained by the Jacobi algorithm (2) by means of two functions f_z^α, g_z^α that substitute the role of the floor function.

In [2], the authors studied the convergence of ternary continued fractions with rational partial quotients, finding infinitely many periodic representations for every cubic root.

Theorem 2.3. *The periodic ternary continued fraction*

$$[\{z, \frac{2z}{d}, \overline{\frac{3dz}{z^3+d^2}, 3z, \frac{3z}{d}}\}, \{0, -\frac{z^2}{d}, \overline{-\frac{3z^2}{z^3+d^2}, -\frac{3dz^2}{z^3+d^2}, -\frac{3z^2}{d}}\}] \tag{6}$$

converges for every integer $z \neq 0$ to the pair of irrationals $(\sqrt[3]{d^2}, \sqrt[3]{d})$, for d integer not cube.

Remark 2.4. A ternary continued fraction with rational partial quotients is clearly determined by sequences of integer numbers. Indeed, given

$$[\{\frac{a_0}{b_0},\frac{a_1}{b_1},...\},\{\frac{c_0}{d_0},\frac{c_1}{d_1},...\}],$$

where a_i,b_i,c_i,d_i integer numbers, for $i=0,1,2,...$, then

$$\begin{pmatrix} a_0d_0 & b_0d_0 & 0 \\ c_0b_0 & 0 & b_0d_0 \\ b_0d_0 & 0 & 0 \end{pmatrix}\cdots\begin{pmatrix} a_nd_n & b_nd_n & 0 \\ c_nb_n & 0 & b_nd_n \\ b_nd_n & 0 & 0 \end{pmatrix}= \tag{7}$$

$$\begin{pmatrix} d_0s_n & d_0b_nd_ns_{n-1} & d_0b_nd_nb_{n-1}d_{n-1}s_{n-2} \\ b_0d_1s'_n & b_0b_nd_ns'_{n-1} & b_0b_nd_nb_{n-1}d_{n-1}s'_{n-2} \\ b_0d_0d_1s''_n & b_0d_0d_1b_nd_ns''_{n-1} & b_0d_0d_1b_nd_nb_{n-1}d_{n-1}s''_{n-2} \end{pmatrix}$$

$$\Updownarrow$$

$$[\{\frac{a_0}{b_0},...,\frac{a_n}{b_n}\},\{\frac{c_0}{d_0},...,\frac{c_n}{d_n}\}]=(\frac{s_n}{b_0d_1s''_n},\frac{s'_n}{d_0s''_n})=(\frac{A_n}{C_n},\frac{B_n}{C_n}),$$

where

$$\begin{cases} s_0=a_0, \quad s_1=a_0a_1d_1+b_0b_1c_1, \quad s_2=a_2d_2s_1+b_2b_1c_2d_1s_0+b_2b_1b_0d_2d_1 \\ s'_0=c_0, \quad s'_1=a_1c_0+b_1d_0, \quad s'_2=b_1b_2c_0c_2+a_1a_2c_0d_2+a_2b_1d_0d_2 \\ s''_0=1, \quad s''_1=a_1, \quad s''_2=b_1b_2c_2+a_1a_2d_2 \end{cases}$$

and

$$\begin{cases} s_n=a_nd_ns_{n-1}+b_nb_{n-1}c_nd_{n-1}s_{n-2}+b_nb_{n-1}b_{n-2}d_nd_{n-1}d_{n-2}s_{n-3} \\ s'_n=a_nd_ns'_{n-1}+b_nb_{n-1}c_nd_{n-1}s'_{n-2}+b_nb_{n-1}b_{n-2}d_nd_{n-1}d_{n-2}s'_{n-3} \\ s''_n=a_nd_ns''_{n-1}+b_nb_{n-1}c_nd_{n-1}s''_{n-2}+b_nb_{n-1}b_{n-2}d_nd_{n-1}d_{n-2}s''_{n-3} \end{cases}, \quad \forall n\geq 3.$$

For these results see [2]. Thus, a ternary continued fraction with rational partial quotients can be represented by matrices with integer entries like

$$\begin{pmatrix} a_id_i & b_id_i & 0 \\ c_ib_i & 0 & b_id_i \\ b_id_i & 0 & 0 \end{pmatrix},$$

which play the same role of the matrices used in (5).

The periodic expansion of Theorem 2.3 has been found starting from the development of

$$(z+\sqrt[3]{d^2})^n=\nu_n^{(0)}+\nu_n^{(1)}\sqrt[3]{d}+\nu_n^{(2)}\sqrt[3]{d^2}, \tag{8}$$

for every integer $z \neq 0$, d integer not cube, and where $\nu_n^{(0)}, \nu_n^{(1)}, \nu_n^{(2)}$ are polynomials such that

$$\lim_{n\to} \frac{\nu_n^{(0)}}{\nu_n^{(2)}} = \sqrt[3]{d^2}, \quad \lim_{n\to} \frac{\nu_n^{(1)}}{\nu_n^{(2)}} = \sqrt[3]{d}$$

These ratios generalize the Rédei rational functions [29]. Indeed Rédei rational functions arise from the development of

$$(z + \sqrt{d})^n = N_n(d, z) + D_n(d, z)\sqrt{d},$$

for every integer $z \neq 0$, d integer not square, and where

$$N_n(d, z) = \sum_{k=0}^{[n/2]} \binom{n}{2k} d^k z^{n-2k}, \quad D_n(d, z) = \sum_{k=0}^{[n/2]} \binom{n}{2k+1} d^k z^{n-2k-1}.$$

The Rédei rational functions are defined as

$$Q_n(d, z) = \frac{N_n(d, z)}{D_n(d, z)}, \quad \forall n \geq 1.$$

The Rédei rational functions are very interesting and useful tools in number theory. Indeed they are permutation functions of finite fields (see, e.g., [22]) and they can be also used in order to generate pseudorandom sequences [35] or to construct a public key cryptographic system [26]. Moreover in [4] and [1], Rédei rational functions are connected to periodic continued fractions with rational partial quotients convergent to square roots. In [2], Rédei rational functions have been generalized in order to obtain periodic representations only for cubic roots. In the next section, we will propose a different generalization of the Rédei rational functions in order to construct periodic ternary continued fractions convergent to any cubic irrationalities.

3 The main case

Let α be a real root of the polynomial $x^3 - px^2 - qx - r$, with $p, q, r \in \mathbb{Q}$. Let us consider

$$(z + \alpha^2)^n = \mu_n^{(0)} + \mu_n^{(1)}\alpha + \mu_n^{(2)}\alpha^2, \tag{9}$$

for z integer number not zero and where the polynomials $\mu_n^{(i)}$ depends on p, q, r, z and we will call it *Cerruti polynomials*, since when $\alpha = \sqrt[3]{d}$ they are the polynomials $\nu_n^{(i)}$ (8) introduced the first time in [2]. Let N be the following fundamental matrix

$$N = \begin{pmatrix} z & r & pr \\ 0 & q+z & pq+r \\ 1 & p & p^2+q+z \end{pmatrix}. \tag{10}$$

Its characteristic polynomial is

$$x^3 - \mathrm{Tr}(N)x^2 + \frac{1}{2}(\mathrm{Tr}(N)^2 - \mathrm{Tr}(N^2))x - \det(N),$$

i.e.,

$$x^3-(p^2+2q+3z)x^2+(q^2-2pr+2p^2z+4qz+3z^2)x-(r^2+q^2z-2prz+p^2z^2+2qz^2+z^3).$$

In the following, we set $I_1(N) = \frac{1}{2}(\mathrm{Tr}(N)^2 - \mathrm{Tr}(N^2))$.

Theorem 3.1. *Let N and $\mu_n^{(i)}$ be the matrix* (10) *and the Cerruti polynomials above defined.*

1. *The characteristic polynomial of N has roots*

$$z + \alpha_1^2, \quad z + \alpha_2^2, \quad z + \alpha_3^2,$$

where $\alpha_1, \alpha_2, \alpha_3$ are the roots of $x^3 - px^2 - qx - r$.

2.
$$N^n = \begin{pmatrix} \mu_n^{(0)} & r\mu_n^{(2)} & r\mu_n^{(1)} + pr\mu_n^{(2)} \\ \mu_n^{(1)} & \mu_n^{(0)} + q\mu_n^{(2)} & (pq+r)\mu_n^{(2)} + q\mu_n^{(1)} \\ \mu_n^{(2)} & \mu_n^{(1)} + p\mu_n^{(2)} & \mu_n^{(0)} + p\mu_n^{(1)} + (p^2+q)\mu_n^{(2)} \end{pmatrix}$$

Proof. 1. Considering that

$$\alpha_1\alpha_2\alpha_3 = r, \quad \alpha_1 + \alpha_2 + \alpha_3 = p, \quad \alpha_1\alpha_2 + \alpha_2\alpha_3 + \alpha_1\alpha_3 = -q,$$

we have

$$\alpha_1^2 + \alpha_2^2 + \alpha_3^2 = p^2 + 2q, \quad \alpha_1^2\alpha_2^2 + \alpha_2^2\alpha_3^2 + \alpha_1^2\alpha_3^2 = q^2 - 2pr.$$

Moreover, expanding the polynomial $(x - (z + \alpha_1^2))(x - (z + \alpha_2^2))(x - (z + \alpha_3^2))$, it is easy to see that the coefficient of x^2, the coefficient of x, and the constant term are

$$-(\alpha_1^2 + \alpha_2^2 + \alpha_3^2 + 3z) = -\mathrm{Tr}(N)$$

$$\alpha_1^2\alpha_2^2 + \alpha_2^2\alpha_3^2 + \alpha_1^2\alpha_3^2 + 2z(\alpha_1^2 + \alpha_2^2 + \alpha_3^2) + 3z^2 = I_1(N)$$

$$-\alpha_1^2\alpha_2^2\alpha_3^2 - z(\alpha_1^2\alpha_2^2 + \alpha_2^2\alpha_3^2 + \alpha_1^2\alpha_3^2) - z^2(\alpha_1^2 + \alpha_2^2 + \alpha_3^2) - z^3 = -\det(N),$$

respectively.

2. By definition of Cerruti polynomials (9), it follows that $\mu_n^{(i)}$'s, for $i = 0, 1, 2$, are linear recurrent sequences of degree 3 whose characteristic polynomial is the minimal polynomial of $z + \alpha^2$ (where α real root of $x^3 - px^2 - qx - r$), i.e., the characteristic polynomial of N.
Thus, we only have to check the initial conditions. We start from

$$(z + \alpha^2)^0 = 1,$$

i.e.

$$\mu_0^{(0)} = 1, \quad \mu_0^{(1)} = 0, \quad \mu_0^{(2)} = 0$$

and since

$$N^0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

the initial condition for $n = 0$ is satisfied. Considering $z+\alpha^2$, it follows that

$$\mu_1^{(0)} = z, \quad \mu_1^{(1)} = 0, \quad \mu_1^{(2)} = 1.$$

Thus,

$$N = \begin{pmatrix} z & r & pr \\ 0 & q+z & pq+r \\ 1 & p & p^2+q+z \end{pmatrix} = \begin{pmatrix} \mu_1^{(0)} & r\mu_1^{(2)} & r\mu_1^{(1)} + pr\mu_1^{(2)} \\ \mu_1^{(1)} & \mu_1^{(0)} + q\mu_1^{(2)} & (pq+r)\mu_1^{(2)} + q\mu_1^{(1)} \\ \mu_1^{(2)} & \mu_1^{(1)} + p\mu_1^{(2)} & \mu_1^{(0)} + p\mu_1^{(1)} + (p^2+q)\mu_1^{(2)} \end{pmatrix}.$$

Finally,

$$(z + \alpha^2)^2 = z^2 + 2\alpha^2 z + \alpha^4 = z^2 + pr + (pq + r)\alpha + (p^2 + q + 2z)\alpha^2$$

and

$$N^2 = \begin{pmatrix} pr + z^2 & p^2r + qr + 2rz & p^3r + 2pqr + r^2 + 2prz \\ pq + r & p^2q + q^2 + pr + 2qz + z^2 & p^3q + 2pq^2 + p^2r + 2qr + 2pqz + 2rz \\ p^2 + q + 2z & p^3 + 2pq + r + 2pz & p^4 + 3p^2q + q^2 + 2pr + 2p^2z + 2qz + z^2 \end{pmatrix}.$$

□

Theorem 3.2. *Let α be a real root largest in modulus of $x^3 - px^2 - qx - r$ and let α_2, α_3 be the remaining roots. Let $\mu_n^{(i)}$ be the Cerruti polynomials (9), then is*

$$\lim_{n\to\infty} \frac{\mu_n^{(0)}}{\mu_n^{(2)}} = \frac{r}{\alpha}, \quad \lim_{n\to\infty} \frac{\mu_n^{(1)}}{\mu_n^{(2)}} = \alpha - p,$$

for any integer z such that $z + \alpha^2$ larger in modulus than $z + \alpha_2^2, z + \alpha_3^2$ and $\mu_n^{(2)} \neq 0$.

Proof. Let

$$\beta_1 = z + \alpha^2, \quad \beta_2 = z + \alpha_2^2, \quad \beta_3 = z + \alpha_3^2$$

be the roots of the characteristic polynomial of N. By the Binet formula

$$\begin{cases} \mu_n^{(0)} = a_1\beta_1^n + a_2\beta_2^n + a_3\beta_3^n \\ \mu_n^{(1)} = b_1\beta_1^n + b_2\beta_2^n + b_3\beta_3^n \\ \mu_n^{(2)} = c_1\beta_1^n + c_2\beta_2^n + c_3\beta_3^n \end{cases}, \quad \forall n \geq 0$$

where the coefficients a_i, b_i, c_i can be obtained by initial conditions, solving the system

$$\begin{cases} a_1 + a_2 + a_3 = 1 \\ a_1\beta_1 + a_2\beta_2 + a_3\beta_3 = z, \\ a_1\beta_1^2 + a_2\beta_2^2 + a_3\beta_3^2 = pr + z^2 \end{cases}$$

and similar systems for the b_i's and c_i's. Since β_1 is larger in modulus than β_2, β_3, we are only interested in

$$\begin{cases} a_1 = \dfrac{\beta_2\beta_3 - z(\beta_2 + \beta_3) + pr + z^2}{(\beta_1 - \beta_2)(\beta_1 - \beta_3)} \\ b_1 = \dfrac{pq + r}{(\beta_1 - \beta_2)(\beta_1 - \beta_3)} \\ c_1 = \dfrac{2z + p^2 + q - (\beta_2 + \beta_3)}{(\beta_1 - \beta_2)(\beta_1 - \beta_3)} \end{cases}.$$

Now,

$$\lim_{n\to\infty} \frac{\mu_n^{(1)}}{\mu_n^{(2)}} = \frac{b_1}{c_1} = \frac{pq + r}{2z + p^2 + q - (2z + \alpha_2^2 + \alpha_3^2)} = \frac{pq + r}{\alpha^2 - q},$$

moreover

$$(\alpha^2 - q)(\alpha - p) = \alpha^3 - p\alpha^2 - q\alpha + pq = pq + r$$

and it is proved that

$$\lim_{n\to\infty} \frac{\mu_n^{(1)}}{\mu_n^{(2)}} = \alpha - p.$$

The proof that $\lim_{n\to\infty} \dfrac{\mu_n^{(0)}}{\mu_n^{(2)}} = \dfrac{r}{\alpha}$ is left to the reader. □

Remark 3.3. It is always possible to find integers z satisfying the condition of Theorem 3.2 and such that $\mu_n^{(2)} \neq 0$.

Theorem 3.4. *Let α be a real root largest in modulus of $x^3 - px^2 - qx - r$ and N the matrix defined in* (10)*, then*

$$[\{z, \frac{2z+p^2+q}{pq+r}, \overline{\frac{(pq+r)\,Tr(N)}{\det(N)}, Tr(N), \frac{Tr(N)}{pq+r}}\},$$
$$\{p, -\frac{z^2+qz+p^2z-pr}{pq+r}, \overline{-\frac{I_1(N)}{\det(N)}, -\frac{(pq+r)I_1(N)}{\det(N)}, -\frac{I_1(N)}{pq+r}}\}] = (\frac{r}{\alpha}, \alpha) \tag{11}$$

for any integer z satisfying the hypothesis of Theorem 3.2.

Proof. By Theorem 3.2, it is sufficient to prove that

$$\frac{A_n}{C_n} = \frac{\mu^{(0)}_{n+1}}{\mu^{(2)}_{n+1}}, \quad \frac{B_n}{C_n} = \frac{\mu^{(1)}_{n+1}}{\mu^{(2)}_{n+1}} + p, \quad \forall n \geq 0,$$

where $\frac{A_n}{C_n}, \frac{B_n}{C_n}$ are the convergents of the ternary continued fraction (11) satisfying Eqs. (4). First of all we prove by induction that

$$A_n = \frac{\mu^{(0)}_{n+1}}{(pq+r)^k \det(N)^{[\frac{n+1}{3}]}}, \quad \forall n \geq 0,$$

where

$$k = \begin{cases} 1 & n \equiv 1 \pmod 3 \\ 0 & \text{otherwise} \end{cases}$$

The inductive basis is straightforward to prove, indeed

$$A_0 = z, \quad A_1 = \frac{pr+z^2}{pq+r}, \quad A_2 = \frac{p^3r + 2pqr + r^2 + 3prz + z^3}{r^2 + q^2z - 2prz + p^2z^2 + 2qz^2 + z^3}.$$

Now, for $n \geq 3$, we consider the cases

$$n \equiv 0 \pmod 3, \quad n \equiv 1 \pmod 3, \quad n \equiv 2 \pmod 3.$$

Let us consider $n \equiv 0 \pmod 3$, then

$$A_n = \mathrm{Tr}(N)A_{n-1} - \frac{(pq+r)I_1(N)}{\det(N)}A_{n-2} + A_{n-3},$$

by inductive hypothesis we have

$$A_n = \mathrm{Tr}\frac{\mu^{(0)}_n}{\det(N)^{[\frac{n}{3}]}} - \frac{(pq+r)I_1(N)}{\det(N)} \cdot \frac{\mu^{(0)}_{n-1}}{(pq+r)\det(N)^{[\frac{n-1}{3}]}} + \frac{\mu^{(0)}_{n-2}}{\det(N)^{[\frac{n-2}{3}]}}.$$

Since $n \equiv 0 \pmod 3$, we have

$$[\frac{n}{3}] = [\frac{n+1}{3}], \quad [\frac{n-1}{3}] = [\frac{n-2}{3}] = [\frac{n+1}{3}] - 1.$$

Using the recurrence relation for the Cerruti polynomials, we obtain

$$A_n = \frac{\mathrm{Tr}(N)\mu_n^{(0)} - I_1(N)\mu_{n-1}^{(0)} + \det(N)\mu_{n-2}^{(0)}}{\det(N)^{[\frac{n+1}{3}]}} = \frac{\mu_{n+1}^{(0)}}{(pq+r)^k \det(N)^{[\frac{n+1}{3}]}}.$$

Let us consider $n \equiv 1 \pmod 3$, then

$$A_n = \frac{\mathrm{Tr(N)}}{pq+r} A_{n-1} - \frac{I_1(N)}{pq+r} A_{n-2} + A_{n-3}$$

and

$$[\frac{n+1}{3}] = [\frac{n}{3}] = [\frac{n-1}{3}] = [\frac{n-2}{3}] + 1.$$

Thus, we easily obtain

$$A_n = \frac{\mathrm{Tr}(N)\mu_n^{(0)} - I_1(N)\mu_{n-1}^{(0)} + \det(N)\mu_{n-2}^{(0)}}{(pq+r)\det(N)^{[\frac{n+1}{3}]}} = \frac{\mu_{n+1}^{(0)}}{(pq+r)^k \det(N)^{[\frac{n+1}{3}]}}.$$

Similarly when $n \equiv 2 \pmod 3$. In a similar way, it is possible to prove that

$$C_n = \frac{\mu_{n+1}^{(2)}}{(pq+r)^k \det(N)^{[\frac{n+1}{3}]}}, \quad \forall n \geq 0.$$

Consequently,

$$\lim_{n\to\infty} \frac{A_n}{C_n} = \lim_{n\to\infty} \frac{\mu_{n+1}^{(0)}}{\mu_{n+1}^{(2)}} = \frac{r}{\alpha}.$$

Finally, let us consider the sequence $(B_n)_{n=0}^{\infty}$. We prove by induction that

$$B_n = \frac{\mu_{n+1}^{(1)} + p\mu_{n+1}^{(2)}}{(pq+r)^k \det(N)^{[\frac{n+1}{3}]}}, \quad \forall n \geq 0.$$

The steps $n = 0, 1, 2$ can be directly checked. Let us consider $n \geq 3$ and $n \equiv 2 \pmod 3$ (similarly the formula can be proved when $n \equiv 0 \pmod 3$ and $n \equiv 1 \pmod 3$). We have

$$B_n = \frac{(pq+r)\mathrm{Tr}(N)}{\det(N)} B_{n-1} - \frac{I_1(N)}{\det(N)} B_{n-2} + B_{n-3} =$$

$$= \frac{(pq+r)\mathrm{Tr}(N)}{\det(N)} \cdot \frac{\mu_n^{(1)} + p\mu_n^{(2)}}{(pq+r)\det(N)^{\frac{n}{3}}} - \frac{I_1(N)}{\det(N)} \cdot \frac{\mu_{n-1}^{(1)} + p\mu_{n-1}^{(2)}}{(pq+r)\det(N)^{\frac{n-1}{3}}} + \frac{\mu_{n-2}^{(1)} + p\mu_{n-2}^{(2)}}{(pq+r)\det(N)^{\frac{n}{3}}}.$$

Since

$$[\frac{n+1}{3}] = [\frac{n}{3}] + 1 = [\frac{n-1}{3}] + 1 = [\frac{n-2}{3}] + 1$$

the formula is proved. Thus,

$$\lim_{n\to\infty} \frac{B_n}{C_n} = \lim_{n\to\infty} \frac{\mu_{n+1}^{(1)}}{\mu_{n+1}^{(2)}} + p = \alpha.$$

□

The previous theorem provides a periodic representation for all the cubic irrationalities α, such that α is the root greatest in modulus of its minimal polynomial. This representation is a ternary continued fraction (1) of period 3 whose partial quotients are given by (11). Clearly, this expansion leads to periodic sequences of integer numbers that represent cubic irrationals. This fact is highlighted by using the matricial approach. In particular, the irrationalities $(\frac{r}{\alpha}, \alpha)$ are represented by a periodic product of matrices (5) whose entries are rational numbers, or equivalently by a periodic product of matrices (7) whose entries are integer numbers.

Example 3.5. Let us consider the cubic polynomial $x^3 - 5x^2 + x - 3$, having a real root

$$\alpha = \frac{1}{3}(5 + \sqrt[3]{44} + \sqrt[3]{242}) \simeq 4.9207,$$

greater in modulus than the complex roots. Theorem 3.4 provides periodic representations of α. If we choose, e.g., $z = 5$ (which satisfies conditions of Theorem 3.2), we have

$$N = \begin{pmatrix} 5 & 3 & 15 \\ 0 & 4 & -2 \\ 1 & 5 & 29 \end{pmatrix}$$

and

$$(\frac{3}{\alpha}, \alpha) = [\{5, -17, \overline{-\frac{19}{141}, 38, -19}\}, \{5, 65, \overline{-\frac{23}{47}, -\frac{46}{47}, 138}\}].$$

Moreover, we can use (7) in order to express the cubic irrationality α as a periodic product of matrices with integer entries

$$\begin{pmatrix} 5 & 1 & 0 \\ 5 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} -17 & 1 & 0 \\ 65 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \overline{\begin{pmatrix} -893 & 6627 & 0 \\ -3243 & 0 & 6627 \\ 6627 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1786 & 47 & 0 \\ 46 & 0 & 47 \\ 47 & 0 & 0 \end{pmatrix} \begin{pmatrix} -19 & 1 & 0 \\ 138 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}}.$$

This periodic representation provides rational approximations for α. For example, the convergents of the ternary continued fraction can be evaluated by using (4) and we obtain rational approximations of $\frac{3}{\alpha}$ and α, respectively:

$$(5, \frac{20}{17}, \frac{88}{127}, \frac{4633}{7447}, \frac{66559}{108838}, ...) = (5, 1.1764, 0.6929, 0.6221, 0.6115, ...)$$

$$(5, \frac{84}{17}, \frac{1251}{254}, \frac{36651}{7447}, \frac{535575}{108838}) = (5, 4.9412, 4.9252, 4.9216, 4.9208, ...)$$

These rational approximations can be obviously obtained by the matricial representation. For example if we set A for the matrix of the pre–period (i.e., the matrix product of the two matrices of the pre–period) and P for the matrix of the period (i.e., the matrix product of the three matrices of the period), then

$$AP = \begin{pmatrix} -147028831 & 10234297 & 388784 \\ -1183085175 & 80962059 & 2763459 \\ -240423142 & 16450423 & 561086 \end{pmatrix}$$

and

$$\frac{-147028831}{-240423142} = \frac{66559}{108838}, \quad \frac{-1183085175}{-240423142} = \frac{535575}{108838}.$$

Example 3.6. Let us consider the cubic polynomial $3x^3 - 12x^2 - 4x + 1$. We can apply Theorem 3.4 with $p = 4, q = 4/3, r = -1/3$. Using $z = 1$, we obtain

$$(-\frac{1}{3\alpha}, \alpha) = [\{1, \frac{58}{15}, \overline{\frac{975}{218}, \frac{65}{3}, \frac{13}{3}}\}, \{4, -\frac{59}{15}, \overline{-\frac{403}{218}, -\frac{2015}{218}, -\frac{403}{45}}\}].$$

that is a periodic representation of α root greatest in modulus of $3x^3 - 12x^2 - 4x + 1$.
If we choose, e.g., $z = -1$, we obtain a different periodic ternary continued fraction convergent to α:

$$(-\frac{1}{3\alpha}, \alpha) = [\{-1, \frac{46}{15}, \overline{\frac{47}{8}, \frac{47}{3}, \frac{47}{15}}\}, \{4, 3, \overline{-\frac{269}{120}, \frac{269}{24}, \frac{269}{45}}\}],$$

where $\alpha \simeq 4.29253$. These periodic representations leads to a periodic product of matrices with integer entries convergent to the irrational α.

4 The remaining cases

When α is the root largest in modulus of a general cubic polynomial $x^3 - px^2 - qx - r$, a periodic representation for α is provided by Theorem 3.4. In this section, we treat all the remaining cases of cubic irrationalities.

If α is the root smallest in modulus of $x^3 - px^2 - qx - r$, Theorem 3.4 does not work. In this case, we consider the reflected polynomial $x^3 + \frac{q}{r}x^2 + \frac{p}{r}x - \frac{1}{r}$ whose roots are $\frac{1}{\alpha}, \frac{1}{\alpha_2}, \frac{1}{\alpha_3}$. In this way $\frac{1}{\alpha}$ is the root greatest in modulus of $x^3 + \frac{q}{r}x^2 + \frac{p}{r}x - \frac{1}{r}$ and by Theorem 3.4 we get a periodic representation for

the pair $(\frac{\alpha}{r}, \frac{1}{\alpha})$. Successively, a periodic representation for α can be derived. We need the following

Theorem 4.1. *Let* $[\{a_0, a_1, \overline{a_2, a_3, a_4}\}, \{b_0, b_1, \overline{b_2, b_3, b_4}\}]$ *be a periodic ternary continued fraction that converges to a pair of real number* (α, β)*, then the periodic ternary continued fraction*

$$[\{ra_0, ra_1, \overline{\frac{a_2}{r^2}, ra_3, ra_4}\}, \{\frac{b_0}{r}, r^2 b_1, \overline{\frac{b_2}{r}, \frac{b_3}{r}, r^2 b_4}\}] \tag{12}$$

converges to the pair of real number $(r\alpha, \frac{1}{r\beta})$*, for* r *rational number.*

Proof. Let $\frac{A_n}{C_n}$ and $\frac{B_n}{C_n}$ be the n–th convergents of the ternary continued fraction of (α, β). Let $\frac{\tilde{A}_n}{\tilde{C}_n}$ and $\frac{\tilde{B}_n}{\tilde{C}_n}$ be the n–th convergents of the ternary continued fractions (12), then

$$\tilde{A}_n = r^{k_1} A_n, \quad \tilde{B}_n = r^{k_2} B_n, \quad \tilde{C}_n = r^{k_3} C_n, \quad \forall n \geq 0$$

where

$$k_1 = \begin{cases} 0, & n \equiv 2 \pmod 3 \\ 1, & n \equiv 0 \pmod 3 \\ 2, & n \equiv 1 \pmod 3 \end{cases}, \quad k_2 = \begin{cases} -1, & n \equiv 2 \pmod 3 \\ 0, & n \equiv 0 \pmod 3 \\ 1, & n \equiv 1 \pmod 3 \end{cases}$$

$$k_3 = \begin{cases} -2, & n \equiv 2 \pmod 3 \\ -1, & n \equiv 0 \pmod 3 \\ 0, & n \equiv 1 \pmod 3 \end{cases}$$

It is straightforward to check these identities for $n = 0, 1, 2$. Let us proceed by induction, considering an integer $m \equiv 0 \pmod 3$. Then

$$\tilde{A}_m = ra_3 \tilde{A}_{m-1} + \frac{b_3}{r} \tilde{A}_{m-2} + \tilde{A}_{m-3} = ra_3 A_{m-1} + b_3 r A_{m-2} + r A_{m-3} = r A_m.$$

Similarly when $m \equiv 1 \pmod 3$ and $m \equiv 2 \pmod 3$, and for the sequences $\tilde{B}_n$ and $\tilde{C}_n$. □

Thus, if α is the root smallest in modulus of $x^3 - px^2 - qx - r$, by Theorem 3.4 we get the periodic ternary continued fraction of $(\frac{\alpha}{r}, \frac{1}{\alpha})$ and by Theorem 4.1 we get the periodic ternary continued fraction of $(\alpha, \frac{1}{r\alpha})$.

Example 4.2. Let us consider the cubic polynomial $x^3 - 2x^2 + x + 1$. It has one real root α whose modulus is smaller than the modulus of the complex

roots. Theorem 3.4 does not work on this polynomial, but we can consider the reflected polynomial x^3+x^2-2x+1 whose roots are the inverse roots of x^3-2x^2+x+1. Thus, $\frac{1}{\alpha}$ is the real root of x^3+x^2-2x+1 largest in modulus and we can apply Theorem 3.4. Posing, e.g., $z=5$, we obtain

$$(-\alpha,\frac{1}{\alpha})=[\{5,-\frac{13}{3},\overline{-\frac{20}{87},20,-\frac{20}{3}}\},\{-1,13,\overline{-\frac{127}{261},\frac{127}{87},\frac{127}{3}}\}].$$

Finally, we multiply by -1 this ternary continued fraction and by Theorem 4.1 we obtain

$$(\alpha,-\frac{1}{\alpha})=[\{-5,\frac{13}{3},\overline{-\frac{20}{87},-20,\frac{20}{3}}\},\{1,13,\overline{\frac{127}{261},-\frac{127}{87},\frac{127}{3}}\}],$$

i.e., we found a periodic representation for α root smallest in modulus of x^3-2x^2+x+1.

Now, we are able to determine a periodic representation for any cubic irrational that is the root largest or smallest in modulus of a cubic polynomial x^3-px^2-qx-r.

Finally, we treat the last case, i.e., α is the intermediate root of a cubic polynomial having three real roots. Let $\alpha_1,\alpha_2,\alpha_3$ be the real root of x^3-px^2-qx-r such that $|\alpha_3|<|\alpha_2|<|\alpha_1|$. Using previous techniques we can get periodic expansions for α_1 and α_3. Moreover, a rational number k can be ever found such that $\alpha_2\pm k$ is the root largest or smallest in modulus of its minimal polynomial $(x-(\alpha_1\pm k))(x-(\alpha_2\pm k))(x-(\alpha_3\pm k))$. The coefficients of $(x-(\alpha_1\pm k))(x-(\alpha_2\pm k))(x-(\alpha_3\pm k))$ can be derived from the coefficients of x^3-px^2-qx-r (see, e.g., Th. 10 and Cor. 11 [5]). Thus, by Theorem 3.4 we know the periodic ternary continued fraction of $(\frac{r'}{\alpha_2\pm k},\alpha_2\pm k)$. Then it is immediate to obtain the periodic expansion of $(\frac{r'}{\alpha_2\pm k},\alpha_2)$ (see (1)).

Example 4.3. Let us consider the Ramanujan cubic polynomial x^3+x^2-2x-1 with three real roots. The roots of this polynomial are quite famous (see, e.g., [38]) and it is well–known that they are

$$\alpha_1=2\cos\frac{2\pi}{7},\quad \alpha_2=2\cos\frac{4\pi}{7},\quad \alpha_3=2\cos\frac{8\pi}{7},$$

where

$$\alpha_1\simeq 1.24698,\quad \alpha_2\simeq -0.445042,\quad \alpha_3\simeq -1.80194.$$

Thus, from Theorem 3.4, for $z=3$, we obtain

$$(\frac{1}{\alpha_3},\alpha_3)=[\{3,-9,\overline{-\frac{2}{13},14,-14}\},\{-1,19,\overline{-\frac{9}{13},\frac{9}{13},63}\}].$$

Considering the polynomial x^3+2x^2-x-1 and $z=1$, we obtain

$$(\alpha_2, \frac{1}{\alpha_2}) = [\{1, -7, -\overline{\frac{9}{13}}, 9, -9\}, \{-2, 8, -\overline{\frac{20}{13}, \frac{20}{13}}, 20\}].$$

Finally, we can consider the minimal polynomial of

$$\alpha_1+1, \quad \alpha_2+1, \quad \alpha_3+1$$

that is x^3-2x^2-x+1, whose root largest in modulus is α_1+1. For $z=2$, by Theorem 3.4, we obtain

$$(-\frac{1}{\alpha_1+1}, \alpha_1+1) = [\{2, 9, \overline{\frac{12}{43}, 12, 12}\}, \{2, -16, -\overline{\frac{41}{43}, -\frac{41}{43}, -41}\}]$$

and

$$(-\frac{1}{\alpha_1+1}, \alpha_1) = [\{2, 9, \overline{\frac{12}{43}, 12, 12}\}, \{1, -16, -\overline{\frac{41}{43}, -\frac{41}{43}, -41}\}].$$

5 The periodic algorithm

An approach to the Hermite problem contemplates the research of a function whose iteration on algebraic irrationalities provides a periodical algorithm. The partial quotients of the ternary continued fraction (11) can be derived from the Jacobi algorithm (2) using two functions f_z^α, g_z^α instead of the floor function.

Definition 1. Let α and $\mathbb{Q}(\alpha)$ be a root of x^3-px^2-qx-r and the algebraic extension of $\mathbb{Q}$, respectively. We define the linear functions $f_z^\alpha, g_z^\alpha : \mathbb{Q}(\alpha) \to \mathbb{Q}$, for $z \in \mathbb{Z}$, such that

1. $f_z^\alpha(q) = g_z^\alpha(q) = q, \quad \forall q \in \mathbb{Q}$
2. $f_z^\alpha\left(\frac{r}{\alpha}\right) = g_z^\alpha\left(\frac{r}{\alpha}\right) = z$
3. $f_z^\alpha(\alpha) = g_z^\alpha(\alpha) = p$
4. $f_z^\alpha(\alpha^2) = 2z+p^2+2q, \quad g_z^\alpha(\alpha^2) = z+p^2+q$

The ternary continued fraction (11) is obtained from the following algorithm:

$$\begin{cases} a_n = f_z^\alpha(x_n) \\ b_n = g_z^\alpha(y_n) \\ x_{n+1} = \dfrac{1}{y_n - b_n} \\ y_{n+1} = \dfrac{x_n - a_n}{y_n - b_n} \end{cases}, \tag{13}$$

for $n = 0, 1, 2, ...$ and $x_0 = \dfrac{r}{\alpha}$, $y_0 = \alpha$, where α root of the polynomial $x^3 - px^2 - qx - r$.

Let us start to use the algorithm (13) with inputs $(\dfrac{r}{\alpha}, \alpha)$, we immediately have

$$a_0 = z, \quad b_0 = p.$$

Now, we evaluate x_1 and y_1:

$$x_1 = \frac{1}{\alpha - p}, \quad y_1 = \frac{\dfrac{r}{\alpha} - z}{\alpha - p}.$$

We need to manipulate x_1 and y_1 in order to find the values of $f_z^\alpha(x_1)$ and $g_z^\alpha(y_1)$. In particular, we will often use that $\alpha^3 = p\alpha^2 + q\alpha + r$. We have

$$x_1 = \frac{1}{\alpha - p} \cdot \frac{\alpha^2 - q}{\alpha^2 - q} = \frac{\alpha^2 - q}{pq + r}$$

$$y_1 = \frac{\dfrac{r}{\alpha} - z}{\alpha - p} \cdot \frac{\alpha^2 - q}{\alpha^2 - q} = \frac{r\alpha - z\alpha^2 + qz - \dfrac{qr}{\alpha}}{pq + r}.$$

Now we can apply the properties of f_z^α and g_z^α and we obtain

$$a_1 = f_z^\alpha(x_1) = \frac{2z + p^2 + q}{pq + r}, \quad b_1 = g_z^\alpha(y_1) = -\frac{z^2 + qz + p^2 z - pr}{pq + r}.$$

Let us continue with

$$x_2 = \frac{1}{y_1 - b_1} = \frac{pq + r}{(\dfrac{r}{\alpha} - z)(\alpha^2 - q) + (z^2 + qz + p^2 z - pr)(\alpha^2 - q)} \cdot \frac{\alpha^2 + z}{\alpha^2 + z} = \frac{(pq + r)(\alpha^2 + z)}{\det(N)},$$

where the last identities follow by using $\alpha^3 = p\alpha^2 + q\alpha + r$. Moreover,

$$y_2 = (x_1 - a_1) \cdot x_2 = \frac{(\alpha^2 - q)(\alpha^2 + z) - (2z + p^2 + q)(\alpha^2 + z)}{\det(N)}$$

and

$$a_2 = \frac{(pq + r)(3z + p^2 + 2q)}{\det(N)} = \frac{(pq + r)\mathrm{Tr}(N)}{\det(N)}, \quad b_2 = -\frac{I_1(N)}{\det(N)}.$$

Then,

$$x_3 = \frac{\det(N)}{(\alpha^2 - q)(\alpha^2 + z) - (2z + p^2 + q)(\alpha^2 + z) + I_1(N)} \cdot \frac{\alpha^2 + z}{\alpha^2 + z} = \frac{\det(N)(\alpha^2 + z)}{\det(N)} = \alpha^2 + z,$$

$$y_3 = \frac{(pq+r)((\alpha^2+z^2)^2 - \mathrm{Tr}(N)(\alpha^2+z))}{\det(N)}$$

and

$$a_3 = 3z + 2q + p^2 = \mathrm{Tr}(N), \quad b_3 = -\frac{(pq+r)I_1(N)}{\det(N)}.$$

Finally,

$$x_4 = \frac{\det(N)}{(pq+r)((\alpha^2+z)^2 - \mathrm{Tr}(N)(\alpha^2+z) + I_1(N))} \cdot \frac{\alpha^2+z}{\alpha^2+z} = \frac{\alpha^2+z}{pq+r},$$

where the last identity is obtained recalling that $\alpha^2 + z$ is the root of the characteristic polynomial of N.

$$y_4 = (\alpha^2 + z - \mathrm{Tr(N)}) \cdot \frac{\alpha^2+z}{pq+r}$$

from which

$$a_4 = \frac{\mathrm{Tr}(N)}{pq+r}, \quad b_4 = -\frac{I_1(N)}{pq+r}.$$

Now we check that $x_5 = x_2$ and $y_5 = y_2$:

$$x_5 = \frac{pq+r}{(\alpha^2+z)^2 - \mathrm{Tr}(N)(\alpha^2+z) + I_1(N)} = \frac{(pq+r)(\alpha^2+z)}{\det(N)} = x_2$$

$$y_5 = \frac{\alpha^2 + z - \mathrm{Tr}(N)}{pq+r} \cdot \frac{(pq+r)(\alpha^2+z)}{\det(N)} = y_2.$$

6 Conclusions

In this paper, given a fixed cubic irrational we have found a periodic representation via ternary continued fraction with rational partial quotients. A periodic representation involving integer numbers can be directly derived from it. A family of algorithms based on Eqs. (13) determines this periodic representation. These algorithms become periodic when the input is a pair $(\frac{r}{\alpha}, \alpha)$, for any cubic irrationality α, whose minimal polynomial is known. The ternary continued fraction (11) is very manageable, since the pre–period has length 2, the period has length 3, and it provides simultaneous rational approximations. Of course, many questions and further developments remain open:

- Generalization of Theorem 3.4 to any algebraic irrationality is the natural development of the present work.

- The problem of the periodicity of the original Jacobi algorithm is still open. It would be possible to use the present work in order to solve this question. Indeed, the sequence of numerator and denominator of the convergents of (11) are linear recurrent sequences. Thus, it could be possible to prove that a ternary continued fraction provided by the Jacobi algorithm and equal to (11) (i.e. converging to the same pair of cubic irrationalities) has numerators and denominators of the convergents that are linear recurrent sequences. Moreover, it could be possible to generalize to ternary continued fraction the result of Lenstra and Shallit [24], which proved that a continued fraction is periodic if and only if numerators (or denominators) of the convergents are linear recurrent sequences.

- The convergence's rate of the ternary continued fraction (11) has not been studied in the present paper and it will be developed in future works, studying the role of the parameter z.

- Cerruti polynomials (9) appear to be very interesting and they could be applied in different fields of number theory. Indeed, they are a generalization of the Rédei rational functions. Since Rédei rational functions are very useful in several fields of number theory, Cerruti polynomials could have many different applications.

7 Acknowledgment

I would like to thank the anonymous referee for the valuable suggestions in the improvement of the presentation of this paper.
Thanks to Marianna, Roberta, Simone, and Vanni.

References

[1] M. Abrate, S. Barbero, U. Cerruti, N. Murru, *Periodic representations and rational approximations of square roots*, Journal of Approximation Theory, Vol. **175**, 83–90, (2013).

[2] M. Abrate, S. Barbero, U. Cerruti, N. Murru, *Periodic representations for cubic irrationalities*, The Fibonacci Quarterly, Vol. **50**, Issue **3**, 252–264, (2012).

[3] S. Assaf, L. C. Chen, T. Cheslack–Postava, B. Cooper, A. Diesl, T. Garrity, M. Lepinski, A. Schuyler, *A dual approach to triangle sequences: a multidimensional continued fraction algorithm*, Integers: Electronic Journal of Combinatorial Number Theory, **5**, A08, (2005).

[4] S. Barbero, U. Cerruti, N. Murru, *Solving the Pell equation via Rédei rational functions*, The Fibonacci Quarterly, Vol. **48**, 348–357, (2010).

[5] S. Barbero, U. Cerruti, N. Murru, *Transforming recurrent sequences by using the Binomial and Invert operators*, Journal of Integer Sequences, Vol. **13**, Article **10.7.7**, (2010).

[6] O. R. Beaver, T. Garrity, *A two–dimensional Minkowsky ?(x) function*, Journal of Number Theory, **107**, 105–134, (2004).

[7] L. Bernstein, *Periodical continued fractions for irrationals n by Jacobi's algorithm*, J. Reine Angew. Math., **213**, 31–38, (1964).

[8] L. Bernstein, *Periodicity of Jacobi's algorithm for a special case of cubic irrationals*, J. Reine Angew. Math., **213**, 137–147, (1964).

[9] L. Bernstein, *New infinite classes of periodic Jacobi–Perron algorithms*, Pacific Journal of Mathematics, Vol. **16**, No. **3**, 439–469, (1965).

[10] L. Bernstein, *The Jacobi–Perron algorithm – its theory and application*, Lectures Notes in Mathematics, Vol. **207**, (1971).

[11] A. J. Brentjes, *Multi–dimensional continued fraction algorithms*, Mathematical Centre Tracts, Amsterdam, (1981).

[12] A. D. Bryuno, V. I. Parusnikov, *Comparison of various generalizations of continued fractions*, Mathematical notes, Vol. **61**, No. **3**, 278–286, (1997).

[13] K. Dasaratha, L. Flapan, T. Garrity, C. Lee, C. Mihaila, N. Neumann-Chun, S. Peluse M. Stoffregen, *Cubic irrationals and periodicity via a family of multi–dimensional continued fraction algorithms*, Available online at: http://www.arxiv.org/abs/1208.4244v2, (2013).

[14] P. H. Daus, *Normal ternary continued fraction expansions for the cube roots of integers*, American Journal of Mathematics, Vol. **44**, No. **4**, 279–296, (1922).

[15] T. Garrity, *On periodic sequences for algebraic numbers*, Journal of Number Theory, Vol. **88**, 86–103, (2001).

[16] A. Gupta, A. Mittal, *Bifurcating continued fractions*, Available online at: http://front.math.ucdavis.edu/math.GM/0002227, (2000).

[17] M.D. Hendy, N.S. Jeans, *The Jacobi–Perron algorithm in integer form*, Mathematics of Computation, Vol. **36**, No. **154**, 565–574, (1981).

[18] C. Hermite, *Extraits de lettres de M. Ch. Hermite a M. Jacobi sur differents objets de la theorie des nombres*, J. Reine Angew. Math. **40**, 286, (1850).

[19] C. G. J. Jacobi, Ges. Werke, Vol. **VI**, 385–426, Berlin Academy, (1891).

[20] O. N. Karpenkov, *Constructing multidimensional periodic continued fractions in the sense of Klein*, Mathematics of Computation, Vol. **78**, No. **267**, 1687–1711, (2009).

[21] O. N. Karpenkov, *Geometry of continued fractions*, Series Algorithms and Computation in Mathematics, Vol. **26**, Springer-Verlag GmbH, Heidelberg, Germany, 2013.

[22] R. Lidl, G. L. Mullen, G. Turnwald, *Dickson polynomials*, Pitman Monogr. Surveys Pure appl. Math. **65**, Longman, (1993).

[23] D. N. Lehmer, *On Jacobi's extension of the continued fraction algorithm*, National Academy of Sciences, Vol. **4**, No. **12**, 360–364, (1918).

[24] H. W. Lenstra, J. O. Shallit, *Continued fractions and linear recurrences*, Mathematics of Computation, Vol. **61**, No. **203**, 351–354, (1993).

[25] G. Martin, *The unreasonable effectualness of continued function expansions*, J. Aust. Math. Soc., Vol. **77**, 305–319, (2004).

[26] R. Nobauer, *Cryptanalysis of the Rédei scheme*, Contributions to General Algebra, **3**, 255–264, (1984).

[27] O. Perron, *Grundlagen fur eine theorie des Jacobischen kettenbruchalgorithmus*, Math. Ann. **64**, 1–76, (1907).

[28] N. S. Raju, *Periodic Jacobi–Perron algorithms and fundamental units*, Pacific Journal of Mathematics, Vol. **64**, No. **1**, 241–251, (1976).

[29] L. Rédei, *Uber eindeuting umkehrbare polynome in endlichen korpen*, Acta Sci. Math. (Szeged), **11**, 85–92, (1946).

[30] F. Schweiger, *The metrical theory of Jacobi–Perron algorithm*, Lectures Notes in Mathematics, Vol. **334**, Springer–Verlag, Berlin, (1973).

[31] F. Schweiger, *Multidimensional continued fractions*, Oxford: Oxford University Press, (2000).

[32] F. Schweiger, *Brun meets Selmer*, Integers, 13, A17, (2013).

[33] J. Tamura, S. Yasutomi, *A new multidimensional continued fraction algorithm*, Mathematics of Computation, **78.268**, 2209–2222, (2009).

[34] J. Tamura, S. Yasutomi, *A new algorithm of continued fractions related to algebraic number fields of degree* ≤ 5, Integers, 11B, A16, (2011).

[35] A. Topuzoglu, A. Winterhof, *Topics in Geometry, Coding Theory and Cryptography*, Algebra and Applications, Vol. **6**, 135–166, (2006).

[36] T. P. Vaughan, *A note on the Jacobi–Perron algorithm*, Pacific Journal of Mathematics, Vol. **72**, No. **1**, 261–271, (1977)

[37] H. C. Williams, G. W. Dueck, *An analogue of the nearest integer continued fraction for certain cubic irrationalities*, Mathematics of computation, Vol. **42**, No. **166**, 683–705, (1984).

[38] R. Witula and D. Slota, *New Ramanujan – Type Formulas and Quasi – Fibonacci Numbers of Order 7*, Journal of Integer Sequences, Vol. **10**, Article **07.5.6**, (2007).