

On the number of distinct exponents in the prime factorization of an integer

*Original*

On the number of distinct exponents in the prime factorization of an integer / Sanna, Carlo. - In: PROCEEDINGS OF THE INDIAN ACADEMY OF SCIENCES. MATHEMATICAL SCIENCES. - ISSN 0253-4142. - STAMPA. - 130:1(2020). [10.1007/s12044-020-0556-y]

*Availability:*

This version is available at: 11583/2802793 since: 2020-05-03T10:42:27Z

*Publisher:*

Springer

*Published*

DOI:10.1007/s12044-020-0556-y

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)



# On the number of distinct exponents in the prime factorization of an integer

CARLO SANNA 

Department of Mathematics, Università degli Studi di Genova, Genoa, Italy  
E-mail: carlo.sanna.dev@gmail.com

MS received 25 February 2019; revised 5 September 2019; accepted 11 November 2019

**Abstract.** Let  $f(n)$  be the number of distinct exponents in the prime factorization of the natural number  $n$ . We prove some results about the distribution of  $f(n)$ . In particular, for any positive integer  $k$ , we obtain that

$$\#\{n \leq x : f(n) = k\} \sim A_k x$$

and

$$\#\{n \leq x : f(n) = \omega(n) - k\} \sim \frac{Bx(\log \log x)^k}{k! \log x},$$

as  $x \rightarrow +\infty$ , where  $\omega(n)$  is the number of prime factors of  $n$  and  $A_k, B > 0$  are some explicit constants. The latter asymptotic extends a result of Aktas and Ram Murty (*Proc. Indian Acad. Sci. (Math. Sci.)* **127**(3) (2017) 423–430) about numbers having mutually distinct exponents in their prime factorization.

**Keywords.** Prime factorization; squarefree numbers; powerful number.

**2010 Mathematics Subject Classification.** Primary: 11N25; Secondary: 11N37, 11N64.

## 1. Introduction

Let  $n = p_1^{a_1} \cdots p_s^{a_s}$  be the factorization of the natural number  $n > 1$ , where  $p_1 < \cdots < p_s$  are prime numbers and  $a_1, \dots, a_s$  are positive integers. Several functions of the exponents  $a_1, \dots, a_s$  have been studied, including their product [17], their arithmetic mean [2, 4, 5, 7], and their maximum and minimum [11, 13, 15, 18]. See also [3, 8] for a more general function.

Let  $f$  be the arithmetic function defined by  $f(1) := 0$  and  $f(n) := \#\{a_1, \dots, a_s\}$  for all natural numbers  $n > 1$ . In other words,  $f(n)$  is the number of distinct exponents in the prime factorization of  $n$ . The first values of  $f(n)$  are listed in sequence A071625 of OEIS [16].

Our first contribution is a quite precise result about the distribution of  $f(n)$ .

**Theorem 1.1.** *There exists a sequence of positive real numbers  $(A_k)_{k \geq 1}$  such that, given any arithmetic function  $\phi$  satisfying  $|\phi(k)| < a^k$  for some fixed  $a > 1$ , we have that the series*

$$M_\phi := \sum_{k=1}^{\infty} A_k \phi(k) \quad (1)$$

converges and

$$\sum_{n \leq x} \phi(f(n)) = M_\phi x + O_{a,\varepsilon}(x^{1/2+\varepsilon}),$$

for all  $x \geq 1$  and  $\varepsilon > 0$ .

From Theorem 1.1, it follows immediately that all the moments of  $f$  are finite and that  $f$  has a limiting distribution. In particular, we highlight the following corollary.

#### COROLLARY 1.1

For each positive integer  $k$ , we have

$$\#\{n \leq x : f(n) = k\} = A_k x + O_\varepsilon(x^{1/2+\varepsilon}),$$

for all  $x \geq 1$  and  $\varepsilon > 0$ .

We also provide a formula for  $A_k$ . Before stating it, we need to introduce some notations. Let  $\psi$  be the Dedekind function defined by

$$\psi(n) := n \prod_{p|n} \left(1 + \frac{1}{p}\right)$$

for each positive integer  $n$ , and let  $(\rho_k)_{k \geq 1}$  be the family of arithmetic functions supported on squarefree numbers and satisfying

$$\rho_1(n) = \begin{cases} 1 & \text{if } n = 1, \\ 0 & \text{if } n > 1, \end{cases} \quad \rho_{k+1}(n) = \begin{cases} 0 & \text{if } n = 1, \\ \frac{1}{n-1} \sum_{\substack{d|n \\ d < n}} \rho_k(d) & \text{if } n > 1, \end{cases}$$

for all squarefree numbers  $n$  and positive integers  $k$ .

**Theorem 1.2.** *We have*

$$A_k = \frac{6}{\pi^2} \sum_{n=1}^{\infty} \frac{\rho_k(n)}{\psi(n)}$$

for each positive integer  $k$ .

Clearly,  $f(n) \leq \omega(n)$  for all positive integers  $n$ , where  $\omega(n)$  denotes the number of prime factors of  $n$ . Motivated by a question of Recamán Santos [14], Aktaş and Ram Murty

[1] studied the natural numbers  $n$  such that all the exponents in their prime factorization are distinct, that is,  $f(n) = \omega(n)$ . They called such numbers *special numbers* (sequence A130091 of OEIS [16]) and they proved the following.

**Theorem 1.3.** *The number of special numbers not exceeding  $x$  is*

$$\frac{Bx}{\log x} + O\left(\frac{x}{(\log x)^2}\right),$$

for all  $x \geq 2$ , where

$$B := \sum_{\ell} \frac{1}{\ell}$$

and the sum of over natural numbers  $\ell$  that are powerful and special.

Let  $g$  be the arithmetic function defined by  $g(n) := \omega(n) - f(n)$  for all positive integers  $n$ . Hence, by the previous observation,  $g$  is a nonnegative function and  $g(n) = 0$  if and only if  $n$  is a special number. We prove the following result about  $g$ , which extends Theorem 1.3 and it is somehow dual to Corollary 1.1.

**Theorem 1.4.** *For each nonnegative integer  $k$ , we have*

$$\#\{n \leq x : g(n) = k\} = \frac{Bx(\log \log x)^k}{k! \log x} \left(1 + O_k\left(\frac{1}{\log \log x}\right)\right),$$

for all  $x \geq 3$ .

*Notation.* We employ the Landau–Bachmann “Big Oh” notation  $O$ , as well as the associated Vinogradov symbol  $\ll$ , with their usual meaning. Any dependence of the implied constants is explicitly stated. We let  $\varepsilon$  denote an arbitrary small positive real number, not necessarily the same at each occurrence. We reserve the letter  $p$  for prime numbers.

## 2. Preliminaries

Recall that a natural number  $n$  is called *powerful* if  $p \mid n$  implies  $p^2 \mid n$ , for all primes  $p$ . For all  $x \geq 1$ , let  $\mathcal{P}(x)$  be the set of powerful numbers not exceeding  $x$ .

*Lemma 2.1.* *We have  $\#\mathcal{P}(x) \ll x^{1/2}$  for every  $x \geq 1$ .*

*Proof.* See [9]. □

*Lemma 2.2.* *We have*

$$\sum_{\substack{\ell \in \mathcal{P} \\ \ell > y}} \frac{1}{\ell} \ll \frac{1}{y^{1/2}}, \quad \sum_{\ell \in \mathcal{P}(y)} \frac{1}{\ell^{1/2}} \ll \log y,$$

for all  $y \geq 2$ .

*Proof.* By Lemma 2.1 and by partial summation, we have

$$\sum_{\substack{\ell \in \mathcal{P} \\ \ell > y}} \frac{1}{\ell} = \frac{\#\mathcal{P}(t)}{t} \Big|_{t=y}^{+\infty} + \int_y^{+\infty} \frac{\#\mathcal{P}(t)}{t^2} dt \ll \int_y^{+\infty} \frac{dt}{t^{1+1/2}} \ll \frac{1}{y^{1/2}}.$$

The proof of the second claim is similar.  $\square$

We need the following upper bound for the number of prime factors of a natural number.

*Lemma 2.3.* We have

$$\omega(n) \ll \frac{\log n}{\log \log n}$$

for all integers  $n \geq 3$ .

*Proof.* See, for example, [6, Proposition 7.10].  $\square$

For every  $x \geq 1$  and every positive integer  $h$ , let  $Q(x; h)$  denote the number of squarefree numbers not exceeding  $x$  and relatively prime with  $h$ .

*Lemma 2.4.* We have

$$Q(x; h) = \frac{6}{\pi^2} \frac{h}{\psi(h)} x + O(4^{\omega(h)}(x^{1/2} + 1))$$

for all  $x \geq 1$  and all positive integers  $h$ .

*Proof.* It follows easily from [10, Eq. 8].  $\square$

For every  $x \geq 1$  and every positive integers  $s, h$ , let  $Q_s(x; h)$  denote the number of squarefree numbers not exceeding  $x$ , having exactly  $s$  prime factors, and relatively prime with  $h$ .

*Lemma 2.5.* We have

$$Q_s(x; h) = \frac{x(\log \log x)^{s-1}}{(s-1)! \log x} \left( 1 + O_{\delta, s} \left( \frac{\log \log \log(h+15)}{\log \log x} \right) \right)$$

for all  $x \geq 3$ ,  $0 < \delta < 1$ , and for all integers  $1 \leq h \leq x^\delta$  and  $s \geq 1$ .

*Proof.* For  $s = 1$ , the claim follows from the Prime Number theorem, while for  $h = 1$ , the claim is a classic result of Landau [12]. Hence, suppose  $s, h > 1$ . Also, we can assume that  $x \geq 3^{1/(1-\delta)}$ . If  $n \leq x$  is a squarefree number having exactly  $s$  prime factors such that  $(n, h) > 1$ , then  $n = pn'$ , where  $p$  is a prime number dividing  $h$  and  $n' \leq x/p$  is a squarefree number having exactly  $s - 1$  prime factor. Therefore,

$$\begin{aligned}
 0 \leq Q_s(x; 1) - Q_s(x; h) &\leq \sum_{p|h} Q_{s-1}\left(\frac{x}{p}, 1\right) \ll_s \sum_{p|h} \frac{x}{p} \frac{(\log \log(x/p))^{s-2}}{\log(x/p)} \\
 &\ll_\delta \frac{x(\log \log x)^{s-2}}{\log x} \sum_{p|h} \frac{1}{p} \ll \frac{x(\log \log x)^{s-1} \log \log \log(h+15)}{\log x \log \log x},
 \end{aligned}$$

where we used the fact that  $p \leq x^\delta$  and the upper bound

$$\sum_{p|h} \frac{1}{p} \leq \sum_{p \leq \omega(h)} \frac{1}{p} \ll \log \log(\omega(h) + 2) \ll \log \log \log(h + 15),$$

which in turn follows from Mertens' second theorem [6, Theorem 4.5] and the simple bound  $\omega(h) \ll \log h$ . Consequently,

$$\begin{aligned}
 Q_s(x; h) &= Q_s(x; 1) + O_{\delta,s} \left( \frac{x(\log \log x)^{s-1} \log \log \log(h+15)}{\log x \log \log x} \right) \\
 &= \frac{x(\log \log x)^{s-1}}{(s-1)! \log x} + O_{\delta,s} \left( \frac{x(\log \log x)^{s-1} \log \log \log(h+15)}{\log x \log \log x} \right),
 \end{aligned}$$

as claimed. □

Finally, we need a lemma about certain sums of powers.

*Lemma 2.6.* *Let  $a_0$  be an integer. For all  $x_1, \dots, x_k > 1$ , we have*

$$\sum_{a_0 < a_1 < \dots < a_k} \frac{1}{x_1^{a_1} \dots x_k^{a_k}} = \frac{1}{(x_1 \dots x_k)^{a_0}} \prod_{j=1}^k \frac{1}{x_j \dots x_k - 1},$$

where the sum is over all integers  $a_1, \dots, a_k$  satisfying  $a_0 < a_1 < \dots < a_k$ .

*Proof.* We proceed by induction on  $k$ . For  $k = 1$ , we have

$$\sum_{a_0 < a_1} \frac{1}{x_1^{a_1}} = \frac{1}{x_1^{a_0+1}} \sum_{d=0}^{\infty} \frac{1}{x_1^d} = \frac{1}{x_1^{a_0}} \frac{1}{x_1 - 1}, \tag{2}$$

as claimed. Suppose that the claim is true for  $k$ , we shall prove it for  $k + 1$ . We have

$$\begin{aligned}
 \sum_{a_0 < \dots < a_{k+1}} \frac{1}{x_1^{a_1} \dots x_{k+1}^{a_{k+1}}} &= \sum_{a_0 < \dots < a_k} \frac{1}{x_1^{a_1} \dots x_k^{a_k}} \sum_{a_k < a_{k+1}} \frac{1}{x_{k+1}^{a_{k+1}}} \\
 &= \sum_{a_0 < \dots < a_{k+1}} \frac{1}{x_1^{a_1} \dots x_{k-1}^{a_{k-1}} (x_k x_{k+1})^{a_k}} \frac{1}{x_{k+1} - 1}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{(x_1 \cdots x_{k+1})^{a_0}} \prod_{j=1}^k \frac{1}{x_j \cdots x_{k+1} - 1} \frac{1}{x_{k+1} - 1} \\
 &= \frac{1}{(x_1 \cdots x_{k+1})^{a_0}} \prod_{j=1}^{k+1} \frac{1}{x_j \cdots x_{k+1} - 1},
 \end{aligned}$$

where we used (2), with  $a_0$  and  $x_1$  replaced respectively by  $a_k$  and  $x_{k+1}$ , and the induction hypothesis. □

### 3. Proof of Theorem 1.1

We begin by proving that for each positive integer  $k$ , there exists  $A_k > 0$  such that

$$N_k(x) := \#\{n \leq x : f(n) = k\} = A_k x + O_\varepsilon(x^{1/2+\varepsilon}), \tag{3}$$

for all  $x \geq 1$  and  $\varepsilon > 0$ . Clearly, every natural number  $n$  can be written in a unique way as  $n = m\ell$ , where  $m$  is a squarefree number,  $\ell$  is a powerful number, and  $(m, \ell) = 1$ . If  $m = 1$ , then  $n = \ell$  is powerful and, by Lemma 2.1, belongs to a set of cardinality  $O(x^{1/2})$ . If  $m > 1$ , then  $f(n) = k$  is equivalent to  $f(\ell) = k - 1$ . Also, for each  $\ell$ , there are exactly  $Q(x/\ell; \ell) - 1$  choices for  $m > 1$ . Therefore, we have

$$N_k(x) = \sum_{\substack{\ell \in \mathcal{P}(x) \\ f(\ell) = k-1}} \left( Q\left(\frac{x}{\ell}; \ell\right) - 1 \right) + O(x^{1/2}), \tag{4}$$

for all  $x \geq 1$ . For each positive integer  $\ell \leq x$ , Lemma 2.3 gives  $4^{\omega(\ell)} \ll_\varepsilon x^\varepsilon$ . Consequently, by Lemma 2.4, we obtain

$$Q\left(\frac{x}{\ell}; \ell\right) = \frac{6}{\pi^2} \frac{x}{\psi(\ell)} + O_\varepsilon\left(\frac{x^{1/2+\varepsilon}}{\ell^{1/2}}\right), \tag{5}$$

for all positive integers  $\ell \leq x$ . By Lemma 2.2, we have

$$\sum_{\substack{\ell \in \mathcal{P} \\ \ell > x}} \frac{1}{\psi(\ell)} < \sum_{\substack{\ell \in \mathcal{P} \\ \ell > x}} \frac{1}{\ell} \ll \frac{1}{x^{1/2}}, \tag{6}$$

for all  $x \geq 1$ . In particular, the series

$$A_k := \frac{6}{\pi^2} \sum_{\substack{\ell \in \mathcal{P} \\ f(\ell) = k-1}} \frac{1}{\psi(\ell)} \tag{7}$$

converges. Also, again by Lemma 2.2, we have

$$\sum_{\ell \in \mathcal{P}(x)} \frac{1}{\ell^{1/2}} \ll \log x \ll_\varepsilon x^\varepsilon. \tag{8}$$

At this point, putting together (4) and (5), and using (6) and (8), we obtain

$$\begin{aligned}
 N_k(x) &= \sum_{\substack{\ell \in \mathcal{P}(x) \\ f(\ell) = k-1}} \left( \frac{6}{\pi^2} \frac{x}{\psi(\ell)} + O_\varepsilon \left( \frac{x^{1/2+\varepsilon}}{\ell^{1/2}} \right) \right) + O(x^{1/2}) \\
 &= A_k x + O \left( \sum_{\substack{\ell \in \mathcal{P} \\ \ell > x}} \frac{x}{\psi(\ell)} \right) + O_\varepsilon \left( \sum_{\ell \in \mathcal{P}(x)} \frac{x^{1/2+\varepsilon}}{\ell^{1/2}} \right) + O(x^{1/2}) \\
 &= A_k x + O_\varepsilon(x^{1/2+\varepsilon}),
 \end{aligned}$$

as desired. Thus (3) is proved.

Now we shall show that

$$A_k \leq \frac{6}{\pi^2} \frac{1}{(k-1)!} \tag{9}$$

for all positive integers  $k$ . For  $k = 1$ , the claim is obvious since  $A_1 = 6/\pi^2$ . Hence, assume  $k \geq 2$ . If  $\ell$  is a powerful number such that  $f(\ell) = k - 1$ , then  $\ell = m_1^{a_1} \cdots m_{k-1}^{a_{k-1}}$  for some integers  $m_1, \dots, m_{k-1} \geq 2$  and  $2 \leq a_1 < \cdots < a_{k-1}$ . Consequently,

$$\begin{aligned}
 \frac{\pi^2}{6} A_k &= \sum_{\substack{\ell \in \mathcal{P} \\ f(\ell) = k-1}} \frac{1}{\psi(\ell)} < \sum_{\substack{\ell \in \mathcal{P} \\ f(\ell) = k-1}} \frac{1}{\ell} < \prod_{j=1}^{k-1} \sum_{m=2}^{\infty} \sum_{a=j+1}^{\infty} \frac{1}{m^a} \\
 &= \prod_{j=1}^{k-1} \sum_{m=2}^{\infty} \frac{1}{m^j(m-1)} \leq \prod_{j=1}^{k-1} \frac{1}{j} = \frac{1}{(k-1)!},
 \end{aligned}$$

where we used the facts that

$$\sum_{m=2}^{\infty} \frac{1}{m(m-1)} = \sum_{m=2}^{\infty} \left( \frac{1}{m-1} - \frac{1}{m} \right) = 1$$

and

$$\begin{aligned}
 \sum_{m=2}^{\infty} \frac{1}{m^j(m-1)} &< \frac{1}{2^j} + \frac{1}{3^j \cdot 2} + \sum_{n=3}^{\infty} \frac{1}{n^{j+1}} \\
 &< \frac{1}{2^j} + \frac{1}{3^j \cdot 2} + \int_2^{+\infty} \frac{dt}{t^{j+1}} = \frac{1}{2^j} + \frac{1}{3^j \cdot 2} + \frac{1}{j2^j} < \frac{1}{j},
 \end{aligned}$$

for all integers  $j \geq 2$ . Thus (9) is proved.

Now let  $\phi$  be an arithmetic function satisfying  $|\phi(k)| < a^k$  for all positive integers  $k$ , where  $a > 1$  is some constant. From (9) it follows that series (1) converges. Define

$$y := 2a + \lfloor C \log x / \log \log(x + 2) \rfloor,$$

where  $C > 0$  is some absolute constant. Since  $f(n) \leq \omega(n)$  for all positive integers  $n$ , by Lemma 2.3, we can choose  $C$  sufficiently large so that  $f(n) \leq y$  for all natural numbers  $n \leq x$ . Moreover, from (9) and  $y \geq 2a$ , we get that

$$\sum_{k > y} A_k \phi(k) \ll \sum_{k > y} \frac{a^k}{(k-1)!} < \frac{a^{y+1}}{y!} \sum_{j=0}^{\infty} \left( \frac{a}{y} \right)^j \ll_a \frac{a^y}{y!} \ll_a \frac{1}{x^{1/2}} \tag{10}$$

and

$$a^y y \ll_{a,\varepsilon} x^\varepsilon, \tag{11}$$

for all  $x \geq 1$ . Therefore, putting together (3), (10) and (11), we have

$$\begin{aligned} \sum_{n \leq x} \phi(f(n)) &= \sum_{k \leq y} N_k(x) \phi(k) = \sum_{k \leq y} (A_k \phi(k)x + O_\varepsilon(\phi(k)x^{1/2+\varepsilon})) \\ &= M_\phi x + O\left(\sum_{k > y} A_k \phi(k)x\right) + O_\varepsilon(a^y y x^{1/2+\varepsilon}) \\ &= M_\phi x + O_{a,\varepsilon}(x^{1/2+\varepsilon}), \end{aligned}$$

for all  $x \geq 1$  and  $\varepsilon > 0$ . The proof is complete.

#### 4. Proof of Theorem 1.2

Recall that  $A_k$  is defined by (7). For  $k = 1$ , the claim is obvious, since  $f(\ell) = 0$  if and only if  $\ell = 1$ . Hence, assume  $k \geq 2$ . If  $\ell$  is a powerful number such that  $f(\ell) = k - 1$ , then  $\ell$  can be written in a unique way as  $\ell = m_1^{a_1} \cdots m_{k-1}^{a_{k-1}}$ , where  $1 < a_1 < \cdots < a_{k-1}$  are integers and  $m_1, \dots, m_{k-1} > 1$  are pairwise coprime squarefree numbers. Therefore, from (7) and Lemma 2.6, we obtain

$$\begin{aligned} \frac{\pi^2}{6} A_k &= \sum_{m_1, \dots, m_{k-1}} \sum_{1 < a_1 < \dots < a_{k-1}} \frac{1}{\psi(m_1^{a_1} \cdots m_{k-1}^{a_{k-1}})} \\ &= \sum_{m_1, \dots, m_{k-1}} \frac{m_1 \cdots m_{k-1}}{\psi(m_1 \cdots m_{k-1})} \sum_{1 < a_1 < \dots < a_{k-1}} \frac{1}{m_1^{a_1} \cdots m_{k-1}^{a_{k-1}}} \\ &= \sum_{m_1, \dots, m_{k-1}} \frac{1}{\psi(m_1 \cdots m_{k-1})} \prod_{j=1}^{k-1} \frac{1}{m_j \cdots m_{k-1} - 1}, \end{aligned}$$

where, here and in the rest of the proof, in summation subscripts  $m_1, \dots, m_{k-1}$  are meant to be pairwise coprime, squarefree and greater than 1. At this point, it is enough to prove that

$$\sum_{n = m_1 \cdots m_{k-1}} \prod_{j=1}^{k-1} \frac{1}{m_j \cdots m_{k-1} - 1} = \rho_k(n)$$

for all squarefree numbers  $n > 1$ . We proceed by induction on  $k$ . For  $k = 2$ , the claim is true since

$$\frac{1}{n-1} = \frac{\rho_1(1)}{n-1} = \frac{1}{n-1} \sum_{\substack{d|n \\ d < n}} \rho_1(d) = \rho_2(n),$$

for all squarefree numbers  $n > 1$ . Assuming that the claim is true for  $k$ , we shall prove it for  $k + 1$ . We have

$$\begin{aligned} \sum_{n=m_1 \cdots m_k} \prod_{j=1}^k \frac{1}{m_j \cdots m_k - 1} &= \frac{1}{n-1} \sum_{m_1 | n} \sum_{n/m_1 = m_2 \cdots m_k} \prod_{j=2}^k \frac{1}{m_j \cdots m_k - 1} \\ &= \frac{1}{n-1} \sum_{m_1 | n} \rho_k(n/m_1) \\ &= \frac{1}{n-1} \sum_{\substack{d | n \\ d < n}} \rho_k(d) = \rho_{k+1}(n), \end{aligned}$$

for all squarefree numbers  $n > 1$ , as desired. The proof is complete.

### 5. Proof of Theorem 1.4

We have to count the number of positive integers  $n \leq x$  such that  $g(n) = k$ . As in the proof of Theorem 1.1, every  $n$  can be written in a unique way as  $n = m\ell$ , where  $m$  is a squarefree number,  $\ell$  is a powerful number, and  $(m, \ell) = 1$ . If  $m = 1$ , then  $n = \ell$  is powerful and by Lemma 2.1, belongs to a set of cardinality  $O(x^{1/2})$ . If  $m > 1$ , then

$$\omega(m) = \omega(n) - \omega(\ell) = g(n) + f(n) - f(\ell) - g(\ell) = k + 1 - g(\ell).$$

In particular,  $1 \leq \omega(m) \leq k + 1$ . Assume  $x$  sufficiently large, and put  $y := (\log x)^2$ . Then, by Lemma 2.2, the number of  $n \leq x$  such that  $\ell > y$  is at most

$$\sum_{\substack{\ell \in \mathcal{P} \\ \ell > y}} \frac{x}{\ell} \ll \frac{x}{y^{1/2}} = \frac{x}{\log x}.$$

Therefore,

$$M_k(x) := \#\{n \leq x : g(n) = k\} = \sum_{s=1}^{k+1} \sum_{\substack{\ell \in \mathcal{P}(y) \\ g(\ell) = k+1-s}} Q_s\left(\frac{x}{\ell}; \ell\right) + O\left(\frac{x}{\log x}\right). \tag{12}$$

For each nonnegative integer  $r$ , put

$$B_r := \sum_{\substack{\ell \in \mathcal{P} \\ g(\ell) = r}} \frac{1}{\ell}.$$

Note that, in light of Lemma 2.2, the series defining  $B_r$  converges and, more precisely,

$$\sum_{\substack{\ell \in \mathcal{P}(y) \\ g(\ell) = r}} \frac{1}{\ell} = B_r + O\left(\frac{1}{y^{1/2}}\right) = B_r + O\left(\frac{1}{\log x}\right). \tag{13}$$

Clearly, we can assume  $x$  sufficiently large so that  $x/y \geq 3$  and  $y \leq x^{\delta/(1+\delta)}$ , for some fixed  $0 < \delta < 1$ . Hence, applying Lemma 2.5, we obtain

$$Q_s\left(\frac{x}{\ell}; \ell\right) = \frac{x(\log \log(x/\ell))^{s-1}}{\ell(s-1)! \log(x/\ell)} \left(1 + O_k\left(\frac{\log \log \log(\ell + 15)}{\log \log(x/\ell)}\right)\right)$$

$$\begin{aligned}
&= \frac{x(\log \log x)^{s-1}}{\ell(s-1)! \log x} \left(1 + O_k \left(\frac{\log \ell}{\log x}\right)\right) \left(1 + O_k \left(\frac{\log \log \log(\ell + 15)}{\log \log x}\right)\right) \\
&= \frac{x(\log \log x)^{s-1}}{\ell(s-1)! \log x} \left(1 + O_k \left(\frac{\log(\ell + 1)}{\log \log x}\right)\right),
\end{aligned}$$

for all positive integers  $s \leq k + 1$  and  $\ell \leq y$ . Consequently,

$$\begin{aligned}
&\sum_{\substack{\ell \in \mathcal{P}(y) \\ g(\ell) = k+1-s}} Q_s \left(\frac{x}{\ell}; \ell\right) \\
&= \frac{x(\log \log x)^{s-1}}{(s-1)! \log x} \sum_{\substack{\ell \in \mathcal{P}(y) \\ g(\ell) = k+1-s}} \frac{1}{\ell} \left(1 + O_k \left(\frac{\log(\ell + 1)}{\log \log x}\right)\right) \\
&= \frac{x(\log \log x)^{s-1}}{(s-1)! \log x} \left(B_{k+1-s} + O\left(\frac{1}{\log x}\right) + O_k \left(\frac{1}{\log \log x}\right)\right) \\
&= \frac{x(\log \log x)^{s-1}}{(s-1)! \log x} \left(B_{k+1-s} + O_k \left(\frac{1}{\log \log x}\right)\right), \tag{14}
\end{aligned}$$

where we used (13) and the fact that the series

$$\sum_{\ell \in \mathcal{P}} \frac{\log(\ell + 1)}{\ell}$$

converges. Thus, putting together (12) and (14), and noting that  $B_0 = B$ , we obtain

$$M_k(x) = \frac{Bx(\log \log x)^k}{k! \log x} \left(1 + O_k \left(\frac{1}{\log \log x}\right)\right),$$

as desired. The proof is complete.

## Acknowledgements

The author is thankful to the anonymous referee for carefully reading the paper and providing useful suggestions. The author is a member of the INdAM group GNSAGA and, during the preparation of this work, was supported by a postdoctoral fellowship of INdAM.

## References

- [1] Aktaş K and Ram Murty M, On the number of special numbers, *Proc. Indian Acad. Sci. (Math. Sci.)* **127(3)** (2017) 423–430
- [2] Cao H Z, On the average of exponents, *Northeast. Math. J.* **10(3)** (1994) 291–296
- [3] Cao H Z, Functions involving the number of prime factors of a natural number, *Acta Math. Sinica (Chin. Ser.)* **39(5)** (1996) 602–608
- [4] De Koninck J-M, Sums of quotients of additive functions, *Proc. Amer. Math. Soc.* **44** (1974) 35–38
- [5] De Koninck J-M and Ivić A, Sums of reciprocals of certain additive functions, *Manuscripta Math.* **30(4)** (1979/80) 329–341
- [6] De Koninck J-M and Luca F, Analytic number theory, Graduate Studies in Mathematics, vol. 134, American Mathematical Society, Providence, RI (2012) Exploring the anatomy of integers

- [7] Duncan R L, On the factorization of integers, *Proc. Amer. Math. Soc.* **25** (1970) 191–192
- [8] Duncan R L, Some applications of the Turán–Kubilius inequality, *Proc. Amer. Math. Soc.* **30** (1971) 69–72
- [9] Golomb S W, Powerful numbers, *Amer. Math. Monthly* **77** (1970) 848–855
- [10] Hazlewood D G, On  $k$ -free integers with small prime factors, *Proc. Amer. Math. Soc.* **52** (1975) 40–44
- [11] Kátai I and Subbarao M V, On the maximal and minimal exponent of the prime power divisors of integers, *Publ. Math. Debrecen* **68(3–4)** (2006) 477–488
- [12] Landau E, Sur quelques problèmes relatifs à la distribution des nombres premiers, *Bull. Soc. Math. France* **28** (1900) 25–38
- [13] Niven I, Averages of exponents in factoring integers, *Proc. Amer. Math. Soc.* **22** (1969) 356–360
- [14] Recamán Santos B, Consecutive numbers with mutually distinct exponents in their canonical prime factorization, <http://mathoverflow.net/questions/201489>
- [15] Sinha K, Average orders of certain arithmetical functions, *J. Ramanujan Math. Soc.* **21(3)** (2006) 267–277
- [16] Sloane N J A, The On-Line Encyclopedia of Integer Sequences, <http://oeis.org>
- [17] Suryanarayana D and Sitaramachandra Rao R, The number of square-full divisors of an integer, *Proc. Amer. Math. Soc.* **34** (1972) 79–80
- [18] Suryanarayana D and Sitaramachandra Rao R, On the maximum and minimum exponents in factoring integers, *Arch. Math. (Basel)* **28(3)** (1977) 261–269

COMMUNICATING EDITOR: Sanoli Gun