

Entropy and logarithm of Kaniadakis calculus expressed by means of an Euler infinite product expansion

Original

Entropy and logarithm of Kaniadakis calculus expressed by means of an Euler infinite product expansion / Sparavigna, A.C.. - ELETTRONICO. - (2020). [10.5281/zenodo.4088214]

Availability:

This version is available at: 11583/2848421 since: 2020-10-14T15:18:02Z

Publisher:

Published

DOI:10.5281/zenodo.4088214

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Entropy and logarithm of Kaniadakis calculus expressed by means of an Euler infinite product expansion

Amelia Carolina Sparavigna

Department of Applied Science and Technology, Politecnico di Torino, Torino, Italy

Here we show how an Euler infinite product expansion can be used to display easily the link of nonadditive Kaniadakis entropy to Shannon entropy. At the same time we can see how the natural logarithm is linked to the expression of κ -logarithm.

Keywords: Nonadditive entropies, κ -calculus

Torino, 14 October 2020, DOI: 10.5281/zenodo.4088214

In 2001, Giorgio Kaniadakis proposed a formulation of entropy based on his new approach to calculation which is today known as κ -calculus [1-3]. This entropy is a nonadditive composable one [4]. In [5], we have considered some further properties of this entropy, with the aim of applying it to the analysis of images [6] and theory of numbers [7], among the many applications of this calculus. Here we aim to show how an Euler infinite product expansion can be used to display easily the link of this entropy to Shannon entropy. At the same time we can see how the natural logarithm is linked to the expression of the κ -logarithm.

Kaniadakis entropy has the discrete form:

$$S_{\kappa} = - \sum_i \frac{p_i^{1+\kappa} - p_i^{1-\kappa}}{2\kappa}$$

Integer i enumerates cases with probability p_i and κ is a real number.

Now, let us consider a term in the sum:

$$\frac{p_i^{1+\kappa} - p_i^{1-\kappa}}{2\kappa} = p_i \frac{p_i^{\kappa} - p_i^{-\kappa}}{2\kappa}$$

Let us use Euler number and logarithm:

$$\frac{p_i}{2\kappa}(p_i^\kappa - p_i^{-\kappa}) = \frac{p_i}{2\kappa}(e^{\kappa \ln p_i} - e^{-\kappa \ln p_i}) = \frac{p_i}{2\kappa}(e^{u_i} - e^{-u_i})$$

where we define $u_i = \kappa \ln p_i$.

In [8] we can find mentioned a useful formula [9], which is an Euler infinite product expansion:

$$e^u - e^{-u} = 2u \left(1 + \frac{u^2}{\pi^2}\right) \left(1 + \frac{u^2}{2^2 \pi^2}\right) \left(1 + \frac{u^2}{3^2 \pi^2}\right) \dots = 2u \prod_{j=1}^{\infty} \left(1 + \frac{u^2}{j^2 \pi^2}\right)$$

Then the entropy can be written as:

$$\begin{aligned} S_\kappa &= - \sum_i \frac{p_i^{1+\kappa} - p_i^{1-\kappa}}{2\kappa} = - \frac{1}{2\kappa} \sum_i 2 p_i u_i \prod_{j=1}^{\infty} \left(1 + \frac{u_i^2}{j^2 \pi^2}\right) \\ &= - \frac{1}{2\kappa} \sum_i 2 p_i \kappa (\ln p_i) \prod_{j=1}^{\infty} \left(1 + \frac{(\kappa \ln p_i)^2}{j^2 \pi^2}\right) = - \sum_i (p_i \ln p_i) \prod_{j=1}^{\infty} \left(1 + \frac{(\kappa \ln p_i)^2}{j^2 \pi^2}\right) \end{aligned}$$

So we can write:

$$S_\kappa = - \sum_i (p_i \ln p_i) \prod_{j=1}^{\infty} \left(1 + \frac{(\kappa \ln p_i)^2}{j^2 \pi^2}\right)$$

And here we can see clearly that Kaniadakis entropy becomes Shannon entropy for $\kappa \rightarrow 0$:

$$S_{Shannon} = - \sum_i p_i \ln p_i$$

Kaniadakis proposed a form of logarithm in the κ -calculus [1-3] so that:

$$S_\kappa = - \sum_i p_i \ln_\kappa p_i$$

where

$$\ln_\kappa p_i = \frac{p_i^\kappa - p_i^{-\kappa}}{2\kappa}$$

As a consequence, using Euler function we can tell that:

$$\ln_{\kappa} p_i = (\ln p_i) \prod_{j=1}^{\infty} \left(1 + \frac{(\kappa \ln p_i)^2}{j^2 \pi^2}\right)$$

In the generalized additivity of Kaniadakis entropy, it appears another function [5]:

$$I_{\kappa} = \sum_i \frac{p_i^{1+\kappa} + p_i^{1-\kappa}}{2}$$

In [8] we find another Euler formula [10]:

$$e^u + e^{-u} = 2 \left(1 + \frac{4u^2}{\pi^2}\right) \left(1 + \frac{4u^2}{3^2 \pi^2}\right) \left(1 + \frac{4u^2}{5^2 \pi^2}\right) \dots = 2 \prod_{j=0}^{\infty} \left(1 + \frac{4u^2}{(2j+1)^2 \pi^2}\right)$$

And then:

$$I_{\kappa} = \sum_i \frac{p_i^{1+\kappa} + p_i^{1-\kappa}}{2} = \sum_i \frac{p_i}{2} \left[2 \prod_{j=0}^{\infty} \left(1 + \frac{4(\kappa \ln p_i)^2}{(2j+1)^2 \pi^2}\right) \right]$$

In the case that $\kappa \rightarrow 0$, we have $I_{\kappa} = 1$.

Let us remember that this function appears in the generalized additivity of Kaniadakis entropy. In the case of two independent systems A and B :

$$S_{\kappa}^{A \cup B} = S_{\kappa}^A I_{\kappa}^B + S_{\kappa}^B I_{\kappa}^A$$

When $\kappa \rightarrow 0$, we have:

$$S_{\kappa}^{A \cup B} = S^A + S^B$$

where the entropies are Shannon entropies.

As a conclusion, using two Euler infinite product expansions we can rewrite Kaniadakis entropy and discuss, in a different manner, its properties with respect to Shannon entropy.

References

- [1] Kaniadakis, G. (2001). Non-linear kinetics underlying generalized statistics. *Physica A: Statistical mechanics and its applications*, 296(3-4), 405-425.
- [2] Kaniadakis, G. (2002). Statistical mechanics in the context of special relativity. *Physical review E*, 66(5), 056125.
- [3] Kaniadakis, G. (2013). Theoretical foundations and mathematical formalism of the power-law tailed statistical distributions. *Entropy*, 15(10), 3983-4010.
- [4] Wada, T. (2004). Thermodynamic stability conditions for nonadditive composable entropies. *Continuum Mechanics and Thermodynamics*, 16(3), 263-267.
- [5] Sparavigna, A. C. (2015). On the generalized additivity of Kaniadakis entropy. *International Journal of Sciences*, 4(02), 44-48.
- [6] Sparavigna, A. C. (2015). Shannon, Tsallis And Kaniadakis Entropies In Bi-level Image Thresholding. *International Journal of Sciences*, 4(02), 35-43.
- [7] Sparavigna, A. C. (2019). Composition Operations of Generalized Entropies Applied to the Study of Numbers. *International Journal of Sciences*, 8(04), 87-92.
- [8] Gabrio Piola. Sul Calcolo degli Integrali Definiti. Pag.120, in *Opuscoli Matematici e Fisici di Diversi Autori*, Tomo Secondo, Milano, presso Paolo Emilio Giusto, 1834.
- [9] Euler. *Introductio in analysin infinitorum*. T. I, num.156.
- [10] Euler. *Introductio in analysin infinitorum*. T. I, num.157.