

An integral containing a Bessel Function and a Modified Bessel Function of the First Kind

Original

An integral containing a Bessel Function and a Modified Bessel Function of the First Kind / Sparavigna, Amelia Carolina.
- In: PHILICA. - ISSN 1751-3030. - 2017:1082(2017). [10.5281/zenodo.4088590]

Availability:

This version is available at: 11583/2848485 since: 2020-10-14T17:53:47Z

Publisher:

PHILICA

Published

DOI:10.5281/zenodo.4088590

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

- [Philica front page](#)
- [Search](#)
- [About Philica](#)
- [Take the tour](#)
- [Publish your work](#)
- [Work needing review](#)
- [Most popular entries](#)
- [Highest-rated entries](#)
- [Recent reviews](#)
- [How to cite Philica](#)
- [FAQs](#)
- [Support Philica](#)
- [Contact us](#)
- [Get confirmed status](#)

An integral containing a Bessel Function and a Modified Bessel Function of the First Kind

Amelia Carolina Sparavigna

Department of Applied Science and Technology, Politecnico di Torino

Published in matho.philica.com

Abstract

Here we discuss the calculation of an integral containing the Bessel function $J_0(r)$ and the modified Bessel function of the first kind $I_1(r)$. The calculus is based on a function of $J_0(r)$, $I_1(r)$ and of their derivatives, having a Wronskian form. The method here described could be useful for training the students in the manipulation of such integrals.

Article body

An integral containing a Bessel Function and a Modified Bessel Function of the First Kind

Amelia Carolina Sparavigna

Politecnico di Torino, Italy

Here we discuss the calculation of an integral containing the Bessel function $J_0(r)$ and the modified Bessel function of the first kind $I_1(r)$. The calculus is based on a function of $J_0(r)$, $I_1(r)$ and of their derivatives, having a Wronskian form. The method here described could be useful for training the students in the manipulation of such integrals.

Here we are proposing a method for the calculus of an integral of the form $\int r^2 J_0(r) I_1(r) dr$. The method is based on some suitable functions, which are the products of the Bessel and modified Bessel functions and their derivatives. The approach, which is coming from the author's experience, could be useful for training the students in the manipulation of integrals containing Bessel functions. The author's experience was concerning a calculus required by a method for determining the thermal diffusivity, based on the measurement of the thermal expansion of a cylindrical sample [1-4]. For this method, I had to consider the problem of the thermoelasticity. In particular, I had to test if the thermal expansion of a cylindrical sample, assumed as the integral of temperature $\theta(r, z, t)$, that is $\Delta = \beta \int_0^1 \theta(r, z, t) dz$, (β is the coefficient of the expansion), was equal to the component u_z of the displacement field \vec{u} found by means of the thermoelasticity equations. These equations are [5]:

$$\mu \nabla^2 \vec{u} + (\lambda + \mu) \text{grad div}(\vec{u}) - (3\lambda + 2\mu) \beta \text{grad} \theta = \rho \vec{u}$$

$$k \nabla^2 \theta - c_v \dot{\theta} - \frac{(c_p - c_v)}{\beta} \text{div}(\vec{u}) = 0$$

λ and μ are the Lamè coefficients. c_p and c_v are the specific heats at constant pressure and volume, k is the thermal conductivity and ρ the density. In the case of a solid material, the second equation reduces to $k \nabla^2 \theta - c_v \dot{\theta} = 0$. The solution of these equations for a cylindrical sample is requiring hyperbolic sines and cosines and Bessel functions $J_0(r)$ and $I_1(r)$, and the related integrals. Here we consider and discuss how to calculate one of these integrals, that of the form: $\int r^2 I_1(r) J_0(r) dr$.

Let us start from the equation of the Bessel function and of the modified Bessel function of the first kind (the variable r is here assumed as dimensionless):

$$r^2 J_0''(r) + r J_0'(r) + r^2 J_0(r) = 0 \quad (1)$$

$$r^2 I_1''(r) + r I_1'(r) - (r^2 + 1) I_1(r) = 0 \quad (2)$$

We can multiply (1) by $I_1(r)$ and (2) by $J_0(r)$ and then subtract the results as follow:

$$r^2 J_0''(r) I_1(r) + r J_0'(r) I_1(r) + r^2 J_0(r) I_1(r) - r^2 I_1''(r) J_0(r) - r I_1'(r) J_0(r) + (r^2 + 1) J_0(r) I_1(r) = 0$$

$$r^2 [J_0''(r) I_1(r) - I_1''(r) J_0(r)] + r [J_0'(r) I_1(r) - I_1'(r) J_0(r)] + (2r^2 + 1) J_0(r) I_1(r) = 0 \quad (3)$$

Let us consider a Wronskian function W and its derivative, defined in the following manner:

$$W = J_0'(r) I_1(r) - J_0(r) I_1'(r)$$

$$W' = J_0''(r) I_1(r) + J_0'(r) I_1'(r) - J_0'(r) I_1'(r) - J_0(r) I_1''(r) = J_0''(r) I_1(r) - J_0(r) I_1''(r)$$

$$r^2 W' + r W + (2r^2 + 1) J_0(r) I_1(r) = 0$$

Therefore, from (3), we have:

$$r^2 W' + 2r W = r W - (2r^2 + 1) J_0(r) I_1(r)$$

A simple integration gives:

$$r^2 W = \int r W dr - 2 \int r^2 J_0(r) I_1(r) dr - \int J_0(r) I_1(r) dr \quad (4)$$

Moreover, we have the derivatives of the Bessel functions [6]:

$$J_0'(r) = -J_1(r) \quad ; \quad J_1'(r) = J_0(r) - \frac{J_1(r)}{r} \quad (5)$$

$$I_0'(r) = I_1(r) \quad ; \quad I_1'(r) = I_0(r) - \frac{I_1(r)}{r}$$

From (4), using (5):

$$-r^2 W = -\int r W dr + 2 \int r^2 J_0(r) I_1(r) dr + \int J_0(r) I_1(r) dr$$

$$= \int r [J_1(r) I_1(r) + J_0(r) I_0(r) - J_0(r) I_1(r)/r] dr + 2 \int r^2 J_0(r) I_1(r) dr + \int J_0(r) I_1(r) dr$$

$$-r^2 W = \int r [J_1(r) I_1(r) + J_0(r) I_0(r)] dr + 2 \int r^2 J_0(r) I_1(r) dr$$

Therefore;

$$\int r^2 J_0(r) I_1(r) dr = -\frac{1}{2} r^2 W - \frac{1}{2} \int r [J_1(r) I_1(r) + J_0(r) I_0(r)] dr \quad (6)$$

Then, to evaluate the integral in the left side of (6), we need the integrals $\int r J_1(r) I_1(r) dr$ and $\int r J_0(r) I_0(r) dr$. These integrals are easy to calculate. Let us consider the two functions $\Pi_1 = r J_0(r) I_1(r)$; $\Pi_2 = r J_1(r) I_0(r)$. Let us evaluate the following:

$$\Pi_1' = J_0(r) I_1(r) - r J_1(r) I_1(r) + r J_0(r) I_0(r) - r J_0(r) I_1(r)/r = -r J_1(r) I_1(r) + r J_0(r) I_0(r)$$

$$\Pi_2' = J_1(r) I_0(r) + r J_0(r) I_0(r) - r J_1(r) I_0(r)/r + r J_1(r) I_1(r) = r J_0(r) I_0(r) + r J_1(r) I_1(r)$$

Adding Π_1' , Π_2' , we obtain:

$$\Pi_1' + \Pi_2' = -r J_1(r) I_1(r) + r J_0(r) I_0(r) + r J_0(r) I_0(r) + r J_1(r) I_1(r) = 2r J_0(r) I_0(r)$$

As a consequence (after an integration):

$$2 \int r J_0(r) I_0(r) dr = \Pi_1 + \Pi_2 \rightarrow \int r J_0(r) I_0(r) dr = \frac{r}{2} [J_0(r) I_1(r) + J_1(r) I_0(r)] \quad (7)$$

Subtracting Π_1' , Π_2' , we obtain:

$$\Pi_1' - \Pi_2' = -r J_1(r) I_1(r) + r J_0(r) I_0(r) - r J_0(r) I_0(r) - r J_1(r) I_1(r) = -2r J_1(r) I_1(r)$$

Therefore, we have:

$$2 \int r J_1(r) I_1(r) dr = -\Pi_1 + \Pi_2 \rightarrow \int r J_1(r) I_1(r) dr = \frac{r}{2} [-J_0(r) I_1(r) + J_1(r) I_0(r)] \quad (8)$$

Using the results (7) and (8):

$$\int r^2 J_0(r) I_1(r) dr = -\frac{1}{2} r^2 [J_0'(r) I_1(r) - J_0(r) I_1'(r)]$$

$$+ \frac{r}{4} [J_0(r) I_1(r) - J_1(r) I_0(r)] - \frac{r}{4} [J_0(r) I_1(r) + J_1(r) I_0(r)]$$

$$= -\frac{1}{2} r^2 [-J_1(r) I_1(r) - J_0(r) I_0(r) + J_0(r) I_1(r)/r] - \frac{r}{2} J_1(r) I_0(r)$$

$$= \frac{1}{2} r^2 [J_0(r) I_0(r) + J_1(r) I_1(r)] - \frac{r}{2} [J_0(r) I_1(r) + J_1(r) I_0(r)]$$

As we have seen in the calculation, to find the integral $\int r^2 J_0(r) I_1(r) dr$ we used functions W , Π_1 , Π_2 and their derivatives. The search for these functions could be intriguing to students, and for this reason, it could stimulate their interest during the study of the integrals of Bessel functions.

References

- [1] Omini, M., Sparavigna, A., & Strigazzi, A. (1990). Dilatometric determination of thermal diffusivity in low conducting materials. Measurement Science and Technology, 1(2), 166.
- [2] Sparavigna, A., Giachello, G., Omini, M., & Strigazzi, A. (1990). High-sensitivity capacitance method for measuring thermal diffusivity and thermal expansion: results on aluminum and copper. International Journal of Thermophysics, 11(6), 1111-1126.
- [3] Omini, M., Sparavigna, A., & Strigazzi, A. (1990). Calibration of capacitive cells for dilatometric measurements of thermal diffusivity. Measurement Science and Technology, 1(11), 1228.
- [4] Sparavigna, A. C. (1990). Nuovo metodo dilatometrico di misura della diffusività termica dei solidi (Tesi di Dottorato).
- [5] Kovalenko, A. D. (1970). Thermoelasticity. Basic theory and applications. Wolters-Noordhoff.
- [6] Abramowitz, M., & Stegun, I. A. (1972). Handbook of mathematical functions, National Bureau of Standards, Applied Mathematics Series - 55, 1972, Chapter 9, Bessel function of Integer Order, F. W. J. Olver.

Information about this Article

This Article was published on 19th July, 2017 at 13:25:53 and has been viewed 593 times.

The full citation for this Article is:
Sparavigna, A. (2017). An integral containing a Bessel Function and a Modified Bessel Function of the First Kind. *PHILICA.COM Article number 1082*.

NEWS: The SOAP Project, in collaboration with CERN, are conducting a [survey on open-access publishing](#). Please take a moment to give them your views