

Special arrangements of lines: Codimension 2 ACM varieties in $P^1 \times P^1 \times P^1$

Original

Special arrangements of lines: Codimension 2 ACM varieties in $P^1 \times P^1 \times P^1$ / Favacchio, G.; Guardo, E.; Picone, B.. - In: JOURNAL OF ALGEBRA AND ITS APPLICATIONS. - ISSN 0219-4988. - STAMPA. - 18:4(2019), pp. 1950073-1-1950073-22. [10.1142/S0219498819500737]

Availability:

This version is available at: 11583/2859984 since: 2021-01-13T09:37:35Z

Publisher:

World Scientific Publishing Co. Pte Ltd

Published

DOI:10.1142/S0219498819500737

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Nature --> vedi Generico

[DA NON USARE] ex default_article_draft

(Article begins on next page)

SPECIAL ARRANGEMENTS OF LINES: CODIMENSION TWO ACM VARIETIES IN $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$

GIUSEPPE FAVACCHIO, ELENA GUARDO, AND BEATRICE PICONE

ABSTRACT. In this paper we investigate special arrangements of lines in multiprojective spaces. In particular, we characterize codimension two arithmetically Cohen-Macaulay (ACM) varieties in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, called *varieties of lines*. We also describe their ACM property from a combinatorial algebra point of view.

INTRODUCTION

Given a variety $X \subseteq \mathbb{P}^{a_1} \times \cdots \times \mathbb{P}^{a_n}$ an interesting problem is the description of the homological invariants of the coordinate ring of X . This problem has been primarily studied for points, although there is not a general answer in this direction. A great difficulty comes from the fact that a set of distinct points $X \subseteq \mathbb{P}^{a_1} \times \cdots \times \mathbb{P}^{a_n}$ is not necessarily arithmetically Cohen-Macaulay (ACM). See, for instance [11, 16, 17, 18, 19, 20] for some results on this topic, and [12, 13] for a recent characterization of the ACM property in $\mathbb{P}^1 \times \cdots \times \mathbb{P}^1$ and, under certain conditions, in $\mathbb{P}^1 \times \mathbb{P}^m$. Recently, multiprojective spaces have received more attention since they arise in many applications. For example, a specific value of the Hilbert function of a collection of (fat) points in a multiprojective space is related to a classical problem of algebraic geometry concerning the dimension of certain secant varieties of Segre varieties (see [2, 4, 5] just to cite some of them). Or in [3, 6], the authors deduce new results about tensors, and in [7], the author focus on the implicitization problem for tensor product surfaces.

In particular, it appears of interest in combinatorial algebraic geometry to study finite arrangements of lines (see [8, 22] for recent developments in \mathbb{P}^2). A line arrangement over an algebraically closed field K is a finite collection $L_1, \dots, L_d \subseteq \mathbb{P}^2$, $d > 1$, of distinct lines in the projective plane and their crossing points (i.e., the points of intersections of the lines). In this paper we investigate special arrangements of lines in multiprojective spaces by focusing on ACM codimension two varieties in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, called *varieties of lines*, since we want to generalize the codimension two ACM property of points in $\mathbb{P}^1 \times \mathbb{P}^1$. Specifically, we study special cases arising from their intersection points (see Theorem 5.6). These varieties can be viewed as special configurations of codimension two linear varieties in \mathbb{P}^5 .

Our paper is structured as follows. In Section 1 we set up our notation and recall known results. In Section 2, we describe a connection between ideals of varieties of lines and some squarefree monomial ideals (Lemma 2.2). We introduce

Date: August 9, 2018.

2010 Mathematics Subject Classification. 13C40, 13F20, 13A15, 14C20, 14M05.

Key words and phrases. Configuration of lines, multiprojective spaces, arithmetically Cohen-Macaulay.

the $\text{Hyp}_n(\star)$ -property (Definition 2.12) to give a combinatorial characterization of ACM varieties of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ using a well known property of chordal graphs (Theorem 2.17). In Section 3 we introduce a numerical way to check the ACM property for any varieties of lines. In Section 4 we describe the Hilbert function of Ferrers varieties of lines, a special case of ACM variety of lines. Finally, in Section 5, we initiate an investigation on varieties of lines whose set of crossing points is a complete intersection of points in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. We also characterize varieties of lines defined by a complete intersection ideal in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ (Theorem 5.10). We end the paper with two possible research topics to explore: (1) the connection between our varieties of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and special configurations of lines of \mathbb{P}^3 and (2) the Hilbert function of any ACM variety of lines (Question 5.13).

Acknowledgement. The authors thank A. Van Tuyl for his comments in a previous version of the paper. They also thank the referee for his/her helpful suggestions. The second author would like to thank GNSAGA and Prin 2015. The software CoCoA [1] was indispensable for all the computations.

1. NOTATION AND BASIC FACTS

Throughout the paper $\mathbb{N} := \{0, 1, 2, \dots\}$ denotes the set of non-negative integers and \preceq denotes the natural partial order on the elements of $\mathbb{N}^3 := \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ defined by $(a_1, a_2, a_3) \preceq (b_1, b_2, b_3)$ in \mathbb{N}^3 if and only if $a_i \leq b_i \forall i = 1, 2, 3$. Let $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ be the standard basis of \mathbb{N}^3 . Let $R := K[x_{1,0}, x_{1,1}, x_{2,0}, x_{2,1}, x_{3,0}, x_{3,1}]$ be the polynomial ring over an algebraically closed field K of characteristic zero. We induce a multi-grading by setting $\deg x_{i,j} = \mathbf{e}_i$ for $i \in \{1, 2, 3\}$. A monomial $m = x_{1,0}^{a_0} x_{1,1}^{a_1} x_{2,0}^{b_0} x_{2,1}^{b_1} x_{3,0}^{c_0} x_{3,1}^{c_1} \in R$ has *tridegree* (or simply, *degree*) $\deg m = (a_0 + a_1, b_0 + b_1, c_0 + c_1)$. We make the convention that 0 has degree $\deg 0 = (i, j, k)$ for all $(i, j, k) \in \mathbb{N}^3$. Note that the elements of the field K all have degree $(0, 0, 0)$. For each $(i, j, k) \in \mathbb{N}^3$, let $R_{i,j,k}$ denote the vector space over K spanned by all the monomials of R of degree (i, j, k) . The polynomial ring R is then a *trigraded ring* because there exists a direct sum decomposition

$$R = \bigoplus_{(i,j,k) \in \mathbb{N}^3} R_{i,j,k}$$

such that $R_{i,j,k} R_{l,m,n} \subseteq R_{i+l, j+m, k+n}$ for all $(i, j, k), (l, m, n) \in \mathbb{N}^3$. An element $F \in R$ is *trihomogeneous* (or simply, *homogeneous*) if $F \in R_{i,j,k}$ for some $(i, j, k) \in \mathbb{N}^3$. An ideal $I = (F_1, F_2, \dots, F_r) \subseteq R$ is a *(tri)homogeneous ideal* if F_i is (tri)homogeneous for all $i = 1, 2, \dots, r$.

Let $I \subseteq R$ be a homogeneous ideal, and we let $I_{i,j,k} := I \cap R_{i,j,k}$ for all $(i, j, k) \in \mathbb{N}^3$. Because I is homogeneous, the quotient ring R/I also inherits a graded ring structure. In particular, we have:

$$R/I = \bigoplus_{(i,j,k) \in \mathbb{N}^3} (R/I)_{i,j,k} = \bigoplus_{(i,j,k) \in \mathbb{N}^3} R_{i,j,k}/I_{i,j,k}.$$

Definition 1.1. Let $I_X \subseteq R$ be the homogeneous ideal defining a variety $X \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. We say that X is *arithmetically Cohen-Macaulay* (ACM) if R/I_X is Cohen-Macaulay, i.e. $\text{depth}(R/I_X) = \text{Krull-dim}(R/I_X)$.

Definition 1.2. We say that a homogeneous ideal J in a polynomial ring S is *Cohen-Macaulay* (CM) if S/J is Cohen-Macaulay.

A point in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ is an ordered set of three points in \mathbb{P}^1 . Say $P := ([a_0, a_1], [b_0, b_1], [c_0, c_1]) \in \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, the defining ideal of P is $I_P := (a_1x_{1,0} - a_0x_{1,1}, b_1x_{2,0} - b_0x_{2,1}, c_1x_{3,0} - c_0x_{3,1})$. Note that I_P is a height three prime ideal generated by homogeneous linear forms of different degree.

Throughout the paper, linear forms are denoted by capital letters. In particular, we use A_i to denote a linear form of degree $(1, 0, 0)$, B_j a linear form of degree $(0, 1, 0)$, and C_k a linear form of degree $(0, 0, 1)$. We denote by $\mathcal{L}(A_i)$, $\mathcal{L}(B_j)$ and $\mathcal{L}(C_k)$ the respective hyperplanes of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, and we say that a hyperplane in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ is of type \mathbf{e}_i if it is defined by a form of degree \mathbf{e}_i .

We recall the following definition (see [20], Definition 2.2).

Definition 1.3. Let $F, G \in R$ be two homogeneous linear forms of different degree. In $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ the variety \mathcal{L} defined by the ideal $(F, G) \subseteq R$ is called a *line* of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and we denote it by $\mathcal{L}(F, G)$.

We say that a line $\mathcal{L}(F, G)$ is of type $\mathbf{e}_i + \mathbf{e}_j$, with $i \neq j$, if $\{\deg F, \deg G\} = \{\mathbf{e}_i, \mathbf{e}_j\}$.

In particular, if $A \in R_{1,0,0}$, $B \in R_{0,1,0}$ and $C \in R_{0,0,1}$, then we denote by $\mathcal{L}(A, B)$ the variety in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ defined by the ideal $(A, B) \subseteq R$ and we call it *line of type* $(1, 1, 0)$. Analogously, we call the variety $\mathcal{L}(A, C)$ *line of type* $(1, 0, 1)$ and the variety $\mathcal{L}(B, C)$ *line of type* $(0, 1, 1)$. We also refer to lines of type $\mathbf{e}_1 + \mathbf{e}_2$, $\mathbf{e}_1 + \mathbf{e}_3$ and $\mathbf{e}_2 + \mathbf{e}_3$ by writing *lines having direction* \mathbf{e}_3 , \mathbf{e}_2 and \mathbf{e}_1 , respectively.

Definition 1.4. We say that $X \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ is a *variety of lines* if it is given by a finite union of distinct lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$.

Definition 1.5. Given $X \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ a variety of lines, we denote by $\mathcal{H}_1(X) := \{\mathcal{L}(A_1), \dots, \mathcal{L}(A_{d_1})\}$, $\mathcal{H}_2(X) := \{\mathcal{L}(B_1), \dots, \mathcal{L}(B_{d_2})\}$ and $\mathcal{H}_3(X) := \{\mathcal{L}(C_1), \dots, \mathcal{L}(C_{d_3})\}$ the hyperplanes of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ containing some lines of X . In particular:

$$X := \bigcup_{(i,j) \in U_3(X)} \mathcal{L}(A_i, B_j) \cup \bigcup_{(i,k) \in U_2(X)} \mathcal{L}(A_i, C_k) \cup \bigcup_{(j,k) \in U_1(X)} \mathcal{L}(B_j, C_k)$$

where $U_3(X) \subseteq [d_1] \times [d_2]$, $U_2(X) \subseteq [d_1] \times [d_3]$ and $U_1(X) \subseteq [d_2] \times [d_3]$ are sets of ordered pairs of integers, with $[n] := \{1, 2, \dots, n\} \subset \mathbb{N}$.

For $i = 1, 2, 3$, we denote by X_i the set of lines of X having direction \mathbf{e}_i and we call $U_i(X)$ the *index set of* X_i .

Thus, the ideal defining X is

$$I_X = \bigcap_{(i,j) \in U_3(X)} (A_i, B_j) \bigcap_{(i,k) \in U_2(X)} (A_i, C_k) \bigcap_{(j,k) \in U_1(X)} (B_j, C_k).$$

In this paper we are interested in a combinatorial characterization of ACM varieties of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and their Hilbert function. In $\mathbb{P}^1 \times \mathbb{P}^1$, to describe combinatorially ACM sets of points, it was crucial the definition of the so called *Ferrers diagram* (see for instance [19]).

Definition 1.6. A tuple $\lambda = (\lambda_1, \dots, \lambda_r)$ of positive integers is a *partition* of an integer s if $\sum_{i=1}^r \lambda_i = s$ and $\lambda_i \geq \lambda_{i+1}$ for every i . We write $\lambda = (\lambda_1, \dots, \lambda_r) \vdash s$.

Definition 1.7. To any partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r) \vdash s$ we can associate the following diagram: on an $r \times \lambda_1$ grid, place λ_1 points on the first horizontal line, λ_2 points on the second, and so on, where the points are left justified. The resulting diagram is called the *Ferrers diagram* of the partition λ .

Definition 1.8. Let Y be a finite set of points in $\mathbb{P}^1 \times \mathbb{P}^1$. We say that Y *resembles a Ferrers diagram* if the set of points looks like a Ferrers diagram, i.e., after relabeling the horizontal and vertical rulings, we can assume that the first horizontal ruling contains the most number of points of Y , the second contains the same number or less of points of Y , and so on.

Applying Lemma 3.17, Theorem 3.21 and Theorem 4.11 in [19], we have

Lemma 1.9. *Let Y be a finite set of points in $\mathbb{P}^1 \times \mathbb{P}^1$. Y is ACM if and only if Y resembles a Ferrers diagram.*

We adapt Definition 1.7 to our context.

Construction 1.10. *Let X be a variety of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and consider the set X_3 of lines of X of type $(1, 1, 0)$ indexed by $U_3(X) \subseteq [d_1] \times [d_2]$. We represent X_3 as a $d_1 \times d_2$ grid, where the horizontal lines are labeled by the $\mathcal{L}(A_i)$'s for $i = 1, \dots, d_1$ and the vertical lines by the $\mathcal{L}(B_j)$'s for $j = 1, \dots, d_2$. By abuse of notation, we denote the horizontal lines by $\mathcal{L}(A_i)$ and the vertical lines by $\mathcal{L}(B_j)$. Then, a line $\mathcal{L}(A_i, B_j) \in X_3$ is drawn as the intersection point of $\mathcal{L}(A_i)$ and $\mathcal{L}(B_j)$ in the grid. Similarly, we can construct a $d_1 \times d_3$ grid representing X_2 and a $d_2 \times d_3$ grid representing X_1 .*

Definition 1.11. Let X be a variety of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and $h \in \{1, 2, 3\}$. We say that X *resembles a Ferrers diagram with respect to the direction h* if the grid representing the lines of X_h , constructed as above, resembles a Ferrers diagram.

Definition 1.12. A finite subset $U = \{(u_i, u_j)\} \subseteq \mathbb{N}^2$ *resembles a Ferrers diagram* if it satisfies the following property:

$$(u_i, u_j) \in U \Rightarrow (u_h, u_k) \in U \quad \forall 1 \leq h \leq i, 1 \leq k \leq j.$$

Remark 1.13. Note that Definition 1.11 is equivalent to say that the index set $U_h(X) \subset \mathbb{N}^2$ resembles a Ferrers diagram as Definition 1.12.

Remark 1.14. Construction 1.10 makes clear the connection between X_h ($h \in \{1, 2, 3\}$) and a set of points in $\mathbb{P}^1 \times \mathbb{P}^1$. X_h is a cone of a set of distinct points on a hyperplane of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. So, we can look at it as a set of points in $\mathbb{P}^1 \times \mathbb{P}^1$ with associated grid as described in the construction.

Example 1.15. Let X be the following variety of 15 lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$:

$$\begin{aligned} X = & \mathcal{L}(A_1, B_2) \cup \mathcal{L}(A_1, B_4) \cup \mathcal{L}(A_1, B_5) \cup \mathcal{L}(A_2, B_2) \cup \mathcal{L}(A_2, B_3) \cup \\ & \cup \mathcal{L}(A_2, B_4) \cup \mathcal{L}(A_2, B_5) \cup \mathcal{L}(A_3, B_1) \cup \mathcal{L}(A_3, B_2) \cup \mathcal{L}(A_3, B_3) \cup \\ & \cup \mathcal{L}(A_3, B_4) \cup \mathcal{L}(A_3, B_5) \cup \mathcal{L}(A_4, B_4) \cup \mathcal{L}(B_1, C_1) \cup \mathcal{L}(B_2, C_2). \end{aligned}$$

Then,

$$\begin{aligned} X_3 = & \{\mathcal{L}(A_1, B_2), \mathcal{L}(A_1, B_4), \mathcal{L}(A_1, B_5), \mathcal{L}(A_2, B_2), \mathcal{L}(A_2, B_3), \mathcal{L}(A_2, B_4), \mathcal{L}(A_2, B_5), \\ & \mathcal{L}(A_3, B_1), \mathcal{L}(A_3, B_2), \mathcal{L}(A_3, B_3), \mathcal{L}(A_3, B_4), \mathcal{L}(A_3, B_5), \mathcal{L}(A_4, B_4)\}. \end{aligned}$$

Using Construction 1.10, X_3 is represented by a 4×5 grid as Figure 1. After renaming, we see that X_3 resembles a Ferrers diagram of type $(5, 4, 3, 1)$. Then, using Lemma 1.9, X_3 is ACM (Figure 2).




FIGURE 1




FIGURE 2

Example 1.16. Let X be the variety of lines as in Example 1.15. We have $X_1 = \{\mathcal{L}(B_1, C_1), \mathcal{L}(B_2, C_2)\}$ and the 2×2 grid representing X_1 does not resemble any Ferrers diagram (Figure 3). Thus X does not resemble a Ferrers diagram with respect to the direction 1. Hence, from Lemma 1.9, X_1 is not ACM.




FIGURE 3

Since Ferrers diagrams play a crucial role in the characterization of the ACM property for a finite set of points in $\mathbb{P}^1 \times \mathbb{P}^1$ (see for instance [20]), it is natural for us to investigate the same property for a variety of lines $X \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ (since X has also codimension 2). In the next section, we will show that the ACM property of X depends on the X_i (see Corollary 2.8), but the ACM-ness of the X_i is not sufficient to ensure that X is also ACM (see Remark 2.9).

2. A COMBINATORIAL CHARACTERIZATION OF ACM VARIETIES OF LINES

In this section, we study the ACM property for varieties of lines from a combinatorial point of view. We refer to [23] for all the introductory material on monomial ideals.

The next lemma can be recovered from [26], Proposition 3.2.

Lemma 2.1. *Let $X \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ be a variety of lines. Then, there exist three forms A, B and C of degree $(1, 0, 0)$, $(0, 1, 0)$ and $(0, 0, 1)$, respectively, such that $(\bar{A}, \bar{B}, \bar{C})$ is a regular sequence in R/I_X .*

Proof. Let $A \in R_{1,0,0}$ be such that $\mathcal{L}(A) \notin \mathcal{H}_1(X)$. We claim that \bar{A} is a nonzero divisor of R/I_X . Indeed, take $F \in R$ a homogeneous form such that $AF \in I_X$. Then $AF \in I_{\mathcal{L}}$, for any line $\mathcal{L} \in X$. Since $I_{\mathcal{L}}$ is a prime ideal and $A \notin I_{\mathcal{L}}$, then we get $F \in I_{\mathcal{L}}$, for any $\mathcal{L} \in X$.

Now we prove the existence of the linear form B . Since X is ACM, then $J := I_X + (A)$ is CM. Moreover, J is homogeneous and its height is 3. Take the primary decomposition of J , say $J = \mathfrak{q}_1 \cap \cdots \cap \mathfrak{q}_t$, and let $\mathfrak{p}_i := \sqrt{\mathfrak{q}_i}$ for $i = 1, \dots, t$.

The set of the nonzero divisors of R/J is then $\bigcup_i \bar{\mathfrak{p}}_i$. In order to prove that there exists an element $B \in R_{0,1,0}$ nonzero divisor of R/J , it is enough to show that $(\bigcup_i \mathfrak{p}_i)_{0,1,0} \subsetneq R_{0,1,0}$. Since $R_{0,1,0}$ is a K -vector space over an infinite field, it is not a union of a finite number of its proper subspaces, and so it is enough to show that $(\mathfrak{p}_i)_{0,1,0} \subsetneq R_{0,1,0}$ for each $i = 1, \dots, t$.

Let $i \in \{1, \dots, t\}$, then we have $I_X \subseteq J \subseteq \mathfrak{p}_i$. Therefore, there exists $\mathcal{L} \in X$ such that $I_X \subseteq I_{\mathcal{L}} \subseteq \mathfrak{p}_i$. This implies $\mathfrak{p}_i = I_{\mathcal{L}} + (A)$. Since $I_{\mathcal{L}} \neq R_{0,1,0}$ we are done.

Analogously we prove the existence of a form $C \in R_{0,0,1}$. \square

We set the notation for this section. Let X be a variety of lines and I_X its defining ideal

$$I_X = \bigcap_{(i,j) \in U_3(X)} (A_i, B_j) \bigcap_{(i,k) \in U_2(X)} (A_i, C_k) \bigcap_{(j,k) \in U_1(X)} (B_j, C_k) \subseteq R.$$

We construct a new polynomial ring in $d_1 + d_2 + d_3$ variables each of them corresponding to a hyperplane containing some lines of X . We denote by $S := K[a_1, \dots, a_{d_1}, b_1, \dots, b_{d_2}, c_1, \dots, c_{d_3}]$ the polynomial ring in $d_1 + d_2 + d_3$ variables and $\deg a_i = (1, 0, 0)$, $\deg b_j = (0, 1, 0)$, $\deg c_k = (0, 0, 1)$. We set

$$J_X = \bigcap_{(i,j) \in U_3(X)} (a_i, b_j) \bigcap_{(i,k) \in U_2(X)} (a_i, c_k) \bigcap_{(j,k) \in U_1(X)} (b_j, c_k) \subseteq S.$$

J_X is a height 2 monomial ideal of S and its associated primes correspond to the components of X .

The next lemma is crucial since, as its consequence, we can connect homological invariants between ACM varieties of lines and some height 2 monomial ideals. Similar arguments were also used in [12] (see proof of Theorem 3.2).

Lemma 2.2. *Let X be a variety of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. Then X is ACM if and only if $J_X \subseteq S$ is CM.*

Proof. Set $T := S[x_{1,0}, x_{1,1}, x_{2,0}, x_{2,1}, x_{3,0}, x_{3,1}]$. Consider J_X as an ideal, say \bar{J}_X , in the ring T . Since J_X is a height 2 monomial ideal in S , then \bar{J}_X , being a cone, continues to be a height 2 monomial ideal. Moreover, \bar{J}_X has the same primary decomposition as J_X . Consider the linear forms $a_i - A_i$, $b_j - B_j$, $c_k - C_k$ and let L be the ideal generated by all these linear forms.

Assume J_X is CM. Thus, in the quotient $T/(\bar{J}_X, L)$ we can view the addition of each linear form in L as a proper hyperplane section. We have that R/I_X and $T/(\bar{J}_X, L)$ both have height 2 and $R/I_X \cong T/(\bar{J}_X, L)$. Then, since J_X is CM, we get X is ACM.

On the other hand, if X is ACM, then, applying Lemma 2.1, there exists a sequence of linear forms $(A, B, C) \subseteq R$ that is regular in the quotient R/I_X . Let $\mathfrak{q} := (A, B, C) \subseteq R$ be the ideal generated by these three linear forms. Consider the ideal $(I_X + \mathfrak{q})/\mathfrak{q} \subseteq R/\mathfrak{q}$, that can be viewed as a codimension 2 monomial ideal in a polynomial ring in three variables. Since a Hilbert-Burch matrix of I_X has the same “structure” as the Hilbert-Burch matrix of a monomial ideal, i.e. it is a matrix with only two non zero entries in each column (see for instance Lemma 3.21 in [14] or Theorem 1.5. in [24]), then I_X is generated by some products among the linear forms defining the lines of X . Since the addition of each linear form in L can be seen as a proper hyperplane section, we also have $R/I_X \cong T/(\bar{J}_X, L)$. Then J_X is CM. \square

Corollary 2.3. *Let X be an ACM variety of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. Then I_X is generated by products of linear forms.*

As a consequence of Lemma 2.2, it is interesting to further investigate the structure of the monomial ideal J_X associated to X . Now we recall only a few definitions we will use in the sequel. We refer to [23, 25] for all preliminaries and for further results on graphs.

A (simple) graph G is a pair $G = (V, E)$, where $V := \{v_1, \dots, v_N\}$ is a set of vertices of G and E is a collection of 2-subsets of V , called the edges of G . The complementary graph of G , denoted by G^c , is the graph $G^c = (V, E^c)$, where $E^c = \{\{v_i, v_j\} \mid \{v_i, v_j\} \notin E\}$. A sequence of vertices of G , (v_1, v_2, \dots, v_t) , is a cycle of length t if $\{v_1, v_2\}, \{v_2, v_3\}, \dots, \{v_t, v_1\} \in E$. A chord is an edge joining two not adjacent vertices in a cycle. A minimal cycle is a cycle without chords. A graph G is called chordal when all its minimal cycles have length three. We associate to a graph $G = (V, E)$ two squarefree monomial ideals in the ring $K[v_1, \dots, v_N]$, the face ideal of G

$$I(G) = (v_i v_j \mid \{v_i, v_j\} \in E)$$

and the cover ideal of G

$$J(G) = \bigcap_{\{v_i, v_j\} \in E} (v_i, v_j).$$

It is a well known fact that $I(G)$ and $J(G)$ are the Alexander dual each of the other. In the sequel we will use the following results:

Theorem 2.4 ([15], Theorem 1). *Let G be a graph. Then $I(G)$ has a linear resolution if and only if G^c is a chordal graph.*

Theorem 2.5 ([9], Theorem 3). *Let G be a graph. Then $I(G)$ has a linear resolution if and only if $J(G)$ is CM.*

Remark 2.6. Let X be a variety of lines. Let $G_X = (V_X, E_X)$ be the graph with vertex set

$$V_X := \{a_1, \dots, a_{d_1}, b_1, \dots, b_{d_2}, c_1, \dots, c_{d_3}\}$$

and edge set

$$\begin{aligned} E_X := & \{ \{a_i, b_j\} \subseteq V_X \mid \mathcal{L}(A_i, B_j) \in X_3 \} \cup \\ & \cup \{ \{a_i, c_k\} \subseteq V_X \mid \mathcal{L}(A_i, C_k) \in X_2 \} \cup \\ & \cup \{ \{b_j, c_k\} \subseteq V_X \mid \mathcal{L}(B_j, C_k) \in X_1 \}. \end{aligned}$$

Then, we note that the monomial ideal J_X is the cover ideal of the graph G_X :

$$J_X = J(G_X) \subseteq S,$$

that is the Stanley-Reisner ideal of the simplicial complex (see Lemma 1.5.4. in [23])

$$\Delta_X := \langle V_X \setminus e \mid e \in E_X \rangle.$$

An useful application of Remark 2.6 is the following lemma.

Lemma 2.7. *Let X be an ACM variety of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and let $\mathcal{H} \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ be a hyperplane containing some lines of X . Then the variety of lines $Y = \{\mathcal{L} \in X \mid \mathcal{L} \not\subseteq \mathcal{H}\}$ is ACM.*

Proof. Let H be the linear form defining \mathcal{H} . Denoted by z the variable of S corresponding to H (the linear form H is one of the forms A_i, B_j, C_k and z is the corresponding variable among a_i, b_j, c_k). We have

$$\text{i) } J_X : z = \bigcap_{\substack{\mathfrak{p} \in \text{ass}(J_X) \\ z \notin \mathfrak{p}}} \mathfrak{p}. \text{ Both are monomial ideals, so the equality easily}$$

follows by checking the inclusions for monomials.

$$\text{ii) } J_X : z \text{ is the Stanley-Reisner ideal of the simplicial complex } \text{link}_{\Delta_X} z \text{ (see sections 1.5.2 and 8.1.1 in [23]). Indeed, the Stanley-Reisner ideal of the link of } z \text{ in } \Delta_X \text{ is generated by monomials corresponding to the elements } F \subseteq V_X \text{ such that } \{z\} \cup F \notin \Delta_X. \text{ All these monomials are in } J_X : z = I_{\Delta_X} : z \text{ and vice versa.}$$

Then, in order to prove the statement, it is enough to show that $J_X : z$ is CM. From Lemma 2.2, we have that J_X is CM, so the statement follows by Corollary 8.1.8 in [23]. \square

Corollary 2.8. *If X is an ACM variety of lines, then X resembles a Ferrers diagram with respect to the direction h , for each $h = 1, 2, 3$.*

Proof. We show that $U_1(X)$ resembles a Ferrers diagram. Analogously, one can show the same for $U_2(X)$ and $U_3(X)$. Let us consider the variety of lines X_1 consisting of the lines of X of type $(0, 1, 1)$. Since $I_{X_1} = I_X \setminus \{\mathcal{L}(A_1), \dots, \mathcal{L}(A_{d_1})\}$, X_1 preserves the ACM property by Lemma 2.7. Moreover, $X_1 = \bigcup_{(j,k) \in U_1(X)} \mathcal{L}(B_j, C_k)$,

i.e., it is a cone of an ACM set of distinct points on a hyperplane of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, see Remark 1.14. A well known characterization, see for instance Theorem 4.11 in [19], shows that this set of points resembles a Ferrers diagram. Using Remark 1.13, $U_1(X)$ resembles a Ferrers diagram. Then, the statement follows from Lemma 1.9. \square

Remark 2.9. From previous corollary, if there exists $i \in \{1, 2, 3\}$ such that X_i is not ACM, then X is not ACM. The following example shows that even if all X_i are ACM X could be not ACM.

Example 2.10. Let us consider the following variety of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$:

$$X = \{\mathcal{L}(A_1, B_1), \mathcal{L}(A_2, B_2), \mathcal{L}(B_3, C_3)\}.$$

It is clear that the sets X_1, X_2 and X_3 resemble a Ferrers diagram, so each of them is ACM. But, in this case, X is not ACM. This follows for instance from Lemma 2.2 and from [23], Lemma 9.1.12.

The next definition introduces a property for varieties of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ in analogy to the known (\star) -property defined for sets of points in $\mathbb{P}^1 \times \mathbb{P}^1$ (see [18]).

Definition 2.11. Let $X \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ be a variety of lines. We say that X has the (\star) -property (or explicitly, *star property*) if given any two lines $L_1, L_2 \in X$, there exists $L_3 \in X$ such that L_1, L_3 and L_2, L_3 are coplanar.

We slight generalize this property for varieties of lines.

Definition 2.12. Let $X \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ be a variety of lines. Let $n \geq 4$, $n \in \mathbb{N}$, we say that X has the n -hyperplanes (\star) property (for short, $\text{Hyp}_n(\star)$ -property) if given n hyperplanes H_1, H_2, \dots, H_n such that $\mathcal{L}(H_i, H_j) \in X$ for any $j \neq i-1, i, i+1$ then $\mathcal{L}(H_u, H_{u+1}) \in X$ for some $u \in \{1, 2, \dots, n\}$, where $H_0 = H_n$ and $H_{n+1} = H_1$.

Remark 2.13. Note that if $n > 6$, then X has the $Hyp_n(\star)$ -property. Indeed, among $n > 6$ hyperplanes there are at least three of the same type and so the condition $\mathcal{L}(H_i, H_j) \in X$ for any $j \neq i - 1, i, i + 1$ (where $H_0 = H_n$ and $H_{n+1} = H_1$) fails to be true.

Remark 2.14. Note that the $Hyp_4(\star)$ -property is equivalent to (\star) -property as Definition 2.11.

Example 2.15. Let us consider the following variety of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$:

$$\begin{aligned} X = & \mathcal{L}(A_1, B_1) \cup \mathcal{L}(A_1, B_2) \cup \mathcal{L}(A_1, B_3) \cup \mathcal{L}(A_2, B_1) \cup \\ & \cup \mathcal{L}(A_2, B_2) \cup \mathcal{L}(A_1, C_1) \cup \mathcal{L}(A_1, C_2) \cup \mathcal{L}(A_2, C_1) \cup \\ & \cup \mathcal{L}(B_1, C_1) \cup \mathcal{L}(B_1, C_2) \cup \mathcal{L}(B_2, C_1) \cup \mathcal{L}(B_3, C_1). \end{aligned}$$

X has the $Hyp_4(\star)$ -property. Indeed, if we take the 4 hyperplanes $\mathcal{L}(A_1), \mathcal{L}(A_2), \mathcal{L}(B_1), \mathcal{L}(B_2)$, we have that $\mathcal{L}(A_1, B_1), \mathcal{L}(A_2, B_2) \in X$ and also $\mathcal{L}(A_1, B_2) \in X$; if we take the 4 hyperplanes $\mathcal{L}(A_1), \mathcal{L}(A_2), \mathcal{L}(B_1), \mathcal{L}(C_1)$, we have that $\mathcal{L}(A_1, B_1), \mathcal{L}(A_2, C_1) \in X$ and also $\mathcal{L}(B_1, C_1) \in X$; and so on, if we take any two lines in X , there exists a third line in X that is coplanar with the other two.

Example 2.16. Let us consider the following variety of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$:

$$\begin{aligned} X = & \mathcal{L}(A_1, B_1) \cup \mathcal{L}(A_1, B_2) \cup \mathcal{L}(A_1, B_3) \cup \mathcal{L}(A_2, B_2) \cup \mathcal{L}(A_1, C_1) \\ & \cup \mathcal{L}(A_1, C_2) \cup \mathcal{L}(A_2, C_1) \cup \mathcal{L}(B_1, C_1) \cup \mathcal{L}(B_3, C_1). \end{aligned}$$

X has the $Hyp_5(\star)$ -property. Indeed, if we take the 5 hyperplanes $\mathcal{L}(A_1), \mathcal{L}(A_2), \mathcal{L}(B_1), \mathcal{L}(B_2), \mathcal{L}(C_1)$ we have that the lines $\mathcal{L}(A_1, B_1), \mathcal{L}(A_1, B_2), \mathcal{L}(A_2, B_2), \mathcal{L}(A_2, C_1), \mathcal{L}(B_1, C_1) \in X$ and also $\mathcal{L}(A_1, C_1) \in X$; if we take the 5 hyperplanes $\mathcal{L}(A_1), \mathcal{L}(A_2), \mathcal{L}(B_3), \mathcal{L}(B_2), \mathcal{L}(C_1)$, we have that $\mathcal{L}(A_1, B_3), \mathcal{L}(A_1, B_2), \mathcal{L}(A_2, B_2), \mathcal{L}(A_2, C_1), \mathcal{L}(B_3, C_1) \in X$ and also $\mathcal{L}(A_1, C_1) \in X$; and so on, if we take any 5 hyperplanes H_1, \dots, H_5 among $\mathcal{L}(A_1), \mathcal{L}(A_2), \mathcal{L}(B_1), \mathcal{L}(B_2), \mathcal{L}(B_3), \mathcal{L}(C_1), \mathcal{L}(C_2)$ such that $\mathcal{L}(H_i, H_j) \in X$ for any $j \neq i - 1, i, i + 1$, then there exists $u \in \{1, \dots, 5\}$ such that $\mathcal{L}(H_u, H_{u+1}) \in X$, where $H_0 = H_5$ and $H_6 = H_1$. Note that if we take $\mathcal{L}(B_1), \mathcal{L}(B_2), \mathcal{L}(B_3)$ among the 5 hyperplanes we choose, the condition $\mathcal{L}(H_i, H_j) \in X$ for any $j \neq i - 1, i, i + 1$ fails to be true and then there is nothing to verify.

The following theorem is the main result of this section.

Theorem 2.17. *Let X be a variety of lines. Then X is ACM if and only if X has the $Hyp_n(\star)$ -property for $n = 4, 5, 6$.*

Proof. Let I_X be the ideal defining the variety of lines $X \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. From Lemma 2.2, X is ACM if and only if $J_X \subseteq S$ is CM. From Remark 2.6, the ideal J_X is the cover ideal of the graph G_X , i.e., $J_X = J(G_X)$. From Theorem 2.5, the face ideal $I(G_X)$ has a linear resolution and then, using Theorem 2.4, G_X^c is a chordal graph, that is, X has the $Hyp_n(\star)$ -property for any n . Remark 2.13 completes the proof. \square

3. A NUMERICAL CHARACTERIZATION OF THE ACM PROPERTY

Since we are interested in the study of the ACM property for varieties of lines X , from now on we assume that $U_h(X)$ resembles a Ferrers diagram for each $h = 1, 2, 3$. In order to give a characterization of the ACM property we introduce the following notation.

Definition 3.1. Let $P = P_{ijk} = \mathcal{L}(A_i) \cap \mathcal{L}(B_j) \cap \mathcal{L}(C_k)$ be a point of a variety of lines X , we call *multiplicity of P* the number of lines of X passing through the point P and we denote it by μ_{ijk} .

Remark 3.2. Since at most three lines of X (one of each type) pass through the point P , $\mu_{ijk} \leq 3$.

Definition 3.3. Given a variety of lines X , we define a 3-dimensional matrix $M_X := (\mu_{ijk}) \in \mathbb{N}^{d_1 \times d_2 \times d_3}$ whose (i, j, k) -entry is the multiplicity of $P_{i,j,k}$. We call it the *matrix of the multiplicities* of X .

We also define

Definition 3.4. $M_X^{(3)} := (\mu_{ij0}) \in \mathbb{N}^{d_1 \times d_2}$, where

$$\mu_{ij0} := \begin{cases} 1 & \text{if } (i, j) \in U_3(X), \text{ i.e., } \mathcal{L}(A_i, B_j) \in X \\ 0 & \text{otherwise.} \end{cases}$$

Analogously, $M_X^{(2)} := (\mu_{i0k}) \in \mathbb{N}^{d_1 \times d_3}$, where $\mu_{i0k} := \begin{cases} 1 & \text{if } \mathcal{L}(A_i, C_k) \in X \\ 0 & \text{otherwise} \end{cases}$

and $M_X^{(1)} := (\mu_{0jk}) \in \mathbb{N}^{d_2 \times d_3}$, where $\mu_{0jk} := \begin{cases} 1 & \text{if } \mathcal{L}(B_j, C_k) \in X \\ 0 & \text{otherwise} \end{cases}$.

Example 3.5. Let us consider $X \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ as Figure 4

$$X = \mathcal{L}(A_1, B_1) \cup \mathcal{L}(A_1, B_2) \cup \mathcal{L}(A_2, B_2) \cup \mathcal{L}(A_1, C_1) \cup \mathcal{L}(A_2, C_1) \\ \cup \mathcal{L}(A_2, C_2) \cup \mathcal{L}(B_1, C_1) \cup \mathcal{L}(B_1, C_2) \cup \mathcal{L}(B_2, C_2).$$

We have

$$\mu_{111} = \mu_{222} = 3, \quad \mu_{121} = \mu_{221} = \mu_{211} = \mu_{112} = \mu_{122} = \mu_{212} = 2$$

and

$$M_X^{(3)} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad M_X^{(2)} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad M_X^{(1)} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$




FIGURE 4. The variety of lines X (in bold).

Now we provide a criterion to establish if X is ACM or not just looking at the matrices of the multiplicities M_X , $M_X^{(1)}$, $M_X^{(2)}$ and $M_X^{(3)}$.

Proposition 3.6. *Let X be a variety of lines. X has the $Hyp_6(\star)$ -property iff for all $a_1, a_2 \in [d_1], b_1, b_2 \in [d_2], c_1, c_2 \in [d_3]$*

$$\text{either } \begin{pmatrix} \mu_{a_1 b_1 c_1} & \mu_{a_1 b_2 c_1} \\ \mu_{a_2 b_1 c_1} & \mu_{a_2 b_2 c_1} \end{pmatrix} \neq \begin{pmatrix} 3 & 2 \\ 2 & 2 \end{pmatrix} \text{ or } \begin{pmatrix} \mu_{a_1 b_1 c_2} & \mu_{a_1 b_2 c_2} \\ \mu_{a_2 b_1 c_2} & \mu_{a_2 b_2 c_2} \end{pmatrix} \neq \begin{pmatrix} 2 & 2 \\ 2 & 3 \end{pmatrix}.$$

Proof. If X does not have the $Hyp_6(\star)$ -property then there exist six planes, say $\mathcal{L}(A_1), \mathcal{L}(A_2), \mathcal{L}(B_1), \mathcal{L}(B_2), \mathcal{L}(C_1), \mathcal{L}(C_2)$, such that the lines $\mathcal{L}(A_1, B_1), \mathcal{L}(A_1, B_2), \mathcal{L}(A_1, C_1), \mathcal{L}(A_2, B_2), \mathcal{L}(A_2, C_1), \mathcal{L}(A_2, C_2), \mathcal{L}(B_1, C_1), \mathcal{L}(B_1, C_2), \mathcal{L}(B_2, C_2)$ belong to X and $\mathcal{L}(A_2, B_1), \mathcal{L}(B_2, C_1), \mathcal{L}(A_1, C_2) \notin X$. Then we have that

$$\begin{pmatrix} \mu_{111} & \mu_{121} \\ \mu_{211} & \mu_{221} \end{pmatrix} = \begin{pmatrix} 3 & 2 \\ 2 & 2 \end{pmatrix} \text{ and } \begin{pmatrix} \mu_{112} & \mu_{122} \\ \mu_{212} & \mu_{222} \end{pmatrix} = \begin{pmatrix} 2 & 2 \\ 2 & 3 \end{pmatrix}.$$

On the other hand if $\begin{pmatrix} \mu_{111} & \mu_{121} \\ \mu_{211} & \mu_{221} \end{pmatrix} = \begin{pmatrix} 3 & 2 \\ 2 & 2 \end{pmatrix}$ and $\begin{pmatrix} \mu_{112} & \mu_{122} \\ \mu_{212} & \mu_{222} \end{pmatrix} = \begin{pmatrix} 2 & 2 \\ 2 & 3 \end{pmatrix}$ then it is easy to check that X does not have the $Hyp_6(\star)$ -property since the lines $\mathcal{L}(A_1, B_1), \mathcal{L}(A_1, B_2), \mathcal{L}(A_1, C_1), \mathcal{L}(A_2, B_2), \mathcal{L}(A_2, C_1), \mathcal{L}(A_2, C_2), \mathcal{L}(B_1, C_1), \mathcal{L}(B_1, C_2), \mathcal{L}(B_2, C_2) \in X$ and $\mathcal{L}(A_2, B_1), \mathcal{L}(B_2, C_1), \mathcal{L}(A_1, C_2) \notin X$. \square

Proposition 3.7. *Let X be a variety of lines. X has the $Hyp_5(\star)$ -property iff for all $a_1, a_2 \in [d_1], b_1, b_2 \in [d_2], c_1, c_2 \in [d_3]$ the following three conditions hold:*

- 1) either $\begin{pmatrix} \mu_{a_1 b_1 c_1} & \mu_{a_1 b_2 c_1} \\ \mu_{a_2 b_1 c_1} & \mu_{a_2 b_2 c_1} \end{pmatrix} \neq \begin{pmatrix} 2 & 1 \\ 2 & 2 \end{pmatrix}$ or $\begin{pmatrix} \mu_{a_1 b_1 0} & \mu_{a_1 b_2 0} \\ \mu_{a_2 b_1 0} & \mu_{a_2 b_2 0} \end{pmatrix} \neq \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$,
- 2) either $\begin{pmatrix} \mu_{a_1 b_1 c_2} & \mu_{a_1 b_2 c_2} \\ \mu_{a_2 b_1 c_2} & \mu_{a_2 b_2 c_2} \end{pmatrix} \neq \begin{pmatrix} 2 & 1 \\ 2 & 2 \end{pmatrix}$ or $\begin{pmatrix} \mu_{a_1 0 c_1} & \mu_{a_1 0 c_2} \\ \mu_{a_2 0 c_1} & \mu_{a_2 0 c_2} \end{pmatrix} \neq \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$,
- 3) either $\begin{pmatrix} \mu_{a_1 b_1 c_1} & \mu_{a_1 b_2 c_1} \\ \mu_{a_1 b_2 c_1} & \mu_{a_1 b_2 c_2} \end{pmatrix} \neq \begin{pmatrix} 2 & 1 \\ 2 & 2 \end{pmatrix}$ or $\begin{pmatrix} \mu_{0 b_1 c_1} & \mu_{0 b_1 c_2} \\ \mu_{0 b_2 c_1} & \mu_{0 b_2 c_2} \end{pmatrix} \neq \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Proof. If X does not have the $Hyp_5(\star)$ -property, we say, without loss of generality, that exist five planes $\mathcal{L}(A_1), \mathcal{L}(A_2), \mathcal{L}(B_1), \mathcal{L}(B_2), \mathcal{L}(C_1)$ such that, among all, only the lines $\mathcal{L}(A_2, B_1), \mathcal{L}(A_1, C_1), \mathcal{L}(B_2, C_1) \notin X$. Then we have

$$\begin{pmatrix} \mu_{111} & \mu_{121} \\ \mu_{211} & \mu_{221} \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 2 & 2 \end{pmatrix} \text{ and } \begin{pmatrix} \mu_{110} & \mu_{120} \\ \mu_{210} & \mu_{220} \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

On the other hand, assume, for instance, we have the following equalities

$$\begin{pmatrix} \mu_{111} & \mu_{121} \\ \mu_{211} & \mu_{221} \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 2 & 2 \end{pmatrix} \text{ and } \begin{pmatrix} \mu_{110} & \mu_{120} \\ \mu_{210} & \mu_{220} \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

From the previous equalities, we get $\mathcal{L}(A_1, C_1) \notin X$ thus, since $\mu_{111} = 2$, we have $\mathcal{L}(A_1, B_1), \mathcal{L}(B_1, C_1) \in X$. Analogously $\mathcal{L}(A_1, B_2) \in X$ and so, since $\mu_{121} = 1$, we have $\mathcal{L}(B_2, C_1) \notin X$. Moreover $\mathcal{L}(A_2, B_2) \in X$ and so, since $\mu_{221} = 2$, we have $\mathcal{L}(A_2, C_1) \in X$. Finally $\mathcal{L}(A_2, B_1) \notin X$ since $\mu_{210} = 0$. So X does not have the $Hyp_5(\star)$ -property. \square

Proposition 3.8. *Let X be a variety of lines. X has the $Hyp_4(\star)$ -property iff for all $a_1, a_2 \in [d_1], b_1, b_2 \in [d_2], c_1, c_2 \in [d_3]$ the following three conditions hold:*

- 1) either $\begin{pmatrix} \mu_{a_1 b_1 c_1} \\ \mu_{a_2 b_1 c_1} \end{pmatrix} \neq \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ or $\begin{pmatrix} \mu_{a_1 b_1 0} \\ \mu_{a_2 b_1 0} \end{pmatrix} \neq \begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

$$2) \text{ either } \begin{pmatrix} \mu_{a_1 b_1 c_1} \\ \mu_{a_1 b_1 c_2} \end{pmatrix} \neq \begin{pmatrix} 1 \\ 1 \end{pmatrix} \text{ or } \begin{pmatrix} \mu_{a_1 0 c_1} \\ \mu_{a_1 0 c_2} \end{pmatrix} \neq \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

$$3) \text{ either } \begin{pmatrix} \mu_{a_1 b_1 c_1} \\ \mu_{a_1 b_2 c_1} \end{pmatrix} \neq \begin{pmatrix} 1 \\ 1 \end{pmatrix} \text{ or } \begin{pmatrix} \mu_{0 b_1 c_1} \\ \mu_{0 b_2 c_1} \end{pmatrix} \neq \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Proof. Suppose that X does not have the $Hyp_4(\star)$ -property. Since we are assuming do not exist four planes $\mathcal{L}(A_1), \mathcal{L}(A_2), \mathcal{L}(B_1), \mathcal{L}(B_2)$ such that $\mathcal{L}(A_1, B_1), \mathcal{L}(A_2, B_2) \in X$ and $\mathcal{L}(A_1, B_2)$ or $\mathcal{L}(A_2, B_1) \notin X$, then, X fails the $Hyp_4(\star)$ -property if, without loss of generality, there exist four planes $\mathcal{L}(A_1), \mathcal{L}(A_2), \mathcal{L}(B_1), \mathcal{L}(C_1)$ such that, among all, only the lines $\mathcal{L}(A_2, B_1), \mathcal{L}(B_1, C_1), \mathcal{L}(A_1, C_1) \notin X$. Then we have

$$\begin{pmatrix} \mu_{111} \\ \mu_{211} \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \text{ and } \begin{pmatrix} \mu_{110} \\ \mu_{210} \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

On the other hand, assume, for instance, we have the following equalities

$$\begin{pmatrix} \mu_{111} \\ \mu_{211} \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \text{ and } \begin{pmatrix} \mu_{110} \\ \mu_{210} \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

From the previous equalities, we get $\mathcal{L}(A_1, B_1) \in X$ thus, since $\mu_{111} = 1$, we have $\mathcal{L}(A_1, C_1), \mathcal{L}(B_1, C_1) \notin X$. Analogously $\mathcal{L}(A_2, B_1) \notin X$ and so, since $\mu_{211} = 1$, we have $\mathcal{L}(A_2, C_1) \in X$. So X does not have the $Hyp_4(\star)$ -property. \square

Example 3.9. Let X be as in Example 3.5 (see Figure 4). We observe that

$$\begin{pmatrix} \mu_{111} & \mu_{121} \\ \mu_{211} & \mu_{221} \end{pmatrix} = \begin{pmatrix} 3 & 2 \\ 2 & 2 \end{pmatrix} \text{ and } \begin{pmatrix} \mu_{112} & \mu_{122} \\ \mu_{212} & \mu_{222} \end{pmatrix} = \begin{pmatrix} 2 & 2 \\ 2 & 3 \end{pmatrix}$$

and then, by Proposition 3.6, we have that X does not have the $Hyp_6(\star)$ -property and so, by Theorem 2.17, X is not ACM.

Example 3.10. Let us consider the variety $W = X \cup \mathcal{L}(A_2, B_1)$, where X is as Example 3.5.




FIGURE 5. The variety of lines W (in bold).

We have $\mu_{111} = \mu_{222} = \mu_{211} = \mu_{212} = 3$, $\mu_{121} = \mu_{221} = \mu_{112} = \mu_{122} = 2$. And for all $a_1, a_2 \in [2]$, $b_1, b_2 \in [2]$, $c_1, c_2 \in [2]$, we have:

$$\begin{pmatrix} \mu_{a_1 b_1 c_1} & \mu_{a_1 b_2 c_1} \\ \mu_{a_2 b_1 c_1} & \mu_{a_2 b_2 c_1} \end{pmatrix} \neq \begin{pmatrix} 3 & 2 \\ 2 & 2 \end{pmatrix}, \begin{pmatrix} \mu_{a_1 b_1 c_1} & \mu_{a_1 b_2 c_1} \\ \mu_{a_2 b_1 c_1} & \mu_{a_2 b_2 c_1} \end{pmatrix} \neq \begin{pmatrix} 2 & 1 \\ 2 & 2 \end{pmatrix},$$

$$\begin{pmatrix} \mu_{a_1 b_1 c_1} & \mu_{a_1 b_1 c_2} \\ \mu_{a_2 b_1 c_1} & \mu_{a_2 b_1 c_2} \end{pmatrix} \neq \begin{pmatrix} 2 & 1 \\ 2 & 2 \end{pmatrix}, \begin{pmatrix} \mu_{a_1 b_1 c_1} & \mu_{a_1 b_1 c_2} \\ \mu_{a_1 b_2 c_1} & \mu_{a_1 b_2 c_2} \end{pmatrix} \neq \begin{pmatrix} 2 & 1 \\ 2 & 2 \end{pmatrix},$$

$$\begin{pmatrix} \mu_{a_1 b_1 c_1} \\ \mu_{a_2 b_1 c_1} \end{pmatrix} \neq \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} \mu_{a_1 b_1 c_1} \\ \mu_{a_1 b_1 c_2} \end{pmatrix} \neq \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} \mu_{a_1 b_1 c_1} \\ \mu_{a_1 b_2 c_1} \end{pmatrix} \neq \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

and then, by Propositions 3.6, 3.7 and 3.8, the variety of lines W has the $Hypp_n(\star)$ -property for $n = 4, 5, 6$ and then, by Theorem 2.17, W is ACM.

4. THE HILBERT FUNCTION OF ACM CODIMENSION TWO VARIETIES IN $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$

In this section we approach the study of the Hilbert function of these varieties. We start from the following specific case.

Definition 4.1. If X is a variety of lines such that the index sets $U_1(X), U_2(X)$ and $U_3(X)$ are Ferrers diagram, then we call X a *Ferrers variety of lines*. That is, after renaming, we assume that if $\mathcal{L}(A_i, B_j) \in U_h(X)$ then $\mathcal{L}(A_{i'}, B_{j'}) \in U_h(X)$ for every $1 \leq i' \leq i, 1 \leq j' \leq j$ and for each direction $h = 1, 2, 3$.

Remark 4.2. As a consequence of Theorem 2.17, note that a Ferrers variety of lines is ACM.

Recall that given a homogeneous ideal $I \subseteq R$, the *Hilbert function of R/I* is the numerical function

$$H_{R/I} : \mathbb{N}^3 \rightarrow \mathbb{N}$$

defined by

$$H_{R/I}(i, j, k) := \dim_K(R/I)_{i,j,k} = \dim_K R_{i,j,k} - \dim_K I_{i,j,k}.$$

The *first difference function of H* , denoted ΔH , is the function $\Delta H : \mathbb{N}^3 \rightarrow \mathbb{N}$ defined by

$$\Delta H(i, j, k) := \sum_{(0,0,0) \leq (l,m,n) \leq (1,1,1)} (-1)^{l+m+n} H(i-l, j-m, k-n).$$

Now, let X be a Ferrers variety of lines and let $X_3 = \bigcup_{(r,s) \in U_3(X)} \mathcal{L}(A_r, B_s)$ be the variety of lines consisting of the lines of X of type $(1, 1, 0)$. Since $U_3(X)$ is a Ferrers diagram, the variety X_3 is ACM (in $\mathbb{P}^1 \times \mathbb{P}^1$) and we can explicitly write out a set of minimal generators of I_{X_3} , see Remark 1.14 and [19]. If $\{(a_{3,i}, b_{3,i})\}$ is the set of the degrees of these minimal generators, we denote by $D_3(X) := \{(a_{3,i}, b_{3,i}, 0)\}$.

Analogously, if we consider the varieties of lines X_1 and X_2 consisting of the lines of X of types $(0, 1, 1)$ and $(1, 0, 1)$, respectively, we obtain the sets of degrees $D_1(X) = \{(0, b_{1,j}, c_{1,j})\}$ and $D_2(X) = \{(a_{2,k}, 0, c_{2,k})\}$. Then we denote by

$$D(X) := \{(\max\{a_{3,i}, a_{2,k}\}, \max\{b_{3,i}, b_{1,j}\}, \max\{c_{1,j}, c_{2,k}\}) \mid \forall (a_{3,i}, b_{3,i}, 0) \in D_3(X), (a_{2,k}, 0, c_{2,k}) \in D_2(X), (0, b_{1,j}, c_{1,j}) \in D_1(X)\}.$$

Finally, we denote by $\hat{D}(X)$ the set of the minimal elements of $D(X)$ with respect to the natural partial order \preceq on the elements of \mathbb{N}^3 .

Theorem 4.3. *Let X be a Ferrers variety of lines. Then I_X is minimally generated by the following set of forms*

$$\left\{ \prod_{i \leq a} A_i \prod_{j \leq b} B_j \prod_{k \leq c} C_k \mid \text{for each } (a, b, c) \in \hat{D}(X) \right\}.$$

Proof. First, we prove that if $(a, b, c) \in \hat{D}(X)$, then $\prod_{i \leq a} A_i \prod_{j \leq b} B_j \prod_{k \leq c} C_k \in I_X$. Indeed $(a, b, c) \in \hat{D}(X)$ implies $\prod_{i \leq a} A_i \prod_{j \leq b} B_j \in I_{X_3}$, $\prod_{j \leq b} B_j \prod_{k \leq c} C_k \in I_{X_1}$, $\prod_{i \leq a} A_i \prod_{k \leq c} C_k \in I_{X_2}$ and they are not necessarily minimal elements of the respective ideal. Thus $\prod_{i \leq a} A_i \prod_{j \leq b} B_j \prod_{k \leq c} C_k \in I_{X_1} \cap I_{X_2} \cap I_{X_3} = I_X$. Now, we show that if $(a, b, c) \in \hat{D}(X)$ and $a > 0$, then $\prod_{i \leq a-1} A_i \prod_{j \leq b} B_j \prod_{k \leq c} C_k \notin I_X$. This fact follows by contradiction. Indeed if $\prod_{i \leq a-1} A_i \prod_{j \leq b} B_j \prod_{k \leq c} C_k \in I_X$, then $(a-1, b, 0)$, $(a-1, 0, c)$, $(0, b, c)$ are degrees of some (not necessarily minimal) elements in the ideal and therefore there is an element in $D(X)$ less than or equal to $(a-1, b, c)$, contradicting the minimality of $(a, b, c) \in \hat{D}(X)$. Analogously, it can be easily showed that if $(a, b, c) \in \hat{D}(X)$ and $b > 0$ (or $c > 0$), then $\prod_{i \leq a} A_i \prod_{j \leq b-1} B_j \prod_{k \leq c} C_k \notin I_X$ (or $\prod_{i \leq a} A_i \prod_{j \leq b} B_j \prod_{k \leq c-1} C_k \notin I_X$). Finally, we claim that I_X is minimally generated by the forms $\prod_{i \leq a} A_i \prod_{j \leq b} B_j \prod_{k \leq c} C_k$ with $(a, b, c) \in \hat{D}(X)$. Take a form $F \in I_X$, without loss of generality we can assume that $F := \prod_{i \in \mathcal{A}} A_i \prod_{j \in \mathcal{B}} B_j \prod_{k \in \mathcal{C}} C_k$ is product of linear forms. By contradiction we assume A_i divides F and A_{i-1} does not divide F . Then $\prod_{i \in \mathcal{A}} A_i \prod_{j \in \mathcal{B}} B_j \in I_{X_3}$. Then $F \in (\prod_{i \leq a'} A_i \prod_{j \leq b'} B_j)$ for some a', b' . Repeating the same argument with respect to the other two directions we get the proof. The minimality come from the minimality of the degrees in $\hat{D}(X)$. \square

The following corollary is an immediate consequence of Theorem 4.3 and the ACM property. Set $\langle \hat{D}(X) \rangle := \{(i, j, k) \mid (i, j, k) \geq (a, b, c), \text{ for some } (a, b, c) \in \hat{D}(X)\}$.

Corollary 4.4. *Let X be a Ferrers variety of lines. Then*

$$\Delta H_X(i, j, k) = \begin{cases} 0 & \text{if } (i, j, k) \in \langle \hat{D}(X) \rangle \\ 1 & \text{otherwise} \end{cases}.$$

Example 4.5. Let us consider the following variety of lines

$$X = \{\mathcal{L}(A_i, B_j) \cup \mathcal{L}(A_i, C_k) \cup \mathcal{L}(B_j, C_k) \mid 1 \leq i \leq 4, 1 \leq j \leq 3, 1 \leq k \leq 2\}.$$

In this case we have $D_3(X) = \{(4, 0, 0), (0, 3, 0)\}$, $D_2(X) = \{(4, 0, 0), (0, 0, 2)\}$ and $D_1(X) = \{(0, 3, 0), (0, 0, 2)\}$. Then $D(X) = \{(4, 3, 2), (4, 3, 0), (4, 0, 2), (0, 3, 2)\}$ and $\hat{D}(X) = \{(4, 3, 0), (4, 0, 2), (0, 3, 2)\}$. Therefore, from Theorem 4.3, a minimal set of generators of I_X is given by:

$$A_1 A_2 A_3 A_4 B_1 B_2 B_3, A_1 A_2 A_3 A_4 C_1 C_2, B_1 B_2 B_3 C_1 C_2$$

and

$$\Delta H_X(i, j, k) = \begin{cases} 0 & \text{if } (i, j, k) \geq (4, 3, 0) \text{ or } (4, 0, 2) \text{ or } (0, 3, 2) \\ 1 & \text{otherwise} \end{cases}.$$

5. CASE STUDY: GRIDS OF LINES AND COMPLETE INTERSECTIONS OF LINES

In the last section we focus on the study of special arrangements of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ having the ACM property. Recall that for a point $P \in \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ there are exactly three lines passing through P , one for each direction. We have the following definition.

Definition 5.1. Let \mathcal{Y} be a finite set of points in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. We call *grid of lines* arising from \mathcal{Y} , and denote it by $X_{\mathcal{Y}}$, the set containing all the lines of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ passing through some point of \mathcal{Y} .

In other words, if \mathcal{Y} is a finite set of points in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, then

$$X_{\mathcal{Y}} := \bigcup_{P_{ijk} \in \mathcal{Y}} \mathcal{L}(A_i, B_j) \cup \mathcal{L}(A_i, C_k) \cup \mathcal{L}(B_j, C_k)$$

where $P_{ijk} := \mathcal{L}(A_i) \cap \mathcal{L}(B_j) \cap \mathcal{L}(C_k)$.

The next example shows that, even if \mathcal{Y} is an ACM set of points, $X_{\mathcal{Y}}$ could be not ACM.

Example 5.2. Suppose $\mathcal{Y} := \{P_{112}, P_{122}, P_{121}, P_{212}\} \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. According to [12], \mathcal{Y} is an ACM set of points. We have $\mathcal{L}(A_2, B_1), \mathcal{L}(B_2, C_1) \in X_{\mathcal{Y}}$ and $\mathcal{L}(A_2, B_2), \mathcal{L}(A_2, C_1), \mathcal{L}(B_1, C_1) \notin X_{\mathcal{Y}}$, that is, $X_{\mathcal{Y}}$ has not the $Hyp_4(\star)$ -property and then $X_{\mathcal{Y}}$ is not ACM.

It is interesting to ask which sets of points $\mathcal{Y} \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ lead to an ACM grid of lines $X_{\mathcal{Y}}$. A special class of CM rings is represented by complete intersections. We recall their definitions and properties.

Definition 5.3. An ideal $I \subset R$ is a *complete intersection* if it is generated by a regular sequence.

As pointed out in [19], Lemma 2.25 a complete intersection is also Cohen-Macaulay.

Definition 5.4. In $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, we say that a set of points \mathcal{C} is a *complete intersection of points of type (a_1, a_2, a_3)* if $I_{\mathcal{C}} = (F_1, F_2, F_3)$ is a complete intersection and $\deg F_i = a_i \mathbf{e}_i$ for $i = 1, 2, 3$.

Note that each F_i in Definition 5.4 is product of linear forms.

Definition 5.5. We say that a variety of lines X is a *complete intersection of lines* in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ if I_X is a complete intersection.

Theorem 5.6. Let $\mathcal{C} \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ be a complete intersection of points of type (a, b, c) . Then $X_{\mathcal{C}}$ is ACM and a trigraded minimal free resolution of $I_{X_{\mathcal{C}}}$ is

$$0 \rightarrow R^2(-a, -b, -c) \rightarrow R(-a, -b, 0) \oplus R(-a, 0, -c) \oplus R(0, -b, -c) \rightarrow I_{X_{\mathcal{C}}} \rightarrow 0.$$

Proof. The grid of lines $X := X_{\mathcal{C}}$ has the $Hyp_n(\star)$ -property for $n = 4, 5, 6$ and then, by Theorem 2.17, X is ACM. Moreover, by Corollary 2.3 the generators of I_X are product of linear forms, so

$$I_X = \left(\prod_{i \in [a]} A_i \prod_{j \in [b]} B_j, \prod_{i \in [a]} A_i \prod_{k \in [c]} C_k, \prod_{j \in [b]} B_j \prod_{k \in [c]} C_k \right).$$

Then a Hilbert-Burch matrix of I_X is

$$\begin{pmatrix} \prod_{i \in [a]} A_i & \prod_{i \in [a]} A_i \\ \prod_{j \in [b]} B_j & 0 \\ 0 & \prod_{k \in [c]} C_k \end{pmatrix}.$$

□

Example 5.7. If $\mathcal{C} \subseteq \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ is a complete intersection of points of type $(2, 3, 2)$, then the grid $X_{\mathcal{C}}$ is formed by 6 lines of type $(1, 1, 0)$, 4 lines of type $(1, 0, 1)$ and 6 lines of type $(0, 1, 1)$:




FIGURE 6. The grid of lines $X_{\mathcal{C}}$ arising from a CI of type $(2, 3, 2)$.

In particular, $I_{X_{\mathcal{C}}}$ has a trigraded minimal free resolution of the following type

$$0 \rightarrow R^2(-2, -3, -2) \rightarrow R(-2, -3, 0) \oplus R(-2, 0, -2) \oplus R(0, -3, -2) \rightarrow I_{X_{\mathcal{C}}} \rightarrow 0.$$

The following example shows that there exists an ACM grid of lines $X_{\mathcal{Y}}$ arising from a not ACM set of points \mathcal{Y} .

Example 5.8. The following set of points $\mathcal{Y} := \{P_{111}, P_{121}, P_{211}, P_{122}, P_{212}, P_{222}\}$ is not an ACM set of points in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ (see [12]). However, $X_{\mathcal{Y}} = X_{\mathcal{C}}$ where $\mathcal{C} := \{P_{ijk} \mid 1 \leq i, j, k \leq 2\}$, and then $X_{\mathcal{Y}}$ is an ACM grid of lines.

From Theorem 5.6, we note that the ideal $I_{X_{\mathcal{C}}}$ is generated by three forms that do not form a regular sequence. That is, even if \mathcal{C} is a complete intersection of points, then its associated $X_{\mathcal{C}}$ variety of lines is not a complete intersection of lines. Thus, it is natural to study which varieties of lines are defined by a complete intersection, i.e., their defining ideal has only two generators. Theorem 5.10 and Remark 5.11 will describe complete intersections of lines in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$.

Remark 5.9. If X is an ACM variety of lines, from Corollary 2.3, I_X is generated by products of linear forms. Then

$$I_X \supseteq \left(\prod_{i \in [a]} A_i \prod_{j \in [b]} B_j, \prod_{i \in [a]} A_i \prod_{k \in [c]} C_k, \prod_{j \in [b]} B_j \prod_{k \in [c]} C_k \right).$$

So any set of minimal generators of I_X contains one element of degree $(a_3, b_3, 0)$, one element of degree $(a_2, 0, c_2)$ and one element of degree $(0, b_1, c_1)$.

Theorem 5.10. *Let X be a variety of lines of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. Then the ideal I_X is a complete intersection if and only if $I_X = (F_1, F_2)$, with $\deg F_1 = ae_i$ and $\deg F_2 = be_j + ce_k$ with $j, k \neq i$, for some $a, b, c \in \mathbb{N}$.*

Proof. One implication is trivial. Let I_X be a complete intersection, i.e. I_X is generated by a regular sequence of length 2, then X is ACM. So, from Remark 5.9, any set of minimal generators of I_X contains one element G_1 of degree $(0, b_1, c_1)$, one element G_2 of degree $(a_2, 0, c_2)$ and one element G_3 of degree $(a_3, b_3, 0)$ for some integers a_i, b_j, c_k . Since I_X is a complete intersection, one of these three generators say, without loss of generality, the one of degree $(0, b_1, c_1)$, is not minimal, i.e. $G_1 \in (G_2, G_3)$. This easily implies $a_2a_3 = 0$. \square

Remark 5.11. From Theorem 5.10, a complete intersection of lines X is then obtained from a grid arising from a complete intersection of points by removing either all the lines having direction \mathbf{e}_i for some i , or all the lines having direction \mathbf{e}_i and \mathbf{e}_j with $i \neq j$. Indeed, from Remark 5.9, we have for instance

$$I_X = \left(\prod_{i \in [a]} A_i, \prod_{j \in [b]} B_j, \prod_{k \in [c]} C_k \right) = \bigcap_{\substack{i \in [a] \\ j \in [b]}} (A_i, B_j) \cap \bigcap_{\substack{i \in [a] \\ k \in [c]}} (A_i, C_k).$$

Example 5.12. Let X be the set of lines of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ obtained by a grid of lines X_C arising from a complete intersection C of type $(4, 3, 2)$ removing all the lines having direction \mathbf{e}_2 :

$$X = \bigcup_{\substack{i \in [4] \\ j \in [3]}} \mathcal{L}(A_i, B_j) \bigcup_{\substack{j \in [3] \\ k \in [2]}} \mathcal{L}(B_j, C_k).$$

Then the ideal I_X is a complete intersection and it is generated by the regular sequence $F_1 = B_1B_2B_3$ and $F_2 = A_1A_2A_3A_4C_1C_2$ of degree $(0, 3, 0)$ and $(4, 0, 2)$, respectively.

We end the paper with two research topics that are still under our investigation.

- (1) Guida, Orecchia and Ramella, in [21], studied the *complete grids* of lines in \mathbb{P}^3 , whose defining ideal is the 1-lifting ideal of a specific monomial ideal J in a polynomial ring S in three variables. In particular, from Example 4.9 in [21] and Corollary 4.4, we noted that the first difference of the Hilbert function of the ideal I_{X_C} of a grid of lines arising from a complete intersection of points of type $(2, 2, 2)$ in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ in degree (i, j, k) is equal to 1 if and only if (i, j, k) belongs to the order ideal $N(J) \subseteq \mathbb{N}^3$ of the specific monomial ideal $J = (x_1^2x_2^2, x_1^2x_3^2, x_2^2x_3^2)$ in S .
- (2) Let us consider the ACM varieties of lines X and the Ferrers variety of lines X' as in Figure 7 and Figure 8, respectively. We have that, for each $h = 1, 2, 3$, X_h and X'_h have the same Hilbert functions. We also get $H_X = H_{X'}$.




FIGURE 7

FIGURE 8

According to many experimental computations using CoCoA, [1], we ask the following question:

Question 5.13. Let X be an ACM variety of lines and X' be a Ferrers variety of lines such that, for $h = 1, 2, 3$, X_h and X'_h have the same Hilbert functions. Is it true that $H_X = H_{X'}$?

REFERENCES

- [1] J. Abbott, A. M. Bigatti, L. Robbiano *CoCoA: a system for doing Computations in Commutative Algebra*. Available at <http://cocoa.dima.unige.it>
- [2] E. Ballico, A. Bernardi, M.V. Catalisano, *Higher Secant Varieties of $\mathbb{P}^n \times \mathbb{P}^1$ embedded in bi-degree (a, b)* . *Commutative Algebra* 40 (2012), pp 3822–3840.
- [3] E. Ballico, A. Bernardi, L. Chiantini, E. Guardo, *Bounds on tensor rank*. (2017) <https://arxiv.org/pdf/1705.02299>.
- [4] M.V. Catalisano, A.V. Geramita, A. Gimigliano, *Ranks of tensors, secant varieties of Segre varieties and fat points*. *Linear Algebra and Appl.* 355 (2002), pp 263–285.
- [5] M.V. Catalisano, A.V. Geramita, A. Gimigliano, *Higher secant varieties of the Segre varieties $\mathbb{P}^1 \times \mathbb{P}^1$* . *J. Pure Appl. Algebra*, 201 (2005), 367–380.
- [6] L. Chiantini, D. Sacchi, *Segre Functions in Multiprojective spaces and tensor analysis*. *Trends Hist. Sci. 8 - From Classical to Modern Algebraic Geometry* (2016), 361–374.
- [7] E. Duarte, *Implicitization of tensor product surfaces in the presence of a generic set of base-points*. *Journal of Algebra and Its Applications*, to appear. <https://arxiv.org/pdf/1610.03820>.
- [8] D. Cook II, B. Harbourne, J. Migliore, U. Nagel, *Line arrangements and configurations of points with an unusual geometric property*. Preprint <https://arxiv.org/pdf/1602.02300>.
- [9] J.A. Eagon, V. Reiner, *Resolutions of Stanley-Reisner rings and Alexander duality*. *Journal of Pure and Applied Algebra*. 1998 Sep 17;130(3):265-75.
- [10] G. Favacchio *The Hilbert function of bigraded algebras in $k[\mathbb{P}^1 \times \mathbb{P}^1]$* . *Journal of Commutative Algebra*. Accepted for publication (2017).
- [11] G. Favacchio, E. Guardo, *The minimal free resolution of fat almost complete intersections in $\mathbb{P}^1 \times \mathbb{P}^1$* . *Canadian Journal of Mathematics* 69(2017), no. 6, 1274-1291.
- [12] G. Favacchio, E. Guardo, J. Migliore. *On the arithmetically Cohen-Macaulay property for sets of points in multiprojective spaces*. *Proceedings of the American Mathematical Society*. Accepted for publication. (2017) DOI: <https://doi.org/10.1090/proc/13981>
- [13] G. Favacchio, J. Migliore. *Multiprojective spaces and arithmetically Cohen Macaulay property*. *Mathematical Proceedings of the Cambridge Philosophical Society*. Accepted for publication. (2018)
- [14] G. Favacchio, A. Ragusa, G. Zappalá, *Tower sets and other configurations with the Cohen-Macaulay property*. *Journal of Pure and Applied Algebra*. 2015 Jun 30;219(6):2260-78.
- [15] R. Fröberg, *On Stanley-Reisner rings*. *Banach Center Publications*. 1990; 26(2):57-70.

- [16] E. Guardo, A. Van Tuyl, *Fat Points in $\mathbb{P}^1 \times \mathbb{P}^1$ and their Hilbert functions*. Canad. J. Math. 56 (2004), no. 4, 716–741.
- [17] E. Guardo, A. Van Tuyl, *ACM sets of points in multiprojective space*. Collect. Math. 59 (2008), no. 2, 191–213.
- [18] E. Guardo, A. Van Tuyl, *Classifying ACM sets of points in $\mathbb{P}^1 \times \mathbb{P}^1$ via separators*. Arch. Math. (Basel) 99 (2012), no. 1, 33–36.
- [19] E. Guardo, A. Van Tuyl, *Arithmetically Cohen-Macaulay sets of points in $\mathbb{P}^1 \times \mathbb{P}^1$* . Springer Briefs in Mathematics (2015).
- [20] E. Guardo, A. Van Tuyl, *On the Hilbert functions of sets of points in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$* . Mathematical Proceedings of the Cambridge Philosophical Society. Vol. 159. No. 01. Cambridge University Press, 2015.
- [21] M. Guida, F. Orecchia, L. Ramella, *Some configurations of lines whose ideals are liftings of monomial ideals*. International Journal of Pure and Applied Mathematics. 2014; 92(5):669–690.
- [22] B. Harbourne, *Asymptotics of linear systems, with connections to line arrangements*. Preprint 2017, <https://arxiv.org/pdf/1705.09946>.
- [23] J. Herzog, T. Hibi, *Monomial ideals*. Graduate Texts in Mathematics. 2011.
- [24] M. Naeem, *Cohen-Macaulay monomial ideals of codimension 2*. Manuscripta Mathematica. 2008 Dec 1;127(4):533–45.
- [25] A. Van Tuyl, *A beginner’s guide to edge and cover ideals*. Monomial ideals, computations and applications, 63–94, Lecture Notes in Math., 2083, Springer, Heidelberg, 2013.
- [26] A. Van Tuyl *The Hilbert functions of ACM sets of points in $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k}$* . Journal of Algebra. 2003 Jun 15;264(2):420–41.

DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIALE A. DORIA, 6 - 95100 - CATANIA, ITALY
E-mail address: favacchio@dmf.unict.it
URL: <http://www.dmf.unict.it/~gfavacchio/>

DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIALE A. DORIA, 6 - 95100 - CATANIA, ITALY
E-mail address: guardo@dmf.unict.it
URL: <http://www.dmf.unict.it/~guardo/>

DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIALE A. DORIA, 6 - 95100 - CATANIA, ITALY
E-mail address: picone@dmf.unict.it