

On the least common multiple of random q-integers

Original

On the least common multiple of random q-integers / Sanna, Carlo. - In: RESEARCH IN NUMBER THEORY. - ISSN 2363-9555. - STAMPA. - 7:1(2021). [10.1007/s40993-021-00242-4]

Availability:

This version is available at: 11583/2872308 since: 2021-02-23T18:53:52Z

Publisher:

Springer

Published

DOI:10.1007/s40993-021-00242-4

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

RESEARCH



On the least common multiple of random q -integers

Carlo Sanna* 

*Correspondence:
carlo.sanna.dev@gmail.com
Department of Mathematical
Sciences, Politecnico di Torino,
Corso Duca degli Abruzzi 24,
10129 Torino, Italy
C. Sanna is a member of GNSAGA
of INdAM and of CryptTO, the
group of Cryptography and
Number Theory of Politecnico di
Torino

Abstract

For every positive integer n and for every $\alpha \in [0, 1]$, let $\mathcal{B}(n, \alpha)$ denote the probabilistic model in which a random set $\mathcal{A} \subseteq \{1, \dots, n\}$ is constructed by picking independently each element of $\{1, \dots, n\}$ with probability α . Cilleruelo, Rué, Šarka, and Zumalacárregui proved an almost sure asymptotic formula for the logarithm of the least common multiple of the elements of \mathcal{A} . Let q be an indeterminate and let $[k]_q := 1 + q + q^2 + \dots + q^{k-1} \in \mathbb{Z}[q]$ be the q -analogue of the positive integer k . We determine the expected value and the variance of $X := \deg \text{lcm}([\mathcal{A}]_q)$, where $[\mathcal{A}]_q := \{[k]_q : k \in \mathcal{A}\}$. Then we prove an almost sure asymptotic formula for X , which is a q -analogue of the result of Cilleruelo et al.

Keywords: Asymptotic formula, Least common multiple, q -analogue, Random set

Mathematics Subject Classification: Primary: 11N37, Secondary: 11B99

1 Introduction

A nice consequence of the Prime Number Theorem is the asymptotic formula

$$\log \text{lcm}(1, 2, \dots, n) \sim n, \quad \text{as } n \rightarrow +\infty, \quad (1)$$

where lcm denotes the least common multiple. Indeed, precise estimates for $\log \text{lcm}(1, \dots, n)$ are equivalent to the Prime Number Theorem with an error term. Thus, a natural generalization is to study estimates for $L_f(n) := \log \text{lcm}(f(1), \dots, f(n))$, where f is a well-behaved function, for instance, a polynomial with integer coefficients. (We ignore terms equal to 0 in the lcm and we set $\text{lcm} \emptyset := 1$.) When $f \in \mathbb{Z}[x]$ is a linear polynomial, the product of linear polynomials, or an irreducible quadratic polynomial, asymptotic formulas for $L_f(n)$ were proved by Bateman et al. [3], Hong et al. [10], and Cilleruelo [6], respectively. In particular, for $f(x) = x^2 + 1$, Rué et al. [15] determined a precise error term for the asymptotic formula. When f is an irreducible polynomial of degree $d \geq 3$, Cilleruelo [6] conjectured that $L_f(n) \sim (d - 1)n \log n$, as $n \rightarrow +\infty$, but this is still an open problem. However, bounds for $L_f(n)$ were proved by Maynard and Rudnick [13], and Sah [16]. Moreover, Rudnick and Zehavi [14] studied the growth of $L_f(n)$ along a shifted family of polynomials.

Another direction of research consists in considering the least common multiple of random sets of positive integers. For every positive integer n and every $\alpha \in [0, 1]$, let

$\mathcal{B}(n, \alpha)$ denote the probabilistic model in which a random set $\mathcal{A} \subseteq \{1, \dots, n\}$ is constructed by picking independently each element of $\{1, \dots, n\}$ with probability α . Cilleruelo et al. [9] studied the least common multiple of the elements of \mathcal{A} and proved the following result (see [1] for a more precise version, and [4, 5, 7, 8, 12, 17–19] for other results of a similar flavor).

Theorem 1.1 *Let \mathcal{A} be a random set in $\mathcal{B}(n, \alpha)$. Then, as $\alpha n \rightarrow +\infty$, we have*

$$\log \text{lcm}(\mathcal{A}) \sim \frac{\alpha \log(1/\alpha)}{1 - \alpha} \cdot n,$$

with probability $1 - o(1)$, where the factor involving α is meant to be equal to 1 for $\alpha = 1$.

Remark 1.1 In the deterministic case $\alpha = 1$, we have $\mathcal{A} = \{1, \dots, n\}$ (surely) and Theorem 1.1 corresponds to (1).

Let q be an indeterminate. The q -analog of a positive integer k is defined by

$$[k]_q := 1 + q + q^2 + \dots + q^{k-1} \in \mathbb{Z}[q].$$

The q -analogs of many other mathematical objects (factorial, binomial coefficients, hypergeometric series, derivative, integral...) have been extensively studied, especially in Analysis and Combinatorics [2, 11]. For every set \mathcal{S} of positive integers, let $[\mathcal{S}]_q := \{[k]_q : k \in \mathcal{S}\}$.

The aim of this paper is to study the least common multiple of the elements of $[\mathcal{A}]_q$ for a random set \mathcal{A} in $\mathcal{B}(n, \alpha)$. Our main results are the following:

Theorem 1.2 *Let \mathcal{A} be a random set in $\mathcal{B}(n, \alpha)$ and put $X := \deg \text{lcm}([\mathcal{A}]_q)$. Then, for every integer $n \geq 2$ and every $\alpha \in [0, 1]$, we have*

$$\mathbb{E}[X] = \frac{3}{\pi^2} \cdot \frac{\alpha \text{Li}_2(1 - \alpha)}{1 - \alpha} \cdot n^2 + O(\alpha n (\log n)^2), \tag{2}$$

where $\text{Li}_2(z) := \sum_{k=1}^{\infty} z^k/k^2$ is the dilogarithm and the factor involving α is meant to be equal to 1 when $\alpha = 1$. In particular,

$$\mathbb{E}[X] \sim \frac{3}{\pi^2} \cdot \frac{\alpha \text{Li}_2(1 - \alpha)}{1 - \alpha} \cdot n^2,$$

as $n \rightarrow +\infty$, uniformly for $\alpha \in [0, 1]$.

Theorem 1.3 *Let \mathcal{A} be a random set in $\mathcal{B}(n, \alpha)$ and put $X := \deg \text{lcm}([\mathcal{A}]_q)$. Then there exists a function $v : (0, 1) \rightarrow \mathbb{R}^+$ such that, as $\alpha n / ((\log n)^3 (\log \log n)^2) \rightarrow +\infty$, we have*

$$\mathbb{V}[X] = (v(\alpha) + o(1)) n^3. \tag{3}$$

Moreover, the upper bound

$$\mathbb{V}[X] \ll \alpha n^3, \tag{4}$$

holds for every positive integer n and every $\alpha \in [0, 1]$.

As a consequence of Theorems 1.2 and 1.3, we obtain the following q -analog of Theorem 1.1.

Theorem 1.4 *Let \mathcal{A} be a random set in $\mathcal{B}(n, \alpha)$. Then, as $\alpha n \rightarrow +\infty$, we have*

$$\deg \text{lcm}([\mathcal{A}]_q) \sim \frac{3}{\pi^2} \cdot \frac{\alpha \text{Li}_2(1-\alpha)}{1-\alpha} \cdot n^2,$$

with probability $1 - o(1)$, where the factor involving α is meant to be equal to 1 for $\alpha = 1$.

Remark 1.2 In the deterministic case $\alpha = 1$, we have (see Lemma 4.1 below)

$$\deg \text{lcm}([1, 2, \dots, n]_q) = \sum_{1 < d \leq n} \varphi(d),$$

and Theorem 1.4 corresponds to the well-known asymptotic formula $\sum_{d \leq n} \varphi(d) \sim \frac{3}{\pi^2} n^2$ (Lemma 3.3 below) for the sum of the first values of the Euler function φ .

Remark 1.3 In Theorem 1.4 the condition $\alpha n \rightarrow +\infty$ is necessary. Indeed, if $\alpha n \leq C$, for some constant $C > 0$, then

$$\mathbb{P}[\mathcal{A} = \emptyset] = (1 - \alpha)^n \geq \left(1 - \frac{C}{n}\right)^n \rightarrow e^{-C}$$

as $n \rightarrow +\infty$, and so no (nontrivial) asymptotic formula for $\deg \text{lcm}([\mathcal{A}]_q)$ can hold with probability $1 - o(1)$.

We conclude this section with some possible questions for further research on this topic. Alsmeyer, Kabluchko, and Marynych [1, Corollary 1.5] proved that, for fixed $\alpha \in [0, 1]$ and for a random set \mathcal{A} in $\mathcal{B}(n, \alpha)$, an appropriate normalization of the random variable $\log \text{lcm}(\mathcal{A})$ converges in distribution to a standard normal random variable, as $n \rightarrow +\infty$. In light of Theorems 1.2 and 1.3, it is then natural to ask whether the random variable

$$\frac{\deg \text{lcm}([\mathcal{A}]_q) - \frac{3}{\pi^2} \cdot \frac{\alpha \text{Li}_2(1-\alpha)}{1-\alpha} \cdot n^2}{\sqrt{v(\alpha)n^3}}$$

converges in distribution to a normal random variable, or to some other random variable.

Another problem could be considering polynomial values, similarly to the results done in the context of integers, and studying $\text{lcm}([f(1)]_q, \dots, [f(n)]_q)$ for $f \in \mathbb{Z}[x]$ or, more generally, $\text{lcm}([f(k)]_q : k \in \mathcal{A})$ with \mathcal{A} a random set in $\mathcal{B}(n, \alpha)$.

2 Notation

We employ the Landau–Bachmann “Big Oh” and “little oh” notations O and o , as well as the associated Vinogradov symbol \ll , with their usual meanings. Any dependence of the implied constants is explicitly stated or indicated with subscripts. For real random variables X and Y , depending on some parameters, we say that “ $X \sim Y$ with probability $1 - o(1)$ ”, as the parameters tend to some limit, if for every $\varepsilon > 0$ we have $\mathbb{P}[|X - Y| > \varepsilon|Y|] = o_\varepsilon(1)$, as the parameters tend to the limit. We let (a, b) and $[a, b]$ denote the greatest common divisor and the least common multiple, respectively, of two integers a and b . As usual, we write $\varphi(n)$, $\mu(n)$, $\tau(n)$, and $\sigma(n)$, for the Euler totient function, the Möbius function, the number of divisors, and the sum of divisors, of a positive integer n , respectively.

3 Preliminaries

In this section we collect some preliminary results needed in later arguments.

Lemma 3.1 *We have*

$$\sum_{m \leq x} \tau(m) \ll x \log x,$$

for every $x \geq 2$.

Proof See, e.g., [20, Part I, Theorem 3.2]. □

Lemma 3.2 *We have*

$$\sum_{[e_1, e_2] > x} \frac{1}{e_1 e_2 [e_1, e_2]} \ll \frac{\log x}{x},$$

for every $x \geq 2$.

Proof From Lemma 3.1 and partial summation, it follows that

$$\begin{aligned} \sum_{m > x} \frac{\tau(m)}{m^2} &= \left[\frac{\sum_{m \leq t} \tau(m)}{t^2} \right]_{t=x}^{+\infty} + 2 \int_x^{+\infty} \frac{\sum_{m \leq t} \tau(m)}{t^3} dt \\ &\ll \int_x^{+\infty} \frac{\log t}{t^2} dt = \left[-\frac{\log t + 1}{t} \right]_{t=x}^{+\infty} \ll \frac{\log x}{x}. \end{aligned}$$

Let $e := (e_1, e_2)$ and $e'_i := e_i/e$ for $i = 1, 2$. Then we have

$$\begin{aligned} \sum_{[e_1, e_2] > x} \frac{1}{e_1 e_2 [e_1, e_2]} &\leq \sum_{e \geq 1} \frac{1}{e^3} \sum_{e'_1 e'_2 > x/e} \frac{1}{(e'_1 e'_2)^2} = \sum_{e \geq 1} \frac{1}{e^3} \sum_{m > x/e} \frac{\tau(m)}{m^2} \\ &\ll \sum_{e \leq x/2} \frac{1}{e^3} \frac{\log(x/e)}{x/e} + \sum_{e > x/2} \frac{1}{e^3} \ll \frac{\log x}{x} + \frac{1}{x^2} \ll \frac{\log x}{x}, \end{aligned}$$

as desired. □

Let us define

$$\Phi(x) := \sum_{n \leq x} \varphi(n) \quad \text{and} \quad \Phi(a_1, a_2; x) := \sum_{n \leq x} \varphi(a_1 n) \varphi(a_2 n),$$

for every $x \geq 1$ and for all positive integers a_1, a_2 .

Lemma 3.3 *We have*

$$\Phi(x) = \frac{3}{\pi^2} x^2 + O(x \log x),$$

for every $x \geq 2$.

Proof See, e.g., [20, Part I, Theorem 3.4]. □

Lemma 3.4 *We have*

$$\Phi(a_1, a_2; x) = C_1(a_1, a_2) x^3 + O(\sigma(a_1) \sigma(a_2) x^2 (\log x)^2), \tag{5}$$

for every $x \geq 2$, where

$$C_1(a_1, a_2) := \frac{a_1 a_2}{3} \sum_{d_1, d_2 \geq 1} \frac{\mu(d_1) \mu(d_2)}{d_1 d_2 [d_1 / (a_1, d_1), d_2 / (a_2, d_2)]} \tag{6}$$

and the series is absolutely convergent.

Proof From the identity $\varphi(n)/n = \sum_{d|n} \mu(d)/d$, it follows that

$$\begin{aligned} \sum_{n \leq x} \frac{\varphi(a_1 n)}{a_1 n} \frac{\varphi(a_2 n)}{a_2 n} &= \sum_{n \leq x} \left(\sum_{d_1 | a_1 n} \frac{\mu(d_1)}{d_1} \sum_{d_2 | a_2 n} \frac{\mu(d_2)}{d_2} \right) \\ &= \sum_{\substack{d_1 \leq a_1 x \\ d_2 \leq a_2 x}} \frac{\mu(d_1)}{d_1} \frac{\mu(d_2)}{d_2} \#\{n \leq x : d_1 | a_1 n \text{ and } d_2 | a_2 n\} \\ &= \sum_{\left[\frac{d_1}{(a_1, d_1)}, \frac{d_2}{(a_2, d_2)} \right] \leq x} \frac{\mu(d_1)}{d_1} \frac{\mu(d_2)}{d_2} \left(\frac{x}{\left[d_1/(a_1, d_1), d_2/(a_2, d_2) \right]} + O(1) \right). \end{aligned}$$

Let $c_i := (a_i, d_i)$ and $e_i := d_i/c_i$, for $i = 1, 2$. On the one hand, we have

$$E_1 := \sum_{\left[\frac{d_1}{(a_1, d_1)}, \frac{d_2}{(a_2, d_2)} \right] \leq x} \frac{1}{d_1 d_2} \leq \sum_{c_1 | a_1} \frac{1}{c_1} \sum_{c_2 | a_2} \frac{1}{c_2} \sum_{e_1 \leq x} \frac{1}{e_1} \sum_{e_2 \leq x} \frac{1}{e_2} \ll \frac{\sigma(a_1) \sigma(a_2)}{a_1 a_2} (\log x)^2.$$

On the other hand, thanks to Lemma 3.2, we have

$$\begin{aligned} E_2 &:= \sum_{\left[\frac{d_1}{(a_1, d_1)}, \frac{d_2}{(a_2, d_2)} \right] > x} \frac{1}{d_1 d_2 \left[d_1/(a_1, d_1), d_2/(a_2, d_2) \right]} \\ &\leq \sum_{c_1 | a_1} \frac{1}{c_1} \sum_{c_2 | a_2} \frac{1}{c_2} \sum_{[e_1, e_2] > x} \frac{1}{e_1 e_2 [e_1, e_2]} \ll \frac{\sigma(a_1) \sigma(a_2) \log x}{a_1 a_2 x}, \end{aligned}$$

which, in particular, implies that the series

$$C_0(a_1, a_2) := \sum_{d_1, d_2 \geq 1} \frac{\mu(d_1) \mu(d_2)}{d_1 d_2 \left[d_1/(a_1, d_1), d_2/(a_2, d_2) \right]}$$

is absolutely convergent. Therefore, we obtain

$$\begin{aligned} \sum_{n \leq x} \frac{\varphi(a_1 n)}{a_1 n} \frac{\varphi(a_2 n)}{a_2 n} &= (C_0(a_1, a_2) + O(E_2)) x + O(E_1) \\ &= C_0(a_1, a_2) x + O\left(\frac{\sigma(a_1) \sigma(a_2)}{a_1 a_2} (\log x)^2 \right). \end{aligned} \tag{7}$$

Now (5) follows from (7) by partial summation and since $C_1(a_1, a_2) = \frac{a_1 a_2}{3} C_0(a_1, a_2)$. \square

Remark 3.1 The obvious bound $\varphi(m) \leq m$ yields $C_1(a_1, a_2) \leq \frac{a_1 a_2}{3}$ (which is not so obvious from (6)).

We end this section with an easy observation that will be useful later.

Remark 3.2 It holds $1 - (1 - x)^k \leq kx$, for all $x \in [0, 1]$ and for all integers $k \geq 0$.

4 Proofs

Henceforth, let \mathcal{A} be a random set in $\mathcal{B}(n, \alpha)$, let $[\mathcal{A}]_q$ be its q -analog, and put $L := \text{lcm}([\mathcal{A}]_q)$ and $X := \text{deg } L$. For every positive integer d , let us define

$$I_{\mathcal{A}}(d) := \begin{cases} 1 & \text{if } d | k \text{ for some } k \in \mathcal{A}; \\ 0 & \text{otherwise.} \end{cases}$$

The following lemma gives a formula for X in terms of $I_{\mathcal{A}}$ and the Euler function.

Lemma 4.1 *We have*

$$X = \sum_{1 < d \leq n} \varphi(d) I_{\mathcal{A}}(d). \tag{8}$$

Proof For every positive integer k , it holds

$$[k]_q = \frac{q^k - 1}{q - 1} = \prod_{\substack{d|k \\ d > 1}} \Phi_d(q),$$

where $\Phi_d(q)$ is the d th cyclotomic polynomials. Since, as it is well known, every cyclotomic polynomial is irreducible over \mathbb{Q} , it follows that L is the product of the polynomials $\Phi_d(q)$ such that $d > 1$ and $d | k$ for some $k \in \mathcal{A}$. Finally, the equality $\deg(\Phi_d(q)) = \varphi(d)$ and the definition of $I_{\mathcal{A}}$ yield (8). \square

Let $\beta := 1 - \alpha$. The next lemma provides two expected values involving $I_{\mathcal{A}}$.

Lemma 4.2 *For all positive integers d, d_1, d_2 , we have*

$$\mathbb{E}[I_{\mathcal{A}}(d)] = 1 - \beta^{\lfloor n/d \rfloor} \tag{9}$$

and

$$\mathbb{E}[I_{\mathcal{A}}(d_1)I_{\mathcal{A}}(d_2)] = 1 - \beta^{\lfloor n/d_1 \rfloor} - \beta^{\lfloor n/d_2 \rfloor} + \beta^{\lfloor n/d_1 \rfloor + \lfloor n/d_2 \rfloor - \lfloor n/[d_1, d_2] \rfloor}.$$

Proof On the one hand, by the definition of $I_{\mathcal{A}}$, we have

$$\mathbb{E}[I_{\mathcal{A}}(d)] = \mathbb{P}[\exists k \in \mathcal{A} : d | k] = 1 - \mathbb{P}\left[\bigwedge_{m \leq \lfloor n/d \rfloor} (dm \notin \mathcal{A})\right] = 1 - \beta^{\lfloor n/d \rfloor},$$

which is (9). On the other hand, by linearity of the expectation and by (9), we have

$$\begin{aligned} \mathbb{E}[I_{\mathcal{A}}(d_1)I_{\mathcal{A}}(d_2)] &= \mathbb{E}[I_{\mathcal{A}}(d_1) + I_{\mathcal{A}}(d_2) - 1 + (1 - I_{\mathcal{A}}(d_1))(1 - I_{\mathcal{A}}(d_2))] \\ &= \mathbb{E}[I_{\mathcal{A}}(d_1)] + \mathbb{E}[I_{\mathcal{A}}(d_2)] - 1 + \mathbb{E}[(1 - I_{\mathcal{A}}(d_1))(1 - I_{\mathcal{A}}(d_2))] \\ &= 1 - \beta^{\lfloor n/d_1 \rfloor} - \beta^{\lfloor n/d_2 \rfloor} + \mathbb{E}[(1 - I_{\mathcal{A}}(d_1))(1 - I_{\mathcal{A}}(d_2))], \end{aligned}$$

where the last expected value can be computed as

$$\begin{aligned} \mathbb{E}[(1 - I_{\mathcal{A}}(d_1))(1 - I_{\mathcal{A}}(d_2))] &= \mathbb{P}[\forall k \in \mathcal{A} : d_1 \nmid k \text{ and } d_2 \nmid k] \\ &= \mathbb{P}\left[\bigwedge_{\substack{k \leq n \\ d_1 | k \text{ or } d_2 | k}} (k \notin \mathcal{A})\right] = \beta^{\lfloor n/d_1 \rfloor + \lfloor n/d_2 \rfloor - \lfloor n/[d_1, d_2] \rfloor}, \end{aligned}$$

and second claim follows. \square

We are ready to compute the expected value of X .

Proof of Theorem 1.2 From Lemmas 4.1 and 4.2, it follows that

$$\mathbb{E}[X] = \sum_{1 < d \leq n} \varphi(d) \mathbb{E}[I_{\mathcal{A}}(d)] = \sum_{1 < d \leq n} \varphi(d)(1 - \beta^{\lfloor n/d \rfloor}). \tag{10}$$

Moreover, since $\lfloor n/d \rfloor = j$ if and only if $n/(j + 1) < d \leq n/j$, we get that

$$\begin{aligned} \sum_{d \leq n} \varphi(d)(1 - \beta^{\lfloor n/d \rfloor}) &= \sum_{j \leq n} (1 - \beta^j) \sum_{n/(j+1) < d \leq n/j} \varphi(d) \\ &= \sum_{j \leq n} (1 - \beta^j) \left(\Phi\left(\frac{n}{j}\right) - \Phi\left(\frac{n}{j+1}\right) \right) \\ &= \alpha \sum_{j \leq n} \beta^{j-1} \Phi\left(\frac{n}{j}\right) \\ &= \frac{3}{\pi^2} \cdot \alpha \sum_{j \leq n} \frac{\beta^{j-1}}{j^2} \cdot n^2 + O\left(\alpha \sum_{j \leq n} \frac{n}{j} \log\left(\frac{n}{j}\right)\right) \\ &= \frac{3}{\pi^2} \cdot \frac{\alpha \operatorname{Li}_2(1 - \alpha)}{1 - \alpha} \cdot n^2 + O(\alpha n(\log n)^2), \end{aligned} \tag{11}$$

where we used Lemma 3.3. Putting together (10) and (11), and noting that, by Remark 3.2, the addend of (11) corresponding to $d = 1$ is $1 - \beta^n = O(\alpha n)$, we get (2). The proof is complete. \square

Now we consider the variance of X .

Proof of Theorem 1.3 From Lemmas 4.1 and 4.2, it follows that

$$\begin{aligned} \mathbb{V}[X] &= \mathbb{E}[X^2] - \mathbb{E}[X]^2 \\ &= \sum_{1 < d_1, d_2 \leq n} \varphi(d_1) \varphi(d_2) \left(\mathbb{E}[I_{\mathcal{A}}(d_1) I_{\mathcal{A}}(d_2)] - \mathbb{E}[I_{\mathcal{A}}(d_1)] \mathbb{E}[I_{\mathcal{A}}(d_2)] \right) \\ &= \sum_{1 < d_1, d_2 \leq n} \varphi(d_1) \varphi(d_2) \beta^{\lfloor n/d_1 \rfloor + \lfloor n/d_2 \rfloor - \lfloor n/[d_1, d_2] \rfloor} (1 - \beta^{\lfloor n/[d_1, d_2] \rfloor}). \end{aligned} \tag{12}$$

Let us define

$$V_n(\alpha) := \frac{1}{n^3} \sum_{d_1, d_2 \leq n} \varphi(d_1) \varphi(d_2) \beta^{\lfloor n/d_1 \rfloor + \lfloor n/d_2 \rfloor - \lfloor n/[d_1, d_2] \rfloor} (1 - \beta^{\lfloor n/[d_1, d_2] \rfloor}).$$

Clearly, we have

$$V_n(\alpha) - \frac{\mathbb{V}[X]}{n^3} \ll \frac{1}{n^3} \sum_{d \leq n} \varphi(d) \beta^n (1 - \beta^{\lfloor n/d \rfloor}) \leq \frac{1}{n^3} \sum_{d \leq n} d \ll \frac{1}{n}.$$

Hence, in order to prove (3), it suffices to show that $V_n(\alpha) = v(\alpha) + o(1)$.

For all vectors $\mathbf{a} := (a_1, a_2)$ and $\mathbf{j} := (j_1, j_2, j_3)$ with components in the set of positive integers, define the quantities

$$\rho_1(\mathbf{a}, \mathbf{j}) := \max\left(\frac{1}{a_1(j_1 + 1)}, \frac{1}{a_2(j_2 + 1)}, \frac{1}{a_1 a_2(j_3 + 1)}\right)$$

and

$$\rho_2(\mathbf{a}, \mathbf{j}) := \min\left(\frac{1}{a_1 j_1}, \frac{1}{a_2 j_2}, \frac{1}{a_1 a_2 j_3}\right).$$

Let $d := (d_1, d_2)$ and $a_i := d_i/d$ for $i = 1, 2$. Then the equalities

$$j_1 = \left\lfloor \frac{n}{d_1} \right\rfloor, \quad j_2 = \left\lfloor \frac{n}{d_2} \right\rfloor, \quad j_3 = \left\lfloor \frac{n}{[d_1, d_2]} \right\rfloor,$$

are equivalent to

$$j_1 \leq \frac{n}{a_1 d} < j_1 + 1, \quad j_2 \leq \frac{n}{a_2 d} < j_2 + 1, \quad j_3 \leq \frac{n}{a_1 a_2 d} < j_3 + 1,$$

which in turn are equivalent to

$$\frac{n}{a_1(j_1 + 1)} < d \leq \frac{n}{a_1 j_1}, \quad \frac{n}{a_2(j_2 + 1)} < d \leq \frac{n}{a_2 j_2}, \quad \frac{n}{a_1 a_2(j_3 + 1)} < d \leq \frac{n}{a_1 a_2 j_3},$$

that is,

$$\rho_1(\mathbf{a}, \mathbf{j}) n < d \leq \rho_2(\mathbf{a}, \mathbf{j}) n.$$

Therefore, letting

$$S_n := \{(\mathbf{a}, \mathbf{j}) \in \mathbb{N}^5 : (a_1, a_2) = 1, \exists d \in \mathbb{N} \text{ s.t. } \rho_1(\mathbf{a}, \mathbf{j}) n < d \leq \rho_2(\mathbf{a}, \mathbf{j}) n\}$$

and

$$S(\mathbf{a}, \mathbf{j}; n) := \frac{1}{n^3} \sum_{\rho_1(\mathbf{a}, \mathbf{j}) n < d \leq \rho_2(\mathbf{a}, \mathbf{j}) n} \varphi(a_1 d) \varphi(a_2 d),$$

we have

$$V_n(\alpha) = \sum_{(\mathbf{a}, \mathbf{j}) \in S_n} \beta^{j_1+j_2-j_3} (1 - \beta^{j_3}) S(\mathbf{a}, \mathbf{j}; n).$$

Now let us define

$$v(\alpha) := \sum_{(\mathbf{a}, \mathbf{j}) \in S_\infty} \beta^{j_1+j_2-j_3} (1 - \beta^{j_3}) D(\mathbf{a}, \mathbf{j}), \tag{13}$$

where

$$S_\infty := \bigcup_{m \geq 1} S_m = \{(\mathbf{a}, \mathbf{j}) \in \mathbb{N}^5 : (a_1, a_2) = 1, \rho_1(\mathbf{a}, \mathbf{j}) < \rho_2(\mathbf{a}, \mathbf{j})\}$$

and

$$D(\mathbf{a}, \mathbf{j}) := C_1(a_1, a_2) (\rho_2(\mathbf{a}, \mathbf{j})^3 - \rho_1(\mathbf{a}, \mathbf{j})^3),$$

recalling that $C_1(a_1, a_2)$ is defined by (6). The convergence of series (13) follows easily from Remark 3.1, $\rho_2(\mathbf{a}, \mathbf{j}) \leq 1/(a_1 a_2 j_3)$, and the fact that $\min(j_1, j_2) \geq j_3$ for all $(\mathbf{a}, \mathbf{j}) \in S_\infty$.

Thanks to Lemma 3.4, for each $(\mathbf{a}, \mathbf{j}) \in S_n$ we have

$$S(\mathbf{a}, \mathbf{j}; n) = D(\mathbf{a}, \mathbf{j}) + O\left(\sigma(a_1) \sigma(a_2) \rho_2(\mathbf{a}, \mathbf{j})^2 \cdot \frac{(\log n)^2}{n}\right).$$

Consequently, we get that

$$V_n(\alpha) = v(\alpha) - \Sigma_1 + O\left(\Sigma_2 \cdot \frac{(\log n)^2}{n}\right), \tag{14}$$

where

$$\Sigma_1 := \sum_{(\mathbf{a}, \mathbf{j}) \in S_\infty \setminus S_n} \beta^{j_1+j_2-j_3} (1 - \beta^{j_3}) D(\mathbf{a}, \mathbf{j})$$

and

$$\Sigma_2 := \sum_{(\mathbf{a}, \mathbf{j}) \in \mathcal{S}_n} \beta^{j_1+j_2-j_3}(1 - \beta^{j_3}) \sigma(a_1) \sigma(a_2) \rho_2(\mathbf{a}, \mathbf{j})^2.$$

Now we have to bound both Σ_1 and Σ_2 .

If $(\mathbf{a}, \mathbf{j}) \in \mathcal{S}_\infty \setminus \mathcal{S}_n$ then $(\rho_2(\mathbf{a}, \mathbf{j}) - \rho_1(\mathbf{a}, \mathbf{j}))n < 1$ and consequently, also by Remark 3.1,

$$D(\mathbf{a}, \mathbf{j}) \ll a_1 a_2 (\rho_2^3 - \rho_1^3) = a_1 a_2 (\rho_1^2 + \rho_1 \rho_2 + \rho_2^2)(\rho_2 - \rho_1) \ll \frac{a_1 a_2 \rho_2^2}{n} \leq \frac{1}{a_1 a_2 j_3^2 n}, \tag{15}$$

where, for brevity, we wrote $\rho_i := \rho_i(\mathbf{a}, \mathbf{j})$ for $i = 1, 2$.

If $(\mathbf{a}, \mathbf{j}) \in \mathcal{S}_\infty$ then, as we already noticed, $\min(j_1, j_2) \geq j_3$ and, moreover,

$$\frac{j_2}{j_3 + 1} < a_1 < \frac{j_2 + 1}{j_3} \quad \text{and} \quad \frac{j_1}{j_3 + 1} < a_2 < \frac{j_1 + 1}{j_3}.$$

Hence, we have

$$\begin{aligned} \sum_{(\mathbf{a}, \mathbf{j}) \in \mathcal{S}_\infty} \frac{\beta^{j_1+j_2-j_3}(1 - \beta^{j_3})}{a_1 a_2 j_3^2} &\leq \sum_{j_3 \geq 1} \frac{1 - \beta^{j_3}}{j_3^2} \sum_{j_1, j_2 \geq j_3} \beta^{j_1+j_2-j_3} \sum_{\substack{j_2/(j_3+1) < a_1 < (j_2+1)/j_3 \\ j_1/(j_3+1) < a_2 < (j_1+1)/j_3}} \frac{1}{a_1 a_2} \\ &\ll \sum_{j_3 \geq 1} \frac{1 - \beta^{j_3}}{j_3^2} \sum_{j_1, j_2 \geq j_3} \beta^{j_1+j_2-j_3} = \frac{1}{\alpha^2} \sum_{j \geq 1} \frac{(1 - \beta^j) \beta^j}{j^2} \\ &\leq \frac{1}{\alpha} \sum_{j \leq 1/\alpha} \frac{1}{j} + \frac{1}{\alpha^2} \sum_{j > 1/\alpha} \frac{1}{j^2} \ll \frac{\log(1/\alpha) + 1}{\alpha}, \end{aligned} \tag{16}$$

where we used the inequality $1 - \beta^j \leq \alpha j$, which follows from Remark 3.2.

On the one hand, from (15) and (16) it follows that

$$\Sigma_1 \ll \frac{\log(1/\alpha) + 1}{\alpha n} = o(1), \tag{17}$$

as $\alpha n / ((\log n)^3 (\log \log n)^2) \rightarrow +\infty$ (actually, $\alpha n / \log n \rightarrow +\infty$ is sufficient).

On the other hand, from $\rho_2(\mathbf{a}, \mathbf{j}) \leq 1/(a_1 a_2 j_3)$, (16), and the bound $\sigma(m) \ll m \log \log m$ (see, e.g., [20, Part I, Theorem 5.7]) it follows that

$$\begin{aligned} \Sigma_2 &\leq \sum_{(\mathbf{a}, \mathbf{j}) \in \mathcal{S}_n} \frac{\beta^{j_1+j_2-j_3}(1 - \beta^{j_3})}{a_1 a_2 j_3^2} \cdot \frac{\sigma(a_1) \sigma(a_2)}{a_1 a_2} \ll \frac{(\log(1/\alpha) + 1)(\log \log n)^2}{\alpha} \\ &= o\left(\frac{n}{(\log n)^2}\right), \end{aligned} \tag{18}$$

as $\alpha n / ((\log n)^3 (\log \log n)^2) \rightarrow +\infty$.

At this point, putting together (14), (17), and (18), we obtain $V_n(\alpha) = v(\alpha) + o(1)$, as desired. The proof of (3) is complete.

It remains only to prove the upper bound (4). From (12) it follows that

$$\begin{aligned} \mathbb{V}[X] &\leq \sum_{[d_1, d_2] \leq n} \varphi(d_1) \varphi(d_2) \beta^{\lfloor n/d_1 \rfloor + \lfloor n/d_2 \rfloor - \lfloor n/[d_1, d_2] \rfloor} (1 - \beta^{\lfloor n/[d_1, d_2] \rfloor}) \\ &\leq \sum_{[d_1, d_2] \leq n} d_1 d_2 \cdot \frac{\alpha n}{[d_1, d_2]} = \alpha n \sum_{[d_1, d_2] \leq n} (d_1, d_2) \leq \alpha n \sum_{d \leq n} d \sum_{a_1 a_2 \leq n/d} 1 \\ &= \alpha n \sum_{d \leq n} d \sum_{m \leq n/d} \tau(m) \ll \alpha n^2 \sum_{d \leq n} \log\left(\frac{n}{d}\right) = \alpha n^2 (n \log n - \log(n!)) < \alpha n^3, \end{aligned}$$

where we used Remark 3.2, Lemma 3.1, and the bound $n! > (n/e)^n$. Thus (4) is proved. \square

Proof of Theorem 1.4 By Chebyshev's inequality, Theorems 1.2 and 1.3, we have

$$\mathbb{P}[|X - \mathbb{E}[X]| > \varepsilon \mathbb{E}[X]] \leq \frac{\mathbb{V}[X]}{(\varepsilon \mathbb{E}[X])^2} \ll \frac{\alpha n^3}{(\varepsilon \alpha n)^2} = \frac{1}{\varepsilon^2 \alpha n} = o_\varepsilon(1),$$

as $\alpha n \rightarrow +\infty$. Hence, using again Theorem 1.2, we get

$$X \sim \frac{3}{\pi^2} \cdot \frac{\alpha \operatorname{Li}_2(1 - \alpha)}{1 - \alpha} \cdot n^2,$$

with probability $1 - o(1)$, as $\alpha n \rightarrow +\infty$. \square

Authors' contributions

The author thanks the anonymous referee, whose careful reading and detailed suggestions led to a considerable improvement of the paper.

Funding Open access funding provided by Politecnico di Torino within the CRUI-CARE Agreement.

Received: 27 December 2020 Accepted: 2 February 2021 Published online: 18 February 2021

References

- Alsmeyer, G., Kabluchko, Z., Marynych, A.: Limit theorems for the least common multiple of a random set of integers. *Trans. Am. Math. Soc.* **372**(7), 4585–4603 (2019)
- Andrews, G.E.: *q-Series: Their Development and Application in Analysis, Number Theory, Combinatorics, Physics, and Computer Algebra*, CBMS Regional Conference Series in Mathematics, vol. 66, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI (1986)
- Bateman, P., Kalb, J., Stenger, A.: Solution to problem 10797: a limit involving least common multiples. *Am. Math. Monthly* **109**(4), 393–394 (2002)
- Bostan, A., Marynych, A., Raschel, K.: On the least common multiple of several random integers. *J. Number Theory* **204**, 113–133 (2019)
- Buraczewski, D., Iksanov, A., Marynych, A.: A Brownian weak limit for the least common multiple of a random m -tuple of integers. <https://arxiv.org/abs/2004.05643>
- Cilleruelo, J.: The least common multiple of a quadratic sequence. *Compos. Math.* **147**(4), 1129–1150 (2011)
- Cilleruelo, J., Guijarro-Ordóñez, J.: Ratio sets of random sets. *Ramanujan J.* **43**(2), 327–345 (2017)
- Cilleruelo, J., Ramana, D.S., Ramaré, O.: Quotient and product sets of thin subsets of the positive integers. *Proc. Steklov Inst. Math.* **296**(1), 52–64 (2017)
- Cilleruelo, J., Rué, J., Šarka, P., Zumalacárregui, A.: The least common multiple of random sets of positive integers. *J. Number Theory* **144**, 92–104 (2014)
- Hong, S., Qian, G., Tan, Q.: The least common multiple of a sequence of products of linear polynomials. *Acta Math. Hungar.* **135**(1–2), 160–167 (2012)
- Kac, V., Cheung, P.: *Quantum Calculus*. Universitext. Springer, New York (2002)
- Mastrostefano, D.: On maximal product sets of random sets, (to appear on *Journal of Number Theory*) <https://arxiv.org/abs/2005.04663>
- Maynard, J., Rudnick, Z.: A lower bound on the least common multiple of polynomial sequences, (to appear on *Rivista di Matematica dell'Università di Parma*) <https://arxiv.org/abs/1910.13218>
- Rudnick, Z., Zehavi, S.: On Cilleruelo's conjecture for the least common multiple of polynomial sequences. <https://arxiv.org/abs/1902.01102>
- Rué, J., Šarka, P., Zumalacárregui, A.: On the error term of the logarithm of the lcm of a quadratic sequence. *J. Théor. Nombres Bordeaux* **25**(2), 457–470 (2013)
- Sah, A.: An improved bound on the least common multiple of polynomial sequences. *J. Théor. Nombres Bordeaux* **32**(3), 891–899 (2020)
- Sanna, C.: On the L.C.M. of shifted Fibonacci numbers <https://arxiv.org/abs/2007.13330>
- Sanna, C.: A note on product sets of random sets. *Acta Math. Hungar.* **162**(1), 76–83 (2020)
- Sanna, C.: On the L.C.M. of random terms of binary recurrence sequences. *J. Number Theory* **213**, 221–231 (2020)
- Tenenbaum, G.: *Introduction to Analytic and Probabilistic Number Theory*, third ed., Graduate Studies in Mathematics, vol. 163, American Mathematical Society, Providence, RI, Translated from the 2008 French edition by Patrick D. F. Ion (2015)

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.