

The hydrodynamic genesis of linear karren patterns

Original

The hydrodynamic genesis of linear karren patterns / Bertagni, M.B., Camporeale, C.. - In: JOURNAL OF FLUID MECHANICS. - ISSN 0022-1120. - ELETTRONICO. - 913:A34(2021). [10.1017/jfm.2021.39]

Availability:

This version is available at: 11583/2872892 since: 2021-03-02T14:38:14Z

Publisher:

Cambridge University Press

Published

DOI:10.1017/jfm.2021.39

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

The hydrodynamic genesis of linear karren patterns

M.B. Bertagni ¹†, and C. Camporeale ²

¹The High Meadows Environmental Institute, Princeton University, Princeton, NJ, USA

²Department of Land, Infrastructure and Environmental Engineering, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10124, Torino, Italy

(Received xx; revised xx; accepted xx)

In karst and alpine areas, the interactions between water and rocks give rise to a large variety of marvelous patterns. In this work, we provide an hydrodynamic model for the formation of dissolutional patterns made of parallel longitudinal channels and commonly referred to as linear karren forms. The model addresses a laminar film of water flowing on a rock that is dissolving. The results show that a transverse instability of the water-rock system leads to a longitudinal channelization responsible for the pattern formation. The instability arises because of a positive feedback within the channels between the higher water flow and the enhanced chemical dissolution. The spatial scales predicted by the linear stability analysis span different orders of magnitude depending on the Reynolds number. This may explain why similar patterns of different sizes are observed on natural rocks. Results also show that the rock solubility affects just the temporal scale of the instability and the rock inclination plays a minor role in the pattern formation. It is eventually discussed how rain is not strictly necessary for the appearance of linear karren patterns, but it may affect some of their features.

Key words: Authors should not enter keywords on the manuscript, as these must be chosen by the author during the online submission process and will then be added during the typesetting process (see <http://journals.cambridge.org/data/relatedlink/jfm-keywords.pdf> for the full list)

1. Introduction

Water is an impressive sculptor. It shapes remarkably regular and beautiful patterns throughout the Earth using different tools, such as sediment mobilization, thermal gradients, and chemistry. In karst environments, water carves through chemical dissolution and precipitation a plethora of peculiar patterns on soluble rocks (e.g. Meakin & Jamtveit 2009; Jamtveit & Hammer 2012). Because of the considerable variety of hydrodynamic and chemical processes involved in karst pattern formation, different criteria may be adopted for their classification, which thus remain an open debate (Ginés *et al.* 2009). At first, we may separate between dissolution and precipitation patterns, even though in many modelling aspects these two reverse processes can be treated in a unified manner (Ford & Williams 2013).

Dissolution occurs when undersaturated water, commonly coming from rain or snow-melt, interacts with a soluble rock. It is the most common process in chemical weath-

† Email address for correspondence: matteobb@princeton.edu



FIGURE 1. Linear karren forms of different spatial scales: wandkarren (a-b), rillenkarrren (c), microrills (d). (a) Wandkarren in limestone, Asturias, Spain. (b) Wandkarren in dolomite, Brenta Dolomites, Italy. (c) Rillenkarrren in dolomite, Brenta Dolomites, Italy. (d) Microrills in limestone, Verzino, Calabria, Italy. Credits: I. Benvenuti Cabral for panel (a); Sauro (2009) for panel (d).

ering and, because the dissolution of some rocks involves the capture of CO_2 from the atmosphere, it has profound implications on the Earth carbon cycle (e.g. Berner *et al.* 1983; Regnier *et al.* 2013). Dissolution patterns are mostly found on exposed limestone, dolomite, gypsum and salt rocks and they are usually referred to with the German word *karren* (Bögli 1960). A wide overview on the state of karren research may be found in the contributions collected in Ginés *et al.* (2009). Dissolution patterns may also arise below-ground, as for example in the case of cave scallops, which are formed on cave walls by turbulent flows of water (Curl 1966; Claudin *et al.* 2017).

Precipitation patterns occur after the water has reached a supersaturated state by percolating through the soil. This happens mainly in caves, where water shapes a great variety of *speleothems*. Because speleothem formation endures millennia, they are precious sources of information on past climates (Fairchild & Baker 2012). Some aspects of speleothems have been characterized by hydrodynamic models in recent years, such as the stalactite shape (Short *et al.* 2005), the ripple-forms appearing on the stalactite surface (Camporeale & Ridolfi 2012; Vesipa *et al.* 2015), and the longitudinal precipitation flutes (Camporeale 2015; Bertagni & Camporeale 2017). Yet, many speleothem features still need to be unveiled (Fairchild & Baker 2012). Exceptionally, precipitation patterns also occur above-ground in geothermal hot springs, where admirable travertine terraces emerge (Goldenfeld *et al.* 2006; Veysey & Goldenfeld 2008).

This work deals with dissolutional karren forms. Because karren are characterized by a multitude of shapes (e.g., circular, linear, polygenetic) and spatial scales (from micrometers to tens of meters), karren classification is still debated (Ginés *et al.* 2009).

Specifically, we deal with karren of linear forms that are hydrodynamically controlled (instead of fracture controlled, see the classification by Ford & Williams 2013, tab. 9.1). These are longitudinal patterns in which the channels originate from the dissolution process driven by the water flow (Fig. 1). Linear karren patterns are usually grouped in relation to their transverse wavelength (distance between two *quasi*-parallel channels), which can span different order of magnitudes: from decimetric wandkarren (Fig. 1a-b) and rinnenkarren, also called runnels, to centimetric rillenkarren (Fig. 1c-d) and millimetric microrills (Fig. 1e), also called rillenstein. The pattern classification is not always straightforward. For example, the wandkarren in panel (b) much recalls the rillenkarren in panel (d) and probably some authors would call them decantation flutings (Ford & Williams 2013). Furthermore, superimposition of one pattern onto another are very common, such as microrills on rillenkarren, or rillenkarren on rinnenkarren (Ginés *et al.* 2009). There is a reason for these impressive similarities: all these patterns share an hydrodynamic common origin. Although this has always been recognized, to date there is not a comprehensive theory that might explain the regular channelization induced by water (Ginés *et al.* 2009). This theoretical lack is the goal of the present paper.

To this aim, we model a laminar film of undersaturated water that flows down a dissolving inclined rock (see Fig. 2 for a graphical sketch). The overall chemistry is reduced to the total concentration $\hat{c}(\hat{\mathbf{x}}, \hat{t})$ of the solute species within the water film (the hat refers to dimensional quantities). The interactions between the flow field and the concentration field define the temporal evolution of the rock surface. From a mathematical point of view, the dynamic water-rock boundary delineates a Stefan problem for the underlying system of partial differential equations. Notably, in recent years Stefan-problems involving fluid motion have been the subject of several experimental investigations (e.g. Mac Huang *et al.* 2015; Wykes *et al.* 2018; Cohen *et al.* 2020; Guérin *et al.* 2020) and theoretical analyses (e.g. Moore 2017; Morrow *et al.* 2019).

A modelling novelty of this work regards the dissolution rate, that we here define by directly addressing the local gradient of concentration at the rock-water interface. This is done by spatially solving the advection-diffusion equation for $\hat{c}(\hat{\mathbf{x}}, \hat{t})$ and it allows us to avoid the empirical formulations for the dissolution rate that are commonly used in the context of karst pattern formation. These are here briefly summarized for completeness. The simplest formulations link the dissolution-precipitation flux to an hydrodynamic quantity, e.g., the depth-averaged velocity (Goldenfeld *et al.* 2006; Veysey & Goldenfeld 2008) or the water depth (Camporeale 2015; Bertagni & Camporeale 2017). Another common approach is to linearly link the dissolution flux to the difference between the saturation concentration and a certain concentration within the water film, e.g., the – uniform– bulk concentration (e.g. Camporeale 2017) or the concentration at the solid wall (e.g. Claudin *et al.* 2017). A more complete formulation for the dissolution flux is the seminal PWP (Plummer-Wigley-Parkhurst) equation (Plummer *et al.* 1978), which has been successfully used within the context of karst pattern formation (e.g. Camporeale & Ridolfi 2012; Vesipa *et al.* 2015). Although these empirical formulations are sustained by experimental and numerical evidences (Buhmann & Dreybrodt 1985; Hammer *et al.* 2008; Dreybrodt 2012), they do not address the local gradients of concentration that may develop within the water film.

The Paper is structured as follows: in Sect. 2, we formulate the problem of linear karren formation; in Sect. 3, we find the mathematical solutions for the hydrodynamics and the solute concentration in the initial case of a flat rock surface; in Sect. 4, we investigate the linear stability of the flat solution to eventually achieve the dispersion relationship that describes incipient karren formation. The results of the linear stability analysis are




FIGURE 2. Sketch of the water film flowing on the soluble rock. The flat state is characterized by the semi-parabolic profile for the longitudinal velocity $u_0(\zeta)$ and the two-dimensional concentration field $c_0(x, \zeta)$. This is the base state of the transverse (z -direction) stability analysis.

reported in Sect. 5 and further discussed, together with some limits of the present model, in Sect. 6. We finally add some concluding remarks in Sect. 7.

2. Formulation of the problem

2.1. Governing equations

A steady water film flowing on a flat surface performs a semi-parabolic velocity profile (Fig. 2). This is commonly referred to as the Nusselt solution and it can be readily derived from momentum conservation (Nusselt 1916). Accordingly, the dimensional film thickness and the surface velocity read

$$\hat{h}_0 = \left(\frac{3\nu\hat{q}}{g\sin\theta} \right)^{1/3}, \quad \hat{u}_s = \left(\frac{9g\hat{q}^2\sin\theta}{8\nu} \right)^{1/3}, \quad (2.1)$$

where $\nu = 10^{-6}$ m²/s is the water kinematic viscosity, g is the gravitational acceleration, θ is the angle with the horizontal, and $\hat{q} = 2\hat{u}_s\hat{h}_0/3$ is the flow rate per unit span. The hat refers to dimensional variables. We use the quantities in eq. (2.1) to scale the governing equations of the problems, which are the Navier-Stokes equations and the advection-diffusion equation for the solute concentration c (scaled with its equilibrium value \hat{c}_{eq} , see Table 1)

$$\nabla \cdot \mathbf{u} = 0, \quad (2.2)$$

$$Re(\mathbf{u} \cdot \nabla \mathbf{u} + \nabla p) = \nabla^2 \mathbf{u} + \mathbf{f}, \quad (2.3)$$

$$Pe \mathbf{u} \cdot \nabla c = \nabla^2 c, \quad (2.4)$$

where $\nabla = (\partial_x, \partial_y, \partial_z)$, $\mathbf{u} = (u, v, w)$ is the velocity field (Fig. 2), p is pressure – scaled with $\rho \hat{u}_s^2$ being ρ the water density – and $\mathbf{f} = (2, -\delta, 0)$ is the gravity term, with $\delta = 2 \cot \theta$. Because of the slow morphological evolution of the rock surface, the *quasi-steady*

approximation ($\partial_t = 0$) is used in eqs. (2.2)-(2.4). The dimensionless numbers appearing in (2.2)-(2.4) are the Reynolds and Peclet numbers, which read

$$Re = \frac{\hat{u}_s \hat{h}_0}{\nu}, \quad Pe = \frac{\hat{u}_s \hat{h}_0}{D} = Re Sc, \quad (2.5)$$

where D is the molecular diffusivity coefficient. For the solute species considered (Tab. 1), an average value of $D \simeq 10^{-9}$ m²/s can be assumed (Ford & Williams 2013). Notice that Pe and Re are linked through the Schmidt number $Sc = \nu/D = 10^3$ ($Pe = 10^3 Re$).

2.2. Hydrodynamic boundary conditions

The boundary conditions are defined on the rock-water interface $\eta(x, z, t)$ and water-air interface $\eta(x, z, t) + h(x, z, t)$ (Fig. 2). It is thus convenient to introduce the vertical coordinate

$$\zeta = \frac{y - \eta(x, z, t)}{h(x, z, t)}, \quad (2.6)$$

so that the water domain is always defined between $\zeta = 0$ (rock-water interface) and $\zeta = 1$ (water-air interface). Notice that y and η are measured relative to the position of the initially flat rock surface and they are scaled with \hat{h}_0 .

At the rock-water interface ($\zeta = 0$), the velocity field satisfies the no-slip and impermeability conditions, which are respectively

$$u = w = 0, \quad v = 0, \quad (2.7)$$

where we neglect the influence of the slow dissolution process in the vertical hydrodynamic velocity. At the water-air interface ($\zeta = 1$), the continuity of the stress is ensured by the so-called dynamic conditions

$$\mathbf{n} \cdot \mathbf{T} \cdot \mathbf{n} = -We \mathcal{K}, \quad \mathbf{n} \cdot \mathbf{T} \cdot \mathbf{t} = 0, \quad (2.8)$$

where $\mathbf{n} = [\nabla(y - \eta - h)]/n$ and \mathbf{t} are the versors normal and tangent to the free-surface ($n = \{1 + [\partial_x(h + \eta)]^2 + [\partial_z(h + \eta)]^2\}^{1/2}$ is the versor normalization). \mathbf{T} is the stress tensor ($\mathbf{T}_{ij} = p \delta_{ij} - (\partial_{x_i} u_j + \partial_{x_j} u_i)/Re$). The term $We \mathcal{K}$ accounts for the normal stress induced by the surface tension σ , where We is the Weber number and \mathcal{K} is the mean curvature of the free-surface (e.g Chang & Demekhin 2002)

$$We = \frac{\sigma}{\rho \hat{h}_0 \hat{u}_s^2} = \frac{Ka}{Re^{5/3} \sin(\theta)}, \quad (2.9)$$

$$\mathcal{K} = \partial_x \left[\frac{\partial_x(h + \eta)}{n} \right] + \partial_z \left[\frac{\partial_z(h + \eta)}{n} \right]. \quad (2.10)$$

From (2.9) we notice that We is a function of Re and the angle θ through the Kapitza number $Ka = 2^{1/3} \sigma / (g^{1/3} \nu^{4/3} \rho)$. Ka only depends on the water properties and is thus constant for our purposes.

The temporal evolution of the free-surface is described by the kinematic condition in $\zeta = 1$

$$\partial_t h = \mathbf{u} \cdot \mathbf{n} = v - u(h + \eta)_x - w(h + \eta)_z, \quad (2.11)$$

where, as in eq. (2.7), we neglected the influence of η_t in the hydrodynamics. Eq. (2.11) ensures the water mass conservation and will be later used as the first solvability equation.

Dissolution reaction		ρ_s (kg/m ³)	\hat{c}_{eq} (kg/m ³)	γ
Calcite	$\text{CaCO}_3 + \text{H}_2\text{O} + \text{CO}_2 \leftrightarrow \text{Ca}^{2+} + 2\text{HCO}_3^-$	2700	0.055	49091
Gypsum	$\text{CaSO}_4 \cdot 2\text{H}_2\text{O} \rightarrow \text{Ca}^{2+} + \text{SO}_4^{2-} + 2\text{H}_2\text{O}$	2300	2.4	958
Halite (salt)	$\text{NaCl} + \text{H}_2\text{O} \rightarrow \text{Na}^+ + \text{Cl}^- + \text{H}^+ + \text{OH}^-$	2300	360	6.4

TABLE 1. Dissolution reaction, density, and solubility of some representative karst minerals. Solubility is given as the amount of mineral that can be dissolved in water at 25° and $p_{\text{CO}_2} = 4 \cdot 10^{-4}$ atm (Dreybrodt 2012; Ford & Williams 2013).

2.3. Concentration boundary conditions

Differently from the hydrodynamic quantities, the concentration c has also a longitudinal dependence due to the downstream solute accumulation within the water film (see Fig. 2). Basically, this makes the concentration field a three-dimensional problem. In fact, c depends on x because of the downstream solute accumulation, on y due to the normal-to-wall effect of the dissolution occurring at the rock-water interface, and on z because of the transverse nature of linear karren forms.

Upstream, as the water film is generally produced by rain or snow-melt, it might be considered free of solute

$$c|_{x=0} = 0. \quad (2.12)$$

With the dissolution of the rock surface, solute molecules are dissociated from the solid through chemical reactions and then transported into the film bulk by diffusion. In general, the slowest between these two processes (surface reaction and transport by diffusion) controls the dissolution kinetics of a karst rock. Very soluble minerals, such as salt, have a rapid dissolution reaction and thus the kinetics is always diffusion-controlled (Ford & Williams 2013). For relatively less soluble minerals, such as gypsum or calcite, numerous studies have demonstrated that in very undersaturated waters, as those commonly found in exposed karst environments, a diffusion-controlled dissolution kinetics prevail (see the review on dissolution kinetics by Morse & Arvidson 2002, and references therein). As equilibrium is approached, there is a transition region to surface-controlled dissolution kinetics. Moreover, for calcite, which dissolves in water enriched of CO_2 , the slow uptake of CO_2 from the atmosphere may also become a limiting factor as equilibrium is approached (Buhmann & Dreybrodt 1985; Kaufmann & Dreybrodt 2007). Notice that CO_2 -conversion does not influence salt and gypsum dissolution as they directly dissociate in pure water (Tab 1).

Since we wish to model incipient karren formation and linear karren forms have been observed to develop upstream first, i.e., where the water is very undersaturated (Glew & Ford 1980; Ginés *et al.* 2009; Slabe *et al.* 2016), we consider a dissolution kinetics that is diffusion-controlled. This implies that surface reactions are fast compared to diffusion and that the solute accumulates in a very thin diffusion boundary layer (DBL) at the solid-liquid interface that is nearly saturated (e.g. Ford & Williams 2013). By also imposing no-flux of solute between water and air, the vertical boundary conditions eventually read

$$\partial_\zeta c|_{\zeta=1} = 0, \quad c|_{\zeta=0} = 1. \quad (2.13)$$

The morphological evolution of the rock surface is thus regulated by the flux of solute that moves into the bulk of the water film through diffusion, i.e. $\rho_s(\hat{V} + \partial_t \hat{\eta}) = D \partial_y \hat{c}$, where ρ_s is the rock density and \hat{V} is the dissolution rate induced by the uniform hydrodynamic

flow ($\hat{V} < 0$). In dimensionless form, the same equation reads

$$\gamma Pe h (V + \partial_t \eta) = \partial_\zeta c, \quad (2.14)$$

where $\gamma = \rho_s / \hat{c}_{eq} \gg 1$ is related to the mineral considered (Tab. 1). Equation (2.14) will be used as the second solvability equation.

3. Flat rock solution

We here find the solution to the problem (2.2)-(2.14) for an initially flat rock surface, i.e., $\eta_0 = \partial_z = 0$ and $h_0 = 1$, where the subscript 0 denotes the absence of karren patterns.

3.1. Hydrodynamics

The Navier-Stokes equations (2.2)-(2.3) reduce to

$$u_0'' = -2, \quad p_0' = -\delta/Re, \quad (3.1)$$

with boundary conditions

$$u_0|_{\zeta=0} = 0, \quad u_0'|_{\zeta=1} = 0, \quad p_0|_{\zeta=1} = 0, \quad (3.2)$$

where ' denotes differentiation in ζ . The solutions are the Nusselt semi-parabolic profile for the longitudinal velocity and the hydrostatic pressure profile

$$u_0 = (2 - \zeta)\zeta, \quad v_0 = w_0 = 0, \quad p_0 = \frac{\delta}{Re}(1 - \zeta). \quad (3.3)$$

3.2. Concentration

The advection-diffusion equation (2.4) reduces to

$$Pe u_0(\zeta) \partial_x c_0 = c_0'', \quad (3.4)$$

where the longitudinal diffusion has been neglected since its effect is smaller than that of longitudinal advection. The initial and boundary conditions (2.12)-(2.13) read

$$c_0|_{x=0} = 0, \quad c_0|_{\zeta=0} = 1, \quad c_0'|_{\zeta=1} = 0. \quad (3.5)$$

Equation (3.4) highlights the coupling between the concentration field c_0 and the velocity profile $u_0(\zeta)$, as well as the x -dependency of the concentration distribution due to the downstream solute accumulation. The solution to this problem is resumed in Polyanin *et al.* (2001, p. 130-132) and it is here more extensively reported. For $x \geq 0$, we seek the solution in the form of the series (Davis 1973)

$$c_0 = 1 - \sum_{m=0}^{\infty} A_m F_m(x) G_m(\zeta), \quad (3.6)$$

where $A_m, F_m(x), G_m(\zeta)$ are unknown functions that must be determined. Substituting the expansion (3.6) into (3.4) and then separating the variables, we obtain

$$\partial_x F_m(x) + \frac{\alpha_m^2}{Pe} F_m(x) = 0, \quad (3.7)$$

$$G_m''(\zeta) + \alpha_m^2 u_0(\zeta) G_m(\zeta) = 0. \quad (3.8)$$

The solution to the x -problem (3.7) is readily given by

$$F_m(x) = \exp\left(-\alpha_m^2 \frac{x}{Pe}\right). \quad (3.9)$$

Instead, eq. (3.8) defines a Sturm–Liouville problem, whose boundary conditions, that are obtained after replacement of (3.6) into the second and third equations in (3.5), are

$$G_m|_{\zeta=0} = 0, \quad G'_m|_{\zeta=1} = 0. \quad (3.10)$$

The solution of the Sturm–Liouville problem for the eigenfunctions G_m (up to a constant factor) is

$$G_m(\zeta) = \exp\left[-\frac{\alpha_m}{2}(1-\zeta)^2\right] \Phi\left(\frac{1}{4} - \frac{\alpha_m}{4}, \frac{1}{2}; \alpha_m(1-\zeta)^2\right), \quad (3.11)$$

where $\Phi(\cdot, \cdot; \cdot)$ is the degenerate hypergeometric function (Abramowitz *et al.* 1966). Substituting (3.11) into the first of the boundary conditions (3.10) provides the transcendental equation for the eigenvalues $\alpha_m > 0$

$$\Phi\left(\frac{1}{4} - \frac{\alpha_m}{4}, \frac{1}{2}; \alpha_m\right) = 0. \quad (3.12)$$

For the definition of the coefficients A_m , the first step is to substitute the series (3.6) into the first equation in (3.5), which yields

$$\sum_{m=0}^{\infty} A_m G_m(\zeta) = 1. \quad (3.13)$$

Multiplying eq. (3.13) by $u_0 G_n$, with $n \neq m$, then integrating between $\zeta = 0$ and $\zeta = 1$, and using the orthogonality condition $\int_0^1 u_0 G_m G_n d\zeta = 0$, one obtains

$$A_m = \frac{\int_0^1 u_0 G_m(\zeta) d\zeta}{\int_0^1 u_0 G_m(\zeta)^2 d\zeta}. \quad (3.14)$$

Notably, instead of solving numerically eq. (3.12) and (3.14), it is possible to evaluate the constants A_m and α_m with the approximated relationships (Polyanin & Nazaikinskii 2015)

$$\alpha_m = 4m + 1.68 \quad (m = 0, 1, 2, \dots), \quad (3.15)$$

$$A_0 = 1.2, \quad A_m = (-1)^m 2.27 \alpha_m^{-7/6} \quad (m = 1, 2, 3, \dots), \quad (3.16)$$

whose maximum error is less than 0.2%. It is also worth mentioning that, for $x/Pe \rightarrow 0$, an easier asymptotic solution for c_0 can be obtained (Polyanin *et al.* 2001)

$$c_0 = \Gamma\left(\frac{1}{3}, \frac{2\zeta^3}{9X}\right) / \Gamma\left(\frac{1}{3}\right) \quad \text{for } x/Pe = X \rightarrow 0, \quad (3.17)$$

where $\Gamma(\cdot)$ and $\Gamma(\cdot, \cdot)$ are the complete and incomplete Gamma functions (Abramowitz *et al.* 1966). $X = x/Pe$ is a convenient scaling that maps the dissolution domain of the problem between $X = 0$, where the water is free of solute, and $X = 1$, where the water film is basically saturated. In fact, some easy algebra shows that the timescale $\hat{\tau}$ needed by water to go from $X = 0$ to $X = 1$ is $\hat{\tau} \sim \hat{h}_0^2/D$, i.e., equivalent to the timescale required by diffusion to involve the full water depth.

The spatial trends of the exact solution (3.6) and the asymptotic solution (3.17) for c_0 are shown in Fig. 3.




FIGURE 3. Vertical profiles of c_0 at different X (log-scale). The solid red lines refers to the exact solution (3.6). The dotted-black lines refer to the asymptotic solution (3.17), which is shown to be valid up to $X \sim 0.1$. The coloured areas evidence the saturation ratio of water in the solute.

3.3. Dissolution

The dissolution rate V can be evaluated by combining the evolution equation for the rock surface (2.14) and the solution for c_0

$$V = \frac{1}{\gamma Pe} c'_0|_{\zeta=0}. \quad (3.18)$$

It is also convenient to define the saturation ratio s as

$$s = \int_0^1 c_0 d\zeta, \quad (3.19)$$

where $s = 0$ indicates pure water and $s = 1$ corresponds to saturated water (ion-pairing is neglected). Because the concentration has been scaled with its saturation value, s is also equivalent to the dimensionless depth-averaged concentration. A graphical evidence of the saturation ratio of the water film is given by the light red areas in the panels of Fig. 3.

The trend $V = V(X)$, obtained after replacing (3.6) in (3.18), is reported in Fig. 4a. Notice that the quantity $\gamma Pe |V|$ is independent of the rock type and the film hydrodynamics. The dissolution rate decreases with X as the water loses chemical aggressivity by increasing its saturation ratio s (right y-axis). Because clear water enhances dissolution, half saturation ($s = 0.5$) is reached in the upstream part of the flow ($X \sim 0.2$) and the rate of solute accumulation progressively decreases downstream. Although at $X = 1$ the flow is not completely saturated (coloured area in Fig. 3d) at those levels of saturation – and even before ($s > 0.9$) – dissolution is known to drop drastically because of the inhibitory action of other chemical species in the rock texture (Kaufmann & Dreybrodt 2007; Ford & Williams 2013).

Between the saturation ratio s and the dissolution rate, there is a nonlinear relationship (Fig. 4b). This nonlinearity, which here arises thanks to the spatially-dependent solution for the concentration, highlights the limits of the common assumption of a diffusive flux linearly related to the saturation ratio, i.e., $V \propto (s - 1)$. Indeed, the dissolution of the rock surface is better specified by the local gradient of concentration at the rock surface ($c'_0|_{\zeta=0}$), rather than by the average concentration gradient along the film thickness ($s - 1$ dimensionally scales with $(\hat{c} - \hat{c}_{\text{eq}})/\hat{h}_0$). Furthermore, it is remarkable to notice that the trend in Fig. 4b is qualitatively similar to experimental evidences for calcite dissolution (Buhmann & Dreybrodt 1985; Kaufmann & Dreybrodt 2007), even though we here consider a simplified dissolution kinetics that is controlled just by diffusion. Kaufmann & Dreybrodt (2007) have in fact shown that two *quasi*-linear regimes (for




FIGURE 4. Dissolution of an initially flat rock. (a) Longitudinal trends of the scaled dissolution rate $\gamma Pe|V|$ (blue) and the saturation ratio s (red). (b) $\gamma Pe|V|$ versus s , from the previous panel, highlighting a nonlinear relationship with a change of dissolution rate around $s \sim 0.3$.

$s < 0.3$ and $s > 0.3$) can be considered in the dissolution rate of calcite. These regimes have linear coefficients separated by an order of magnitude and they have been qualitatively plotted with dashed lines in Fig. 4b.

4. Linear Stability Analysis

We here evaluate the stability of the flat-rock solution reported in the previous section to a small transverse perturbation written in normal modes

$$(\mathbf{u}, p, c, h, \eta) = (\mathbf{u}_0(\zeta), p_0(\zeta), c_0(\zeta, X), 1, 0) + \varepsilon(\mathbf{u}_1(\zeta), p_1(\zeta), c_1(\zeta, X), h_1, \eta_1) e^{\omega t + ikz}, \quad (4.1)$$

where ε is an infinitesimal parameter, k is the transverse wavenumber, and ω is the growth rate. Due to the transverse invariance of the problem, ω is a real number, i.e., there is no angular phase. When $\omega > 0$, the base-state is unstable and the perturbation grows in time to eventually form periodic longitudinal patterns.

The ansatz (4.1) follows the assumption of longitudinal invariance ($\partial_x = 0$) just for the hydrodynamic quantity (\mathbf{u}, p, h) and not for the concentration field, which has a longitudinal structure given by the solute accumulation within the water film. The variable X defines the transverse section wherein the linear stability analysis is performed.

4.1. Hydrodynamics

Since the hydrodynamic problem is longitudinally invariant, the continuity equation (2.2) at the linear order (ε) imposes $v'_1 + ik w_1 = 0$. Therefore, we can introduce the scalar stream function φ , such that

$$v_1 = -ik\varphi, \quad w_1 = \varphi'. \quad (4.2)$$

After some algebraic manipulations, the Navier-Stokes equations (2.2)-(2.3) are reduced to the Orr-Sommerfeld equation for a domain longitudinally invariant (e.g. Chang & Demekhin 2002; Kalliadasis *et al.* 2011)

$$\varphi^{iv} - 2k^2\varphi'' + k^4\varphi = 0, \quad (4.3)$$

with boundary conditions

$$\varphi' = \varphi = 0 \quad \text{in } \zeta = 0, \quad (4.4)$$

$$\varphi'' + k^2\varphi = \varphi^{(3)} - 3k^2\varphi' - ik(\eta_1 + h_1)(k^2 Re We + \delta) = 0 \quad \text{in } \zeta = 1. \quad (4.5)$$

We remind that equations (4.4) are the no-slip and impermeability conditions at the rock-water interface (corresponding to eqs. (2.7)), while equations (4.5) are the linearized dynamic conditions at the water-air interface, (eqs. (2.8)). The advantage of the Orr-Sommerfeld approach is that the transverse hydrodynamic problem is reduced to the biharmonic equation (4.3), the solution thereof is

$$\varphi = i (\delta + k^2 Re We) (\eta_1 + h_1) f(\zeta), \quad (4.6)$$

where

$$f(\zeta) = \frac{(k\zeta \coth(k\zeta) - 1)(k \sinh k + \cosh k) - k^2 \zeta \cosh k}{k(2k^2 + \cosh(2k) + 1)} \sinh(k\zeta). \quad (4.7)$$

Combining (4.2) and (4.6) readily provides the solution for v_1 and w_1 (where the latter can be shown to be much larger than the former). Because of the longitudinal invariance of the linear problem, u_1 is not explicitly related to φ . Thus, for its evaluation, we directly address the conservation of longitudinal momentum (2.3), which at order ε reads

$$u_1'' - k^2 u_1 = -k^2 u_0' (\eta_1 + \zeta h_1) + Re u_0' v_1 + 2u_0'' h_1, \quad (4.8)$$

with boundary conditions

$$u_1|_{\zeta=0} = 0, \quad u_1'|_{\zeta=1} = 0. \quad (4.9)$$

The terms in the left side of eq. (4.8) are the vertical and transverse diffusion of momentum, respectively. The terms in the right side arise from advection ($Re u_0' v_1$) and the change of the vertical coordinate (from y to ζ , see eq. (2.6)).

By solving the equation (4.8) through the method of variation of parameters (e.g. Bender & Orszag 2013), we eventually obtain

$$u_1 = 2 \operatorname{sech} k [(\eta_1 + h_1) \sinh(k\zeta)/k - \eta_1 \cosh(k - k\zeta)] + 2(1 - \zeta)(\eta_1 + \zeta h_1), \quad (4.10)$$

where the contribution of v_1 in (4.10) has been neglected as it would considerably complicate the solution without any perceptible numerical change.

4.2. Concentration

At $\mathcal{O}(\varepsilon)$, the linearized advection-diffusion equation (2.4) reads

$$c_1'' - k^2 c_1 + \Upsilon(\zeta, X) = u_0(\zeta) \partial_X c_1 \quad (4.11)$$

where Υ is the inhomogeneous term

$$\Upsilon(\zeta, X) = k^2 c_0' (\eta_1 + \zeta h_1) - Pe c_0' v_1 - 2h_1 c_0'' - u_1 \partial_X c_0, \quad (4.12)$$

and the initial and boundary conditions (2.12)-(2.13) are

$$c_1|_{X=0} = 0, \quad c_1|_{\zeta=0} = 0, \quad c_1'|_{\zeta=1} = 0. \quad (4.13)$$

The solution to the problem (4.11)-(4.13) is (Polyanin & Nazaikinskii 2015)

$$c_1 = \int_0^X \int_0^1 \Upsilon(\zeta_m, X_m) \mathcal{G}(\zeta, \zeta_m, X - X_m) d\zeta_m dX, \quad (4.14)$$

where ζ_m and X_m are dummy variables, and \mathcal{G} is the modified Green function

$$\mathcal{G}(\zeta, \zeta_m, X) = \sum_{m=0}^{\infty} \frac{\mathcal{H}_m(\zeta) \mathcal{H}_m(\zeta_m)}{\|\mathcal{H}_m^2\|} \mathcal{F}_m(X), \quad \|\mathcal{H}_m^2\| = \int_0^1 u_0 \mathcal{H}_m^2 d\zeta. \quad (4.15)$$

$\mathcal{H}_m(\zeta)$ and $\mathcal{F}_m(X)$ are the solutions associated to the homogeneous part of the equation (4.11). They can be obtained, after separation of variables, following the procedure used

to solve the base-state problem (3.4)-(3.5) for c_0 . The X -problem leads to

$$\mathcal{F}_m(X) = \exp(-\lambda_m^2 X). \quad (4.16)$$

The ζ -problem defines the Sturm–Liouville problem

$$\mathcal{H}_m''(\zeta) + (\lambda_m^2 u_0(\zeta) - k^2) \mathcal{H}_m(\zeta) = 0, \quad (4.17)$$

$$\mathcal{H}_m|_{\zeta=0} = 0, \quad \mathcal{H}'_m|_{\zeta=1} = 0, \quad (4.18)$$

whose solution for the eigenfunctions \mathcal{H}_m (up to a constant factor) is

$$\mathcal{H}_m(\zeta) = \exp\left[-\frac{\lambda_m}{2}(1-\zeta)^2\right] \Phi\left(\frac{k^2 + \lambda_m - \lambda_m^2}{4\lambda_m}, \frac{1}{2}; \lambda_m(1-\zeta)^2\right). \quad (4.19)$$

Substituting (4.19) into the first of the boundary conditions (4.18) provides the transcendental equation for the eigenvalues $\lambda_m > 0$

$$\Phi\left(\frac{k^2 + \lambda_m - \lambda_m^2}{4\lambda_m}, \frac{1}{2}; \lambda_m\right) = 0, \quad (4.20)$$

which highlights how the eigenvalues λ_m are function of the wavenumber k .

We remind that to evaluate the rock dissolution, we only need the concentration gradient at the rock-water interface (see eq. (2.14)). Using the solution (4.14) and separating the terms related to h_1 and η_1 , the gradient at the rock-water interface reads

$$c'_1|_{\zeta=0} = \mathcal{I}_{h_1} h_1 + \mathcal{I}_{\eta_1} \eta_1, \quad (4.21)$$

where the expressions for the integral terms \mathcal{I}_{h_1} and \mathcal{I}_{η_1} are reported in Appendix A.

4.3. Dispersion relationship

At this point, (2.11) and (2.14) are adopted as solvability equations. These describe the temporal evolution of the rock-water and water-air interfaces and, at order ε , they read

$$\omega h_1 = ik\varphi|_{\zeta=1} \quad (4.22)$$

$$\omega \eta_1 = \frac{1}{\gamma Pe} c'_1|_{\zeta=0} - h_1 V. \quad (4.23)$$

Substituting the solutions for the stream function (4.6) and the concentration gradient (4.21), we obtain the linear system

$$\omega \begin{pmatrix} h_1 \\ \eta_1 \end{pmatrix} = \begin{pmatrix} a_1 & a_2 \\ a_3 & a_4 \end{pmatrix} \cdot \begin{pmatrix} h_1 \\ \eta_1 \end{pmatrix}, \quad (4.24)$$

where

$$a_1 = a_2 = \frac{(k^2 Re We + \delta)(k - \cosh k \sinh k)}{k(2k^2 + \cosh(2k) + 1)}, \quad (4.25)$$

$$a_3 = \frac{\mathcal{I}_{h_1}}{\gamma Pe} - V, \quad a_4 = \frac{\mathcal{I}_{\eta_1}}{\gamma Pe}. \quad (4.26)$$

The system (4.24) admits two non-trivial solutions for the eigenvalues

$$\omega_{\eta,h} = \frac{(a_1 + a_4) \pm \sqrt{(a_1 + a_4)^2 - 4a_1(a_4 - a_3)}}{2}, \quad (4.27)$$

which we discriminate with the subscripts h and η , as it will be shown in the following that the former is related to the free surface and the latter to the rock surface.

By assuming the *Rigid Lid Approximation* (RLA) on the free surface i.e., imposing $h_t = 0$ in the kinematic condition (2.11), the first equation in the system (4.24) reduces to $h_1 = -\eta_1$. In this case, the problem provides just one eigenvalue

$$\omega_{\text{RLA}} = V + \frac{(\mathcal{I}_{\eta_1} - \mathcal{I}_{h_1})}{\gamma Pe}. \quad (4.28)$$

By substituting $h_1 = -\eta_1$ into the stream function (4.6), it is also evident that the RLA induces zero spanwise and vertical velocities, i.e., $v_1 = w_1 = 0$. The existence of a secondary flow in the cross-section plane is in fact strictly related to the free surface dynamics. Instead, the streamwise velocity perturbation (4.10) is still non-null as it is triggered by the bottom perturbation. In the next Section, we will discuss on the validity and the convenience of this approximate solution in the context of karren formation.

5. Results

The dispersion relationship (4.27) links the two growth rates, ω_h and ω_η , to the transverse wavenumber k and the control parameters that embody the physics of the problem. In particular, ω_h is associated to development of hydrodynamic waves on the free-surface (rivulets) and ω_η is related to the growth of patterns on the rock surface (karren). Hence, our focus is on the morphological eigenvalue ω_η . The control parameters are: the Reynolds number Re , which is related to the flow rate; the longitudinal coordinate X , which is a proxy for the saturation ratio s (Fig. 4a); the angle of the rock inclination θ ; the parameter γ , which is associated to the mineral type (Tab. 1).

Summarizing, when $\omega_\eta > 0$, the perturbation grows in time (instability) generating the karren pattern. On the opposite, $\omega_\eta < 0$ indicates that the perturbation decays in time (stability) restoring the base-state solution. The condition $\omega_\eta = 0$ discriminates between stable and unstable domains, and its trend in the parameter space evidences the *neutral stability curve*. Figure 5a shows that the rock-water interface is unstable to a band of wavenumbers between 0 and a cutoff value k_c for any Re , i.e., there is no critical Reynolds number for karren formation. The cutoff wavenumber k_c is independent of Re (vertical line in Fig. 5a) meaning that, dimensionally speaking, k_c scales with the flow depth. For a fixed value of Re , the behaviour of the morphological eigenvalue ω_η versus the wavenumber is reported in Figure 5b. The maximum of the growth rate, i.e., the fastest growing mode, is indicated with k_m . As commonly done in linear stability analyses (Cross & Hohenberg 1993), we assume k_m to be pattern wavenumber. Yet, as karren instability shows a band of wavenumbers (indicated with Δk) with growth rates very close to the one of k_m , we will later add some considerations on the validity of this assumption (Sect. 6.2).

Figure 5b also reports the trends of the hydrodynamic eigenvalue ω_h and the approximated morphological eigenvalue ω_{RLA} , obtained in the rigid lid approximation. The hydrodynamic eigenvalue is always stable ($\omega_h < 0$) for the angles of karren formation ($\theta < \pi/2$) due to the stabilizing effects of gravity and surface tension, which prevent any transverse wave (rivulet) to form. It is interesting to notice that gravity becomes instead destabilizing in overhanging conditions ($\theta > \pi/2$), creating water rivulets that initiate longitudinal precipitation patterns (Camporeale 2015; Bertagni & Camporeale 2017).

The morphological eigenvalue ω_{RLA} well reproduces the overall system dynamics when the free-surface responds very quickly to the rock evolution ($|\omega_h| \gg \omega_\eta$) and thus it can be considered flat ($\partial_t h = 0$). This is indeed the case for the wavenumbers involved in karren formation. In fact, $\omega_{\text{RLA}} = \omega_\eta$ for the fastest growing mode k_m . However, ω_{RLA} loses reliability for perturbations with wavelengths so long that hydrodynamic and morphological timescales are comparable ($|\omega_h| \sim \omega_\eta$). Although these very long




FIGURE 5. Karren instability. (a) Contour plot of the least stable growth rate ω_η in the $\{k, Re\}$ plane ($X = 0.05$, $\theta = \pi/4$, calcite). The neutral stability curve is reported with a black solid line. The cutoff k_c and fastest growing k_m wavenumbers are highlighted. The numbers on the contours are the values of ω_η . (b) Growth rates versus the wavenumber k ($X = 0.01$, $Re = 1$). ω_h and ω_η (red solid lines) are the hydrodynamic-stable and morphological-unstable eigenvalues, respectively, from eq. (4.27). Δk is the interval of unstable wavenumbers with growth rate very close to the one of k_m (see Sect. 6.2). ω_{RLA} (dotted light line) is the morphological growth rate obtained in the rigid lid approximation, from eq. (4.28). (c) Role of mass ($\partial_z^2 c$) and momentum ($\partial_z^2 u$) diffusion on ω_η . The light-blue line is ω_η evaluated including v_1 in the solution for u_1 and c_1 (it is indistinguishable from the red line for $k > 10^{-4}$).

wavelengths are outside the range of karren patterns, to neglect them by freezing the free-surface dynamics prevents to mathematically obtain the fastest growing mode k_m . We also notice that the rigid lid approximation introduces an error in the mass conservation ($\omega_{RLA} \neq 0$ for $k=0$). In fact, in the particular case $k=0$, which corresponds to a spatially uniform perturbation of the base state – the so-called Goldstone mode (e.g., Kalliadasis *et al.* 2011) – the base state is vertically and uniformly shifting, and the water depth must preserve its length to assure mass conservation. So the correct solution should be $h_1 = 0$ instead of $h_1 = -\eta_1$ as imposed by the RLA. Yet, the latter approximation may be adopted, for example, for a quicker evaluation of the timescale of the instability as $\omega_{RLA}(k_m) = \omega_\eta(k_m)$.

In Figure 5c, we show that the stabilization of the high wavenumbers (short wavelengths) is induced by a coupling of the transverse diffusion of longitudinal momentum – $\partial_z^2 u$ in eq. (2.3) – and solute – $\partial_z^2 c$ in eq. (2.4) –. In particular, the dashed green and orange lines are the eigenvalues obtained without the transverse diffusion of solute and momentum, respectively. Instead, the dash-dot line is the eigenvalue without both diffusion processes ($\partial_z^2 c$ and $\partial_z^2 u$) that highlights how diffusion is needed for a cutoff mode k_c and a fastest growing mode k_m to arise.

Furthermore, we may notice by observing the dotted line in Fig. 5c that including the vertical velocity v_1 in the solutions of u_1 (4.10) and c_1 (4.14) complicates their analytical expressions (which are not reported here) and causes numerical issues when solving the integrals in (4.21) for small k , but overall it does not affect karren instability (the dotted and solid lines in Fig. 5c are indistinguishable for $k > 10^{-4}$).

The influence of the other control parameters (X , θ , γ) on the morphological growth rate ω_η is shown in Fig. 6. The contourplot 6a in the plane $\{k, X\}$ shows two maxima: one very close to the water inlet ($X < 10^{-2}$) and one further downstream ($X \sim 0.5$). This




FIGURE 6. Influences of X , the rock angle θ and the rock mineral on karren instability for a fixed water flow ($Re = 1$). (a) Contourplot of the morphological growth rate ω_η in the $\{k, X\}$ plane (calcite, $\theta = \pi/4$). The dashed line marks the fastest growing modes. (b) Morphological growth rate ω_η versus k for several angles θ ($\theta = \pi/8, \pi/4, 3\pi/8, \pi/2$, calcite and $X = 0.01$). The stars mark the fastest growing modes. (c) Morphological growth rate ω_η versus k for several mineral types ($\theta = \pi/4$, $X = 0.01$ and the γ from Tab. 1 for the three minerals.)

suggests that the pattern may first develop at two different longitudinal locations. Yet, a fully nonlinear analysis could probably reveal some influence of the upstream part of the growing pattern to the downstream dissolution process. The wavenumber k_m (dashed line) is not strongly affected by X .

For a fixed Re , the angle θ seems to have very little influence on karren instability, besides a stabilizing effect of long wavelengths (Fig. 6b). In contrast, the solubility of the mineral is shown to boost the instability (Fig. 6c), so that the timescale of karren formation with respect to the rock type is salt < gypsum < limestone, in agreement with field observations (Mottershead & Lucas 2001).

6. Discussion

The positive feedback responsible for karren instability can be understood observing the longitudinal velocity and concentration fields in perturbed conditions (Fig. 7). In correspondence of the rock trough, the deeper and faster flow (panel a) more efficiently transports away the solute concentration. This leads to a lower concentration of solute into the water (panels b and c) and to an increased dissolution, which further deepens the rock trough. The opposite happens on the rock crest, where dissolution is dampened. This positive feedback mechanism generates linear karren forms. The spatial scale of the instability is influenced by the transverse diffusive processes of longitudinal velocity and solute (Fig. 5c). In Fig. 7, we may also notice that the free-surface is basically flat, further indicating the validity of the rigid lid approximation for the wavenumber of karren formation ($k \sim k_m$).

6.1. Dimensional features

The fastest growing mode k_m of the linear stability analysis provides the dimensional wavelength $\hat{L} = 2\pi/k_m$ and an indication of the timescale of the instability $\hat{T} = 1/\hat{\omega}_\eta(k_m)$. Figure 8 explores the dependence of these dimensional quantities on the control parameters Re and X .

The water flow rate (Re) is shown to have a crucial role in karren formation. In fact, \hat{L} increase with the Reynolds number, spanning several orders of magnitude: from millimetric to decimetric values (Fig. 8a). This result supports the so far speculative idea




FIGURE 7. Instability mechanism. Transverse sections (vertically exaggerated) showing the response of the longitudinal velocity u_1 (a) and the concentration c_1 (b-c) to a small perturbation of the rock surface (calcite, $Re = 1$, $\theta = \pi/4$, $\varepsilon = 0.05$, $k = k_m$, $X = 0.01$ for panel b and $X = 0.1$ for panel c). c_1 varies along X inheriting the longitudinal trend of the concentration c_0 at base-state (see Fig. 3).




FIGURE 8. Contourplots in the $\{X, Re\}$ plane for the dimensional wavelength \hat{L} (a) and timescale of the instability \hat{T} (b) from the linear stability analysis (calcite and $\theta = \pi/4$). The upper x -axis reports the saturation ratio s .

that is the flow rate to discriminate the linear karren type: very thin water films ($Re \sim 10^{-3}$) originated by dew or sea spray give rise to millimetric microrills; thicker water films ($Re \sim 10^{-1} - 1$) produced by rainwater are responsible for the formation of centimetric rillenkarren; and decimetric wandkarren are formed by high water flows ($Re > 10$), which develop downstream of a basin where rain or snow-melt water accumulates (e.g., on a

hill slope). Even \hat{T} is highly dependent on Re , ranging from tens of minutes to weeks (Fig. 8b). These timescales seem pretty short. Indeed, they are closer to experimental evidences of karren formation under a constant water flow (Gu erin *et al.* 2020), rather than to field observations (Mottershead & Lucas 2001), where the intermittent nature of rains prolongs the karren development. Furthermore, the reason for the low values of \hat{T} may be twofold: the diffusion-controlled dissolution kinetics may overestimate the karren-formation rate as the temporal transients of the slow chemical reactions are neglected; the linear stability analysis provides just an indication of the timescale of the instability, while nonlinear effects are expected to slow down the pattern formation (by roughly an order of magnitude if nonlinearities are included into a classical Landau equation (Cross & Hohenberg 1993)).

The wavelength and the timescale very weakly depend on the longitudinal coordinate X , and thus on the level of saturation of water (the values of s are reported on the upper x-axis). In particular, the timescale \hat{T} shows a maximum at $X \sim 0.5$ (for a fixed Re), which is due to the second maximum in the growth rate ω_η shown in Fig. 6a.

6.2. Open issues

The water that shapes some linear karren forms, such as rillenkarren, normally comes from rain events (Gin es *et al.* 2009). For this reason, experimental efforts have mainly focused on reproducing rillenkarren by dissolving blocks of minerals under artificial rain (Glew & Ford 1980; Slabe *et al.* 2016). Although rain might affect some features of karren formation (as we discuss in the following), our model shows that a water flow – without falling raindrops being involved – is sufficient to generate linear karren forms. This is in agreement with some notable recent experiments by Gu erin *et al.* (2020), who observed initially millimetric rills forming on plaster blocks dissolving under the action of a wall-induced turbulent flow ($200 < Re < 700$)[†]. One experimental run was in the laminar regime ($Re \sim 54$) and the same millimetric rills were first observed to arise. The experiments have also shown that, as the dissolution process endures and the effect of the nonlinearities grow, the rills coalesce in time to eventually form wider karren patterns. The range of Reynolds number that was experimentally investigated and the narrow width of the plaster blocks (10 cm) limit the possibility of a deeper comparison with our theory, which is developed under the assumption of a laminar flow and predicts a linear wavenumber, for $Re \sim 54$, that is the same order of the block width. Further research, both at the theoretical and experimental levels, might shed more light on the complex dynamics that create these fascinating patterns.

Here we further discuss on how rain might affect the present model and the results of the linear stability analysis. A first interesting aspect regards rain stochasticity, that implies that the water flow is intermittent and highly variable (Re changes in time). This discloses several questions on the influence of stochasticity on rillenkarren formation, e.g., are extreme events more efficient? Does it exist a constant flow that would create the same rock morphology produced by the stochastic sequence of rain events? The answers remain undisclosed for further numerical or experimental research. A second aspect regards rain spatial extension, which makes it a diffuse source of water: raindrops falling along the karren length add pure water to the flow, introducing X -dependencies in several parameters (e.g., Re raises, the saturation ratio s decreases, and the penetration length \hat{L} increases).

Rain might also affect some outputs of the linear stability analysis, such as the selected

[†] The experimental Reynolds number (Re_{exp}) is defined through the depth-averaged velocity instead of the free-surface velocity, i.e., $Re_{exp} \sim 2 Re/3$

wavenumber k_m . In fact, the classical assumption of the fastest growing mode as the pattern wavenumber is based on the hypotheses that: i) the rock-water system is initially flat; ii) all modes are decoupled as they are initially perturbed with an infinitesimal perturbation; iii) the fastest growing mode overcomes the others. However, Fig. 5b shows that there is an interval of unstable wavenumbers, indicated with Δk , with growth rates very close to the one associated to k_m . This might indicate that, when more realistic finite perturbations are taken into account, any mode within the interval Δk may overcome the others (k_m included) and become the pattern wavenumber. In this perspective, raindrops, which have a characteristic (sub-)millimetric length-scale, may act as a random spatial forcing on the water film and boost the growth of a mode with a comparable length-scale. In a similar way, defects in the rock texture, moss, decantation wells and preferential channels for the water flow upstream of the rock, may act as spatial forcings and affect the selection of the pattern wavenumber.

7. Summary and Conclusion

In this manuscript, we have framed the pivotal role of water hydrodynamics into a theoretical model that explains the genesis of linear karren forms. The model addresses a film of water that flows on a dissolving rock under some simplifying assumptions: i) the water film is laminar; ii) the overall chemistry is reduced to the solute concentration; iii) diffusion is considered as the limiting process in the dissolution kinetics.

The linear stability analysis of the flat solution has revealed that a transverse instability is responsible for the appearance of longitudinal parallel channels on the rock surface. The instability mechanism is a positive feedback between the increased flow rate within the channels and the enhanced dissolution (Fig. 7). The results show that the transverse diffusive processes of solute and longitudinal velocity stabilize the high wavenumbers and affect the wavenumber selection (Fig. 5c). Furthermore, the flow rate is shown to be a discriminating factor in the transverse wavelength of the pattern (Fig. 8), possibly explaining why many similar linear karren of different sizes are observed in karst areas (Fig. 1). The results also suggest that, even though rain might affect some features of karren formation (as discussed in Sec. 6.2), a water film without falling raindrops is sufficient for the pattern occurrence.

Although the linear stability analysis provides encouraging results, we shall remind that these are strictly valid just at the pattern genesis, i.e., before nonlinear effects induced by the finite size of the pattern become preponderant in the overall dynamics. Hence, a (fully) nonlinear analysis could greatly benefit to the present formulation, providing further insights on several features of the pattern formation: such as the nonlinear wavelength selection and the amplitude growth.

We thank the research group at the Paris Diderot University (M. Berhanu, A. Guérin, J. Derr, S. Courrech du Pont) for interesting discussions on karren formation. We are grateful to Jost von Hardenberg for his support in the numerical coding. We also acknowledge the anonymous Referees for their valuable review work.

Appendix A. Integral terms

The terms in eq. (4.21) read

$$\mathcal{I}_{h_1} = \sum_{m=0}^{\infty} \mathcal{H}'_m(0) \int_0^X \int_0^1 \Upsilon_{h_1}(\zeta_m, X_m) \frac{\mathcal{H}_m(\zeta_m)}{\|\mathcal{H}_m^2\|} \mathcal{F}_m(X - X_m) d\zeta_m dX_m,$$

$$\mathcal{I}_{\eta_1} = \sum_{m=0}^{\infty} \mathcal{H}'_m(0) \int_0^X \int_0^1 \Upsilon_{\eta_1}(\zeta_m, X_m) \frac{\mathcal{H}_m(\zeta_m)}{\|\mathcal{H}_m^2\|} \mathcal{F}_m(X - X_m) d\zeta_m dX_m,$$

where the inhomogeneous term (4.12) has been divided into $\Upsilon = \Upsilon_{h_1} h_1 + \Upsilon_{\eta_1} \eta_1$ and

$$\Upsilon_{h_1}(\zeta, X) = \left[k^2 \zeta \partial_\zeta c_0 - 2\partial_\zeta^2 c_0 - 2\partial_X c_0 \left(\frac{\operatorname{sech} k \sinh k\zeta}{k} + \zeta(1 - \zeta) \right) \right], \quad (\text{A } 1)$$

$$\Upsilon_{\eta_1}(\zeta, X) = \left\{ k^2 \partial_\zeta c_0 - 2\partial_X c_0 \left[1 - \zeta - \operatorname{sech} k \left(\cosh(k - k\zeta) - \frac{\sinh(k\zeta)}{k} \right) \right] \right\}. \quad (\text{A } 2)$$

Notice that the contribution of v_1 in Υ has been neglected as it would considerably complicate the solution without any perceptible numerical change

REFERENCES

- ABRAMOWITZ, M., STEGUN, I.A. & OTHERS 1966 Handbook of mathematical functions. *Appl. Math. Series* **55**, 62.
- BENDER, C.M. & ORSZAG, S.A. 2013 *Advanced mathematical methods for scientists and engineers I: Asymptotic methods and perturbation theory*. Springer Science & Business Media.
- BERNER, R A, LASAGA, A C & GARRELS, R M 1983 The carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years. *American Journal of Science* **283**, 641–683.
- BERTAGNI, M.B. & CAMPOREALE, C. 2017 Nonlinear and subharmonic stability analysis in film-driven morphological patterns. *Phys. Rev. E* **96** (5), 053115.
- BÖGLI, A. 1960 Kalklösung und karrenbildung. *Zeitschrift für Geomorphologie* (Supplementband 2), 4–21.
- BUHMANN, D. & DREYBRODT, W. 1985 The kinetics of calcite dissolution and precipitation in geologically relevant situations of karst areas: 1. Open system. *Chemical geology* **48** (1-4), 189–211.
- CAMPOREALE, C. 2015 Hydrodynamically locked morphogenesis in karst and ice flutings. *J. Fluid Mech.* **778**, 89–119.
- CAMPOREALE, C. 2017 An asymptotic approach to the crenulation instability. *J. Fluid Mech.* **826**, 636–652.
- CAMPOREALE, C. & RIDOLFI, L. 2012 Hydrodynamic-driven stability analysis of morphological patterns on stalactites and implications for cave paleoflow reconstructions. *Phys. Rev. Lett.* **108** (23), 238501.
- CHANG, H. & DEMEKHIN, E. A. 2002 *Complex wave dynamics on thin films*. Elsevier.
- CLAUDIN, P., DURÁN, O. & ANDREOTTI, B. 2017 Dissolution instability and roughening transition. *J. Fluid Mech.* **832**.
- COHEN, CAROLINE, BERHANU, MICHAEL, DERR, JULIEN & DU PONT, SYLVAIN COURRECH 2020 Buoyancy-driven dissolution of inclined blocks: Erosion rate and pattern formation. *Physical Review Fluids* **5** (5), 053802.
- CROSS, M. & HOHENBERG, P.C. 1993 Pattern formation outside of equilibrium. *Rev. Mod. Phys.* **65** (3), 851.
- CURL, R.L. 1966 Scallops and flutes .
- DAVIS, E J 1973 Exact solutions for a class of heat and mass transfer problems. *The Canadian Journal of Chemical Engineering* **51** (5), 562–572.
- DREYBRODT, W. 2012 *Processes in karst systems: physics, chemistry, and geology*, , vol. 4. Springer Science & Business Media.
- FAIRCHILD, I J & BAKER, A 2012 *Speleothem science: from process to past environments*, , vol. 3. John Wiley & Sons.
- FORD, D & WILLIAMS, P D 2013 *Karst hydrogeology and geomorphology*. John Wiley & Sons.
- GINÉS, A, KNEZ, M, SLABE, T & DREYBRODT, W 2009 *Karst Rock Features. Karren sculpturing*, , vol. 9. Založba ZRC.

- GLEW, JR & FORD, DC 1980 A simulation study of the development of rillenkarren. *Earth Surface Processes* **5** (1), 25–36.
- GOLDENFELD, N., CHAN, P. Y. & VEYSEY, J. 2006 Dynamics of precipitation pattern formation at geothermal hot springs. *Phys. Rev. Lett.* **96** (25), 254501.
- GUÉRIN, ADRIEN, DERR, JULIEN, DU PONT, SYLVAIN COURRECH & BERHANU, MICHAEL 2020 Streamwise dissolution patterns created by a flowing water film. *Physical Review Letters* **125** (19), 194502.
- HAMMER, Ø, DYSTHE, DK, LELU, B, LUND, H, MEAKIN, P & JAMTVEIT, B 2008 Calcite precipitation instability under laminar, open-channel flow. *Geochimica et Cosmochimica Acta* **72** (20), 5009–5021.
- JAMTVEIT, B & HAMMER, Ø 2012 Sculpting of rocks by reactive fluids. *Geochemical Perspectives* **1** (3), 341–342.
- KALLIADASIS, S., RUYER-QUIL, C., SCHEID, B. & VELARDE, M. G. 2011 *Falling liquid films*, vol. 176. Springer Science & Business Media.
- KAUFMANN, G. & DREYBRODT, W. 2007 Calcite dissolution kinetics in the system $\text{CaCO}_3\text{-H}_2\text{O-CO}_2$ at high undersaturation. *Geochim. Cosmochim. Acta* **71** (6), 1398–1410.
- MAC HUANG, JINZI, MOORE, M NICHOLAS J & RISTROPH, LEIF 2015 Shape dynamics and scaling laws for a body dissolving in fluid flow. *Journal of Fluid Mechanics* **765**.
- MEAKIN, P. & JAMTVEIT, B. 2009 Geological pattern formation by growth and dissolution in aqueous systems. *Proc R Soc Lond A Math Phys Sci* p. rspa20090189.
- MOORE, M NICHOLAS J 2017 Riemann-hilbert problems for the shapes formed by bodies dissolving, melting, and eroding in fluid flows. *Communications on Pure and Applied Mathematics* **70** (9), 1810–1831.
- MORROW, LIAM C, KING, JOHN R, MORONEY, TIMOTHY J & MCCUE, SCOTT W 2019 Moving boundary problems for quasi-steady conduction limited melting. *SIAM Journal on Applied Mathematics* **79** (5), 2107–2131.
- MORSE, J W & ARVIDSON, R S 2002 The dissolution kinetics of major sedimentary carbonate minerals. *Earth-Science Reviews* **58** (1-2), 51–84.
- MOTTERSHEAD, D & LUCAS, G 2001 Field testing of Glew and Ford’s model of solution flute evolution. *Earth Surf Proc Land* **26** (8), 839–846.
- NUSSELT, W. 1916 Die oberflächenkondensation des wasserdampfes the surface condensation of water. *Zetschr. Ver. Deutch. Ing.* **60**, 541–546.
- PLUMMER, LN, WIGLEY, TML & PARKHURST, DL 1978 The kinetics of calcite dissolution in co 2-water systems at 5 degrees to 60 degrees c and 0.0 to 1.0 atm co 2. *American journal of science* **278** (2), 179–216.
- POLYANIN, A D, KUTEPOV, AM, KAZENIN, DA & VYAZMIN, AV 2001 *Hydrodynamics, mass and heat transfer in chemical engineering*, , vol. 14. CRC Press.
- POLYANIN, A D & NAZAIKINSKII, V E 2015 *Handbook of linear partial differential equations for engineers and scientists*. CRC press.
- REGNIER, P, FRIEDLINGSTEIN, P, CIAIS, P, MACKENZIE, F T, GRUBER, N, JANSSENS, I A, LARUELLE, G G, LAUERWALD, R, LUYSSAERT, S, ANDERSSON, A J & OTHERS 2013 Anthropogenic perturbation of the carbon fluxes from land to ocean. *Nat. Geosci.* **6** (8), 597–607.
- SAURO, U 2009 I Karren: le forme scolpite sulla roccia.
- SHORT, M. B., BAYGENTS, J. C. & GOLDSTEIN, R. E. 2005 Stalactite growth as a free-boundary problem. *Phys. Fluids (1994-present)* **17** (8), 083101.
- SLABE, T, HADA, A & KNEZ, M 2016 Laboratory modeling of karst phenomena and their rock relief on plaster: subsoil karren, rain flutes karren and caves. *Acta Carsologica* **45** (2).
- VESIPA, R, CAMPOREALE, C & RIDOLFI, L 2015 Thin-film-induced morphological instabilities over calcite surfaces. *Proc R Soc Lond A Math Phys Sci* **471** (2176), 20150031.
- VEYSEY, J & GOLDENFELD, N 2008 Watching rocks grow. *Nat. Phys.* **4** (4), 310–313.
- WYKES, MEGAN S DAVIES, MAC HUANG, JINZI, HAJJAR, GEORGE A & RISTROPH, LEIF 2018 Self-sculpting of a dissolvable body due to gravitational convection. *Physical Review Fluids* **3** (4), 043801.