

N = 2 supersymmetric S-folds

*Original*

N = 2 supersymmetric S-folds / Guarino, A., Sterckx, C., Trigiante, M.. - In: JOURNAL OF HIGH ENERGY PHYSICS. - ISSN 1029-8479. - ELETTRONICO. - 2020:4(2020). [10.1007/JHEP04(2020)050]

*Availability:*

This version is available at: 11583/2885494 since: 2021-04-08T10:29:28Z

*Publisher:*

Springer

*Published*

DOI:10.1007/JHEP04(2020)050

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

## $\mathcal{N} = 2$ supersymmetric S-folds

---

Adolfo Guarino,<sup>a,b</sup> Colin Sterckx<sup>c</sup> and Mario Trigiante<sup>d</sup>

<sup>a</sup>*Departamento de Física, Universidad de Oviedo,  
Avda. Federico García Lorca 18, 33007 Oviedo, Spain*

<sup>b</sup>*Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA),  
Calle de la Independencia 13, 33004 Oviedo, Spain*

<sup>c</sup>*Université Libre de Bruxelles (ULB) and International Solvay Institutes,  
Service de Physique Théorique et Mathématique,  
Campus de la Plaine, CP 231, B-1050, Brussels, Belgium*

<sup>d</sup>*Dipartimento di Fisica, Politecnico di Torino, Corso Duca degli Abruzzi 24,  
I-10129 Turin, Italy and INFN, Sezione di Torino, Italy*

*E-mail:* [adolfo.guarino@uniovi.es](mailto:adolfo.guarino@uniovi.es), [colin.sterckx@ulb.ac.be](mailto:colin.sterckx@ulb.ac.be),  
[mario.trigiante@polito.it](mailto:mario.trigiante@polito.it)

**ABSTRACT:** Multi-parametric families of  $\text{AdS}_4$  vacua with various amounts of supersymmetry and residual gauge symmetry are found in the  $[\text{SO}(1,1) \times \text{SO}(6)] \times \mathbb{R}^{12}$  maximal supergravity that arises from the reduction of type IIB supergravity on  $\mathbb{R} \times \text{S}^5$ . These provide natural candidates to holographically describe new strongly coupled three-dimensional CFT's which are localised on interfaces of  $\mathcal{N} = 4$  super-Yang-Mills theory. One such  $\text{AdS}_4$  vacua features a symmetry enhancement to  $\text{SU}(2) \times \text{U}(1)$  while preserving  $\mathcal{N} = 2$  supersymmetry. Fetching techniques from the  $\text{E}_{7(7)}$  exceptional field theory, its uplift to a class of  $\mathcal{N} = 2$  S-folds of type IIB supergravity of the form  $\text{AdS}_4 \times \text{S}^1 \times \text{S}^5$  involving S-duality twists of hyperbolic type along  $\text{S}^1$  is presented.

**KEYWORDS:** Flux compactifications, String Duality, Supergravity Models, Superstring Vacua

ARXIV EPRINT: [2002.03692](https://arxiv.org/abs/2002.03692)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>AdS<sub>4</sub> vacua of [SO(1, 1) × SO(6)] × ℝ<sup>12</sup> maximal supergravity</b>	<b>3</b>
2.1	The $\mathcal{N} = 8$ theory: gauging and scalar potential	4
2.2	$\mathbb{Z}_2^3$ invariant sector	5
2.3	New families of AdS <sub>4</sub> vacua	6
2.3.1	$\mathcal{N} = 0$ vacua with $SU(2) \rightarrow SU(2) \times U(1)^2 \rightarrow SU(3) \times U(1) \rightarrow SO(6)$ symmetry	7
2.3.2	$\mathcal{N} = 1$ vacua with $U(1)^2 \rightarrow SU(2) \times U(1) \rightarrow SU(3)$ symmetry	7
2.3.3	$\mathcal{N} = 2$ vacua with $U(1)^2 \rightarrow SU(2) \times U(1)$ symmetry	8
2.3.4	$\mathcal{N} = 4$ vacuum with SO(4) symmetry	9
<b>3</b>	<b>S-folds with <math>\mathcal{N} = 2</math> supersymmetry</b>	<b>9</b>
3.1	Type IIB uplift using E <sub>7(7)</sub> -EFT	10
3.2	S-fold interpretation	16
3.3	Connection with Janus-like solutions	17
<b>4</b>	<b>Conclusions</b>	<b>18</b>

---

## 1 Introduction


Electromagnetic duality in four-dimensional maximal supergravity has provided a very rich phenomenology as far as the existence of new gaugings and vacuum solutions are concerned. The prototypical example is the dyonically-gauged SO(8) supergravity where the action of electromagnetic duality on the gauging generates a one-parameter family of inequivalent theories parameterised by a continuous parameter  $c \in [0, \sqrt{2} - 1]$  [1]. Setting the parameter to  $c = 0$  then the standard (electric) SO(8) supergravity of de Wit and Nicolai [2] is recovered which is known to arise upon dimensional reduction of eleven-dimensional supergravity on a seven-sphere  $S^7$ . The various AdS<sub>4</sub> vacua of the  $c = 0$  theory [3] (see also [4] for an updated encyclopedic reference) get generalised to one-parameter families of vacua when turning on  $c$  and, more importantly, new and genuinely dyonic AdS<sub>4</sub> vacua also appear which do not have a well defined (electric)  $c \rightarrow 0$  limit [1, 5–7]. Other types of four-dimensional solutions, like domain-walls [8, 9] or black holes [10–12], have also been investigated using instead a phase-like parameterisation  $\omega = \arg(1 + ic) \in [0, \pi/8]$  of the electromagnetic deformation parameter. However, and despite the rich structure of new solutions at  $c \neq 0$ , the question about the eleven-dimensional interpretation of the electromagnetic parameter  $c$  remains elusive and various no-go theorems have been stated against the existence of such a higher dimensional origin [13, 14]. Also, for the new supersymmetric

AdS<sub>4</sub> vacua at  $c \neq 0$ , the holographic interpretation of the deformation parameter remains obscure from the perspective of the AdS<sub>4</sub>/CFT<sub>3</sub> correspondence.

Unlike for the SO(8) theory, much more is by now known about the dyonically-gauged ISO(7) supergravity that arises from the reduction of massive IIA supergravity on a six-sphere S<sup>6</sup> [15]. In this case the electromagnetic deformation parameter is a discrete (on/off) deformation, namely, it can be set to  $c = 0$  or  $1$  without loss of generality [16]. Various AdS<sub>4</sub> [17–19], domain-wall [19, 20], and black hole [21–24] solutions have been constructed which necessarily require a non-zero electromagnetic deformation parameter  $c$ . Within this massive IIA context, the electromagnetic parameter is identified with the Romans mass parameter  $\hat{F}_0$  of the ten-dimensional theory [25], and has a holographic interpretation as the Chern-Simons level  $k$  of a three-dimensional super-Chern-Simons dual theory [26].

The role of the electromagnetic deformation  $c$  has been much less investigated in the context of type IIB supergravity. The relevant dyonically-gauged supergravity in this case is the  $[\text{SO}(1,1) \times \text{SO}(6)] \ltimes \mathbb{R}^{12}$  theory which arises from the reduction of type IIB supergravity on the product  $\mathbb{R} \times \text{S}^5$  [27]. As for the ISO(7) theory, the electromagnetic deformation is again a discrete (on/off) deformation, namely,  $c = 0$  or  $1$  [16]. This four-dimensional supergravity has been shown to contain various types of AdS<sub>4</sub> vacua preserving different amounts of supersymmetry as well as of residual gauge symmetry. In particular, an  $\mathcal{N} = 4$  and SO(4) symmetric solution was reported in [28] and subsequently, in [27], uplifted to a class of AdS<sub>4</sub>  $\times$  S<sup>1</sup>  $\times$  S<sup>5</sup> S-fold backgrounds of type IIB supergravity using the E<sub>7(7)</sub> exceptional field theory (E<sub>7(7)</sub>-EFT). These S-folds involve S-duality twists  $A_{(k)}$  ( $k \geq 3$ ) that induce  $\text{SL}(2, \mathbb{Z})_{\text{IIB}}$  monodromies  $\mathfrak{M}(k) = -\mathcal{ST}^k$  of hyperbolic type along S<sup>1</sup>, and can be systematically constructed as quotients of degenerate Janus-like solutions of the type IIB theory [29, 30] where the string coupling  $g_s$  diverges at infinity. Together with the  $\mathcal{N} = 4$  & SO(4) solution, additional  $\mathcal{N} = 0$  & SO(6) [31] and  $\mathcal{N} = 1$  & SU(3) [32] solutions have been found and uplifted to similar S-fold backgrounds of type IIB supergravity with hyperbolic monodromies in [32]. From a holographic perspective, these AdS<sub>4</sub> vacua describe new strongly coupled three-dimensional CFT's, referred to as  $J$ -fold CFT's in [33] (see also [34, 35] and [36]), which are localised on interfaces of  $\mathcal{N} = 4$  super-Yang-Mills theory (SYM) [37]. In the  $\mathcal{N} = 4$  case [33], a hyperbolic monodromy  $J = -\mathcal{ST}^k \in \text{SL}(2, \mathbb{Z})_{\text{IIB}}$  was shown to introduce a Chern-Simons level  $k$  in the dual  $J$ -fold CFT which, in turn, is constructed from the  $T(\text{U}(N))$  theory [38] upon suitable gauging of flavour symmetries. A diagram illustrating this type IIB construction is depicted in figure 1.

On the other hand, a classification of interface SYM theories was performed in [39] (see also [40]) in correspondence to the various amounts of supersymmetry, as well as the largest possible global symmetry, preserved by the interface operators. Three supersymmetric cases were identified: interfaces with  $\mathcal{N} = 4$  & SO(4) symmetry,  $\mathcal{N} = 2$  & SU(2)  $\times$  U(1) symmetry and  $\mathcal{N} = 1$  & SU(3) symmetry. While the S-folds in [27] and [32] respectively match the symmetries of the  $\mathcal{N} = 4$  and  $\mathcal{N} = 1$  cases, the gravity duals of the would be  $\mathcal{N} = 2$   $J$ -fold CFT's localised on the interface with SU(2)  $\times$  U(1) symmetry remain missing. In this work we fill this gap and present a new family of AdS<sub>4</sub>  $\times$  S<sup>1</sup>  $\times$  S<sup>5</sup> S-folds with  $\mathcal{N} = 2$  supersymmetry, SU(2)  $\times$  U(1) symmetry and, as in the previous cases, involving S-duality twists that induce monodromies of hyperbolic type along S<sup>1</sup>.



**Figure 1.** Type IIB S-folds with hyperbolic monodromies  $\mathfrak{M}(k) = -ST^k$  along  $S^1$  and connection with three-dimensional  $J$ -fold CFT's.

The paper is organised as follows. In section 2 we perform a study of multi-parametric families of  $AdS_4$  vacua in the  $[SO(1, 1) \times SO(6)] \times \mathbb{R}^{12}$  maximal supergravity. We find four families of vacua, one of them being  $\mathcal{N} = 2$  supersymmetric and containing a vacuum with a residual symmetry enhancement to  $SU(2) \times U(1)$ . In section 3, by implementing a generalised Scherk-Schwarz (S-S) ansatz in  $E_{7(7)}$ -EFT, we uplift such an  $AdS_4$  vacuum to a class of  $AdS_4 \times S^1 \times S^5$   $\mathcal{N} = 2$  S-folds of type IIB supergravity with  $SU(2) \times U(1)$  symmetry and a non-trivial hyperbolic monodromy along  $S^1$ . In section 4 we present our conclusions and discuss future directions.

## 2 $AdS_4$ vacua of $[SO(1, 1) \times SO(6)] \times \mathbb{R}^{12}$ maximal supergravity

We continue the study of  $AdS_4$  vacua initiated in [31], and further investigated in [28] and [32], for the dyonically-gauged maximal supergravity with non-abelian gauge group

$$G = [SO(1, 1) \times SO(6)] \times \mathbb{R}^{12}. \tag{2.1}$$

We will show how the  $AdS_4$  vacua of [28, 31, 32] actually correspond to very special points (featuring residual symmetry enhancements) within multi-parametric families of solutions. Each of these families preserves a given amount supersymmetry, namely,  $\mathcal{N} = 0, 1, 2$  or 4. More specifically we find:

- A three-parameter family of  $\mathcal{N} = 0$  &  $SU(2)$  symmetric  $AdS_4$  vacua with symmetry enhancements to  $SU(2) \times U(1)^2$ ,  $SU(3) \times U(1)$  and  $SO(6) \sim SU(4)$  at specific values of the three arbitrary parameters.
- A two-parameter family of  $\mathcal{N} = 1$  &  $U(1)^2$  symmetric  $AdS_4$  vacua with symmetry enhancements to  $SU(2) \times U(1)$  and  $SU(3)$  at specific values of the two arbitrary parameters.
- A one-parameter family of  $\mathcal{N} = 2$  &  $U(1)^2$  symmetric  $AdS_4$  vacua with a symmetry enhancement to  $SU(2) \times U(1)$  at a special value of the arbitrary parameter.
- A single  $\mathcal{N} = 4$  &  $SO(4)$  symmetric  $AdS_4$  vacuum.

The  $\mathcal{N} = 2$  family of AdS<sub>4</sub> vacua is new and we will uplift the solution with  $SU(2) \times U(1)$  enhanced residual symmetry to a new and analytic family of S-fold backgrounds of type IIB supergravity in section 3.

### 2.1 The $\mathcal{N} = 8$ theory: gauging and scalar potential

We follow the conventions and notation of [32], which slightly differ from those of [27], to describe the dyonically-gauged maximal supergravity with gauge group  $G$  in (2.1). For the purposes of this work, i.e. the study of AdS<sub>4</sub> vacua, we set to zero all the vector and (auxiliary [41]) tensor fields of the theory, so that the bosonic Lagrangian reduces to the following one

$$\mathcal{L}_{\mathcal{N}=8} = \left( \frac{R}{2} - V_{\mathcal{N}=8} \right) * 1 + \frac{1}{96} \text{Tr} (dM \wedge *dM^{-1}), \quad (2.2)$$

which describes the scalar fields  $M_{MN}$  coupled to Einstein gravity in the presence of a scalar potential. The scalar fields serve as coordinates on the coset space of maximal supergravity

$$M_{MN} = \mathcal{V} \mathcal{V}^t \in \frac{E_{7(7)}}{SU(8)}, \quad (2.3)$$

with  $M = 1, \dots, 56$  being a fundamental index of  $E_{7(7)}$ . The coset representative  $\mathcal{V}$  is constructed by direct exponentiation of the 70 non-compact generators  $t_A{}^B$  (with  $t_A{}^A = 0$ ) and  $t_{ABCD} = t_{[ABCD]}$  generators of  $E_{7(7)}$  in the  $SL(8)$  basis.<sup>1</sup> The scalar potential in (2.2), which survives our truncation to the Einstein-scalar sector, is induced by the gauging of the group  $G$  in (2.1) within the maximal theory and has the following general form:

$$V_{\mathcal{N}=8} = \frac{g^2}{672} X_{MN}{}^R X_{PQ}{}^S M^{MP} \left( M^{NQ} M_{RS} + 7 \delta_R^Q \delta_S^N \right), \quad (2.4)$$

which depends on the gauge coupling  $g$ , the scalar matrix  $M_{MN}$  (and its inverse  $M^{MN}$ ) and on a constant *embedding tensor*  $X_{MN}{}^P$  living in the **912** of  $E_{7(7)}$  [43]. This tensor codifies how the gauge group  $G$  is embedded into the  $E_{7(7)}$  duality group of maximal supergravity. Moreover, it also specifies the gauge connection which involves both electric and magnetic vector fields transforming under the  $Sp(56)$  group of electromagnetic transformations of the theory (for reviews see [44, 45]).

Under  $SL(8) \subset E_{7(7)}$  the index  $M$  decomposes into antisymmetric pairs  $M = ([AB], [AB])$  where  $A = 1, \dots, 8$  denotes a fundamental index of  $SL(8)$ . This implies that, for gaugings of subgroups of  $SL(8)$ , the non-vanishing electric and magnetic components of the embedding tensor are given by [31]

$$\begin{aligned} \text{electric :} \quad & X_{[AB][CD]}{}^{[EF]} = -X_{[AB]}{}^{[EF]}{}_{[CD]} = -8 \delta_{[A}^{[E} \eta_{B][C} \delta_{D]}^F], \\ \text{magnetic :} \quad & X^{[AB]}{}_{[CD]}{}^{[EF]} = -X^{[AB][EF]}{}_{[CD]} = -8 \delta_{[C}^{[A} \tilde{\eta}^{B][E} \delta_{D]}^F], \end{aligned} \quad (2.5)$$

in terms of two symmetric matrices  $\eta_{AB}$  and  $\tilde{\eta}^{AB}$ . For the gauging of  $G \subset SL(8)$  in (2.1) these are

$$\eta_{AB} = \text{diag}(0, \mathbb{I}_6, 0) \quad \text{and} \quad \tilde{\eta}^{AB} = c \text{diag}(-1, 0_6, 1). \quad (2.6)$$

---

<sup>1</sup>We adopt the conventions in the appendix of [42] for the explicit form of the  $t_A{}^B$  and  $t_{ABCD}$  matrices.

As stated in the introduction, the magnetic part of the embedding tensor in (2.5) allows for an (on/off) electromagnetic parameter  $c$  so that  $\tilde{\eta}^{AB} \propto c$ .

## 2.2 $\mathbb{Z}_2^3$ invariant sector

In order to efficiently search for extrema of the scalar potential (2.4), we will now construct a  $\mathbb{Z}_2^3$  invariant sector of the  $[\text{SO}(1,1) \times \text{SO}(6)] \ltimes \mathbb{R}^{12}$  maximal supergravity. This sector can be recast as a minimal  $\mathcal{N} = 1$  supergravity coupled to seven chiral multiplets  $z_i$  with  $i = 1, \dots, 7$ . The same invariant sector has recently been explored in the dyonically-gauged ISO(7) theory [19] and the purely electric SO(8) theory [46], and it originally appeared in the context of type II orientifold compactifications with generalised fluxes [47, 48].

To describe this sector of the maximal theory, we first focus on a four-element Klein subgroup of SL(8). Its action on the fundamental index  $A$  is given by

$$\begin{aligned} \mathbb{Z}_2^{(1)} : (x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8) &\rightarrow (x_1, x_2, x_3, -x_4, -x_5, -x_6, -x_7, x_8), \\ \mathbb{Z}_2^{(2)} : (x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8) &\rightarrow (x_1, -x_2, -x_3, x_4, x_5, -x_6, -x_7, x_8), \end{aligned} \tag{2.7}$$

together with the remaining generators  $\mathbb{I}$  and  $\mathbb{Z}_2^{(1)}\mathbb{Z}_2^{(2)}$ . In addition, we will also require invariance under an extra  $\mathbb{Z}_2^*$  generator acting as

$$\mathbb{Z}_2^* : (x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8) \rightarrow (x_1, -x_2, x_3, -x_4, x_5, -x_6, x_7, -x_8). \tag{2.8}$$

The resulting  $\mathbb{Z}_2^3$  invariant sector describes  $\mathcal{N} = 1$  supergravity coupled to seven chiral multiplets (and no vector multiplets)

$$z_i = -\chi_i + ie^{-\varphi_i} \quad \text{with} \quad i = 1, \dots, 7. \tag{2.9}$$

The fourteen real spinless fields are associated with generators  $t_A^B$  (scalars) and  $t_{[ABCD]}$  (pseudo-scalars) of  $E_{7(7)}$  in the SL(8) basis. The former have associated generators of the form

$$\begin{aligned} g_{\varphi_1} &= -t_1^1 - t_2^2 - t_3^3 + t_4^4 + t_5^5 + t_6^6 + t_7^7 - t_8^8, \\ g_{\varphi_2} &= -t_1^1 + t_2^2 + t_3^3 - t_4^4 - t_5^5 + t_6^6 + t_7^7 - t_8^8, \\ g_{\varphi_3} &= -t_1^1 + t_2^2 + t_3^3 + t_4^4 + t_5^5 - t_6^6 - t_7^7 - t_8^8, \\ g_{\varphi_4} &= t_1^1 - t_2^2 + t_3^3 + t_4^4 - t_5^5 + t_6^6 - t_7^7 - t_8^8, \\ g_{\varphi_5} &= t_1^1 + t_2^2 - t_3^3 - t_4^4 + t_5^5 + t_6^6 - t_7^7 - t_8^8, \\ g_{\varphi_6} &= t_1^1 + t_2^2 - t_3^3 + t_4^4 - t_5^5 - t_6^6 + t_7^7 - t_8^8, \\ g_{\varphi_7} &= t_1^1 - t_2^2 + t_3^3 - t_4^4 + t_5^5 - t_6^6 + t_7^7 - t_8^8, \end{aligned} \tag{2.10}$$

whereas the latter correspond with generators given by

$$\begin{aligned} g_{\chi_1} &= t_{1238}, & g_{\chi_4} &= t_{2578}, \\ g_{\chi_2} &= t_{1458}, & g_{\chi_5} &= t_{4738}, & g_{\chi_7} &= t_{8246}, \\ g_{\chi_3} &= t_{1678}, & g_{\chi_6} &= t_{6358}, \end{aligned} \tag{2.11}$$

Exponentiating (2.10) and (2.11) with coefficients  $\varphi_i$  and  $\chi_i$  as

$$\mathcal{V} = \text{Exp} \left[ -12 \sum_{i=1}^7 \chi_i g_{\chi_i} \right] \text{Exp} \left[ \frac{1}{4} \sum_{i=1}^7 \varphi_i g_{\varphi_i} \right], \quad (2.12)$$

yields a parameterisation of an  $M_{MN} = \mathcal{V}\mathcal{V}^t \in [\text{SL}(2)/\text{SO}(2)]^7$  subspace of the coset space in (2.3). The kinetic terms in the resulting  $\mathcal{N} = 1$  sector follow from (2.2) and (2.12), and are given by

$$\mathcal{L}_{kin} = -\frac{1}{4} \sum_{i=1}^7 [(\partial\varphi_i)^2 + e^{2\varphi_i}(\partial\chi_i)^2]. \quad (2.13)$$

These match the standard kinetic terms  $\mathcal{L}_{kin} = -(\partial_{z_i, \bar{z}_j}^2 K) dz_i \wedge *d\bar{z}_j$  for a set of seven chiral fields  $z_i$  with Kähler potential

$$K = -\sum_{i=1}^7 \log[-i(z_i - \bar{z}_i)]. \quad (2.14)$$

Lastly, when restricted to the  $\mathbb{Z}_2^3$  invariant sector entering (2.12), the scalar potential, as computed from (2.4), can be recovered from a holomorphic superpotential

$$W = 2g[z_1 z_5 z_6 + z_2 z_4 z_6 + z_3 z_4 z_5 + (z_1 z_4 + z_2 z_5 + z_3 z_6) z_7] + 2gc(1 - z_4 z_5 z_6 z_7), \quad (2.15)$$

using the standard  $\mathcal{N} = 1$  formula

$$V_{\mathcal{N}=1} = e^K \left[ K^{z_i \bar{z}_j} D_{z_i} W D_{\bar{z}_j} \bar{W} - 3W\bar{W} \right], \quad (2.16)$$

where  $D_{z_i} W \equiv \partial_{z_i} W + (\partial_{z_i} K)W$  is the Kähler derivative and  $K^{z_i \bar{z}_j}$  is the inverse of the Kähler metric  $K_{z_i \bar{z}_j} \equiv \partial_{z_i \bar{z}_j}^2 K$ . Note that only the last term in the superpotential (2.15) turns out to be sensitive to the electromagnetic parameter  $c$ .

### 2.3 New families of AdS<sub>4</sub> vacua

A thorough study of the structure of extrema of the scalar potential (2.4), restricted to the  $\mathbb{Z}_2^3$  invariant sector, reveals a rich structure of (fairly) symmetric AdS<sub>4</sub> vacua. We find four families of vacua preserving  $\mathcal{N} = 0, 1, 2$  or 4 supersymmetry as well as various residual gauge symmetries ranging from  $U(1)^2$  to  $SO(6) \sim SU(4)$ . The three supersymmetric families are also supersymmetric within the  $\mathcal{N} = 1$  model with seven chirals presented in the previous section, and therefore satisfy the F-flatness conditions

$$D_{z_i} W = 0, \quad (2.17)$$

that follow from the superpotential (2.15) and Kähler potential (2.14). Importantly, all the AdS<sub>4</sub> vacua we will present in this section are genuinely dyonic, namely, they disappear if taking the limit  $c \rightarrow 0$  to a purely electric gauging of G in (2.1).

**2.3.1  $\mathcal{N} = 0$  vacua with  $SU(2) \rightarrow SU(2) \times U(1)^2 \rightarrow SU(3) \times U(1) \rightarrow SO(6)$  symmetry**

There is a three-parameter family of  $\mathcal{N} = 0$  solutions that preserves  $SU(2)$  and is located at

$$z_{1,2,3} = c \left( -\chi_{1,2,3} + i \frac{1}{\sqrt{2}} \right) \quad \text{and} \quad z_4 = z_5 = z_6 = z_7 = i, \quad (2.18)$$

with  $\chi_{1,2,3}$  being arbitrary (real) parameters. This family of solutions has a vacuum energy given by

$$V_0 = -2\sqrt{2}g^2c^{-1}, \quad (2.19)$$

and a spectrum of normalised scalar masses of the form

$$\begin{aligned} m^2L^2 = & 6(\times 2), \quad -3(\times 2), \quad 0(\times 28), \\ & -\frac{3}{4} + \frac{3}{2}\chi^2(\times 2), \\ & -\frac{3}{4} + \frac{3}{2}(\chi - 2\chi_i)^2(\times 2) \quad i = 1, 2, 3, \\ & -\frac{3}{4} + \frac{3}{2}\chi_i^2(\times 4) \quad i = 1, 2, 3, \\ & -3 + 6\chi_i^2(\times 2) \quad i = 1, 2, 3, \\ & -3 + \frac{3}{2}(\chi_i \pm \chi_j)^2(\times 2) \quad i < j, \end{aligned} \quad (2.20)$$

where  $\chi \equiv \chi_1 + \chi_2 + \chi_3$  and  $L^2 = -3/V_0$  is the  $AdS_4$  radius. This family of solutions is perturbatively unstable due to the mass eigenvalue  $-3$  lying below the Breitenlohner-Freedman bound for stability in  $AdS_4$  [49]. The computation of the vector masses yields

$$\begin{aligned} m^2L^2 = & 0(\times 3), \quad 6(\times 1), \\ & \frac{9}{4} + \frac{3}{2}\chi_i^2(\times 4) \quad i = 1, 2, 3, \\ & \frac{3}{2}(\chi_i \pm \chi_j)^2(\times 2) \quad i < j. \end{aligned} \quad (2.21)$$

Note that a generic solution in this family preserves an  $SU(2)$  symmetry as three vectors are generically massless. Therefore, out of the 28 massless scalars in (2.20), only 3 of them correspond to physical directions in the scalar potential. An additional  $U(1)^2$  factor appears when imposing a pairwise identification between the free axions  $\chi_{1,2,3}$ , thus resulting in a symmetry enhancement to  $SU(2) \times U(1)^2$ . A further identification  $\chi_1 = \chi_2 = \chi_3 \neq 0$  implies a symmetry enhancement to  $SU(3) \times U(1)$ . Lastly, setting  $\chi_{1,2,3} = 0$  enhances the symmetry to  $SU(4) \sim SO(6)$ . This  $SO(6)$  symmetric solution was originally studied in [29] from a ten-dimensional perspective and, more recently, connected with a family of type IIB S-fold backgrounds in [32].

**2.3.2  $\mathcal{N} = 1$  vacua with  $U(1)^2 \rightarrow SU(2) \times U(1) \rightarrow SU(3)$  symmetry**

There is a two-parameter family of  $\mathcal{N} = 1$  supersymmetric  $AdS_4$  solutions that preserves  $U(1)^2$  and is located at

$$z_{1,2,3} = c \left( -\chi_{1,2,3} + i \frac{\sqrt{5}}{3} \right) \quad \text{and} \quad z_4 = z_5 = z_6 = z_7 = \frac{1}{\sqrt{6}}(1 + i\sqrt{5}), \quad (2.22)$$

subject to the constraint

$$\chi_1 + \chi_2 + \chi_3 = 0. \quad (2.23)$$

This family of AdS<sub>4</sub> solutions has a vacuum energy given by

$$V_0 = -\frac{162}{25\sqrt{5}}g^2c^{-1}, \quad (2.24)$$

and a spectrum of normalised scalar masses of the form

$$\begin{aligned} m^2L^2 = & 0(\times 28), \quad 4 \pm \sqrt{6}(\times 2), \quad -2(\times 2), \\ & -\frac{14}{9} + 5\chi_i^2 \pm \frac{1}{3}\sqrt{4 + 45\chi_i^2}(\times 2) \quad i = 1, 2, 3, \\ & -\frac{14}{9} + \frac{5}{4}\chi_i^2 \pm \frac{1}{6}\sqrt{16 + 45\chi_i^2}(\times 2) \quad i = 1, 2, 3, \\ & \frac{7}{9} + \frac{5}{4}\chi_i^2(\times 2) \quad i = 1, 2, 3, \\ & -2 + \frac{5}{4}(\chi_i - \chi_j)^2(\times 2) \quad i < j, \end{aligned} \quad (2.25)$$

where  $L^2 = -3/V_0$  is the AdS<sub>4</sub> radius. The computation of the vector masses yields

$$\begin{aligned} m^2L^2 = & 0(\times 2), \quad 6(\times 1), \quad 2(\times 1), \\ & \frac{16}{9} + \frac{5}{4}\chi_i^2 \pm \frac{1}{6}\sqrt{64 + 45\chi_i^2}(\times 2) \quad i = 1, 2, 3, \\ & \frac{25}{9} + \frac{5\chi_i^2}{4}(\times 2) \quad i = 1, 2, 3, \\ & \frac{5}{4}(\chi_i - \chi_j)^2(\times 2) \quad i < j. \end{aligned} \quad (2.26)$$

Note that a generic solution in this family preserves  $U(1)^2$  as only two vectors are generically massless. Therefore, out of the 28 massless scalars in (2.25), only 2 of them correspond to physical directions in the potential. The residual symmetry gets enhanced to  $SU(2) \times U(1)$  when imposing a pairwise identification between the axions  $\chi_{1,2,3}$  so that a total of four vectors become massless. Finally there is a symmetry enhancement to  $SU(3)$  when setting  $\chi_{1,2,3} = 0$  so that a total of eight vectors become massless. The  $SU(3)$  symmetric solution was recently uplifted to a ten-dimensional family of type IIB S-fold backgrounds in [32].

### 2.3.3 $\mathcal{N} = 2$ vacua with $U(1)^2 \rightarrow SU(2) \times U(1)$ symmetry

There is a one-parameter family of  $\mathcal{N} = 2$  supersymmetric AdS<sub>4</sub> solutions that preserves  $U(1)^2$  and is located at

$$z_1 = -\bar{z}_3 = c \left( -\chi + i\frac{1}{\sqrt{2}} \right), \quad z_2 = ic, \quad z_4 = z_6 = i \quad \text{and} \quad z_5 = z_7 = \frac{1}{\sqrt{2}}(1 + i). \quad (2.27)$$

This family of AdS<sub>4</sub> solutions has a vacuum energy given by

$$V_0 = -3g^2c^{-1}, \quad (2.28)$$

and a spectrum of normalised scalar masses of the form

$$\begin{aligned}
 m^2 L^2 = & 0(\times 30), \quad 3 \pm \sqrt{17}(\times 2), \quad -2(\times 4), \quad 2(\times 6), \quad -2 + 4\chi^2(\times 6) \\
 & -1 + 4\chi^2 \pm \sqrt{16\chi^2 + 1}(\times 2), \quad \chi^2 \pm \sqrt{\chi^2 + 2}(\times 8),
 \end{aligned}
 \tag{2.29}$$

where  $L^2 = -3/V_0$  is the  $\text{AdS}_4$  radius. The computation of the vector masses yields

$$\begin{aligned}
 m^2 L^2 = & 0(\times 2), \quad 6(\times 2), \quad 4(\times 2), \quad 2(\times 4), \\
 & 4\chi^2(\times 2), \quad 2 + \chi^2 \pm \sqrt{\chi^2 + 2}(\times 8).
 \end{aligned}
 \tag{2.30}$$

Note that a generic solution in this family preserves  $U(1)^2$  as only two vectors are generically massless. Therefore, out of the 30 massless scalars in (2.29), only 4 of them correspond to physical directions in the scalar potential. However, the residual symmetry gets enhanced to  $SU(2) \times U(1)$  when  $\chi = 0$  and two additional vectors become massless. This special  $\text{AdS}_4$  vacuum will be uplifted to a ten-dimensional family of type IIB S-fold backgrounds in section 3.

### 2.3.4 $\mathcal{N} = 4$ vacuum with $SO(4)$ symmetry

There is an  $\mathcal{N} = 4$  supersymmetric  $\text{AdS}_4$  solution that preserves  $SO(4)$  and is located at

$$z_1 = z_2 = z_3 = ic \quad \text{and} \quad z_4 = z_5 = z_6 = -\bar{z}_7 = \frac{1}{\sqrt{2}}(1 + i).
 \tag{2.31}$$

This  $\text{AdS}_4$  solution has a vacuum energy given by

$$V_0 = -3g^2 c^{-1},
 \tag{2.32}$$

as for the previous solution, and a spectrum of normalised scalar masses of the form

$$m^2 L^2 = 0(\times 48), \quad 10(\times 1), \quad 4(\times 10), \quad -2(\times 11),
 \tag{2.33}$$

where  $L^2 = -3/V_0$  is the  $\text{AdS}_4$  radius. The computation of the vector masses yields

$$m^2 L^2 = 0(\times 6), \quad 6(\times 7), \quad 2(\times 15),
 \tag{2.34}$$

thus reflecting the  $SO(4)$  residual symmetry at the  $\text{AdS}_4$  solution. Therefore, out of the 48 massless scalars in (2.33), only 26 of them correspond to physical directions in the scalar potential. This  $\mathcal{N} = 4$  solution was first reported in [28], and then uplifted to a ten-dimensional family of type IIB S-fold backgrounds in [27].

## 3 S-folds with $\mathcal{N} = 2$ supersymmetry

From this moment on we will set

$$g = c = 1,
 \tag{3.1}$$

without loss of generality. From (2.18), (2.22), (2.27) and (2.31) it becomes clear that varying  $c$  amounts to a rescaling of the vacuum expectation values of  $z_{1,2,3} \propto c$  at the  $\text{AdS}_4$

vacua. After  $c$  has been set to unity, varying  $g$  simply corresponds to a rescaling of the vacuum energy  $V_0 \propto g^2 c^{-1}$  and thus to a redefinition of the AdS<sub>4</sub> radius  $L^2 = -3/V_0$ . Let us emphasise again that all the AdS<sub>4</sub> vacua in section 2.3 are genuinely dyonic as they do not survive the limit  $c \rightarrow 0$  to implement a purely electric gauging. In this limit one has that  $\text{Im}(z_{1,2,3}) \rightarrow 0$  or, by virtue of (2.9), a runaway behaviour towards the boundary of moduli space  $\varphi_{1,2,3} \rightarrow \infty$ .

Going back to the goal of this section, the  $\mathcal{N} = 2$  family of solutions in section 2.3.3 is new and preserves a  $U(1)^2$  symmetry. It is a one-parameter family of AdS<sub>4</sub> vacua and, in the special case of the parameter vanishing  $\chi = 0$ , there is an enhancement of symmetry to  $SU(2) \times U(1)$ . Following [27], and implementing a generalised S-S ansatz in E<sub>7(7)</sub>-EFT [50], we will uplift such an  $\mathcal{N} = 2$   $SU(2) \times U(1)$  symmetric AdS<sub>4</sub> vacuum to a class of ten-dimensional S-fold backgrounds of type IIB supergravity of the form  $\text{AdS}_4 \times S^1 \times S^5$  with an S-duality hyperbolic monodromy along  $S^1$ .

### 3.1 Type IIB uplift using E<sub>7(7)</sub>-EFT

Generalised Scherk-Schwarz (S-S) reductions of exceptional field theory (EFT) have proved a very efficient method to perform consistent truncations of eleven-dimensional and type IIB supergravity on spheres and hyperboloids [51]. Here we are interested in the uplift of an AdS<sub>4</sub> vacuum of a four-dimensional gauged maximal supergravity, which thus selects the E<sub>7(7)</sub>-EFT of [50] as the natural framework to carry out this mission.

The E<sub>7(7)</sub>-EFT lives in an extended space-time that consists of an external four-dimensional space with coordinates  $x^\mu$  ( $\mu = 0, \dots, 3$ ) and a 56-dimensional generalised internal space with coordinates  $Y^M$  ( $M = 1, \dots, 56$ ) in the fundamental representation **56** of E<sub>7(7)</sub>, subject to the action of the E<sub>7(7)</sub>-covariant generalised diffeomorphisms. In order to uplift an AdS<sub>4</sub> vacuum amongst those in section 2.3 to a ten-dimensional background of type IIB supergravity, the relevant field content of E<sub>7(7)</sub>-EFT reduces to the external metric  $g_{\mu\nu}(x, Y)$  and the internal generalised metric  $\mathcal{M}_{MN}(x, Y)$  (vector and tensor fields are consistently set to zero). These are connected with the metric  $g_{\mu\nu}(x)$  and the scalar fields  $M_{MN}(x)$  of the four-dimensional maximal supergravity in (2.2) via a generalised S-S ansatz [51]

$$\begin{aligned} g_{\mu\nu}(x, Y) &= \rho^{-2}(Y)g_{\mu\nu}(x), \\ \mathcal{M}_{MN}(x, Y) &= U_M^K(Y)U_N^L(Y)M_{KL}(x). \end{aligned} \tag{3.2}$$

The entire dependence on the  $Y^M$  coordinates is then encoded in a twist matrix  $U_M^K(Y)$  and a scaling function  $\rho(Y)$  satisfying

$$\begin{aligned} (U^{-1})_M^P(U^{-1})_N^Q \partial_P U_Q^K \Big|_{\mathbf{912}} &= \frac{1}{7} \rho X_{MN}^K, \\ \partial_N (U^{-1})_M^N - 3\rho^{-1} \partial_N \rho (U^{-1})_M^N &= 2\rho \vartheta_M, \end{aligned} \tag{3.3}$$

where  $X_{MN}^K$  is the embedding tensor specifying the gauging in the four-dimensional supergravity,  $\vartheta_M$  is a constant scaling tensor and  $\Big|_{\mathbf{912}}$  denotes projection onto the **912** irreducible representation of E<sub>7(7)</sub> where the embedding tensor lives.

For the dyonic gauging of  $G \subset \text{SL}(8)$  in (2.1) the non-vanishing components of the embedding tensor were given in (2.5) and the tensor  $\vartheta_M$  vanishes identically. The generalised S-S ansatz depends on six physical coordinates  $(y^i, \tilde{y}) \in Y^M$ : five of them are electric  $y^i$  ( $i = 2, \dots, 6$ ) and one is magnetic  $\tilde{y}$ . Considering the electric-magnetic splitting of generalised coordinates  $Y^M = (Y^{AB}, Y_{AB})$  under  $\text{SL}(8) \subset \text{E}_{7(7)}$ , one has

$$y^i = Y^{i7} \in Y^{AB} \quad \text{and} \quad \tilde{y} = Y_{18} \in Y_{AB}. \quad (3.4)$$

In terms of the physical coordinates  $(y^i, \tilde{y})$  the scaling function  $\rho$  in (3.2)–(3.3) reads

$$\rho(y^i, \tilde{y}) = \hat{\rho}(y^i) \check{\rho}(\tilde{y}), \quad (3.5)$$

where the two factors in (3.5) are given by

$$\hat{\rho}^4 = 1 - |\vec{y}|^2 \quad \text{and} \quad \check{\rho}^4 = 1 + \tilde{y}^2, \quad (3.6)$$

and  $\vec{y} \equiv (y^i)$ . On the other hand, the generalised twist matrix  $(U^{-1})_M{}^N$  in (3.2)–(3.3) is  $\text{SL}(8)$ -valued and possesses a block diagonal structure

$$(U^{-1})_M{}^N = \begin{pmatrix} (U^{-1})_{[AB]}{}^{[CD]} & 0 \\ 0 & (U^{-1})^{[AB]}{}_{[CD]} = U_{[CD]}{}^{[AB]} \end{pmatrix}, \quad (3.7)$$

with components

$$(U^{-1})_{[AB]}{}^{[CD]} = 2(U^{-1})_{[A}{}^{[C}(U^{-1})_{B]}{}^{D]}, \quad (3.8)$$

and

$$(U^{-1})_A{}^B = \left(\frac{\check{\rho}}{\hat{\rho}}\right)^{\frac{1}{2}} \begin{pmatrix} 1 & 0 & 0 & \check{\rho}^{-2}\tilde{y} \\ 0 & \delta^{ij} + \hat{K}y^i y^j & \hat{\rho}^2 y^i & 0 \\ 0 & \hat{\rho}^2 y^j \hat{K} & \hat{\rho}^4 & 0 \\ \check{\rho}^{-2}\tilde{y} & 0 & 0 & \check{\rho}^{-4}(1 + \tilde{y}^2) \end{pmatrix}. \quad (3.9)$$

The twist matrix in (3.9) also depends on a function  $\hat{K}(y^i)$  which is given in this case by a hypergeometric function [27]

$$\hat{K} = -{}_2F_1\left(1, 2, \frac{1}{2}; 1 - |\vec{y}|^2\right). \quad (3.10)$$

Using the dictionary between the fields of type IIB supergravity and those of  $\text{E}_{7(7)}$ -EFT [52, 53], together with the S-S ansatz (3.2) involving generalised twist parameters (3.5)–(3.9), one arrives at the final uplift formulae

$$\begin{aligned} G^{mn} &= G^{\frac{1}{2}} \mathcal{M}^{mn}, \\ \mathbb{B}_{mn}{}^\alpha &= G^{\frac{1}{2}} G_{mp} \epsilon^{\alpha\beta} \mathcal{M}^p{}_{n\beta}, \\ C_{klmn} - \frac{3}{2} \epsilon_{\alpha\beta} \mathbb{B}_{k[l}{}^\alpha \mathbb{B}_{mn]}{}^\beta &= -\frac{1}{2} G^{\frac{1}{2}} G_{kp} \mathcal{M}^p{}_{lmn}, \\ m_{\alpha\beta} &= \frac{1}{6} G \left( \mathcal{M}^{mn} \mathcal{M}_{m\alpha n\beta} + \mathcal{M}^m{}_{k\alpha} \mathcal{M}^k{}_{m\beta} \right), \end{aligned} \quad (3.11)$$

for the purely internal components of the type IIB fields: (inverse) metric  $G^{mn}$ , two-form potentials  $\mathbb{B}^\alpha = (B_2, C_2)$  with  $\alpha = 1, 2$ , four-form potential  $C_4$  and axion-dilaton  $m_{\alpha\beta}$ . The various blocks  $\mathcal{M}^{mn}$ ,  $\mathcal{M}^p_{n\beta}$ ,  $\mathcal{M}^p_{lmn}$  and  $\mathcal{M}_{m\alpha n\beta}$  entering the r.h.s. of (3.11) can be extracted from the internal generalised metric  $\mathcal{M}_{MN}(x, y^i, \tilde{y})$  by performing the group-theoretical decomposition that is relevant for the embedding of type IIB supergravity into  $E_{7(7)}$ -EFT:

$$\begin{aligned}
 E_{7(7)} &\supset && GL(6) \times SL(2)_{\text{IIB}} \times \mathbb{R}^+ \\
 \mathbf{56} &\rightarrow && (\mathbf{6}, \mathbf{1})_{+2} + (\mathbf{6}', \mathbf{2})_{+1} + (\mathbf{20}, \mathbf{1})_0 + (\mathbf{6}, \mathbf{2})_{-1} + (\mathbf{6}', \mathbf{1})_{-2} \\
 Y^M &\rightarrow && y^m + y_{m\alpha} + y^{mnp} + y^{m\alpha} + y_m
 \end{aligned}
 \tag{3.12}$$

The physical coordinates are identified as  $y^m = (y^i, \tilde{y})$ , with  $m = (i, 7)$  and  $i = 2, \dots, 6$ , which implies a further group-theoretical branching  $GL(6) \rightarrow GL(1) \times GL(5)$  compatible with the  $\mathbb{R}(\text{or } S^1) \times S^5$  factorisation of the geometry we are behind of. The various mappings between coordinates discussed above are summarised as

$$\begin{array}{c|c|c|c|c}
 y^m & & y_{m\alpha} & & y^{mnp} & & y^{m\alpha} & & y_m \\
 \hline
 y^i & y^7 & y_{i\alpha} & y_{7\alpha} & y^{ijk} & y^{ij7} & y^{i\alpha} & y^{7\alpha} & y_i & y_7 \\
 Y^{i7} & Y_{18} & Y_{i\alpha} & \epsilon_{\alpha\beta} Y^{\beta 7} & \epsilon^{ijkj'k'} Y_{j'k'} & Y^{ij} & Y^{i\alpha} & \epsilon^{\alpha\beta} Y_{\beta 7} & Y_{i7} & Y^{18}
 \end{array}
 \tag{3.13}$$

We refer the reader to the original works [52, 53] (and also [27, 32]) for more details on the generalised S-S reductions of  $E_{7(7)}$ -EFT and their connection with the gauged maximal supergravities.

We now move to the uplift of the  $AdS_4$  vacuum with  $\mathcal{N} = 2 \& SU(2) \times U(1)$  symmetry discussed in section 2.3.3 to a ten-dimensional background of type IIB supergravity using (3.11). We have explicitly verified that the ten-dimensional equations of motion and Bianchi identities of type IIB supergravity are satisfied.<sup>2</sup>

**Ten-dimensional metric.** We adopt the conventions of [32] to describe the geometry of the round five-sphere  $S^5$ . Using coordinates  $y^i$  ( $i = 2, \dots, 6$ ) to parameterise  $S^5$ , the metric and its inverse are given by

$$\hat{G}_{ij} = \delta_{ij} + \frac{\delta_{ik}\delta_{jl}y^k y^l}{1 - y^m \delta_{mn} y^n} \quad \text{and} \quad \hat{G}^{ij} = \delta^{ij} - y^i y^j.
 \tag{3.14}$$

However it will also be convenient to introduce a set of embedding coordinates  $\mathcal{Y}_{\underline{m}}$  on  $\mathbb{R}^6$  ( $\underline{m} = 2, \dots, 7$ ) of the form

$$\mathcal{Y}_{\underline{m}} = \left\{ y^i, \mathcal{Y}_7 \equiv (1 - |\tilde{y}|^2)^{\frac{1}{2}} \right\} \quad \text{with} \quad \delta^{\underline{m}\underline{n}} \mathcal{Y}_{\underline{m}} \mathcal{Y}_{\underline{n}} = 1,
 \tag{3.15}$$

so that the Killing vectors on  $S^5$  are constructed as

$$\mathcal{K}_{\underline{mn}}{}^i \equiv \hat{G}^{ij} \partial_j \mathcal{Y}_{[\underline{m}} \mathcal{Y}_{\underline{n}]} = \delta^i_{[\underline{m}} \mathcal{Y}_{\underline{n}]} .
 \tag{3.16}$$

---

<sup>2</sup>We adopt the type IIB conventions in the appendix B of [32].

Following the derivation of [27], the internal part of the ten-dimensional metric has components in (3.11) given by

$$\begin{aligned}
 G^{11} &= \Delta \hat{\rho}^4 M_{1818} = 2\Delta(1 + \tilde{y}^2), \\
 G^{1k} &= \Delta \hat{\rho}^2 \mathcal{K}_{ij}{}^k M_{18}^{ij} = 0, \\
 G^{ij} &= \Delta \mathcal{K}_{kl}{}^i \mathcal{K}_{mn}{}^j M^{klmn} = \Delta (\hat{G}^{ij} + L^{ij}),
 \end{aligned} \tag{3.17}$$

where  $M_{18}^{ij} = 0$  as a consequence of having set  $\chi = 0$  in the  $\mathcal{N} = 2$  AdS<sub>4</sub> vacuum, and where we have defined

$$L^{ij} = \begin{pmatrix}
 \mathcal{Y}_4^2 + \mathcal{Y}_5^2 + \mathcal{Y}_6^2 & -\mathcal{Y}_6\mathcal{Y}_7 & -\mathcal{Y}_2\mathcal{Y}_4 - \mathcal{Y}_2\mathcal{Y}_5 & -\mathcal{Y}_2\mathcal{Y}_6 \\
 -\mathcal{Y}_6\mathcal{Y}_7 & \mathcal{Y}_4^2 + \mathcal{Y}_5^2 + \mathcal{Y}_7^2 & -\mathcal{Y}_3\mathcal{Y}_4 - \mathcal{Y}_3\mathcal{Y}_5 & \mathcal{Y}_2\mathcal{Y}_7 \\
 -\mathcal{Y}_2\mathcal{Y}_4 & -\mathcal{Y}_3\mathcal{Y}_4 & 1 - \mathcal{Y}_4^2 - \mathcal{Y}_4\mathcal{Y}_5 & -\mathcal{Y}_4\mathcal{Y}_6 \\
 -\mathcal{Y}_2\mathcal{Y}_5 & -\mathcal{Y}_3\mathcal{Y}_5 & -\mathcal{Y}_4\mathcal{Y}_5 & 1 - \mathcal{Y}_5^2 - \mathcal{Y}_5\mathcal{Y}_6 \\
 -\mathcal{Y}_2\mathcal{Y}_6 & \mathcal{Y}_2\mathcal{Y}_7 & -\mathcal{Y}_4\mathcal{Y}_6 - \mathcal{Y}_5\mathcal{Y}_6 & \mathcal{Y}_4^2 + \mathcal{Y}_5^2 + \mathcal{Y}_7^2
 \end{pmatrix}. \tag{3.18}$$

The warping factor  $\Delta$  in (3.17) is nowhere vanishing and reads

$$\Delta = (\det G)^{\frac{1}{2}} \rho^2 = \frac{1}{\sqrt{2}} (1 + \mathcal{Y}_4^2 + \mathcal{Y}_5^2)^{-\frac{1}{4}}. \tag{3.19}$$

The six-dimensional internal metric becomes more transparent if first introducing a new variable for the magnetic coordinate

$$\tilde{y} = \sinh \eta \quad \text{with} \quad \eta \in (-\infty, \infty), \tag{3.20}$$

and then a set of angular variables for S<sup>5</sup> of the form

$$\begin{aligned}
 y^2 &= \cos \theta \cos \left( \frac{\beta}{2} \right) \cos \left( \frac{\alpha + \gamma}{2} \right), & y^3 &= \cos \theta \cos \left( \frac{\beta}{2} \right) \sin \left( \frac{\alpha + \gamma}{2} \right), \\
 y^4 &= \cos \phi \sin \theta, & y^5 &= \sin \phi \sin \theta, \\
 y^6 &= \cos \theta \sin \left( \frac{\beta}{2} \right) \cos \left( \frac{\alpha - \gamma}{2} \right),
 \end{aligned} \tag{3.21}$$

with ranges given by

$$\theta \in \left[ 0, \frac{\pi}{2} \right], \quad \phi \in [0, 2\pi], \quad \alpha \in [0, 2\pi], \quad \beta \in [0, \pi], \quad \gamma \in [0, 2\pi]. \tag{3.22}$$

In this manner, and upon introducing a set of SU(2) left-invariant one-forms

$$\begin{aligned}
 \sigma_1 &= \frac{1}{2} (-\sin \alpha d\beta + \cos \alpha \sin \beta d\gamma), \\
 \sigma_2 &= \frac{1}{2} (\cos \alpha d\beta + \sin \alpha \sin \beta d\gamma), \\
 \sigma_3 &= \frac{1}{2} (d\alpha + \cos \beta d\gamma),
 \end{aligned} \tag{3.23}$$

the internal six-dimensional metric takes a simple  $\mathbb{R} \times S^5$  form

$$ds_6^2 = \frac{1}{2} \Delta^{-1} [d\eta^2 + ds_{S^2}^2 + \cos^2 \theta ds_{S^3}^2], \quad (3.24)$$

with a warping factor

$$\Delta^{-1} = (6 - 2 \cos(2\theta))^{\frac{1}{4}}, \quad (3.25)$$

and where we have introduced  $S^2$  and (squashed)  $S^3$  metrics to describe the deformation of the internal  $S^5$ . These metrics are explicitly given by

$$ds_{S^2}^2 = d\theta^2 + \sin^2 \theta d\phi^2 \quad \text{and} \quad ds_{S^3}^2 = \sigma_2^2 + 8\Delta^4 (\sigma_1^2 + \sigma_3^2). \quad (3.26)$$

Bringing together (3.24) and the external  $\text{AdS}_4$  part of the geometry, one obtains a ten-dimensional metric of the form<sup>3</sup>

$$ds^2 = \frac{1}{2} \Delta^{-1} [ds_{\text{AdS}_4}^2 + d\eta^2 + ds_{S^2}^2 + \cos^2 \theta ds_{S^3}^2]. \quad (3.27)$$

This metric has an  $\text{SU}(2) \times \text{U}(1)_\phi \times \text{U}(1)_\sigma$  symmetry, where  $\text{U}(1)_\sigma$  acts as a rotation on the  $(\sigma_1, \sigma_3)$ -plane. Finally, our choice of *undeformed* frames for the metric (3.27) is

$$\begin{aligned} ds_{\text{AdS}_4}^2 : \quad & \hat{e}^0 = \frac{L}{r} dr, \quad \hat{e}^i = \frac{L}{r} dx^i \quad (i = 1, 2, 3) \quad \text{and} \quad \eta_{ij} = (-1, 1, 1) \\ ds_{\mathbb{R}}^2 : \quad & \hat{e}^4 = d\eta \\ ds_{S^2}^2 : \quad & \hat{e}^5 = d\theta, \quad \hat{e}^6 = \sin \theta d\phi \\ ds_{S^3}^2 : \quad & \hat{e}^7 = \sigma_1, \quad \hat{e}^8 = \sigma_2, \quad \hat{e}^9 = \sigma_3 \end{aligned} \quad (3.28)$$

with  $L^2 = -3/V_0 = 1$  being the  $\text{AdS}_4$  radius at the four-dimensional  $\mathcal{N} = 2$   $\text{SU}(2) \times \text{U}(1)$  symmetric  $\text{AdS}_4$  vacuum.

**$B_2$  and  $C_2$  potentials.** The two-form potentials  $\mathbb{B}^\alpha = (B_2, C_2)$  in (3.11) transform as a doublet under the global S-duality group  $\text{SL}(2, \mathbb{R})_{\text{IIB}}$  of type IIB supergravity. An explicit computation along the lines of [27] shows that

$$\begin{aligned} \mathbb{B}_{1j}^\alpha &= 0, \\ \mathbb{B}_{ij}^\alpha &= \Delta G_{ik} \mathcal{K}_{kl}^k \partial_j \mathcal{Y}^m \epsilon^{\alpha\beta} (A^{-1})^\gamma_\beta M^{kl}_{m\gamma}, \end{aligned} \quad (3.29)$$

in terms of a local  $\text{SO}(1, 1) \subset \text{SL}(2, \mathbb{R})_{\text{IIB}}$  twist matrix

$$A^\alpha_\beta \equiv \begin{pmatrix} \cosh \eta & \sinh \eta \\ \sinh \eta & \cosh \eta \end{pmatrix}, \quad (A^{-1})^\gamma_\beta \equiv \begin{pmatrix} \cosh \eta & -\sinh \eta \\ -\sinh \eta & \cosh \eta \end{pmatrix}. \quad (3.30)$$

---

<sup>3</sup>Restoring the explicit dependence of the warping factor (3.25) on the parameter  $c$  one finds  $\Delta \propto c$ , so the (electric) limit  $c \rightarrow 0$  of the metric (3.27) becomes pathological. In other words, the ten-dimensional solution is genuinely dyonic, namely, it requires  $c \neq 0$ , as for its associated  $\text{AdS}_4$  vacuum in (2.27) with  $\chi = 0$ .

This matrix encodes the dependence of the two-form potentials on the direction  $\eta$ . Using the scalar block  $M^{kl}_{m\gamma}$  at the  $\mathcal{N} = 2$  AdS<sub>4</sub> vacuum under consideration, and using differential form notation, one finds

$$\mathbb{B}^\alpha = A^\alpha_\beta \mathfrak{b}^\beta, \quad (3.31)$$

with

$$\begin{aligned} \mathfrak{b}^1 &= \frac{1}{\sqrt{2}} \cos \theta \left[ \left( \cos \phi d\theta + \frac{1}{2} \sin(2\theta) d(\cos \phi) \right) \wedge \sigma_2 + \cos \phi \frac{4 \sin(2\theta)}{6 - 2 \cos(2\theta)} \sigma_1 \wedge \sigma_3 \right], \\ \mathfrak{b}^2 &= -\frac{1}{\sqrt{2}} \cos \theta \left[ \left( \sin \phi d\theta + \frac{1}{2} \sin(2\theta) d(\sin \phi) \right) \wedge \sigma_2 + \sin \phi \frac{4 \sin(2\theta)}{6 - 2 \cos(2\theta)} \sigma_1 \wedge \sigma_3 \right]. \end{aligned} \quad (3.32)$$

The two-form potentials in (3.32) preserve  $SU(2) \times U(1)_\sigma$  but break the  $U(1)_\phi$  factor due to the explicit dependence on the coordinate  $\phi$ .

**C<sub>4</sub> potential.** The internal component of the four-form potential  $C_4$  can be explicitly obtained from the third uplift formula in (3.11). Computing the associated (purely internal) five-form field strength, and imposing ten-dimensional self-duality, one gets

$$\begin{aligned} \tilde{F}_5 &= dC_4 - \frac{1}{2} \epsilon_{\alpha\beta} \mathbb{B}^\alpha \wedge \mathbb{H}^\beta \\ &= (1 + \star) \left[ 6\sqrt{2} \Delta^{5/2} \text{vol}_{M_5} \right. \\ &\quad \left. - 4\Delta^4 \sin \theta \cos^3 \theta d\eta \wedge \left( \cos(2\phi) d\theta - \frac{1}{2} \sin(2\theta) \sin(2\phi) d\phi \right) \wedge \sigma_1 \wedge \sigma_2 \wedge \sigma_3 \right], \end{aligned} \quad (3.33)$$

where

$$\text{vol}_{M_5} = \sqrt{2} \Delta^{3/2} \sin \theta \cos^3 \theta d\theta \wedge d\phi \wedge \sigma_1 \wedge \sigma_2 \wedge \sigma_3, \quad (3.34)$$

denotes the volume of the deformed five-sphere. Note that  $U(1)_\phi$  is also broken by  $\tilde{F}_5$  due to its explicit dependence on the coordinate  $\phi$ .

**Axion-dilaton.** The axion-dilaton matrix  $m_{\alpha\beta}$  can be obtained from the last equation in (3.11). Transforming linearly under S-duality, a direct computation shows an explicit dependence of  $m_{\alpha\beta}$  on the  $A$ -twist in (3.30) of the form

$$m_{\alpha\beta} = \frac{1}{\text{Im}\tau} \begin{pmatrix} |\tau|^2 & -\text{Re}\tau \\ -\text{Re}\tau & 1 \end{pmatrix} = (A^{-t})_\alpha^\gamma \mathfrak{m}_{\gamma\delta} (A^{-1})^\delta_\beta, \quad (3.35)$$

with  $\tau = C_0 + ie^{-\Phi}$  and

$$\mathfrak{m}_{\gamma\delta} = 2\Delta^2 \begin{pmatrix} 1 + \sin^2 \theta \cos^2 \phi & -\frac{1}{2} \sin^2 \theta \sin(2\phi) \\ -\frac{1}{2} \sin^2 \theta \sin(2\phi) & 1 + \sin^2 \theta \sin^2 \phi \end{pmatrix}. \quad (3.36)$$

Again  $U(1)_\phi$  is broken by the explicit dependence of (3.36) on the angle  $\phi$ . This concludes the uplift of the  $AdS_4$  vacuum with  $\mathcal{N} = 2$  and  $SU(2) \times U(1)$  symmetry discussed in section 2.3.3 to a ten-dimensional background of type IIB supergravity. It is worth emphasising that, if trivialising the  $A$ -twist in (3.30), i.e.  $A^\alpha_\beta = \delta^\alpha_\beta$ , then the ten-dimensional equations of motion of type IIB supergravity are no longer satisfied.

### 3.2 S-fold interpretation

The dependence of the full type IIB solution on the coordinate  $\eta$  along the  $\mathbb{R}$  direction of the geometry (3.27) is totally encoded in the local  $SL(2, \mathbb{R})_{IIB}$   $A$ -twist in (3.30). This twist matrix is of hyperbolic type and thus induces a non-trivial monodromy

$$\mathfrak{M}_{S^1} = A^{-1}(\eta)A(\eta + T) = \begin{pmatrix} \cosh T & \sinh T \\ \sinh T & \cosh T \end{pmatrix}, \quad (3.37)$$

when forcing the  $\eta$  coordinate to be periodic  $\eta \rightarrow \eta + T$  with period  $T$ , namely, when replacing  $\mathbb{R} \rightarrow S^1$  in the geometry. Generalising the  $A$ -twist in (3.30) to a discrete  $k$ -family ( $k \in \mathbb{N}$  with  $k \geq 3$ ) of new ones

$$A_{(k)} = Ag(k) \quad \text{with} \quad g(k) = \begin{pmatrix} \frac{(k^2 - 4)^{\frac{1}{4}}}{\sqrt{2}} & 0 \\ \frac{k}{\sqrt{2}(k^2 - 4)^{\frac{1}{4}}} & \frac{\sqrt{2}}{(k^2 - 4)^{\frac{1}{4}}} \end{pmatrix}, \quad (3.38)$$

the monodromy (3.37) gets generalised to a  $k$ -family of  $SL(2, \mathbb{Z})_{IIB}$  hyperbolic monodromies

$$\mathfrak{M}(k) = A_{(k)}^{-1}(\eta)A_{(k)}(\eta + T(k)) = \begin{pmatrix} k & 1 \\ -1 & 0 \end{pmatrix}, \quad k \geq 3, \quad (3.39)$$

with  $T(k) = \log(k + \sqrt{k^2 - 4}) - \log(2)$  and  $\text{Tr}\mathfrak{M}(k) > 2$ . Therefore, as discussed in [27] (see also [32]), these backgrounds can be interpreted as locally geometric compactifications on  $S^1 \times S^5$  involving a  $k$ -family of S-duality monodromies (3.39). These monodromies can be written as

$$\mathfrak{M}(k) = -S\mathcal{T}^k \quad \text{with} \quad S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \mathcal{T} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad (3.40)$$

and thus define a  $k$ -family of S-fold backgrounds. Moreover, the argument wielded in [27] for the straightforward uplift of the four-dimensional supersymmetries to ten dimensions relied on the monodromy (3.37) being in the hyperbolic conjugacy class of  $SL(2, \mathbb{R})_{IIB}$ . This is still our case, so the S-folds presented here preserve  $\mathcal{N} = 2$  supersymmetry.

Lastly, various holographic aspects of both  $\mathcal{N} = 4$  [27] and  $\mathcal{N} = 1$  [32, 36] S-folds with hyperbolic monodromies have respectively been investigated in [33–35] and [36] within the

context of three-dimensional quiver theories involving  $\mathcal{N} = 4$   $T(U(N))$  theories [38], and their potential generalisation to  $\mathcal{N} = 1$  SCFT's. It would be interesting to extend these holographic studies to the  $\mathcal{N} = 2$  S-folds with hyperbolic monodromies (3.39) presented in this work.

### 3.3 Connection with Janus-like solutions

The type IIB solution with  $\mathcal{N} = 2$   $SU(2) \times U(1)$  symmetry we just obtained can be mapped to a new (but equivalent) solution with a linear dilaton profile along the coordinate  $\eta$  upon performing a global  $\Lambda \in SL(2, \mathbb{R})_{\text{IIB}}$  transformation, equivalently a change of duality frame, based on the matrix element

$$\Lambda = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}. \tag{3.41}$$

The composed action of  $\Lambda A^{-1}(\eta)$  on (3.36) yields a shift of the form  $\Phi \rightarrow \Phi - 2\eta$ . Therefore, a degenerate Janus-like behaviour with a linear dilaton  $\Phi$  running from  $-\infty$  to  $\infty$  becomes manifest

$$g_s = e^\Phi \propto e^{-2\eta}, \tag{3.42}$$

giving rise to a varying string coupling  $g_s$  that interpolates between the singular values 0 and  $\infty$ .

Upon performing the  $\Lambda \in SL(2, \mathbb{R})_{\text{IIB}}$  transformation (3.41) on the original solution found in section 3.1, a new type IIB background is generated. The metric and self-dual five-form flux are  $SL(2, \mathbb{R})_{\text{IIB}}$  singlets and are not affected by the transformation. Therefore, they take the same form as in (3.27) and (3.33), namely,

$$ds^2 = \frac{1}{2} \Delta^{-1} [ds_{\text{AdS}_4}^2 + d\eta^2 + d\theta^2 + \sin^2 \theta d\phi^2 + \cos^2 \theta (\sigma_2^2 + 8\Delta^4 (\sigma_1^2 + \sigma_3^2))],$$

$$\tilde{F}_5 = 4\Delta^4 \sin \theta \cos^3 \theta (1 + \star) \left[ 3d\theta \wedge d\phi \wedge \sigma_1 \wedge \sigma_2 \wedge \sigma_3 - d\eta \wedge \left( \cos(2\phi) d\theta - \frac{1}{2} \sin(2\theta) \sin(2\phi) d\phi \right) \wedge \sigma_1 \wedge \sigma_2 \wedge \sigma_3 \right]. \tag{3.43}$$

The axion-dilaton matrix  $m_{\alpha\beta}$  in (3.35) transforms linearly under  $SL(2, \mathbb{R})_{\text{IIB}}$ . Reading off the new components of  $\tau$  one finds

$$\Phi = -2\eta + \log \left[ \frac{1}{2} \Delta^2 (5 - \cos(2\theta) - 2 \sin^2 \theta \sin(2\phi)) \right],$$

$$C_0 = -2e^{2\eta} \frac{\cos(2\phi) \sin^2 \theta}{5 - \cos(2\theta) - 2 \sin^2 \theta \sin(2\phi)}. \tag{3.44}$$

The two-form potentials  $\mathbb{B}^\alpha = (B_2, C_2)$  in (3.31)–(3.32) transform as an  $\text{SL}(2, \mathbb{R})_{\text{IIB}}$  doublet and take the new form<sup>4</sup>

$$\begin{aligned}
 B_2 &= e^{-\eta} \left[ \frac{1}{2} \cos \theta ((\cos \phi + \sin \phi) d\theta + \frac{1}{2} \sin(2\theta) (\cos \phi - \sin \phi) d\phi) \wedge \sigma_2 \right. \\
 &\quad \left. + 2\Delta^4 \cos \theta \sin(2\theta) (\cos \phi + \sin \phi) \sigma_1 \wedge \sigma_3 \right], \\
 C_2 &= e^\eta \left[ \frac{1}{2} \cos \theta ((\cos \phi - \sin \phi) d\theta - \frac{1}{2} \sin(2\theta) (\cos \phi + \sin \phi) d\phi) \wedge \sigma_2 \right. \\
 &\quad \left. + 2\Delta^4 \cos \theta \sin(2\theta) (\cos \phi - \sin \phi) \sigma_1 \wedge \sigma_3 \right].
 \end{aligned}
 \tag{3.45}$$

The nowhere vanishing warping factor still reads

$$\Delta^{-4} = 6 - 2 \cos(2\theta).
 \tag{3.46}$$

In the asymptotic region at  $\eta \rightarrow -\infty$  one has that  $g_s$  in (3.42) diverges (strong coupling) and  $B_2$  dominates over other gauge potentials, e.g.,  $C_0 \rightarrow 0$  and  $C_2 \rightarrow 0$ . On the contrary, in the asymptotic region at  $\eta \rightarrow \infty$ , the solution becomes dominated by  $C_0$  and  $C_2$  whereas  $g_s \rightarrow 0$  (weak coupling) and  $B_2 \rightarrow 0$ . At intermediate values of the coordinate  $\eta$  one has an interpolating behaviour between these two regimes. Finally, it is also worth noticing that, unlike for the  $\mathcal{N} = 4$  [27] and  $\mathcal{N} = 1$  [32] S-folds, there is no  $\text{SL}(2, \mathbb{R})_{\text{IIB}}$  frame in which the axion  $C_0$  (and thus the dual  $\theta$ -angle) vanishes identically or becomes independent of the coordinate  $\eta$ .

## 4 Conclusions

In this work we have extended the study of  $\text{AdS}_4$  vacua in [28, 31, 32] for the dyonically-gauged  $[\text{SO}(1, 1) \times \text{SO}(6)] \ltimes \mathbb{R}^{12}$  maximal supergravity and found multi-parametric families of new  $\text{AdS}_4$  vacua. Within one such families, all the solutions preserve the same amount of supersymmetry but, importantly, residual symmetry enhancements occur at particular values of the parameters. The previously known  $\mathcal{N} = 0$  &  $\text{SO}(6)$  [31],  $\mathcal{N} = 1$  &  $\text{SU}(3)$  [32] and  $\mathcal{N} = 4$  &  $\text{SO}(4)$  [28]  $\text{AdS}_4$  vacua are shown to correspond to the points of largest symmetry enhancement within their respective families. This is in line with the analysis of (global) symmetry breaking patterns of three-dimensional interface SYM theories presented in [39].

In the second part of the paper we focused on the new family of  $\mathcal{N} = 2$  supersymmetric  $\text{AdS}_4$  vacua and, more concretely, on the vacuum within this family featuring the largest possible residual symmetry, which turns to be  $\text{SU}(2) \times \text{U}(1)$ . By implementing a generalised

---

<sup>4</sup>The two terms in  $B_2$  and  $C_2$  which are proportional to  $\sigma_1 \wedge \sigma_3$  can be eliminated by means of a gauge transformation of the form

$$\begin{aligned}
 B_2 &\rightarrow B_2 - d(2\sqrt{2}\Delta^4 e^{-\eta} \sin(2\theta) \cos \theta \cos \psi \sigma_2), \\
 C_2 &\rightarrow C_2 + d(2\sqrt{2}\Delta^4 e^\eta \sin(2\theta) \cos \theta \sin \psi \sigma_2),
 \end{aligned}$$

where we have shifted the coordinate  $\phi \rightarrow \psi + \frac{\pi}{4}$ . However, since these terms are generated by the generalised S-S ansatz discussed in section 3.1, we will retain them here.

S-S ansatz in  $E_{7(7)}$ -EFT, we uplifted the  $\text{AdS}_4$  vacuum to a new family of  $\text{AdS}_4 \times \text{S}^1 \times \text{S}^5$  S-folds of type IIB supergravity with hyperbolic monodromies  $\mathfrak{M}(k) = -\mathcal{ST}^k$  (with  $k \geq 3$ ) along  $\text{S}^1$ . The residual  $\text{SU}(2) \times \text{U}(1)$  symmetry and  $\mathcal{N} = 2$  supersymmetry of the  $\text{AdS}_4$  vacuum are realised on the S-folds: the internal  $\text{S}^5$  is deformed into a product of  $\text{S}^2$  and (squashed)  $\text{S}^3$  with  $\text{SU}(2) \times \text{U}(1)_\sigma \times \text{U}(1)_\phi$  isometries and a warping factor, whereas the background fluxes break the  $\text{U}(1)_\phi$  factor explicitly by introducing a dependence on the coordinate  $\phi$ . In many aspects, the realisation of symmetries is much alike the  $\text{AdS}_5 \times \text{S}^5$  background by Pilch and Warner [54] that uplifts the  $\mathcal{N} = 2$  and  $\text{SU}(2) \times \text{U}(1)$  symmetric  $\text{AdS}_5$  vacuum of the five-dimensional  $\text{SO}(6)$  maximal supergravity presented in [55].

Finally it would be interesting to investigate the brane setups underlying the families of S-folds presented here (and in [32]), especially due to the non-trivial  $\text{SL}(2, \mathbb{Z})_{\text{IIB}}$  hyperbolic monodromies  $\mathfrak{M}(k) = -\mathcal{ST}^k$ . It would also be interesting to investigate holographic aspects of such  $\mathcal{N} = 2$  and  $\mathcal{N} = 1$  S-folds (in the spirit of the  $J$ -fold CFT's of [33–36] with  $J = -\mathcal{ST}^k$ ), as well as to study holographic RG flows by explicitly constructing domain-wall solutions interpolating between the various families of  $\text{AdS}_4$  vacua presented in this work. Lastly, since the S-folds here and in [32] display  $\text{SU}(2)$  isometries in the internal geometry, it would also be interesting to apply non-abelian T-duality in order to generate new analytic type IIA backgrounds. We plan to address these and related issues in the future.

## Acknowledgments

We are grateful to Nikolay Bobev, Gianluca Inverso, Yolanda Lozano and Henning Samtleben for interesting conversations. The work of AG is supported by the Spanish government grant PGC2018-096894-B-100 and by the Principado de Asturias through the grant FC-GRUPIN-IDI/2018/000174. The research of CS is supported by IISN-Belgium (convention 4.4503.15). CS is a Research Fellow of the F.R.S.-FNRS (Belgium).

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] G. Dall'Agata, G. Inverso and M. Trigiante, *Evidence for a family of  $\text{SO}(8)$  gauged supergravity theories*, *Phys. Rev. Lett.* **109** (2012) 201301 [[arXiv:1209.0760](https://arxiv.org/abs/1209.0760)] [[INSPIRE](#)].
- [2] B. de Wit and H. Nicolai,  *$N = 8$  Supergravity*, *Nucl. Phys. B* **208** (1982) 323 [[INSPIRE](#)].
- [3] N.P. Warner, *Some New Extrema of the Scalar Potential of Gauged  $N = 8$  Supergravity*, *Phys. Lett.* **128B** (1983) 169 [[INSPIRE](#)].
- [4] I.M. Comsa, M. Firsching and T. Fischbacher,  *$\text{SO}(8)$  Supergravity and the Magic of Machine Learning*, *JHEP* **08** (2019) 057 [[arXiv:1906.00207](https://arxiv.org/abs/1906.00207)] [[INSPIRE](#)].
- [5] A. Borghese, A. Guarino and D. Roest, *All  $G_2$  invariant critical points of maximal supergravity*, *JHEP* **12** (2012) 108 [[arXiv:1209.3003](https://arxiv.org/abs/1209.3003)] [[INSPIRE](#)].

- [6] A. Borghese, G. Dibitetto, A. Guarino, D. Roest and O. Varela, *The SU(3)-invariant sector of new maximal supergravity*, *JHEP* **03** (2013) 082 [[arXiv:1211.5335](#)] [[INSPIRE](#)].
- [7] A. Borghese, A. Guarino and D. Roest, *Triality, Periodicity and Stability of SO(8) Gauged Supergravity*, *JHEP* **05** (2013) 107 [[arXiv:1302.6057](#)] [[INSPIRE](#)].
- [8] A. Guarino, *On new maximal supergravity and its BPS domain-walls*, *JHEP* **02** (2014) 026 [[arXiv:1311.0785](#)] [[INSPIRE](#)].
- [9] J. Tarrío and O. Varela, *Electric/magnetic duality and RG flows in AdS<sub>4</sub>/CFT<sub>3</sub>*, *JHEP* **01** (2014) 071 [[arXiv:1311.2933](#)] [[INSPIRE](#)].
- [10] A. Anabalón and D. Astefanesei, *Black holes in  $\omega$ -deformed gauged N = 8 supergravity*, *Phys. Lett. B* **732** (2014) 137 [[arXiv:1311.7459](#)] [[INSPIRE](#)].
- [11] H. Lü, Y. Pang and C.N. Pope, *An  $\omega$  deformation of gauged STU supergravity*, *JHEP* **04** (2014) 175 [[arXiv:1402.1994](#)] [[INSPIRE](#)].
- [12] S.-Q. Wu and S. Li, *Thermodynamics of Static Dyonically AdS Black Holes in the  $\omega$ -Deformed Kaluza-Klein Gauged Supergravity Theory*, *Phys. Lett. B* **746** (2015) 276 [[arXiv:1505.00117](#)] [[INSPIRE](#)].
- [13] B. de Wit and H. Nicolai, *Deformations of gauged SO(8) supergravity and supergravity in eleven dimensions*, *JHEP* **05** (2013) 077 [[arXiv:1302.6219](#)] [[INSPIRE](#)].
- [14] K. Lee, C. Strickland-Constable and D. Waldram, *New Gaugings and Non-Geometry*, *Fortsch. Phys.* **65** (2017) 1700049 [[arXiv:1506.03457](#)] [[INSPIRE](#)].
- [15] A. Guarino and O. Varela, *Consistent  $\mathcal{N} = 8$  truncation of massive IIA on S<sup>6</sup>*, *JHEP* **12** (2015) 020 [[arXiv:1509.02526](#)] [[INSPIRE](#)].
- [16] G. Dall'Agata, G. Inverso and A. Marrani, *Symplectic Deformations of Gauged Maximal Supergravity*, *JHEP* **07** (2014) 133 [[arXiv:1405.2437](#)] [[INSPIRE](#)].
- [17] A. Guarino and O. Varela, *Dyonically  $\mathit{ISO}(7)$  supergravity and the duality hierarchy*, *JHEP* **02** (2016) 079 [[arXiv:1508.04432](#)] [[INSPIRE](#)].
- [18] A. Guarino, J. Tarrío and O. Varela, *Halving  $\mathit{ISO}(7)$  supergravity*, *JHEP* **11** (2019) 143 [[arXiv:1907.11681](#)] [[INSPIRE](#)].
- [19] A. Guarino, J. Tarrío and O. Varela, *Flowing to  $\mathcal{N} = 3$  Chern-Simons-matter theory*, *JHEP* **03** (2020) 100 [[arXiv:1910.06866](#)] [[INSPIRE](#)].
- [20] A. Guarino, J. Tarrío and O. Varela, *Romans-mass-driven flows on the D2-brane*, *JHEP* **08** (2016) 168 [[arXiv:1605.09254](#)] [[INSPIRE](#)].
- [21] A. Guarino and J. Tarrío, *BPS black holes from massive IIA on S<sup>6</sup>*, *JHEP* **09** (2017) 141 [[arXiv:1703.10833](#)] [[INSPIRE](#)].
- [22] A. Guarino, *BPS black hole horizons from massive IIA*, *JHEP* **08** (2017) 100 [[arXiv:1706.01823](#)] [[INSPIRE](#)].
- [23] S.M. Hosseini, K. Hristov and A. Passias, *Holographic microstate counting for AdS<sub>4</sub> black holes in massive IIA supergravity*, *JHEP* **10** (2017) 190 [[arXiv:1707.06884](#)] [[INSPIRE](#)].
- [24] F. Benini, H. Khachatryan and P. Milan, *Black hole entropy in massive Type IIA*, *Class. Quant. Grav.* **35** (2018) 035004 [[arXiv:1707.06886](#)] [[INSPIRE](#)].
- [25] L.J. Romans, *Massive N=2a Supergravity in Ten-Dimensions*, *Phys. Lett.* **169B** (1986) 374 [[INSPIRE](#)].

- [26] A. Guarino, D.L. Jafferis and O. Varela, *String Theory Origin of Dyonically  $N = 8$  Supergravity and Its Chern-Simons Duals*, *Phys. Rev. Lett.* **115** (2015) 091601 [[arXiv:1504.08009](#)] [[INSPIRE](#)].
- [27] G. Inverso, H. Samtleben and M. Trigiante, *Type II supergravity origin of dyonic gaugings*, *Phys. Rev. D* **95** (2017) 066020 [[arXiv:1612.05123](#)] [[INSPIRE](#)].
- [28] A. Gallerati, H. Samtleben and M. Trigiante, *The  $\mathcal{N} > 2$  supersymmetric AdS vacua in maximal supergravity*, *JHEP* **12** (2014) 174 [[arXiv:1410.0711](#)] [[INSPIRE](#)].
- [29] D. Bak, M. Gutperle and S. Hirano, *A dilatonic deformation of AdS<sub>5</sub> and its field theory dual*, *JHEP* **05** (2003) 072 [[hep-th/0304129](#)] [[INSPIRE](#)].
- [30] E. D'Hoker, J. Estes and M. Gutperle, *Ten-dimensional supersymmetric Janus solutions*, *Nucl. Phys. B* **757** (2006) 79 [[hep-th/0603012](#)] [[INSPIRE](#)].
- [31] G. Dall'Agata and G. Inverso, *On the Vacua of  $N = 8$  Gauged Supergravity in 4 Dimensions*, *Nucl. Phys. B* **859** (2012) 70 [[arXiv:1112.3345](#)] [[INSPIRE](#)].
- [32] A. Guarino and C. Sterckx, *S-folds and (non-)supersymmetric Janus solutions*, *JHEP* **12** (2019) 113 [[arXiv:1907.04177](#)] [[INSPIRE](#)].
- [33] B. Assel and A. Tomasiello, *Holographic duals of 3d S-fold CFTs*, *JHEP* **06** (2018) 019 [[arXiv:1804.06419](#)] [[INSPIRE](#)].
- [34] I. Garozzo, G. Lo Monaco and N. Mekareeya, *The moduli spaces of S-fold CFTs*, *JHEP* **01** (2019) 046 [[arXiv:1810.12323](#)] [[INSPIRE](#)].
- [35] I. Garozzo, G. Lo Monaco, N. Mekareeya and M. Sacchi, *Supersymmetric Indices of 3d S-fold SCFTs*, *JHEP* **08** (2019) 008 [[arXiv:1905.07183](#)] [[INSPIRE](#)].
- [36] N. Bobev, F.F. Gautason, K. Pilch, M. Suh and J. Van Muiden, *Janus and J-fold Solutions from Sasaki-Einstein Manifolds*, *Phys. Rev. D* **100** (2019) 081901 [[arXiv:1907.11132](#)] [[INSPIRE](#)].
- [37] A.B. Clark, D.Z. Freedman, A. Karch and M. Schnabl, *Dual of the Janus solution: An interface conformal field theory*, *Phys. Rev. D* **71** (2005) 066003 [[hep-th/0407073](#)] [[INSPIRE](#)].
- [38] D. Gaiotto and E. Witten, *S-duality of Boundary Conditions In  $N = 4$  Super Yang-Mills Theory*, *Adv. Theor. Math. Phys.* **13** (2009) 721 [[arXiv:0807.3720](#)] [[INSPIRE](#)].
- [39] E. D'Hoker, J. Estes and M. Gutperle, *Interface Yang-Mills, supersymmetry and Janus*, *Nucl. Phys. B* **753** (2006) 16 [[hep-th/0603013](#)] [[INSPIRE](#)].
- [40] D. Gaiotto and E. Witten, *Janus Configurations, Chern-Simons Couplings, And The theta-Angle in  $N = 4$  Super Yang-Mills Theory*, *JHEP* **06** (2010) 097 [[arXiv:0804.2907](#)] [[INSPIRE](#)].
- [41] B. de Wit, H. Samtleben and M. Trigiante, *Magnetic charges in local field theory*, *JHEP* **09** (2005) 016 [[hep-th/0507289](#)] [[INSPIRE](#)].
- [42] A. Guarino, *CSO<sub>c</sub> superpotentials*, *Nucl. Phys. B* **900** (2015) 501 [[arXiv:1508.05055](#)] [[INSPIRE](#)].
- [43] B. de Wit, H. Samtleben and M. Trigiante, *The maximal  $D = 4$  supergravities*, *JHEP* **06** (2007) 049 [[arXiv:0705.2101](#)] [[INSPIRE](#)].
- [44] H. Samtleben, *Lectures on Gauged Supergravity and Flux Compactifications*, *Class. Quant. Grav.* **25** (2008) 214002 [[arXiv:0808.4076](#)] [[INSPIRE](#)].

- [45] M. Trigiante, *Gauged Supergravities*, *Phys. Rept.* **680** (2017) 1 [[arXiv:1609.09745](#)] [[INSPIRE](#)].
- [46] N. Bobev, T. Fischbacher and K. Pilch, *Properties of the new  $\mathcal{N} = 1$   $AdS_4$  vacuum of maximal supergravity*, *JHEP* **01** (2020) 099 [[arXiv:1909.10969](#)] [[INSPIRE](#)].
- [47] G. Aldazabal, P.G. Camara, A. Font and L.E. Ibáñez, *More dual fluxes and moduli fixing*, *JHEP* **05** (2006) 070 [[hep-th/0602089](#)] [[INSPIRE](#)].
- [48] G. Aldazabal, P.G. Camara and J.A. Rosabal, *Flux algebra, Bianchi identities and Freed-Witten anomalies in F-theory compactifications*, *Nucl. Phys. B* **814** (2009) 21 [[arXiv:0811.2900](#)] [[INSPIRE](#)].
- [49] P. Breitenlohner and D.Z. Freedman, *Positive Energy in anti-de Sitter Backgrounds and Gauged Extended Supergravity*, *Phys. Lett.* **115B** (1982) 197 [[INSPIRE](#)].
- [50] O. Hohm and H. Samtleben, *Exceptional field theory. II.  $E_{7(7)}$* , *Phys. Rev. D* **89** (2014) 066017 [[arXiv:1312.4542](#)] [[INSPIRE](#)].
- [51] O. Hohm and H. Samtleben, *Consistent Kaluza-Klein Truncations via Exceptional Field Theory*, *JHEP* **01** (2015) 131 [[arXiv:1410.8145](#)] [[INSPIRE](#)].
- [52] A. Baguet, O. Hohm and H. Samtleben,  *$E_{6(6)}$  Exceptional Field Theory: Review and Embedding of Type IIB*, *PoS(CORFU2014)133* (2015) [[arXiv:1506.01065](#)] [[INSPIRE](#)].
- [53] A. Baguet, O. Hohm and H. Samtleben, *Consistent Type IIB Reductions to Maximal 5D Supergravity*, *Phys. Rev. D* **92** (2015) 065004 [[arXiv:1506.01385](#)] [[INSPIRE](#)].
- [54] K. Pilch and N.P. Warner, *A new supersymmetric compactification of chiral IIB supergravity*, *Phys. Lett. B* **487** (2000) 22 [[hep-th/0002192](#)] [[INSPIRE](#)].
- [55] A. Khavaev, K. Pilch and N.P. Warner, *New vacua of gauged  $N = 8$  supergravity in five-dimensions*, *Phys. Lett. B* **487** (2000) 14 [[hep-th/9812035](#)] [[INSPIRE](#)].