

Exact response theory and Kuramoto dynamics

*Original*

Exact response theory and Kuramoto dynamics / Amadori, D., Colangeli, M., Correa, A., Rondoni, L.. - In: PHYSICA D-NONLINEAR PHENOMENA. - ISSN 0167-2789. - STAMPA. - 429:133076(2022), pp. 1-11.  
[10.1016/j.physd.2021.133076]

*Availability:*

This version is available at: 11583/2942252 since: 2021-12-10T14:03:41Z

*Publisher:*

Elsevier

*Published*

DOI:10.1016/j.physd.2021.133076

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

Elsevier postprint/Author's Accepted Manuscript

© 2022. This manuscript version is made available under the CC-BY-NC-ND 4.0 license  
<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:  
<http://dx.doi.org/10.1016/j.physd.2021.133076>

(Article begins on next page)

# EXACT RESPONSE THEORY AND KURAMOTO DYNAMICS

DEBORA AMADORI, MATTEO COLANGELI, ASTRID CORREA, AND LAMBERTO RONDONI

ABSTRACT. The dynamics of Kuramoto oscillators is investigated in terms of the exact response theory based on the Dissipation Function, which has been introduced in the field of nonequilibrium molecular dynamics. While linear response theory is a cornerstone of nonequilibrium statistical mechanics, it does not apply, in general, to systems undergoing phase transitions. Indeed, even a small perturbation may in that case result in a large modification of the state. An exact theory is instead expected to handle such situations. The Kuramoto dynamics, which undergoes synchronization transitions, is thus investigated *analytically and numerically* as a testbed for the exact theory mentioned above. A comparison between the two approaches shows how the linear theory fails, while the exact theory yields the correct response.

## 1. INTRODUCTION

The response of a system with many degrees of freedom to an external stimulus is a central topic in nonequilibrium statistical mechanics. Its investigation has greatly progressed with the works of Callen, Green, Kubo, and Onsager, in particular, who contributed to the development of linear response theory [32, 36]. In the '90s, the derivation of the Fluctuation Relations [19, 21, 26] provided the framework for a more general response theory, applicable to both Hamiltonian as well as dissipative deterministic particle systems [8, 10, 12, 13, 14, 24, 36, 40]. The study of response in stochastic processes, with a special focus on diffusion and Markov jump processes, has also been inspired by fluctuation relations, and has been studied *e.g.* in [2, 4, 7, 11, 15]. Moreover, the role of causality, expressed by the Kramers-Kronig relations, in nonlinear extensions of the linear response theory has been discussed in [35].

The introduction of the Dissipation Function, first made explicit in [22], and developed as the observable of interest in Fluctuation Relations [23, 41], paved the way to an exact response theory. A theory expected to hold in presence of arbitrarily large perturbations and modifications of states, which allows the study of the relaxation of particle systems to equilibrium or non-equilibrium steady states.

In this work we present and apply the Dissipation Function formalism to the Kuramoto model [33, 34], which is considered a prototype of many particle systems exhibiting *synchronization*, a phenomenon familiar in many physical and biological contexts [25, 27, 31, 37, 42, 43]. Furthermore, the Kuramoto model provides the stage for a large

---

*Date:* November 5, 2021.

research endeavor, in applied mathematics, control theory and statistical physics [1, 3, 5, 18, 28, 39, 43]. See [29, 16] for recent reviews on the subject.

In this paper, our aim is two-fold. On the one hand, we probe the exact response theory on a dissipative system with many degrees of freedom undergoing nonequilibrium phase transitions, which is in fact a challenging open problem. On the other hand, while a vast mathematical literature exists on the Kuramoto model, it is interesting to analyze it from a new statistical mechanical perspective, in which some known results are reinterpreted, cf. *e.g.* Refs.[6, 17].

Our conclusion is that, while the linear response theory cannot characterize the Kuramoto synchronization process, the exact theory does. In particular, we obtain synchronization within the formalism of the Dissipation Function, thus showing how such a behaviour is captured by the exact response theory, while it is not evidenced by the linear theory. Synchronization corresponds indeed to the maximum value of the Dissipation Function, which we prove is attained in time. When the number of oscillators  $N$  is large, this maximum value is proportional to the oscillators coupling constant  $K$ .

This paper is organized as follows. In Sec. 2 we review some basic properties of the Kuramoto dynamics. In Sec. 3 we illustrate the main ingredients of the Dissipation Function response theory. In Sec. 4 we study the response theory for the Kuramoto dynamics of identical oscillators. In Sec. 5 we review the linear response theory, and we compare it, [analytically and numerically](#), with the exact response formalism. We draw our conclusions in Sec. 6.

## 2. THE KURAMOTO SYSTEM

The Kuramoto dynamics is defined on the  $N$ -dimensional torus,  $\mathcal{T}^N = (\mathbb{R}/(2\pi\mathbb{Z}))^N$ , with  $N \geq 1$ , by the following set of coupled first order ODEs, for the phases  $\theta_i(t)$ :

$$\dot{\theta}_i = \omega_i + \frac{K}{N} \sum_{j=1}^N \sin(\theta_j - \theta_i) \quad i = 1, \dots, N \quad (2.1)$$

where  $K > 0$  is a constant, and the natural frequencies  $\omega_i \in \mathbb{R}$  are drawn from some given distribution  $g(\omega)$ .

The  $N$  oscillators are represented by points rotating on the unit circle centered at the origin of the complex plane, more precisely by  $e^{i\theta_j}$  with  $j = 1, \dots, N$ . By introducing the polar coordinates of the barycenter,

$$Re^{i\Phi} = \frac{1}{N} \sum_{j=1}^N e^{i\theta_j} \quad (2.2)$$

with  $R \in [0, 1]$  and  $\Phi \in \mathbb{R}$  (defined if  $R > 0$ ), one can rewrite Eq.(2.1) as follows:

$$\dot{\theta}_i = \omega_i + KR \sin(\Phi - \theta_i), \quad i = 1, \dots, N \quad (2.3)$$

where  $R = R(\theta(t))$  is the *order parameter* and  $\Phi = \Phi(\theta(t))$  the *collective phase*, with  $\theta = (\theta_1, \dots, \theta_N) \in \mathcal{M} = \mathcal{T}^N$ , and  $\mathcal{M}$  the phase space. It is to be remarked that the

Kuramoto dynamics (2.3) can be written as a gradient flow:

$$\dot{\theta} = -\nabla f(\theta) \quad (2.4)$$

with potential

$$f(\theta) = -\sum_{i=1}^N \omega_i \theta_i + \frac{K}{2N} \sum_{i,j=1}^N \left(1 - \cos(\theta_j - \theta_i)\right) \quad (2.5)$$

that is analytic in  $\theta$ .

A *complete frequency synchronization* occurs when the differences  $\theta_i(t) - \theta_j(t)$  tend to a constant for all  $i$  and  $j$ , and  $R(\theta(t))$  tends to a given  $R^\infty \in (0, 1]$ , as  $t \rightarrow +\infty$ . In case  $R^\infty = 1$ , all the  $N$  terms of the sum in (2.2) coincide, hence the Kuramoto system undergoes *phase synchronization*.

**Identities for the order parameter.** We remark that equation (2.2) leads to the following identities,

$$R = \frac{1}{N} \sum_{i=1}^N \cos(\Phi - \theta_i), \quad (2.6)$$

$$0 = \frac{1}{N} \sum_{i=1}^N \sin(\Phi - \theta_i), \quad (2.7)$$

$$R \sin(\Phi - \theta_i) = \frac{1}{N} \sum_{j=1}^N \sin(\theta_j - \theta_i), \quad i = 1, \dots, N \quad (2.8)$$

$$R \cos(\Phi - \theta_i) = \frac{1}{N} \sum_{j=1}^N \cos(\theta_j - \theta_i), \quad i = 1, \dots, N. \quad (2.9)$$

From the identities (2.6) and (2.9), we find that:

$$R^2 = \frac{1}{N^2} \sum_{i,j=1}^N \cos(\theta_j - \theta_i). \quad (2.10)$$

For  $\theta \in \mathcal{M}$ , we can rewrite Eq.(2.3) as:

$$\dot{\theta} = W + V(\theta) = V_K(\theta) \quad (2.11)$$

where  $W = (\omega_1, \dots, \omega_N)$  is interpreted as an *equilibrium* vector field made of  $N$  natural frequencies, while  $V$  represents a *nonequilibrium* perturbation with components:

$$V_i(\theta) = \frac{K}{N} \sum_{j=1}^N \sin(\theta_j - \theta_i) = KR \sin(\Phi - \theta_i), \quad i = 1, \dots, N. \quad (2.12)$$

For later use, we prove the following identity.

**Lemma 2.1.** *The divergence of the Kuramoto vector field  $V_K$  of Eq.(2.11), i.e. the associated phase space volumes variation rate  $\Lambda$ , satisfies:*

$$\Lambda := \operatorname{div}_\theta V = K(1 - NR^2). \quad (2.13)$$

*Proof.* By means of (2.12), for  $i = 1, \dots, N$  one has

$$\begin{aligned} \partial_{\theta_i} V_i &= \frac{K}{N} \partial_{\theta_i} \left( \sum_{i \neq j=1}^N \sin(\theta_j - \theta_i) \right) \\ &= -\frac{K}{N} \left( \sum_{i \neq j=1}^N \cos(\theta_j - \theta_i) \right) = -\frac{K}{N} \left( \sum_{j=1}^N \cos(\theta_j - \theta_i) - 1 \right) \\ &= -KR \cos(\Phi - \theta_i) + \frac{K}{N} \end{aligned}$$

where we used (2.9). Summing over  $i$ , and using (2.6), Eq.(2.13) follows.  $\square$

Therefore, the Kuramoto dynamics do not preserve the phase space volumes, and  $\Lambda$  actually varies in time, since  $R$  is a function of the dynamical variables  $\theta(t)$ .

### 3. MATHEMATICAL FRAMEWORK OF RESPONSE THEORY

Let us summarize the mathematical framework of exact response theory, as described in [8, 24, 30, 41]. The starting point is a flow  $S^t : \mathcal{M} \rightarrow \mathcal{M}$ , with phase space  $\mathcal{M} \subset \mathbb{R}^N$ ,  $N \geq 1$ , that is usually determined by an ODE system

$$\dot{\theta} = V(\theta), \quad \theta \in \mathcal{M} \quad (3.1)$$

with  $V$  defined on  $\mathcal{M}$ . Let  $S^t \theta$  denote the solution at time  $t \in \mathbb{R}$ , with initial condition  $\theta$ , of such ODEs. The second ingredient is a probability measure  $d\mu_0(\theta) = f_0(\theta)d\theta$  on  $\mathcal{M}$ , with positive and continuously differentiable density  $f_0$ . A time evolution is induced on the simplex of probabilities on  $\mathcal{M}$ , defining the probability at a time  $t \in \mathbb{R}$  as:

$$\mu_t(E) = \mu_0(S^{-t}E)$$

for each measurable set  $E \subset \mathcal{M}$ . This amounts to consider probability in a phase space, like the mass of a fluid in real space. The corresponding continuity equation for the probability densities is the (generalized) Liouville equation:

$$\frac{\partial f}{\partial t} + \operatorname{div}_{\theta}(fV) = 0 \quad (3.2)$$

Denoting by  $f_t$  the solution of Eq.(3.2) with initial datum  $f_0$ , we can write  $d\mu_t = f_t d\theta$ .

Introducing the *Dissipation Function*  $\Omega^{f,V}$  [30, 41]:

$$\Omega^{f,V}(\theta) := -\Lambda(\theta) - V(\theta) \cdot \nabla \log f(\theta) \quad (3.3)$$

with  $\nabla = (\partial_{\theta_1}, \dots, \partial_{\theta_N})$  and

$$\Lambda = \operatorname{div}_{\theta} V,$$

the Euler version (3.2) of the Liouville equation may also be written as:

$$\frac{\partial f}{\partial t} = f \Omega^{f,V} \quad (3.4)$$

Also, the equation (3.2) can be cast in the Lagrangian form

$$\frac{df}{dt} = -f \Lambda, \quad (3.5)$$

where  $\frac{d}{dt} = \frac{\partial}{\partial t} + V \cdot \nabla_\theta$  is the total derivative along the flow (3.1).

Direct integration of Eq.(3.5) yields

$$f_{s+t}(S^t\theta) = \exp \left\{ -\Lambda_{0,t}(\theta) \right\} f_s(\theta), \quad \forall t, s \geq 0 \quad (3.6)$$

where we used the notation

$$\mathcal{O}_{s,t}(\theta) := \int_s^t \mathcal{O}(S^\tau\theta) d\tau \quad (3.7)$$

for the phase functions, or *observables*,  $\mathcal{O} : \mathcal{M} \rightarrow \mathbb{R}$ , so that, in particular,  $\Lambda_{0,t}(\theta) = \int_0^t \Lambda(S^\tau\theta) d\tau$ . In the following Proposition, this notation is used with the observable  $\mathcal{O} = \Omega^{f,V}(\theta)$  given in (3.3), so that the time integral in (3.7) will correspondingly be denoted by  $\Omega_{s,t}^{f,V}$ .

**Proposition 3.1.** *For all  $t, s \in \mathbb{R}$ , the following identity holds:*

$$f_{s+t}(\theta) = \exp \left\{ \Omega_{-t,0}^{f_s,V}(\theta) \right\} f_s(\theta). \quad (3.8)$$

*Proof.* We start by claiming that

$$\Omega_{0,s}^{f_t,V}(\theta) = \log \frac{f_t(\theta)}{f_t(S^s\theta)} - \Lambda_{0,s}(\theta). \quad (3.9)$$

Indeed, one has:

$$V(S^u\theta) \cdot \nabla \log f_t(S^u\theta) = \frac{d}{du} \log f_t(S^u\theta) \quad (3.10)$$

because  $t$  is fixed and  $f_t$  does not depend explicitly on  $u$ , hence Eqs.(3.3) and (3.10) imply:

$$\begin{aligned} \Omega_{0,s}^{f_t,V}(\theta) &= - \int_0^s [\Lambda(S^u\theta) + V \cdot \nabla \log f_t(S^u\theta)] du \\ &= -\Lambda_{0,s}(\theta) - \int_0^s \frac{d}{du} \log f_t(S^u\theta) d\theta = -\Lambda_{0,s}(\theta) - \log \frac{f_t(S^s\theta)}{f_t(\theta)} \end{aligned}$$

which leads to Eq.(3.9). Next, Eqs.(3.6) and Eq.(3.9) lead to

$$\exp \left\{ \Omega_{s,s+t}^{f_s,V}(\theta) \right\} f_s(S^{s+t}\theta) = \exp \left\{ -\Lambda_{s,s+t}(\theta) \right\} f_s(S^s\theta) = f_{s+t}(S^{s+t}\theta) \quad (3.11)$$

which yields (3.8).  $\square$

As a consequence of Proposition 3.1, a probability density  $f$  is *invariant* under the dynamics if and only if  $\Omega^{f,V}$  identically vanishes:

$$\Omega^{f,V}(\theta) = 0, \quad \forall \theta \in \mathcal{M}. \quad (3.12)$$

In the sequel, we shall use the notation

$$\langle \mathcal{O} \rangle_t := \int_{\mathcal{M}} \mathcal{O}(\theta) f_t(\theta) d\theta \quad (3.13)$$

to denote the average of an observable with respect to the probability measure  $\mu_t = f_t d\theta$ . The response theory states that the average  $\langle \mathcal{O} \rangle_t$  can be expressed in terms of the initial density  $f_0$ , which is known. More precisely:

**Lemma 3.1. (Exact response):** *Given  $\{S^t\}_{t \in \mathbb{R}}$  and an integrable observable  $\mathcal{O} : \mathcal{M} \rightarrow \mathbb{R}$ , the following identity holds:*

$$\langle \mathcal{O} \rangle_t = \langle \mathcal{O} \rangle_0 + \int_0^t \langle (\mathcal{O} \circ S^\tau) \Omega_V^{f_0} \rangle_0 d\tau. \quad (3.14)$$

*Proof.* First of all,  $f_0$  is smooth as a function of  $\theta$  by assumption, and evolves according to the Liouville equation. Therefore,  $f_t$  is also smooth with respect to  $\theta$  and  $t$  for every finite time  $t$ . In turn,  $\Omega^{f_t, V}(\theta)$  is differentiable with respect to  $\theta$  and  $t$ , if  $f_0$  (that depends only on  $\theta$ ) is differentiable with respect to  $\theta$ . These conditions are immediately verified for differentiable  $f_0$ , and smooth dynamics on a compact manifold. Therefore two identities can be derived for integrable  $\mathcal{O}$ :

$$\begin{aligned} \mathcal{O}_{0,s}(\theta) &= \int_0^s \mathcal{O}(S^u \theta) du = \int_\tau^{s+\tau} \mathcal{O}(S^{u-\tau} \theta) du = \int_\tau^{s+\tau} \mathcal{O}(S^{-\tau} S^u \theta) du \\ &= \mathcal{O}_{\tau, s+\tau}(S^{-\tau} \theta), \end{aligned}$$

which is valid for every  $\tau \in \mathbb{R}$ , and

$$\begin{aligned} \langle \mathcal{O} \rangle_{t+s} &= \int \mathcal{O}(\theta) f_{t+s}(\theta) d\theta = \int \mathcal{O}(S^s(S^{-s}\theta)) f_{t+s}(S^s(S^{-s}\theta)) \left| \frac{\partial \theta}{\partial (S^{-s}\theta)} \right| d(S^{-s}\theta) \\ &= \int \mathcal{O}(S^s(S^{-s}\theta)) f_{t+s}(S^s(S^{-s}\theta)) \exp \left\{ \Lambda_{-s,0}(\theta) \right\} d(S^{-s}\theta) \\ &= \int \mathcal{O}(S^s(S^{-s}\theta)) f_{t+s}(S^s(S^{-s}\theta)) \exp \left\{ \Lambda_{0,s}(S^{-s}\theta) \right\} d(S^{-s}\theta) \\ &= \int \mathcal{O}(S^s \theta) f_{t+s}(S^s \theta) \exp \left\{ \Lambda_{0,s}(\theta) \right\} d\theta = \int \mathcal{O}(S^s \theta) f_t(\theta) d\theta \\ &= \langle \mathcal{O} \circ S^s \rangle_t \end{aligned} \quad (3.15)$$

to obtain [30]:

$$\frac{d}{ds} \langle \mathcal{O} \rangle_s = \langle \mathcal{O} (\Omega^{f_r, V} \circ S^{r-s}) \rangle_s \quad (3.16)$$

which holds  $\forall r \in \mathbb{R}^+$ . Note that in Eq. (3.15) we used the relation

$$\left| \frac{\partial \theta}{\partial (S^{-s}\theta)} \right| = \exp \left\{ \Lambda_{-s,0}(\theta) \right\}, \quad (3.17)$$

which is discussed in Appendix B, see Eq. (B.3). Choosing  $r = 0$  one finds

$$\frac{d}{ds} \langle \mathcal{O} \rangle_s = \langle \mathcal{O} (\Omega^{f_0, V} \circ S^{-s}) \rangle_s = \langle (\mathcal{O} \circ S^s) \Omega^{f_0, V} \rangle_0 \quad (3.18)$$

where we used (3.15). Then, integrating over time from 0 to  $t$ , Eq.(3.18) yields (3.14).  $\square$

It may be useful to endow the density  $f_0$  with physical meaning. To this aim, we assume that, for  $t \in (-\infty, 0)$ , the dynamics is determined by a (non dissipative) vector field  $V_0$ ,

$$\dot{\theta} = V_0(\theta) \quad (3.19)$$

and that it has reached an equilibrium state, described by the probability measure  $d\mu_0 = f_0 d\theta$ , long before the time  $t = 0$ . In other words, we assume that  $\mu_0$  is invariant under the dynamics (3.19), which we call *unperturbed* or *reference* dynamics. At time  $t = 0$ , the dynamics (3.19) is perturbed and the perturbation remains in place for  $t \geq 0$ . In general, the density  $f_0$  will not be invariant under the perturbed vector field  $V$ , Eq.(3.1) say. Therefore, it will evolve as prescribed by Eq.(3.4) into a different density,  $f_t$ , at time  $t > 0$ . Nevertheless, Eq.(3.14) expresses the average  $\langle \mathcal{O} \rangle_t$  in terms of a correlation function computed with respect to  $f_0$ , the non-invariant density, which is only invariant under the unperturbed dynamics.

The full range of applicability of this theory is still to be identified. However, it obviously applies to smooth dynamics on smooth compact manifolds, such as the Kuramoto dynamics (2.1), which has  $\mathcal{M} = \mathcal{T}^N$ . One advantage of using the Dissipation Function, compared to other possible exact approaches to response, apart from molecular dynamics efficiency, is that it corresponds to a physically measurable quantity, *e.g.* proportional to a current, that is adapted to the initial state of the system of interest. Moreover, it provides necessary and sufficient conditions for relaxation of ensembles, as well as sufficient conditions for the single system relaxation, known as T-mixing [30, 41].

Henceforth, we shall denote by  $\Omega^{f,0}$  the Dissipation Function (3.3) defined in terms of the unperturbed flow  $V_0$ . The analysis of the response theory for a specific example of the Kuramoto model is discussed in the next Section.

#### 4. RESPONSE THEORY FOR IDENTICAL OSCILLATORS

Let us focus on the case of identical oscillators, namely the Kuramoto dynamics in which all the natural frequencies  $\omega_i$  in Eq.(2.1) equal the same constant  $\omega \in \mathbb{R}$ . In particular, let the unperturbed dynamics be defined by the vector field  $V_0(\theta) = W = (\omega, \dots, \omega)$ , which corresponds to  $K = 0$  in Eq.(2.1), *i.e.* to decoupled oscillators, equipped with same natural frequency. Such dynamics are conservative, since  $\text{div}_\theta V_0 = 0$ . The corresponding steady state can then be considered an equilibrium state. At time  $t = 0$  the perturbation  $V$  is switched on, and we can write:

$$\dot{\theta} = \begin{cases} W & t < 0 \\ W + V(\theta) & t > 0 \end{cases} \quad (4.1)$$

The perturbed dynamics corresponds to the Kuramoto dynamics (2.1), which is not conservative, cf. Eq.(2.13). As an initial probability density, invariant under the unperturbed dynamics, we may take the factorized density:

$$f_0(\theta) = (2\pi)^{-N} \quad (4.2)$$

which, indeed, yields:

$$\Omega_0^{f_0} = -(\operatorname{div} V_0 + V_0 \cdot \nabla \log f_0) \equiv 0, \quad \text{and} \quad \frac{\partial f}{\partial t} = 0. \quad (4.3)$$

After the perturbation, the Dissipation Function takes the form:

$$\Omega_V^{f_0} = -(\operatorname{div}_\theta V + V \cdot \nabla \log f_0) = K(NR^2 - 1) = \frac{K}{N} \sum_{i,j=1}^N \cos(\theta_j - \theta_i) - K \quad (4.4)$$

and the density evolves as:

$$f_t(\theta) = \frac{1}{(2\pi)^N} \exp \left[ -K(t - NR_{-t,0}^2(\theta)) \right] \quad (4.5)$$

where  $R_{-t,0}$  denotes the integral of  $R$  from time  $-t$  to 0, cf. Eq.(3.7).

**Remark 4.1.** *The Dissipation Function Eq.(4.4) is of class  $C^\infty$ .*

Using the formula (3.14) to compute the response for the observable  $\mathcal{O} = \Omega_V^{f_0}$ , we obtain:

$$\langle \Omega_V^{f_0} \rangle_t = \langle \Omega_V^{f_0} \rangle_0 + \int_0^t \langle (\Omega_V^{f_0} \circ S^\tau) \Omega_V^{f_0} \rangle_0 d\tau \quad (4.6)$$

that is

$$\int_{\mathcal{M}} \Omega_V^{f_0}(\theta) f_t(\theta) d\theta = (2\pi)^{-N} \int_{\mathcal{M}} \Omega_V^{f_0}(\theta) d\theta + (2\pi)^{-N} \int_0^t \int_{\mathcal{M}} \Omega_V^{f_0}(S^\tau(\theta)) \Omega_V^{f_0}(\theta) d\theta d\tau.$$

Moreover:

$$\langle R^2 \rangle_0 = \frac{1}{N}, \quad \text{hence} \quad \langle \Omega_V^{f_0} \rangle_0 = K(N\langle R^2 \rangle_0 - 1) = 0 \quad (4.7)$$

as expected.

**Remark 4.2.** *Note that the scalar field  $\Omega_0^{f_0}$  is identically 0, while  $\Omega_V^{f_0}$  is not, see (4.4). However, the phase space average  $\langle \Omega_V^{f_0} \rangle_0$  vanishes.*

Therefore, using Eqs.(3.14) and (4.4) we can write:

$$\begin{aligned} \langle \Omega_V^{f_0} \rangle_t &= \int_0^t \langle (\Omega_V^{f_0} \circ S^\tau) \Omega_V^{f_0} \rangle_0 d\tau \\ &= KN \int_0^t \langle \Omega_V^{f_0} [R^2 \circ S^\tau] \rangle_0 d\tau - K \int_0^t \langle \Omega_V^{f_0} \rangle_0 d\tau \\ &= KN \int_0^t \langle \Omega_V^{f_0} [R^2 \circ S^\tau] \rangle_0 d\tau \\ &= K^2 N^2 \int_0^t \langle R^2 [R^2 \circ S^\tau] \rangle_0 d\tau - K^2 N \int_0^t \langle R^2 \circ S^\tau \rangle_0 d\tau \end{aligned}$$

For the second integral we have:

$$\begin{aligned} \int_0^t \langle R^2 \circ S^\tau \rangle_0 d\tau &= \frac{1}{(2\pi)^N} \int_0^t \int_{\mathcal{M}} R^2(S^\tau \theta) d\theta d\tau \\ &= \frac{1}{(2\pi)^N} \int_0^t \int_{\mathcal{M}} R^2(S^\tau \theta) \left| \frac{\partial \theta}{\partial S^\tau \theta} \right| dS^\tau \theta d\tau \\ &= \frac{1}{(2\pi)^N} \int_0^t \int_{\mathcal{M}} R^2(S^\tau \theta) e^{\Lambda_{0,\tau}(\theta)} dS^\tau \theta \end{aligned}$$

Explicit calculations can be carried out for  $N = 2$  and will be discussed in Sec. 4.1, while the study of the general case with  $N > 2$  is deferred to Sec. 4.2.

**4.1. The case with two oscillators.** For  $N = 2$  and  $\omega \geq 0$ , consider the system for two oscillators:

$$\begin{cases} \dot{\theta}_1 = \frac{\omega}{2} + \frac{K}{2} \sin(\theta_2 - \theta_1) \\ \dot{\theta}_2 = -\frac{\omega}{2} + \frac{K}{2} \sin(\theta_1 - \theta_2). \end{cases} \quad (4.8)$$

For  $\omega = 0$  we say that the two oscillators are *identical* (in the sense that their natural frequencies coincide). Setting  $\psi = \theta_1 - \theta_2$ , we obtain the following equation:

$$\frac{d\psi}{dt} = \omega - K \sin(\psi). \quad (4.9)$$

With a slight abuse of notation, in the following we denote by  $S^t \theta$ ,  $S^t \psi$  the flows corresponding to (4.8), (4.9) respectively, with initial data  $\theta = (\theta_1, \theta_2)$ .

The solution of (4.9) can be explicitly expressed as ([9, Appendix D]) :

$$\psi(t) = \psi \circ S^t = 2 \arctan(g(\psi, t)) \quad (4.10)$$

where

$$g(\psi, t) = \begin{cases} \frac{\sqrt{K^2 - \omega^2}}{\omega} \left[ \tanh \left( h_1(\psi) - \frac{t\sqrt{K^2 - \omega^2}}{2} \right) + \frac{K}{\sqrt{K^2 - \omega^2}} \right] & \text{if } K > \omega \\ \frac{\sqrt{\omega^2 - K^2}}{\omega} \tan \left( \frac{t\sqrt{\omega^2 - K^2}}{2} + h_2(\psi) \right) + \frac{K}{\omega} & \text{if } 0 \leq K < \omega \\ e^{-Kt} \tan\left(\frac{\psi}{2}\right) & \text{if } \omega = 0 \end{cases}$$

and the functions  $h_1$  and  $h_2$  are given by:

$$\begin{aligned} h_1(\psi) &= \tanh^{-1} \left( \frac{\omega}{\sqrt{K^2 - \omega^2}} \tan \left( \frac{\psi}{2} \right) - \frac{K}{\sqrt{K^2 - \omega^2}} \right), \\ h_2(\psi) &= \arctan \frac{\omega \tan \left( \frac{\psi}{2} \right) - K}{\sqrt{\omega^2 - K^2}}. \end{aligned}$$

Recalling Eq.(2.10), we find that  $(R^2 \circ S^t)$  can be written as

$$R^2(S^t \theta) = \frac{1}{2} [1 + \cos(S^t \psi)] = \frac{1}{g^2(S^t \psi) + 1}. \quad (4.11)$$

Restricting our analysis to the identical oscillator case,  $\omega = 0$ , yields:

$$R^2(S^t\theta) = \left( \tan^2\left(\frac{\psi}{2}\right) e^{-2Kt} + 1 \right)^{-1} \quad (4.12)$$

and

$$S^t\psi \rightarrow 0 \quad \text{for } t \rightarrow +\infty, \quad \text{if } |\psi| \neq \pi$$

while

$$|S^t\psi| \rightarrow \pi \quad \text{for } t \rightarrow -\infty, \quad \text{if } \psi \neq 0.$$

In particular, for  $\theta_1 \neq \theta_2$  and  $\theta_1, \theta_2 \in [0, 2\pi)$ , then  $t \rightarrow -\infty$  yields  $S^t\psi \rightarrow -\pi$  if  $\theta_1 < \pi$ , and  $S^t\psi \rightarrow \pi$  if  $\theta_1 > \pi$ .

Then, the set

$$E_\infty = \{(\theta_1, \theta_2) \in \mathcal{T}^2 : \theta_1 = \theta_2\}$$

is invariant and attracting for the Kuramoto dynamics, while the set

$$E_{-\infty} = \{(\theta_1, \theta_2) \in \mathcal{T}^2 : |\theta_1 - \theta_2| = \pi\}$$

is invariant and repelling. This also implies that:

$$R^2(S^t\theta) \rightarrow 0, \quad \Omega_V^{f_0} \rightarrow -K, \quad \text{for } \psi \neq 0, \quad t \rightarrow -\infty$$

while

$$R^2(S^t\theta) \rightarrow 1, \quad \Omega_V^{f_0} \rightarrow K, \quad \text{for } |\psi| \neq \pi, \quad t \rightarrow \infty.$$

Consequently, Eq.(4.5) shows that the probability piles up on the zero Lebesgue measure sets  $E_\infty$  and  $E_{-\infty}$ , respectively for  $t \rightarrow \infty$  and  $t \rightarrow -\infty$ .

For  $\tau \geq 0$ , the following relations also hold:

$$\langle R^2 \circ S^\tau \rangle_0 = \frac{1}{(2\pi)^2} \int_{\mathcal{M}} \frac{1}{\tan^2\left(\frac{\theta_1 - \theta_2}{2}\right) e^{-2K\tau} + 1} d\theta = \frac{1}{e^{-K\tau} + 1} \quad (4.13)$$

and

$$\langle R^2(R^2 \circ S^\tau) \rangle_0 = \frac{1}{8\pi^2} \int_{\mathcal{M}} \frac{1 + \cos(\theta_1 - \theta_2)}{\tan^2\left(\frac{\theta_1 - \theta_2}{2}\right) e^{-2K\tau} + 1} d\theta = \frac{2e^{-K\tau} + 1}{2(e^{-K\tau} + 1)^2}$$

which then yields

$$\int_0^t \langle R^2 \circ S^\tau \rangle_0 d\tau = t + \frac{\ln(e^{-Kt} + 1)}{K} - \frac{\ln(2)}{K}$$

and

$$\int_0^t \langle R^2(R^2 \circ S^\tau) \rangle_0 d\tau = \frac{t}{2} + \frac{1}{2K} \left[ \frac{3}{2} + \ln\left(\frac{e^{-Kt} + 1}{2}\right) - \frac{2}{e^{Kt} + 1} - \frac{1}{e^{-Kt} + 1} \right].$$

Thus, we finally obtain the explicit expressions

$$\langle \Omega_V^{f_0} \rangle_t = K \tanh\left(\frac{Kt}{2}\right) \quad (4.14)$$

and

$$\langle (\Omega_V^{f_0} \circ S^t) \Omega_V^{f_0} \rangle_0 = \frac{K^2}{1 + \cosh(Kt)}. \quad (4.15)$$




FIGURE 1. Behavior of  $\langle \Omega_V^{f_0} \rangle_t$  and  $\langle (\Omega_V^{f_0} \circ S^t) \Omega_V^{f_0} \rangle_0$  as functions of time, for  $N = 2$ ,  $K = 1$  and  $\omega = 0$ . Disks and solid lines correspond to the numerical and analytical results, respectively. The averages were taken over a set of 5000 trajectories with initial data sampled from the uniform distribution on  $[0, 2\pi)$ .

In the limit  $t \rightarrow +\infty$ , we thus find the asymptotic values

$$\left\langle \Omega_V^{f_0} \right\rangle_t \rightarrow K \quad \text{and} \quad \left\langle (\Omega_V^{f_0} \circ S^t) \Omega_V^{f_0} \right\rangle_0 \rightarrow 0 \quad (4.16)$$

In particular, the two-time autocorrelation of  $\Omega_V^{f_0}$  is monotonic as also shown in the two panels of Fig.1. Indeed, Eq.(4.15) yields:

$$\frac{d}{dt} \left\langle (\Omega_V^{f_0} \circ S^t) \Omega_V^{f_0} \right\rangle_0 = -K^2 \frac{\sinh Kt}{(1 + \cosh Kt)^2}$$

which is  $\leq 0$  for  $t \geq 0$ .

**4.2. The general case.** In this Subsection we assume  $N \geq 2$  and  $\omega = 0$ , considering the following dynamics:

$$\dot{\theta}_i = \frac{K}{N} \sum_{j=1}^N \sin(\theta_j - \theta_i) = KR \sin(\Phi - \theta_i), \quad i = 1, \dots, N. \quad (4.17)$$

where  $R$  and  $\Phi$  are defined in Eq.(2.2). We are going to prove that the observable  $\left\langle \Omega_V^{f_0} \right\rangle_t$  is a monotonic function of time, and we can estimate the asymptotic value it attains in the large time limit. We start by proving the following result.

**Lemma 4.1.** *For every  $t > 0$ , the time derivative of the expectation of the Dissipation Function obeys:*

$$\frac{d}{dt} \left( \Omega_V^{f_0}(S^t \theta) \right) \geq 0 \quad \text{and} \quad \frac{d}{dt} \left\langle \Omega_V^{f_0} \right\rangle_t = \left\langle (\Omega_V^{f_0} \circ S^t) \Omega_V^{f_0} \right\rangle_0 \geq 0. \quad (4.18)$$

*Proof.* First, we note that by setting  $\mathcal{O} = \Omega_V^{f_0}$  in Eq. (3.18), we find:

$$\frac{d}{dt} \left\langle \Omega_V^{f_0} \right\rangle_t = \left\langle (\Omega_V^{f_0} \circ S^t) \Omega_V^{f_0} \right\rangle_0. \quad (4.19)$$

Moreover, Eq. (3.15) with  $t = 0$  and  $\mathcal{O} = \Omega_V^{f_0}$  yields:

$$\left\langle \Omega_V^{f_0} \right\rangle_t = \left\langle \Omega_V^{f_0} \circ S^t \right\rangle_0. \quad (4.20)$$

Therefore, we can write:

$$\begin{aligned} \frac{d}{dt} \left\langle \Omega_V^{f_0} \circ S^t \right\rangle_0 &= \frac{d}{dt} \int_{\mathcal{M}} \Omega_V^{f_0}(S^t\theta) f_0(\theta) d\theta = \\ &= \int_{\mathcal{M}} \frac{d}{dt} \left( \Omega_V^{f_0}(S^t\theta) \right) f_0(\theta) d\theta = \left\langle \frac{d}{dt} \left( \Omega_V^{f_0}(S^t\theta) \right) \right\rangle_0. \end{aligned} \quad (4.21)$$

Then, using Eq.(2.5) in Ref.[6] we find:

$$\frac{d}{dt} R^2(S^t\theta) = \frac{2K}{N} R^2(S^t\theta) \sum_{j=1}^N \sin^2(S^t\theta_j - \Phi(S^t\theta)) \quad (4.22)$$

where  $S^t\theta_j$  denotes the  $j$ -th element of  $S^t\theta$ , and then

$$\frac{d}{dt} \left( \Omega_V^{f_0}(S^t\theta) \right) = 2K^2 R^2(S^t\theta) \left[ \sum_{j=1}^N \sin^2(S^t\theta_j - \Phi(S^t\theta)) \right] \geq 0 \quad (4.23)$$

for all  $\theta \in \mathcal{M}$ . By integrating over  $\mathcal{M}$  we obtain (4.18). This completes the proof.  $\square$

**Remark 4.3.** *Unlike stationary current autocorrelations, that may fluctuate between positive and negative values, the two times  $\Omega_V^{f_0}$  autocorrelation, computed with respect to the initial probability measure, is non-negative.*

Theorem 2.4 of Ref.[6] shows that non stationary solutions of the system (4.17) converge, as  $t \rightarrow +\infty$ , either to a complete frequency synchronized state  $\Theta^*$ , *i.e.* to a state denoted by  $(N, 0)$ , that takes the form:

$$\Theta^* = (\varphi^*, \dots, \varphi^*) \quad (4.24)$$

in which all phases are equal; or to a state denoted by  $(N - 1, 1)$ , that takes the form:

$$\Theta^\dagger = (\varphi^* + k_1\pi, \varphi^* + k_2\pi, \varphi^* + k_3\pi, \varphi^* + k_4\pi, \dots, \varphi^* + k_N\pi) \quad (4.25)$$

where  $k_i \in \{-1, +1\}$  for a single  $i \in \{1, 2, \dots, N\}$ , and all  $k_j = 0$  with  $j \neq i$ . This can be understood also in terms of the Dissipation Function. In the first place, without loss of generality, let us consider a fixed point  $\bar{\theta}$  of type  $(N - 1, 1)$  whose antipodal is in the  $N$ -component, *i.e.*

$$\bar{\theta} = (\varphi^*, \dots, \varphi^*, (\varphi^* + \pi) \bmod 2\pi) \quad (4.26)$$

for a  $\varphi^* \in [0, 2\pi)$ . Then, the following holds:

**Proposition 4.1.** *The set of initial data such that the solution to (4.17) reaches a stationary  $(N - 1, 1)$ -state for  $t \rightarrow +\infty$  has 0-measure.*

*Proof.* For  $V(\theta)$  as in (2.11), the Jacobian matrix  $A(\theta) \doteq \nabla V(\theta)$  is given by

$$A_{ij} = \begin{cases} \frac{\partial V_j}{\partial \theta_i} = \frac{1}{N} \cos(\theta_i - \theta_j), & i \neq j \\ \frac{\partial V_j}{\partial \theta_j} = -\frac{1}{N} \sum_{k \neq j}^N \cos(\theta_j - \theta_k) \end{cases}$$

For the fixed point  $\bar{\theta}$  set in (4.26) we obtain a symmetric matrix  $\bar{A} = A(\bar{\theta})$  whose entries are

$$\bar{A}_{ij} = \begin{cases} \frac{1}{N} & i \neq j \text{ and } i, j \neq N \\ -\frac{1}{N} & i \neq j \text{ and } i = N \text{ or } j = N \\ -\frac{N-3}{N} & i = j < N \\ \frac{N-1}{N} & i = j = N \end{cases} \quad (4.27)$$

By the symmetry of  $\bar{A}$ , the extremal representation of the eigenvalues  $\{\lambda_k\}_{k=1}^N$  of  $\bar{A}$  are given by the optimization problem:

$$\max_{1 \leq k \leq N} \lambda_k = \max_{\|x\|=1} \{x' \nabla \bar{A} x\}, \quad \min_{1 \leq k \leq N} \lambda_k = \min_{\|x\|=1} \{x' \bar{A} x\}$$

Setting  $x$  to be the standard-basis vectors  $\mathbf{e}_i$ , where  $\mathbf{e}_i$  denotes the vector with a 1 in the  $i$ th coordinate and 0's elsewhere, we see that

$$\min_{1 \leq k \leq N} \lambda_k \leq \min_{1 \leq i \leq N} \{\bar{A}\}_{ii} = -\frac{N-3}{N} < 0, \quad 0 < \frac{N-1}{N} = \max_{1 \leq i \leq N} \{\bar{A}\}_{ii} \leq \max_{1 \leq k \leq N} \lambda_k.$$

Therefore, there exists at least one positive eigenvalue and at least one negative eigenvalue. Indeed, the matrix  $\bar{A}$  has the eigenvalues  $\lambda_- = -(N-2)/N$  with algebraic multiplicity  $N-2$ ,  $\lambda_2 = 0$  and  $\lambda_3 = 1$  with algebraic multiplicity 1. This can be checked considering the proposed subspaces of the center, stable and unstable subspace of the linearized system at  $\bar{\theta}$

$$E^c = \left\{ \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ \vdots \\ 1 \end{bmatrix} \right\}, \quad E^s = \left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \dots, \begin{bmatrix} -1 \\ 0 \\ \vdots \\ 0 \\ 1 \\ 0 \end{bmatrix} \right\} \quad \text{and} \quad E^u = \left\{ \begin{bmatrix} -1 \\ -1 \\ \vdots \\ -1 \\ -1 \\ N-1 \end{bmatrix} \right\}.$$

Then, the Center Manifold Theorem [38, p.116] yields the existence of an  $(N-2)$ -dimensional stable manifold  $W^s(\bar{\theta})$  tangent to the stable subspace  $E^s$ , and the existence of a 1-dimensional unstable manifold  $W^u(\bar{\theta})$ , and 1-dimensional center manifold  $W^c(\bar{\theta})$  tangents to the  $E^u$  and  $E^c$  subspaces respectively. Consequently, the dimension of the center manifold conjoint with the stable manifold is smaller than  $N$ , which implies a null Lebesgue measure in  $\mathbb{R}^n$ .  $\square$

Moreover, we have:




FIGURE 2. Behavior of  $\langle \Omega_V^{f_0} \rangle_t$  (left panel) and  $\langle (\Omega_V^{f_0} \circ S^t) \Omega_V^{f_0} \rangle_0$  (right panel), both rescaled by  $(N - 1)$ , as functions of time, for  $K = 1$ ,  $\omega = 0$  and for different values of  $N$ . The curves on the right panel represent the time derivative of those in the left panel. In particular,  $t = 0$  in the right panel represents  $K^2/N$ , cf. Eq.(4.31).

**Lemma 4.2. (Synchronization):** *Irrespective of the initial condition  $\theta \in \mathcal{T}$ , the Dissipation Function obeys:*

$$\lim_{t \rightarrow \infty} \Omega_V^{f_0}(S^t \theta) = \begin{cases} K(N-1), & \text{for } \theta \neq \Theta^\dagger \\ K(N-1) \left( \frac{N-4}{N} \right) & \text{for } \theta = \Theta^\dagger \end{cases} \quad (4.28)$$

where  $K(N-1)$ , the maximum of  $\Omega_V^{f_0}$  in  $\mathcal{T}^N$ , corresponds to  $(N, 0)$  synchronization.

*Proof.* Because of Theorem 2.4 in Ref.[6] and of the continuity of  $\Omega_V^{f_0}$ , the long time limit of  $\Omega_V^{f_0} \circ S^t$  in the case  $\theta \neq \Theta^\dagger$  is given by  $\Omega_V^{f_0}(\Theta^*)$ . Then, Eq.(2.10) and Eq.(4.4), yield the first line of Eq.(4.28). The case  $\theta = \Theta^\dagger$ , gives, instead:

$$R^* e^{i\varphi^*} = \frac{1}{N} \left( (N-1)e^{i\varphi^*} + e^{i(\varphi^* + \pi)} \right) = \frac{N-2}{N} e^{i\varphi^*}, \quad (4.29)$$

Substituting in Eq.(4.4) we obtain the second line of (4.28).  $\square$

**Remark 4.4.** Equation (4.28) implies that

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \frac{\Omega_V^{f_0}(S^t \theta)}{N} = K. \quad (4.30)$$

In other words, the large  $t$  limit followed by the large  $N$  limit implies that the coupling constant  $K$ , which drives the synchronization process in the Kuramoto dynamics (2.1), equals the average Dissipation per oscillator. For fixed  $N$ , synchronization is also evident from the fact that Eq.(4.23) must converge to 0, for  $\Omega_V^{f_0}$  to become constant.

This also implies  $R^2(S^t \theta) \rightarrow 1$ , as  $t \rightarrow \infty$ . It suffices to consider the definition (4.4) of  $\Omega_V^{f_0}$ . For different values of  $N$ , Fig. 2 illustrates the behavior of  $\langle \Omega_V^{f_0} \rangle_t$  and of its time derivative, which is  $\langle (\Omega_V^{f_0} \circ S^t) \Omega_V^{f_0} \rangle_0$ , as functions of time. The initial growth of

the autocorrelation may look unusual, since autocorrelations are commonly found to decrease. However, unlike standard calculations that rely on an invariant distribution,<sup>1</sup> our autocorrelation is computed with respect to the transient probability measure  $\mu_0$ . The figure portrays the result of numerical simulations. The right panel of Fig. 2, shows that for sufficiently large  $N$  the autocorrelation function  $\langle (\Omega_V^{f_0} \circ S^t) \Omega_V^{f_0} \rangle_0$  reaches a maximum before it decreases, as required for convergence to a steady state. An interesting result is the following.

**Lemma 4.3.** *For  $N \geq 2$ , the derivative of the time dependent average of  $\Omega_V^{f_0}$ , computed at time  $t = 0$  obeys:*

$$\frac{d}{dt} \langle \Omega_V^{f_0} \rangle_t \Big|_{t=0} = \left\langle \left( \Omega_V^{f_0} \right)^2 \right\rangle_0 = K^2 \frac{N-1}{N}. \quad (4.31)$$

Note that the derivative of the mean Dissipation Function equals its autocorrelation function, as expressed by Eq.(4.18). Therefore, Eq.(4.31) gives the value of this autocorrelation function at  $t = 0$ , as shown in the right panel of Fig. 2.

*Proof.* Using (4.20), (4.21) and (4.23) we find that

$$\frac{d}{dt} \langle \Omega_V^{f_0} \rangle_t = \frac{d}{dt} \langle \Omega_V^{f_0} \circ S^t \rangle_0 = 2K^2 \left\langle R^2(S^t \theta) \sum_{j=1}^N \sin^2 \left( S^t \theta_j - \Phi(S^t \theta) \right) \right\rangle_0. \quad (4.32)$$

Thus, at  $t = 0$ , the integrand of (4.32) reads

$$\begin{aligned} R^2(\theta) \sum_{j=1}^N \sin^2(\Phi - \theta_j) &= \frac{1}{N^2} \sum_{j=1}^N \left( \sum_{l=1}^N \sin(\theta_l - \theta_j) \right)^2 \\ &= \frac{1}{N^2} \sum_{j=1}^N \left[ \sum_{l=1}^N \sin^2(\theta_j - \theta_l) + \sum_{l=1}^N \sum_{\substack{k=1 \\ k \neq l}}^N \sin(\theta_l - \theta_j) \sin(\theta_k - \theta_j) \right]. \end{aligned} \quad (4.33)$$

Furthermore, we have:

$$\begin{aligned} \int_0^{2\pi} \int_0^{2\pi} \sin(\theta_l - \theta_j) \sin(\theta_k - \theta_j) d\theta_l d\theta_k \\ = \int_0^{2\pi} \sin(\theta_l - \theta_j) d\theta_l \int_0^{2\pi} \sin(\theta_k - \theta_j) d\theta_k = 0. \end{aligned} \quad (4.34)$$

---

<sup>1</sup>In linear response the initial distribution is considered invariant to first order in the perturbation.

Therefore, considering (4.33) and (4.34) over (4.32) at time  $t = 0$  we have:

$$\begin{aligned}
\left. \frac{d}{dt} \langle \Omega_V^{f_0} \rangle_t \right|_{t=0} &= 2K^2 \int_{\mathcal{M}} R^2(\theta) \sum_{j=1}^N \sin^2(\Phi - \theta_j) f_0(\theta) d\theta \\
&= 2 \frac{K^2}{N^2} \frac{1}{(2\pi)^N} \int_{\mathcal{M}} \sum_{j=1}^N \sum_{l=1}^N \sin^2(\theta_j - \theta_l) d\theta \\
&= 2 \frac{K^2}{(2\pi)^2} \frac{N-1}{N} \int_0^{2\pi} \int_0^{2\pi} \sin^2(\theta_1 - \theta_2) d\theta_1 d\theta_2 \\
&= K^2 \frac{N-1}{N}.
\end{aligned}$$

This completes the proof of (4.31).  $\square$

## 5. COMPARISON WITH LINEAR RESPONSE

In this Section we compare the foregoing exact response formalism with the standard linear response [20]. Consider a perturbed vector field  $V_\varepsilon$ , defined as

$$V_\varepsilon(\theta) = V_0(\theta) + \varepsilon V_p(\theta) \quad (5.1)$$

where the parameter  $\varepsilon$  expresses the strength of the perturbation. Following Section 4, we identify  $\varepsilon$  with  $K$ , and define:

$$V_0(\theta) = \omega \quad (5.2)$$

$$V_{p,j}(\theta) = R \sin(\Phi - \theta_j), \quad j = 1, \dots, N \quad (5.3)$$

Correspondingly, we denote by  $S_\varepsilon^t$  and  $S_0^t$  the perturbed and unperturbed flows, respectively. From Eq. (3.3), we obtain:

$$\Omega_\varepsilon^{f_0} = \Omega_0^{f_0} + \varepsilon \Omega_p^{f_0} = \varepsilon \Omega_p^{f_0} \quad (5.4)$$

where  $\Omega_0^{f_0}$  and  $\Omega_p^{f_0}$  denote the Dissipation Function (4.4) evaluated in terms of the vector fields  $V_0$  and  $V_p$ , respectively. In particular, we have:

$$\Omega_p^{f_0} = \frac{1}{N} \sum_{i,j=1}^N \cos(\theta_j - \theta_i) - 1 \quad (5.5)$$

The last equality in Eq.(5.4) derives from the fact that  $\Omega_0^{f_0} \equiv 0$  if, as assumed,  $f_0$  is invariant under the unperturbed dynamics, cf. Eq.(4.3). We may then write the *exact* response Eq.(3.14) as:

$$\langle \mathcal{O} \rangle_{t,\varepsilon} = \langle \mathcal{O} \rangle_0 + \varepsilon \int_0^t \langle (\mathcal{O} \circ S_\varepsilon^\tau) \Omega_p^{f_0} \rangle_0 d\tau, \quad (5.6)$$

where  $\mathcal{O} \circ S_\varepsilon^t$  denotes the observable  $\mathcal{O}$  composed with the perturbed flow. Because this formula is exact, the parameter  $\varepsilon$  in it does not need to be small, and it appears both

as a factor multiplying the integral and as a subscript indicating the perturbed flow  $S_\varepsilon^t$ . Next, using Eq. (3.8), we can write

$$f_t(\theta) = \exp \left\{ \varepsilon \int_{-t}^0 \Omega_p^{f_0}(S_\varepsilon^\tau \theta) d\tau \right\} f_0(\theta). \quad (5.7)$$

which can be expanded about  $\varepsilon = 0$ , and truncated to first order, to obtain the linear approximation of the evolving probability density:

$$\bar{f}_t(\theta; \varepsilon) = f_0(\theta) \left( 1 + \varepsilon \frac{d}{d\varepsilon} \exp \left\{ \varepsilon \int_{-t}^0 \Omega_p^{f_0}(S_\varepsilon^\tau \theta) d\tau \right\} \Big|_{\varepsilon=0} \right) \quad (5.8)$$

$$\begin{aligned} &= f_0(\theta) \left( 1 + \varepsilon \int_{-t}^0 \Omega_p^{f_0}(S_0^\tau \theta) d\tau \right) \\ &= f_0(\theta) \left( 1 + \varepsilon \int_0^t \Omega_p^{f_0}(S_0^{-\tau} \theta) d\tau \right). \end{aligned} \quad (5.9)$$

Note that the expansion in the variable  $\varepsilon$  of the exponential in Eq.(5.7), requires computing the derivatives with respect to  $\varepsilon$  of the time integral in it. This, in turn, requires the derivatives of the Dissipation Function  $\Omega_p^{f_0}(S_\varepsilon^\tau \theta)$ , and of the evolved trajectory points  $S_\varepsilon^\tau \theta$ . Because both the Dissipation Function and the dynamics are smooth on a compact manifold, their derivatives are bounded, and their integral up to any time  $t$  computed at  $\varepsilon = 0$  is also bounded. Multiplied by  $\varepsilon$ , this integral gives a vanishing contribution to the first derivative of the exponential in Eq.(5.7). There only remain the exponential and the integral computed at  $\varepsilon = 0$ , multiplied by the increment  $\varepsilon$ , which is the brackets in Eq.(5.9). We then define:

$$\overline{\langle \mathcal{O} \rangle}_{t,\varepsilon} = \int_{\mathcal{M}} \mathcal{O}(\theta) \bar{f}_t(\theta; \varepsilon) d\theta = \langle \mathcal{O} \rangle_0 + \varepsilon \int_0^t \left\langle \mathcal{O} \left( \Omega_p^{f_0} \circ S_0^{-\tau} \right) \right\rangle_0 d\tau \quad (5.10)$$

which is the linear response result. At the same time, the invariance of the correlation function under time translations of the unperturbed dynamics, which is proven in Appendix B, yields:

$$\overline{\langle \mathcal{O} \rangle}_{t,\varepsilon} = \langle \mathcal{O} \rangle_0 + \varepsilon \int_0^t \left\langle (\mathcal{O} \circ S_0^\tau) \Omega_p^{f_0} \right\rangle_0 d\tau \quad (5.11)$$

It is interesting to note that, unlike the Green-Kubo formulae, which are obtained from small Hamiltonian perturbations, here the perturbation is not Hamiltonian. Therefore, we may call (5.11) a *generalized* GK formula. It is worth comparing it with the exact response formula (5.6), as follows:

$$\langle \mathcal{O} \rangle_{t,\varepsilon} - \overline{\langle \mathcal{O} \rangle}_{t,\varepsilon} = \varepsilon \int_0^t \left\langle \left[ (\mathcal{O} \circ S_\varepsilon^\tau) - (\mathcal{O} \circ S_0^\tau) \right] \Omega_p^{f_0} \right\rangle_0 d\tau \quad (5.12)$$

which shows that the two formulae tend to be the same, in the small  $\varepsilon$  limit, as expected. Thanks to the use of the Dissipation Function, their difference lies only in the use of the perturbed rather than the unperturbed flow inside  $\mathcal{O}$ .

Let us dwell on the response of two relevant observables, in the case in which  $V_0 = \omega = 0$ , hence  $S_0^t$  is the identity operator, Id. First, taking  $\mathcal{O} = \Omega_\varepsilon^{f_0} = \varepsilon \Omega_p^{f_0}$ , we find

$$\begin{aligned} \langle \Omega_\varepsilon^{f_0} \rangle_{t,\varepsilon} - \overline{\langle \Omega_\varepsilon^{f_0} \rangle}_{t,\varepsilon} &= \int_0^t \left\langle \left[ \left( \Omega_\varepsilon^{f_0} \circ S_\varepsilon^\tau \right) - \left( \Omega_\varepsilon^{f_0} \circ S_0^\tau \right) \right] \Omega_\varepsilon^{f_0} \right\rangle_0 d\tau \\ &= \int_0^t \left[ \left\langle \left( \Omega_\varepsilon^{f_0} \circ S_\varepsilon^\tau \right) \Omega_\varepsilon^{f_0} \right\rangle_0 - \left\langle \left( \Omega_\varepsilon^{f_0} \right)^2 \right\rangle_0 \right] d\tau \end{aligned} \quad (5.13)$$

where we used the identity  $\left( \Omega_\varepsilon^{f_0} \circ S_0^\tau \right) = \Omega_\varepsilon^{f_0}$ , which derives from the fact that  $S_0^t = \text{Id}$ , and which yields, cf. Eq.(4.31):

$$\left\langle \left( \Omega_\varepsilon^{f_0} \right)^2 \right\rangle_0 = \varepsilon^2 \frac{N-1}{N} \quad (5.14)$$

For  $N = 2$ , we can also use the explicit expression (4.15) for the autocorrelation function:

$$\left\langle \left( \Omega_\varepsilon^{f_0} \circ S_\varepsilon^\tau \right) \Omega_\varepsilon^{f_0} \right\rangle_0 = \frac{\varepsilon^2}{1 + \cosh(\varepsilon\tau)} \quad (5.15)$$

which leads to:

$$\langle \Omega_\varepsilon^{f_0} \rangle_{t,\varepsilon} = \varepsilon \tanh\left(\frac{\varepsilon t}{2}\right), \quad \text{and} \quad \overline{\langle \Omega_\varepsilon^{f_0} \rangle}_{t,\varepsilon} = \frac{\varepsilon^2 t}{2} \quad (5.16)$$

so that

$$\langle \Omega_\varepsilon^{f_0} \rangle_{t,\varepsilon} = \overline{\langle \Omega_\varepsilon^{f_0} \rangle}_{t,\varepsilon} + o(\varepsilon^2)t \quad (5.17)$$

In other words, for any  $\varepsilon > 0$ , the difference of the two responses is small at small times, but it diverges linearly as time passes.

As a second instance, let us take  $\mathcal{O} = \psi = \theta_1 - \theta_2$ . From (4.4) and (4.11) we have:

$$\Omega_\varepsilon^{f_0} = 2\varepsilon R^2(\psi) - \varepsilon = \frac{2\varepsilon}{\tan^2\left(\frac{\psi}{2}\right) + 1} - \varepsilon = \varepsilon \cos(\psi) \quad (5.18)$$

Moreover, Eq.(4.10) yields:

$$(\psi \circ S_\varepsilon^t) = 2 \arctan \left[ \tan\left(\frac{\psi}{2}\right) e^{-\varepsilon t} \right] \quad (5.19)$$

and we can write:

$$\begin{aligned} \langle \psi \rangle_{t,\varepsilon} - \overline{\langle \psi \rangle}_{t,\varepsilon} &= \int_0^t \left[ \left\langle (\psi \circ S_\varepsilon^\tau) \Omega_\varepsilon^{f_0} \right\rangle_0 - \left\langle (\psi \circ S_0^\tau) \Omega_\varepsilon^{f_0} \right\rangle_0 \right] d\tau \\ &= \int_0^t \left[ \left\langle (\psi \circ S_\varepsilon^\tau) \Omega_\varepsilon^{f_0} \right\rangle_0 - \left\langle \psi \Omega_\varepsilon^{f_0} \right\rangle_0 \right] d\tau \end{aligned} \quad (5.20)$$

where we used  $S_0^t = \text{Id}$ , which implies  $(\psi \circ S_0^\tau) \equiv \psi$ . Therefore, using (5.18) and (5.19) in (5.20), we obtain:

$$\begin{aligned} \langle \psi \rangle_{t,\varepsilon} - \overline{\langle \psi \rangle}_{t,\varepsilon} &= \frac{1}{(2\pi)^2} \int_0^t \int_{\mathcal{M}} 2 \arctan \left[ \tan\left(\frac{\theta_1 - \theta_2}{2}\right) e^{-\varepsilon\tau} \right] \cos(\theta_1 - \theta_2) d\theta d\tau \\ &\quad - \frac{1}{(2\pi)^2} \int_0^t \int_{\mathcal{M}} (\theta_1 - \theta_2) \cos(\theta_1 - \theta_2) d\theta d\tau = 0 \end{aligned} \quad (5.21)$$

The last equality follows from the fact that the integrands in Eq. (5.21) are odd continuous and periodic functions, that are integrated over a whole period, so that one actually:

$$\langle \psi \rangle_{t,\varepsilon} = \overline{\langle \psi \rangle}_{t,\varepsilon} \equiv 0 \quad , \quad \forall t > 0. \quad (5.22)$$

Clearly, there are observables for which the difference of responses is irrelevant, since they do not evolve in time, and others for which the difference is substantial, even under small perturbations. In any event, the exact response characterizes the synchronization transition, while the linear response does not.

## 6. CONCLUDING REMARKS

We investigated the Kuramoto dynamics for identical oscillators through the statistical mechanics framework of response theory. As a reference (unperturbed) dynamics we took a system of uncoupled oscillators, with statistical properties given by a factorized  $N$ -body distribution with uniform marginal densities. Next, we interpreted the classical Kuramoto mean-field dynamics as a perturbation of the reference one. For any finite number  $N$  of oscillators, we then derived an exact response formula whose validity holds for arbitrarily large perturbations, and we computed, both analytically and numerically, the asymptotic value of the Dissipation Function. The latter is indeed the main ingredient of the exact response theory. Explicit analytical results are given for  $N = 2$ . We also investigated the autocorrelation function of the Dissipation Function, and highlighted its non-monotonic behavior, for sufficiently large  $N$ . Finally, we compared the exact response formalism with the linear response regime. We found that the two responses differ substantially, even for very small perturbations, and that only the exact response describes the transition to synchronized states.

This indicates that the exact response theory, which by definition must be capable of describing even systems undergoing non-equilibrium phase transitions, may actually be used in practice. Synchronization phenomena, which are ubiquitous in Nature, are indeed of that kind.

## ACKNOWLEDGEMENTS

L. R. acknowledges partial support from Ministero dell'Istruzione e Ministero dell'Università e della Ricerca Grant Dipartimenti di Eccellenza 2018-2022 (E11G18000350001).

## APPENDIX A. UNSTABLE FIXED POINTS FOR THE IDENTICAL CASE

In this section we show explicitly the existence of unstable points in any neighborhood of a fixed point of  $(N - 1, 1)$  type.

**Proposition A.1.** *Let  $\bar{\theta}$  be the stationary type  $(N - 1, 1)$  solution set in (4.26) and  $\delta > 0$ . If  $\theta = (\theta_1, \dots, \theta_N)$  satisfy*

$$|\theta_j - \varphi^*| \leq \delta^2, \quad j = [1, \dots, N - 1] \quad (A.1)$$

$$\theta_N = \varphi^* + \pi + \delta \quad (A.2)$$

then there exists a  $\delta_0$  such that for any  $0 < |\delta| < \delta_0$  one has:

$$R^2(\theta) > \left(\frac{N-2}{N}\right)^2 \quad (\text{A.3})$$

and therefore  $R(S^t\theta) \rightarrow 1$  as  $t \rightarrow \infty$ .

*Proof.* From the equation (2.10) we have that

$$\begin{aligned} R^2(\theta) - \left(\frac{N-2}{N}\right)^2 &= \frac{1}{N^2} \sum_{i,j=1}^N \cos(\theta_i - \theta_j) - \sum_{i,j=1}^{N-1} 1 - 2 \sum_{i=1}^{N-1} 1 + 1 \\ &= \frac{1}{N^2} \left[ \underbrace{\sum_{i,j=1}^{N-1} [\cos(\theta_i - \theta_j) - 1]}_{I_1} + 2 \underbrace{\sum_{j=1}^{N-1} [\cos(\theta_N - \theta_j) + 1]}_{I_2} \right] \\ &= \frac{1}{N^2} (I_1 + I_2) \end{aligned}$$

Next, we estimate the lower bounds of  $I_1$  and  $I_2$ . We use the elementary inequality  $\frac{x^2}{4} \leq 1 - \cos(x) \leq \frac{x^2}{2}$ , which is valid for  $|x| \leq x_0$  where  $x_0 \in (0, \frac{5\pi}{6})$ . Then, by using (A.1), for  $I_1$  we get

$$\begin{aligned} I_1 &= \sum_{i,j=1}^{N-1} \cos(\theta_i - \theta_j) - 1 \geq -\frac{1}{2} \sum_{i,j=1}^{N-1} (\theta_i - \theta_j)^2 \\ &= -\frac{1}{2} \sum_{i,j=1}^N [(\theta_i - \varphi^*) + (\varphi^* - \theta_j)]^2 \geq -2\delta^4(N-1)^2 \end{aligned} \quad (\text{A.4})$$

if  $2\delta^2 \leq x_0$ . On the other hand, for  $I_2$  we first observe that for  $1 \leq j \leq N-1$

$$\begin{aligned} |\theta_N - \pi - \theta_j| &\leq |\theta_N - \pi - \varphi^*| + |\varphi^* - \theta_j| \\ &\leq \delta^2 + |\delta| \\ &\leq 2|\delta| \end{aligned}$$

if we take  $|\delta| \leq 1$ .

Then we can use the inequality (A.1) to obtain that

$$\begin{aligned} \cos(\theta_N - \theta_j) + 1 &= 1 - \cos(\theta_N - \pi - \theta_j) \\ &\geq \frac{1}{4}(\delta^2 + |\delta|)^2 = \frac{1}{4}\delta^2(1 + |\delta|)^2 \\ &\geq \frac{\delta^2}{4} \end{aligned}$$

where we consider that  $\delta^2 + |\delta| \leq 2|\delta| \leq x_0$ . Therefore

$$I_2 \geq (N-1) \frac{\delta^2}{2} \quad (\text{A.5})$$

and it follows from the equations (A.4) and (A.5) that

$$\begin{aligned} R^2(\theta) - \left(\frac{N-2}{N}\right)^2 &\geq \frac{1}{N^2} \left\{ \frac{N-1}{2} \delta^2 - 2\delta^4(N-1)^2 \right\} \\ &= \delta^2 \frac{N-1}{N^2} \left[ \frac{1}{2} - 2\delta^2(N-1) \right] \\ &> 0 \end{aligned}$$

for  $\delta^2 < \frac{1}{2(N-1)}$ . In summary, if we choose  $\delta_0 = \min\{1, \frac{x_0}{2}, \frac{1}{2\sqrt{N-1}}\}$ , then (A.3) holds.

Finally, to prove that  $R(S^t\theta) \rightarrow 1$  as  $t \rightarrow +\infty$ , we use the fact that  $t \rightarrow R(S^t\theta)$  is not decreasing and converges to a value  $(N-2k)/N > 0$  for some integer  $k \geq 0$ .

By (A.3) and the monotonicity we deduce that  $R(S^t\theta) > \frac{N-2k}{N}$  for all  $k \geq 1$  and all  $t \geq 0$ , and therefore we conclude that, necessarily, the limiting value has  $k = 0$ . The proof is complete.  $\square$

## APPENDIX B. STATIONARY CORRELATION FUNCTIONS

Given a vector field  $V_0$ , let  $f_0$  be an invariant probability density under the flow  $S_0^t$  generated by  $V_0$ . With the notation set by Eq.(3.7), let  $\Lambda_{0,t}^0$  be the time integral over a trajectory segment, from time 0 to time  $t$ , of the phase space volume variation rate  $\Lambda$ , which is the divergence of the vector field  $V_0$ . Two-time correlation functions between two generic observables  $\mathcal{A}, \mathcal{B} : \mathcal{M} \rightarrow \mathbb{R}$ , evaluated with the density  $f_0$ , are invariant under the time translations determined by  $S_0^t$ . This can be shown as follows. First we note that, proceeding as in Eq. (3.9), one finds

$$\begin{aligned} \Omega_{-t,0}^{f_s,0} &= \int_{-t}^0 \Omega_0^{f_s}(S_0^\tau\theta) d\tau = -\Lambda_{-t,0}^0 - \int_{-t}^0 \frac{d}{d\tau} (\log f_s(S_0^\tau\theta)) d\tau \\ &= -\Lambda_{-t,0}^0 - \log \frac{f_s(\theta)}{f_s(S_0^{-t}\theta)} \end{aligned} \quad (\text{B.1})$$

Upon setting  $s = 0$  in (B.1) and using Eq.(3.12), we find  $\Omega_{-t,0}^{f_0,0} \equiv 0$ , from which we obtain the following useful relation

$$f_0(\theta) = \exp \left\{ -\Lambda_{-t,0}^0(\theta) \right\} f_0(S_0^{-t}\theta) \quad (\text{B.2})$$

where the exponential term is related to the Jacobian determinant of the dynamics as [30]:

$$\left| \frac{\partial (S_0^{-t}\theta)}{\partial \theta} \right| = \exp \left\{ -\Lambda_{-t,0}^0(\theta) \right\} \quad (\text{B.3})$$

Let us look, next, at time correlation functions of the form

$$\langle (\mathcal{A} \circ S_0^{s+\tau}) (\mathcal{B} \circ S_0^t) \rangle_0 = \int_{\mathcal{M}} \mathcal{A}(S_0^{s+\tau}\theta) \mathcal{B}(S_0^t\theta) f_0(\theta) d\theta$$

for any  $s, t, \tau \in \mathbb{R}$ . By a change of variables, one finds

$$\begin{aligned}
\langle (\mathcal{A} \circ S_0^{s+\tau}) (\mathcal{B} \circ S_0^t) \rangle_0 &= \int_{\mathcal{M}} \mathcal{A}(S_0^s \theta) \mathcal{B}(S_0^{t-\tau} \theta) f_0(S_0^{-\tau} \theta) d(S_0^{-\tau} \theta) \\
&= \int_{\mathcal{M}} \mathcal{A}(S_0^s \theta) \mathcal{B}(S_0^{t-\tau} \theta) f_0(S_0^{-\tau} \theta) \left| \frac{\partial (S_0^{-\tau} \theta)}{\partial \theta} \right| d\theta \\
&= \int_{\mathcal{M}} \mathcal{A}(S_0^s \theta) \mathcal{B}(S_0^{t-\tau} \theta) e^{-\Lambda_{-\tau,0}^0} f_0(S_0^{-\tau} \theta) d\theta \\
&= \int_{\mathcal{M}} \mathcal{A}(S_0^s \theta) \mathcal{B}(S_0^{t-\tau} \theta) f_0(\theta) d\theta \\
&= \langle (\mathcal{A} \circ S_0^s) (\mathcal{B} \circ S_0^{t-\tau}) \rangle_0
\end{aligned} \tag{B.4}$$

where we used (B.3) and, in the last line, the formula (B.2).

## REFERENCES

- [1] J. Acebrón, L. Bonilla, C. Pérez, F. Ritort, and R. Spigler. The Kuramoto model: A simple paradigm for synchronization phenomena. *Rev. Mod. Phys.*, 77:137–185, 2005.
- [2] G. S. Agarwal. Fluctuation-Dissipation Theorems for Systems in Non-Thermal Equilibrium and Applications. *Z. Physik*, 252:25–38, 1972.
- [3] A. Arenas, A. Díaz-Guilera, J. Kurths, Y. Moreno, and C. Zhou. Synchronization in complex networks. *Physics Reports*, 469(3):93–153, 2008.
- [4] M. Baiesi, C. Maes, and B. Wynants. Nonequilibrium Linear Response for Markov Dynamics, I: Jump Processes and Overdamped Diffusions. *J. Stat. Phys.*, 137(5):1094, 2009.
- [5] N. Balmforth and R. Sassi. A shocking display of synchrony. *Physica D: Nonlinear Phenomena*, 143(1):21–55, 2000.
- [6] D. Benedetto, E. Caglioti, and U. Montemagno. On the complete phase synchronization for the Kuramoto model in the mean-field limit. *Commun. Math. Sci.*, 13(7):1775–1786, 2015.
- [7] T. Bodineau, B. Derrida, and J. L. Lebowitz. A diffusive system driven by a battery or by a smoothly varying field. *J. Stat. Phys.*, 140:648–675, 2010.
- [8] S. Caruso, C. Giberti, and L. Rondoni. Dissipation Function: Nonequilibrium Physics and Dynamical Systems. *Entropy*, 22:835, 2020.
- [9] Y. Choi, S. Ha, S. Jung, and Y. Kim. Asymptotic formation and orbital stability of phase-locked states for the Kuramoto model. *Physica D: Nonlinear Phenomena*, 241(7):735–754, 2012.
- [10] M. Colangeli and V. Lucarini. Elements of a unified framework for response formulae. *J. Stat. Mech. Theory Exp.*, 2014:P01002, 2014.
- [11] M. Colangeli, C. Maes, and B. Wynants. A meaningful expansion around detailed balance. *J. Phys. A*, 44(9):095001, 13, 2011.
- [12] M. Colangeli and L. Rondoni. Equilibrium, fluctuation relations and transport for irreversible deterministic dynamics. *Physica D: Nonlinear Phenomena*, 241(6):681–691, 2012.
- [13] M. Colangeli, L. Rondoni, and A. Vulpiani. Fluctuation-dissipation relation for chaotic non-Hamiltonian systems. *J. Stat. Mech. Theory Exp.*, 2012:L04002, 2012.
- [14] S. Dal Cengio and L. Rondoni. Broken versus non-broken time reversal symmetry: irreversibility and response. *Symmetry*, 8(8):Art. 73, 20, 2016.
- [15] B. Derrida. Non-equilibrium steady states: fluctuations and large deviations of the density and of the current. *J. Stat. Mech. Theory Exp.*, 2007(7):P07023, 45, 2007.
- [16] H. Dietert and B. Fernandez. The mathematics of asymptotic stability in the Kuramoto model. *Proc. R. Soc. A.*, 474(2220):20180467, 20, 2018.

- [17] J.-G. Dong and X. Xue. Synchronization analysis of Kuramoto oscillators. *Commun. Math. Sci.*, 11(2):465–480, 2013.
- [18] F. Dörfler and F. Bullo. Synchronization in complex networks of phase oscillators: A survey. *Automatica*, 50(6):1539–1564, 2014.
- [19] D.J. Evans, E.G.D. Cohen, and G.P. Morriss. Probability of second law violations in shearing steady flows. *Phys. Rev. Lett.*, 71:2401, 1993.
- [20] D.J. Evans and G. Morriss. *Statistical Mechanics of Nonequilibrium Liquids*. Cambridge University Press, 2008.
- [21] D.J. Evans and D.J. Searles. Equilibrium microstates which generate second law violating steady states. *Phys. Rev. E*, 50:1645–1648, 1994.
- [22] D.J. Evans and D.J. Searles. The Fluctuation Theorem. *Advances in Physics*, 51(7):1529–1585, 2002.
- [23] D.J. Evans, D.J. Searles, and L. Rondoni. Application of the Gallavotti–Cohen fluctuation relation to thermostated steady states near equilibrium. *Phys. Rev. E*, 71:056120, 2005.
- [24] D.J. Evans, D.J. Searles, and S.R. Williams. On the fluctuation theorem for the dissipation function and its connection with response theory. *J. Chem. Phys.*, 128(014504), 2008.
- [25] J. Fell and N. Axmacher. The role of phase synchronization in memory processes. *Nat. Rev. Neurosci.*, 12(2):105–118, 2011.
- [26] G. Gallavotti and E.G.D. Cohen. Dynamical ensembles in stationary states. *J. Statist. Phys.*, 80:931–970, 1995.
- [27] L. Glass. Synchronization and rhythmic processes in physiology. *Nature*, 410(6825):277–284, 2001.
- [28] S. Gupta, A. Campa, and S. Ruffo. *Statistical physics of synchronization*. Springer, 2018.
- [29] S. Ha, D. Ko, J. Park, and X. Zhang. Collective synchronization of classical and quantum oscillators. *EMS Surv. Math. Sci.*, 3(2):209–267, 2016.
- [30] O.G. Jepps and L. Rondoni. A dynamical-systems interpretation of the dissipation function, T-mixing and their relation to thermodynamic relaxation. *J. Phys. A: Math. Theor.*, 49:154002, 2016.
- [31] P. Jiruska, M. De Curtis, J. Jefferys, C. Schevon, S. Schiff, and K. Schindler. Synchronization and desynchronization in epilepsy: controversies and hypotheses. *J. Physiol.*, 591(4):787–797, 2013.
- [32] R. Kubo. The fluctuation-dissipation theorem. *Rep. Prog. Phys.*, 29:255–284, 1966.
- [33] Y. Kuramoto. Self-entrainment of a population of coupled non-linear oscillators. In *International Symposium on Mathematical Problems in Theoretical Physics*, pages 420–422. Springer-Verlag, 1975.
- [34] Y. Kuramoto. *Chemical oscillations, waves, and turbulence*. Springer Series in Synergetics. Springer-Verlag, Berlin, 1984.
- [35] V. Lucarini and M. Colangeli. Beyond the linear fluctuation-dissipation theorem: the role of causality. *J. Stat. Mech. Theory Exp.*, 2012:P05013, 2012.
- [36] U.B.M. Marconi, A. Puglisi, L. Rondoni, and A. Vulpiani. Fluctuation–dissipation: Response theory in statistical physics. *Physics Reports*, 461:111–195, 2008.
- [37] A. Motter, S. Myers, M. Anghel, and T. Nishikawa. Spontaneous synchrony in power-grid networks. *Nature Physics*, 9(3):191–197, 2013.
- [38] L. Perko. *Differential equations and dynamical systems*. Springer-Verlag New York, 2006.
- [39] A. Pikovsky, M. Rosenblum, and J. Kurths. *Synchronization: A Universal Concept in Nonlinear Sciences*. Cambridge University Press, 2001.
- [40] D. Ruelle. General linear response formula in statistical mechanics, and the fluctuation-dissipation theorem far from equilibrium. *Physics Letters A*, 245:220–224, 1998.
- [41] D.J. Searles, L. Rondoni, and D.J. Evans. The steady state fluctuation relation for the dissipation function. *J. Stat. Phys.*, 128(6):1337–1363, 2007.
- [42] W. Singer. Neuronal synchrony: a versatile code review for the definition of relations. *Neuron*, 24(24):49–64, 1999.
- [43] S. Strogatz. From Kuramoto to Crawford: exploring the onset of synchronization in populations of coupled oscillators. *Physica D: Nonlinear Phenomena*, 143(1):1–20, 2000.

DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL'INFORMAZIONE E MATEMATICA (DISIM), UNIVERSITY OF L'AQUILA – L'AQUILA, ITALY

*Email address:* `debora.amadori@univaq.it`

DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL'INFORMAZIONE E MATEMATICA (DISIM), UNIVERSITY OF L'AQUILA – L'AQUILA, ITALY

*Email address:* `matteo.colangeli1@univaq.it`

DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL'INFORMAZIONE E MATEMATICA (DISIM), UNIVERSITY OF L'AQUILA – L'AQUILA, ITALY

*Email address:* `astridherminia.correaluces@graduate.univaq.it`

DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY, AND INFN, SEZIONE DI TORINO, VIA P. GIURIA 1, 10125 TORINO, ITALY, ORCID ID: 0000-0002-4223-6279

*Email address:* `lamberto.rondoni@polito.it`