

Diversity of the Bacterial Community of Three Soils Revealed by Illumina-Miseq Sequencing of 16S rRNA Gene in the South of Brazzaville, Congo

Original

Diversity of the Bacterial Community of Three Soils Revealed by Illumina-Miseq Sequencing of 16S rRNA Gene in the South of Brazzaville, Congo / Mabila, S.T., Goma-Tchimbakala, J., Goma-Tchimbakala, E.J.C.D., Lebonguy, A.A., Banga, A.B.. - In: AMERICAN JOURNAL OF MICROBIOLOGICAL RESEARCH. - ISSN 2328-4137. - STAMPA. - 8:4(2020), pp. 141-149. [10.12691/ajmr-8-4-4]

Availability:

This version is available at: 11583/2953736 since: 2022-01-27T12:25:27Z

Publisher:

Science and Education Publishing

Published

DOI:10.12691/ajmr-8-4-4

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Diversity of the Bacterial Community of Three Soils Revealed by Illumina-Miseq Sequencing of 16S rRNA Gene in the South of Brazzaville, Congo

Shaloom Teresa MABIALA², Joseph GOMA-TCHIMBAKALA^{1,2,*},
Emerance Jessica Claire D'Assise GOMA-TCHIMBAKALA^{2,3},
Augustin Aimé LEBONGUY², Alvychelle Benith BANGA²

¹Ecole Nationale Supérieure d'Agronomie et de Foresterie BP 69 Brazzaville, Congo

²Institut Nationale de Recherche en Sciences Exactes et Naturelles (IRSEN) BP 2400 Brazzaville, Congo

³Politecnico di Torino, Corso Duca Degli Abruzzi 24, Turin, Italy

*Corresponding author: goma.tchimbakala@gmail.com

Received November 10, 2020; Revised December 11, 2020; Accepted December 18, 2020

Abstract The study was conducted on three soils (MFILOU, SNR and ORSTOM) at the south of Brazzaville in Congo. The aim of the work was to study the composition and diversity of the soil microbial community. Microbial diversity was assessed using the Miseq-Sequencing. The results showed that microbial diversity was represented by 1289 OTUs assigned to the Bacteria (1282 OTUs), Fungi (2 OTUs), Viridiplantae (3 OTUs) and metazoa (20TUs) domain with 97% of similarity. The 1282 bacterial OTUs were affiliated with 12 phyla, 31 classes, 59 orders, 122 families, 288 genera et 521 species for all sites. The Proteobacteria (45.59%-29.92%), Firmicutes (27.27%-7.35%), Acidobacteria (16.74%-10.15%), Actinobacteria (12.35%-5.07%) and Nitrospirae are the most abundant common phyla for the three sites. The most dominant common classes were Alphaproteobacteria (38.87%-24.77%), Bacilli (27.20%-6.95%), Acidobacteria (16.47%-7.59%), Actinobacteria (12.35%-5.07%) and Nitrospira (9.81%-1.78%). The most abundant common orders for the 3 sites are represented by Rhizobiales (33.96%-20.38%), Bacillales (27.20%-6.95%), Acidobacteriales (16.49%-7.61%), Actinomycetales (10.20%-3.6%) and Nitrospirales (9.81%-1.78%). Bacillaceae (25.37%-5.60%), Acidobacteriaceae (16.49%-7.61%), Bradyrhizobiaceae (10.81%-4.61%), Nitrospiraceae (9.81%-1.78%) and Chitinophagaceae (4.24%-1.59%) were the best distributed common families in the microbial community of the three sites. Bacillus (25.27%-5.56%), yhe most abundant and common genera were Rhodoplanes (15.48%-5.30%), Bradyrhizobium (10.74%-4.39%), Nitrospira (9.81%-1.78%) and Acidobacterium (8.49%-6.51%). At species level, *Rhodoplanes spp.* (15.26%-5.10%), *Bradyrhizobium spp.* (10.59%-4.35%) and *Acidobacterium spp.* (8.49%-6.51%) were the most common and abundant in the three soils.

Keywords: soil microbial community, microbial diversity, Illumina-Miseq

Cite This Article: Shaloom Teresa MABIALA, Joseph GOMA-TCHIMBAKALA, Emerance Jessica Claire D'Assise GOMA-TCHIMBAKALA, Augustin Aimé LEBONGUY, and Alvychelle Benith BANGA, "Diversity of the Bacterial Community of Three Soils Revealed by Illumina-Miseq Sequencing of 16S rRNA Gene in the South of Brazzaville, Congo." *American Journal of Microbiological Research*, vol. 8, no. 4 (2020): 141-149. doi: 10.12691/ajmr-8-4-4.

1. Introduction

Microbial communities of soil play a key role in ecosystems. Bacteria, fungi and archaea are closely related to nutrient and energy cycling, which support the structure and functional stability of ecosystem [1,2]. Kuczynski et al [3] have reported that composition and diversity of soil microbial communities are influenced by different soil characteristics and vegetation types under land-use change (soil carbon, nutrient depletion and reduced water holding capacity). Soil microbes play important roles in plant growth and health. The higher abundance of beneficial microbes (Bacillus, Agromyces, Micromonospora, Pseudonocardia,

Acremonium, Lysobacter, Mesorhizobium, Microvirga, Bradyrhizobium, Acremonium, Chaetomium) are positively related to the higher soil quality, including better plant growth, lower disease incidence, and higher nutrients contents, soil enzyme activities and soil pH. Several authors found that soil texture represents one of the most important factors influencing the structure of microbial communities as well as, pH, cation exchange capacity, and organic matter content. Soil texture can affect microbial community structure directly by providing a suitable habitat for specific microorganisms which in turn making a maximum degradation process [4]. In other hand some works have found that soil water content is a main driver of soil microbial community structure rather than soil pH [5,6]. Zhang et al., [7]

showed that different vegetation types had different effects on soil bacterial community structure. Wang et al., [8] claimed that diversity and catabolic activity of the bacterial community are greatly impacted by the composition of different vegetation types. Several studies found that the microbial community composition and alpha-diversity varied significantly over time, whereas the change in beta-diversity was relatively small [9,10]. Among soil microbes, bacteria play a pivotal role in crop production, supplying nutrients to crops, stimulating plant growth, controlling the activity of plant pathogens, and improving soil structure [11]. Bacteria are integral components of soil, where their community structure and diversity have been found to be linked to many soil environmental characteristics, such as the physical and chemical properties of the soil [12,13].

In the Congo, apart from the studies by Lebonguy et al [14], no study on the composition and diversity of soil microorganisms in different ecosystems is available. These kinds of studies are needed for improving the assessment of the status of microbial community and diversity of different ecosystems of Congo. Thus, the present study was carried out to improve the knowledge of the diversity and composition of the bacterial community in the rhizospheric soil of different types of legumes. The aim of the study was to know how the characteristics of these soils impact the diversity and composition of the bacterial community under the legumes. For this study, we used Illumina-MiSeq sequencing of PCR amplified 16S rRNA genes from total DNA to assess bacterial diversity, community structure, and the relative abundance of bacterial taxa within three sites from the south of Brazzaville in Congo.

2. Material and Methods

2.1. Study Site and Soil Sampling

The study was conducted in south of Brazzaville in three sites: MFILOU (4°15'50,7600" S, 15°14'6,1512" E), SNR (4°16'41,4944" S, 15°15'5,7341" E) and ORSTOM (4°16'42,1439" S, 15°14'24,6538" E). The climate of Brazzaville is characterized by a rainy season from October to May and a dry and cool season from June to September. The average annual temperature is 26 °C in the rainy season and 20 °C in the dry season. The average annual rainfall is 1200mm. The soil is sandy loam. The vegetation of MFILOU and SNR was characterized by legumes and non-legumes while at ORSTOM, the main species were *Panicum maximum*, *Indigofera hirsuta*, *Mimosa pudica* and *Milletia laurentii*.

The rhizospheric soil samples were collected at five locations in each site. The soil samples were taken in the 0-20 cm horizon using a 5 cm diameter auger. The five soil samples of each site were put together to form one composite soil. The soil was then transported to the laboratory in iceboxes. At the laboratory, the stones, various debris and roots were removed from the soil by sieving with a sieve of 2mm mesh. Each sample was homogenized then 10 g of soils were put in the sterilized tubes and stored at -80°C for DNA extraction. The

remaining soils were used for an analysis of the physicochemical properties of the soil.

2.2. Soil Physicochemical Properties

The soil pH was measured with a « OHAUS STARTER 3000 » pH meter using soil suspension with deionized distilled water (1:25 w/v). Total organic carbon content of soil samples was determined using Walkley-Black method [15]. Total nitrogen was determined using the Kjeldahl method described by Bremner and Mulvaney (1982) [16]. The ammoniacal nitrogen is extracted over 10g of soil after stirring for 30 minutes in 50ml of a 1N KCL solution on Shaker. After decantation and filtration on filter paper, the assay was carried out on a 10 ml aliquot of supernatant supplemented with 1 mL of Nessler's reagent [16]. The phosphorus was determined using the Olsen method [17]. Total iron was determined using DEB method [18]. The magnesium was extracted on 20g of soil with 150 mL of ammonium acetate at pH7. The mixture is stirred for 1 hour then filtered through a filter paper without ash. Then, the soil extract was used to determine spectrophotometrically the magnesium.

2.3. DNA Extraction, illumina-MiSeq Sequencing and Bioinformatic Analyses

DNA extraction, MiSeq sequencing and bioinformatic analyses were performed by Mr DNA laboratories in USA.

Genomic DNA was extracted from 0.5 g of soil samples (dry weight) using the PowerSoil kit (MOBIO Laboratories, Carlsbad, CA, USA). The concentration of the extracted DNA was determined using a Nanodrop 2000C Spectrophotometer (Thermo Scientific, Wilmington, DE, USA). The DNA extract from the soil samples was kept at 80°C until it was used.

The V4 variable region of the bacterial 16S rRNA gene were amplified with primers 515F (5'-GTGCCAGCMGCCGCGGTAA-3') and 806R (5'-GGACTACHVGGTWTCTAAT-3'). The following program was used to conduct the PCR reaction: 3 min of denaturation at 94°C, 30-35 cycles of 30 s at 94°C for amplification, 40 s at 53°C for annealing, 1min at 72°C for elongation, and a final extension at 72°C for 5 min.

After amplification, PCR products are checked in 2% agarose gel to determine the success of amplification and the relative intensity of bands. Multiple samples are pooled together (e.g., 100 samples) in equal proportions based on their molecular weight and DNA concentrations. Pooled samples are purified using calibrated Ampure XP beads. Then the pooled and purified PCR product is used to prepare Illumina DNA library. Sequencing was performed at MR DNA (www.mrdnalab.com, Shallowater, TX, USA) on a MiSeq following the manufacturer's guidelines. Sequence data were processed using MR DNA analysis pipeline (MR DNA, Shallowater, TX, USA). In summary, sequences were joined, depleted of barcodes then sequences <150bp removed, sequences with ambiguous base calls removed. Sequences were denoised, OTUs generated and chimeras removed. Operational taxonomic units (OTUs) were defined by clustering at 3% divergence (97% similarity). Final OTUs were

taxonomically classified using BLASTn against a curated database derived from RDPH and NCBI (www.ncbi.nlm.nih.gov, http://rdp.cme.msu.edu).

2.4. Statistical Analysis

One-way analyses of variance (ANOVA) and relative abundance of phyla and classes were carried out using Excel 2013. For alpha diversity, all analyses were based on the OTU clusters with a cutoff of 3% dissimilarity. Chao1, Shannon and Simpson indexes, equitability J' were calculated using PAST software (v.3.26) to estimate the richness and diversity of the bacterial community of each site. Rarefaction curves with the average number of observed OTUs were generated using PAST software (v.3.01) to compare relative levels of bacterial OTU diversity across the three different sites. The dendrogram of site similarity index was plotted from the Bray-Curtis association matrix. The principal Component Analysis (PCA) and Redundancy analysis (RDA) were performed by CANOCO software (v 4.56) to assess the beta diversity. At the end, Venn diagrams were constructed to visualize shared and unique genera between sites.

3. Results

3.1. Soil Physicochemical Properties

Table 1 shows the results of physicochemical properties. In the three sites sand was the main content of the soil. While Clay was the lower soil component of these three sites. The pH of soil varied from 6.89% in the soil of ORSTOM to 7.21% in the soil of MFILOU. The highest carbon contents (16.2‰), total nitrogen (1.7‰), phosphorus (0.4‰), iron (0.37%), NH₄⁺ (0.8‰) and magnesium (1.0‰) were observed in the soil of ORSTOM site. These chemical properties showed the lowest content in the two other soils (MFILOU and SNR).

Table 1. Soil characteristics

Sites	Clay	Slit	Sand	pH	C	N	P	Fe	NH ₄ ⁺	Mg
	%									
MFILOU	5.50	21.91	67.52	7.21	10.5	1.0	0.3	2.5	0.02	0.5
SNR	4	15.33	78.74	5.01	14.2	1.2	0.2	3.3	0.02	0.7
ORSTOM	5.77	19.54	73.19	6.89	16.2	1.7	0.4	3.7	0.08	1.0

3.2. Bacterial Community Composition

The sequences obtained from 16S rARN sequencing of the soils were grouped with 97% of similarity sequence, in 1289 valid OTUs. Among these OTUs, 99.45% (1282 OTUs) have been assigned to the domain of Bacteria and 0.55% was attributed to Fungi, Metazoa and Viridiplantae. The bacterial OTUs was assigned to 12 phyla, 31 classes, 59 orders, 122 families, 288 genera et 521 species for all sites. ANOVA applied to phyla, classes, orders, families, genera and species does not show any significant difference between soils of the three prospected sites (p>0,05). From the 1282 bacterial OTUs, 960 OTUs were affiliated to MFILOU (with 10463 individuals), 825 OTUs to SNR (with 8062 individuals) and 855 OTUs to ORSTOM (with 11133 individuals). The dendrogram of

similarity (Figure 1) between soils shows two clusters: the first was formed by ORSTOM, SNR and MFILOU were combined in the second cluster. The value of Bray-Curtis similarity between SNR and MFILOU was 0.59. The rarefaction curves (Figure 2) plotted with Shannon index and number of individuals showed that MFILOU has the greatest diversity followed by SNR and ORSTOM. On the other hand ORSTOM had the highest number of individuals than MFILOU and SNR.




Figure 1. Dendrogram UPMGA of the three soils (Bray-Curtis Similarity and OTUs)




Figure 2. Rarefaction curve plotted against Shannon index and individuals




Figure 3. Relative abundance of the dominant phyla in the three sites

3.3. Relative Abundance of Phyla

Among the 12 phyla founded in the present study, only 7 for ORSTOM have a relative abundance $\geq 1\%$ while this number was 9 for MFILOU and SNR; Proteobacteria is the most dominant phylum in all sites (Figure 3).

At MFILOU, the most representative phyla are: Proteobacteria (36.25%), followed by Chloroflexi (19.44%), Acidobacteria (10.15%), Firmicutes (10.13%), Nitrospirae (7.93%), Actinobacteria (5.55%), Bacteroidetes (4.26%), Planctomycetes (3.54%) and Cyanobacteria (1.12%). At SNR, the most dominant phyla are: Proteobacteria (45.59%), Chloroflexi (11.51%), Acidobacteria (10.58%), Nitrospirae (9.81%), Firmicutes (7.35%), Actinobacteria (5.07%), Bacteroidetes (4.10%), Planctomycetes (2.46%) and Verrucomicrobia (2%). At ORSTOM, Proteobacteria (29.92%) is followed by Firmicutes (27.27%), Acidobacteria (16.74%), Actinobacteria (12.35%), Bacteroidetes (6.06%), Verrucomicrobia (3.62%) and Nitrospirae (1.78%). Chao1 indicated that phyla richness were the same in the 3 sites (Table 2). According to Shannon and Simpson indexes the phyla diversity of MFILOU ($H' = 1.89$; $1-D = 0.79$) was higher than in SNR ($H' = 1.79$; $1-D = 0.74$) and ORSTOM ($H' = 1.74$; $1-D = 0.78$). The phyla equitability (J') showed that at MFILOU ($J' = 0.76$) phyla were more equitably distributed than in SNR ($J' = 0.72$) and ORSTOM ($J' = 0.70$).

Table 2. Alpha diversity indexes and richness at phylum level

Alpha diversity index	MFILOU	SNR	ORSTOM
S (number of phylum)	12	12	12
N (number of individuals)	10463	8062	11133
Simpson index (1-D)	0.7975	0.7472	0.7873
Shannon index (H')	1.899	1.793	1.749
Equitability (e ^{H/S})	0.5565	0.5004	0.4792
Equitability (J')	0.7641	0.7214	0.7039
Chao-1	12	12	12

PCA plotted with phyla (Figure 4) showed that Nitrospirae, Proteobacteria and Gemmatimonadetes are abundant at SNR while Cyanobacteria, Deinococcus_thermus, Planctomycetes and Chloroflexi are abundant at MFILOU. On the other hand, Firmicutes, Bacteroidetes, Actinobacteria, Acidobacteria and Verrucomicrobia are abundant at ORSTOM. The two first axis of PCA explained 100% of variations observed between soils at phylum level. Axis 1 explained 87.3% of variation while axis 2 explained 12.7% of variation. By taking into account axis 1, Nitrospirae, Proteobacteria and Gemmatimonadetes are negatively correlated to SNR while Cyanobacteria, Deinococcus_thermus, Planctomycetes and Chloroflexi are negatively correlated to MFILOU. On the other hand, Firmicutes, Bacteroidetes, Actinobacteria, Acidobacteria and Verrucomicrobia are positively correlated to ORSTOM. By taking into account axis 2, Nitrospirae, Proteobacteria and Gemmatimonadetes are negatively correlated to SNR while Actinobacteria, Acidobacteria and Verrucomicrobia are negatively correlated to ORSTOM. Cyanobacteria, Deinococcus_thermus, Planctomycetes, Chloroflexi, Firmicutes and Bacteroidetes are positively correlated to MFILOU.




Figure 4. PCA plotted with the bacterial phyla




Figure 5. Relative abundances of the dominant classes in the three sites

3.4. Relative abundance of Classes

In this study 31 classes were obtained from the 3 sites. At MFILOU 11 classes were retrieved, 14 at SNR and 9 at ORSTOM with a relative abundance $\geq 1\%$ (Figure 5). At MFILOU Alphaproteobacteria (30.73%) was the most dominant class followed by Chloroflexia (16.36%), Bacilli (10%), Acidobacteriia (8.63%), Nitrospira (7.93%), Actinobacteria (5.55%), Planctomycetia (3.49%), Sphingobacteriia (3.47%), Anaerolineae (3.03%), Betaproteobacteria (2.48%), Gammaproteobacteria (2.18%). While at SNR the most dominant class was Alphaproteobacteria (38.87%) followed by Nitrospira (9.81%), Chloroflexia (8.23%), Acidobacteriia (7.59%), Bacilli (6.95%), Actinobacteria (5.07%), Sphingobacteriia (3.87%), Anaerolineae (3.12%), Betaproteobacteria (2.91%), Holophagae (2.86%), Planctomycetia (2.45%), Gammaproteobacteria (2.1%), Verrucomicrobiae (1.77%) and Deltaproteobacteria (1.69%). However, at ORSTOM Bacilli (27.20%) was the most dominant class followed by

Alphaproteobacteria (24.77%), Acidobacteriia (16.47%), Actinobacteria (12.35%), Sphingobacteriia (5.86%), Verrucomicrobiae (3.33%), Betaproteobacteria (2.62%), Gammaproteobacteria (2.11%), Nitrospira (1.78%).

3.5. Relative Abundance of Orders

The 1282 OTUs were grouped into 59 taxonomic orders. Among these orders, 13, 15 and 12 were represented with a relative abundance $\geq 1\%$ respectively at MFILOU, SNR and ORSTOM (Figure 6). At MFILOU, Rhizobiales (22.85%) is the most represented order followed by Chloroflexales (16.25%), Bacillales (10%), Acidobacteriales (8.65%), Nitrospirales (7.93%), Rhodospirillales (4.11%), Actinomycetales (4.10%), Planctomycetales (3.49%), Sphingobacteriales (3.47%), Sphingomonadales (3.45%), Anaerolineales (3.03%), Burkholderiales (1.96%) and Xanthomonadales (1.51%). At SNR, Rhizobiales (33.96%) is the most represented order followed by Nitrospirales (9.81%), Chloroflexales (8.22%), Acidobacteriales (7.61%), Bacillales (6.95%), Sphingobacteriales (3.87%), Rhodospirillales (3.7%), Actinomycetales (3.6%), Anaerolineales (3.12%), Holophagales (2.86%), Planctomycetales (2.45%), Burkholderiales (2.40%), Verrucomicrobiales (1.77%), Xanthomonadales (1.63%) and Sphingomonadales (1%). At ORSTOM Bacillales (27.20%) is the most represented order followed by Rhizobiales (20.38%), Acidobacteriales (16.49%), Actinomycetales (10.20%), Sphingobacteriales (5.86%), Verrucomicrobiales (3.33%), Sphingomonadales (2.85%), Burkholderiales (2.49%), Xanthomonadales (2.03%), Solirubrobacterales (1.96%), Nitrospirales (1.78%) and Rhodospirillales (1.15%).




Figure 6. Relative abundance of the dominant bacterial order in the three sites

3.6. Relative abundance of Families

A total of 122 bacterial families were obtained in the tested soils. However, only 15, 17 and 16 families

respectively at MFILOU, SNR and ORSTOM have a relative abundance $\geq 1\%$ (Figure 7). At MFILOU, Chloroflexaceae (16.09%) is the most represented family followed by Hyphomicrobiaceae (10.56%), Acidobacteriaceae (8.65%), Bacillaceae (8.35%), Nitrospiraceae (7.93%), Bradyrhizobiaceae (4.61%), Rhodospirillaceae (4.05%), Sphingomonadaceae (3.35%), Anaerolineaceae (3.03%), Methylobacteriaceae (2.69%), Rhodobiaceae (2.21%), Sphingobacteriaceae (1.86%), Chitinophagaceae (1.59%), Sinobacteraceae (1.36%) and Rhizobiaceae (1.20%). At SNR, Hyphomicrobiaceae (19.75%) is the most represented family followed by Nitrospiraceae (9.81%), Bradyrhizobiaceae (8.12%), Chloroflexaceae (8.11%), Acidobacteriaceae (7.61%), Bacillaceae (5.60%), Rhodospirillaceae (3.58%), Anaerolineaceae (3.12%), Holophagaceae (2.86%), Sphingobacteriaceae (2.05%), Chitinophagaceae (1.81%), Rhodobiaceae (1.68%), Rhizobiaceae (1.66%), Verrucomicrobiaceae (1.6%), Sinobacteraceae (1.42%), Burkholderiaceae (1.17%) and Sphingomonadaceae (1%). At ORSTOM the most represented family is Bacillaceae (25.37%) followed by Acidobacteriaceae (16.49%), Bradyrhizobiaceae (10.81%), Hyphomicrobiaceae (6.40%), Chitinophagaceae (4.23%), Pseudonocardiaceae (4.20%), Verrucomicrobiaceae (3.26%), Sphingomonadaceae (2.78%), Nitrospiraceae (1.78%), Sphingobacteriaceae (1.63%), Streptomycetaceae (1.4%), Paenibacillaceae (1.30%), Sinobacteraceae (1.29%), Burkholderiaceae (1.16%), Conexibacteraceae (1.14%) and Methylobacteriaceae (1.10%).




Figure 7. Relative abundance of the main family in the three sites

3.7. Relative Abundance of Genera

In this study 288 taxonomic genera have been founded and the most representative of them (with a relative abundance $\geq 1\%$) are presented in Figure 8. The most representative genera are Acidobacterium (8.49%)

followed by *Bacillus* (8.19%), *Nitrospira* (7.93%), *Rhodoplanes* (6.54%), *Bradyrhizobium* (4.39%), *Skermanella* (2.85%), *Microvirga* (2.63%), *Sphingomonas* (2.37%), *Rhodobium* (2.21%), *Hyphomicrobium* (2.10%), *Pirellula* (2.03%), *Sphingobacterium* (1.84%), *Bellilinea* (1.77%), *Longilinea* (1.26%), *Steroidobacter* (1.19%) and *Pedomicrobium* (1.01%) for MFILOU. While at SNR, *Rhodoplanes* (15.48%) is followed by *Nitrospira* (9.81%), *Bradyrhizobium* (7.93%), *Acidobacterium* (6.51%), *Bacillus* (5.56%), *Holophaga* (2.51%), *Hyphomicrobium* (2.30%), *Skermanella* (2.25%), *Bellilinea* (2.18%), *Sphingobacterium* (2.04%), *Rhodobium* (1.68%), *Prostheco bacter* (1.48%), *Pirellula* (1.42%), *Steroidobacter* (1.40%) and *Pedomicrobium* (1.04%). On the other hand, at ORSTOM *Bacillus* (25.27%) is followed by *Bradyrhizobium* (10.74%), *Acidobacterium* (7.10%), *Rhodoplanes* (5.30%), *Terriglobus* (4.39%), *Pseudonocardia* (3.58%), *Flavisolibacter* (3.36%), *Prostheco bacter* (3.17%), *Candidatus koribacter* (3.16%), *Nitrospira* (1.78%), *Sphingobacterium* (1.61%), *Sphingomonas* (1.56%), *Steroidobacter* (1.29%), *Streptomyces* (1.23%), *Edaphobacter* (1.22%), *Conexibacter* (1.14%) et *Oxalophagus* (1.01%).




Figure 8. Relative abundances of the dominant genus in the three sites

3.8. Relative abundance of species

The bacterial OTU are distributed in 521 species and the most abundant are reported in Figure 9. The 5 most dominant species are *Chloroflexus spp.* (16.09%), *Acidobacterium spp.* (8.49%), *Nitrospira spp.* (7.93%), *Rhodoplanes spp.* (6.42%) and *Bradyrhizobium spp.* (4.35%) for MFILOU; *Rhodoplanes spp.* (15.26%), *Nitrospira spp.* (9.81%), *Chloroflexus spp.* (8.11%),

Bradyrhizobium spp. (7.87%) and *Acidobacterium spp.* (6.51%) for SNR; *Bacillus sp.* (11.17%), *Bradyrhizobium spp.* (10.59%), *Acidobacterium spp.* (7.07%), *Rhodoplanes spp.* (5.10%) and *Bacillus spp.* (4.86%) for ORSTOM.




Figure 9. Relative abundance of the 18 most abundant species of the 3 sites

3.9. Alpha and Beta Diversity

Diversity index and richness were also calculated at genera level (Table 3). Chao-1 indicated that the genera richness were highest at MFILOU (280) than SNR (270.4) and ORSTOM (240.8). According to Shannon and Simpson indexes the diversity at MFILOU ($H' = 3.66$, $1-D = 0.942$) and SNR ($H' = 3.61$, $1-D = 0.941$) were similar and higher than ORSTOM ($H' = 3.37$, $1-D = 0.90$). The genera equitability (J') was similar at MFILOU (0.672) and SNR (0.673). Venn diagram (Figure 10) showed that the three soils shared 152 genera. MFILOU and ORSTOM shared 24 genera while SNR and MFILOU 35 genera. On the other hand SNR and ORSTOM shared 17 genera. The number of genera specific to MFILOU, SNR and ORSTOM was respectively 23, 11 and 24.

Table 3. Alpha Diversity index and richness at genera level

Alpha diversity index	MFILOU	SNR	ORSTOM
S (nombre de genres)	234	215	217
N (nombre d'individus)	10463	8062	11133
Simpson index (1-D)	0,9421	0,9413	0,908
Shannon index (H')	3,667	3,618	3,37
Equitability (e ^{H/S})	0,1672	0,1732	0,134
Equitability (J')	0,6722	0,6736	0,6265
Chao-1	281	270,4	240,8




Figure 10. Venn Diagram of genus between the three sites

PCA was plotted with the 10 most abundant genera for all soils (Figure 11). *Bacillus* and *Bradyrhizobium* are abundant at ORSTOM while *Acidobacterium*, *Sphingomonas*, *Nitrospira*, *Hyphomicrobium*, *Rhodobium*, *Skermanella*, and *Microvirga* are abundant at MFILOU; *Rhodoplanes* is abundant at SNR. Axis 1 and axis 2 explained 100% of variations observed between soil at genera level. Axis 1 explained 92.9% of variations while axis 2 explained 7.1% of variations. By taking in account axis 1, *Bacillus* and *Bradyrhizobium* are positively correlated to ORSTOM. *Acidobacterium* and *Sphingomonas* are positively correlated to MFILOU while *Microvirga*, *Nitrospira*, *Hyphomicrobium*, *Rhodobium* and *Skermanella* are negatively correlated to MFILOU. On the other hand, *Rhodoplanes* are negatively correlated to SNR. By taking in account axis 2, *Bradyrhizobium* is negatively correlated to ORSTOM while *Bacillus* is positively correlated to ORSTOM. *Sphingomonas*, *Acidobacterium*, *Microvirga*, *Nitrospira*, *Hyphomicrobium*, *Rhodobium* and *Skermanella* are positively correlated to MFILOU. *Rhodoplanes* is negatively correlated to SNR.




Figure 11. PCA plotted with the 10 most abundant genera of bacterial community

In this study, RDA was used to explore relationships between environmental factors and soil microbial community structure. RDA was plotted with the 10 most abundant genera and the physicochemical characteristics of soil. According to the RDA analysis the 12 environmental parameters (Table 3) accounted for 100%

of the variation in the observed bacterial community structure. Axis 1 explained 92.9% of this variation and axis 2 explained the other 7.1% (Figure 12). We found that sand and kind of nitrogen components, and carbon were the driver of bacterial communities respectively at SNR for *Rhodoplanes* and ORSTOM for *Bradyrhizobium* and *Bacilli* while clay, pH and silt impacted the distribution of bacteria at MFILOU for *Acidobacterium*, *Sphingomonas*, *Microvirga*, *Rhodobium*, *Skermanella*, *Nitrospira*.




Figure 12. RDA plotted with the 10 most abundant genera and physicochemical properties of soils

4. Discussion

Soil is an extremely rich environment which harbors a multitude of microorganisms, among which bacteria are the most abundant in term of biomass and taxonomic diversity. Bacteria are the most plentiful and diverse group of microorganisms in the soil, and by number of species found per gram of soil [19]. These assertions agreed with our result in the fact that 99,45% of OTU found were attributed to bacteria. For the three sites prospected, at phylum level, all the bacteria were related to Proteobacteria, Chloroflexi, Actinobacteria, Acidobacteria, Planctomycetes, Nitrospira, Verrucomicrobia, Bacteroidetes, Gemmatimonadetes, Firmicutes, Deinococcus_thermus, Cyanobacteria. We found that Proteobacteria was the most dominant phylum in all sites (45,59% - 29,92%). Alami et al. [20] found in their study on the rhizospheric soil of *Coptis chinensis* that Proteobacteria, Actinobacteria, Chloroflexi, Acidobacteria, Bacteroidetes, Gemmatimonadetes were the dominant bacterial phyla. Among these phyla, Proteobacteria was the most dominant phylum as in our study. This kind of result was also found by Sun et al [21] in a tropical forest soil in China, where they found that Proteobacteria, Planctomycetes, Nitrospirae, Verrucomicrobia, Bacteroidetes, Chloroflexi, Acidobacteria and Actinobacteria constituted the major phyla. This assertion was assumed by several authors in their works, Wang et al [8] on the soil covered by halophytic vegetation in China, Zang et al

[22] in soil of mixed bamboo and broad-leaved forest in China, Wang et al [23] in a Mineral Sandy loam Soil with a lower pH in the Desert of Maine (USA), Goma-Tchimbakala and Lebonguy [14] in Sandy-Loam Soil Polluted by hydrocarbons in Congo. The present study and all the previous cited above have shown that the bacterial community of soils at phylum level is similar although belonging to different environments and having different characteristics.

Several studies found that the microbial community composition and alpha-diversity varied significantly over time, whereas the change in beta-diversity was relatively small [9] and [10]. In this study, alpha diversity represented by richness (S observed and Chao1) and diversity index (Shannon and Simpson indexes, equitability) declined from the cultivated soils of MFILOU and SNR to the soil of ORSTOM that was exempt of culture. This result is in accordance with several other studies in which it was demonstrated that continuous cultivation resulted in decrease of bacterial diversity [20]. According to Alami et al. [20] and Kennedy et al. [24], microbial richness and diversity have a crucial role in soil quality, health and ecosystem sustainability. These authors claimed that the reduction of soil microbial richness and diversity could contribute to alter plant performance and insufficient resistance against disease and pest in the continuous cultures. Hamarashid et al [25] have shown that bacterial population in loamy sand was higher than in sandy loam soil. In the present study, the result showed that in ORSTOM and MFILOU in which soils were sandy loam bacterial number was higher than in loamy sand soil of SNR. The dendrogram (UPMGA) has shown two clusters: the first formed by ORSTOM and the second by SNR and MFILOU. This result can be explained by the fact that SNR and MFILOU are agricultural soil therefore subject to several practices such as green manure amendment for soil enrichment while ORSTOM is a forest and was not subjected to any agricultural practices. Many previous studies showed that soil microbial communities structure and diversity are affected by many factors, including plant species, soil types, organic breeding, and agricultural management (e.g., fertilizer application, periodic outflow, and crop rotation) [19,26,27,28]. Venn diagram and the rarefaction curves plotted against Shannon index have shown that bacterial community of SNR and MFILOU is more diverse than of ORSTOM but bacterial community of ORSTOM is greater in term of number of bacteria than SNR and MFILOU. The great diversity and low number of bacteria of MFILOU and SNR can be explained by agricultural practices. The choice of cultivation method is essential to increase crop yield and suppress disease, and it may cause changes in microbial communities as different plant species secrete different types of root exudates, which can alter the structure of the soil bacterial community [29]. On the other hand, Chen et al [30] have shown that the microbial biomass in the soil with legumes is higher than soil with grass. At ORSTOM around soil sampling point (within a radius of 1m) the vegetation were composed by *Indigofera hirsuta* which is a spontaneous leguminous plant while at SNR and MFILOU vegetation were mixt comprising legumes and non-legume plant.

In this study, RDA showed that soil pH, clay and silt have an influence on distribution of Acidobacterium, Sphingomonas, Microvirga, Rhodobium, Nitrospirae that found to be most related with soil of MFILOU while in soil of ORSTOM the main components that impacted bacteria distribution of Bradyrhizobium and Bacillus were phosphorus, total nitrogen, ammonium, and magnesium. Soil pH has been demonstrated to be one of the main factors determining soil microbial composition and diversity and has also been found to be correlate with soil bacterial phylogenetic diversity, with peak diversity near neutral pH [12]. Some other studies showed that soil pH was closely correlated with bacterial diversity and soil microbial communities [31,32,33]. At SNR, bacterial distribution of Rhodoplanes was greatly related with the sand content. The findings of our study were in accordance with Lauber et al. [12] and Fierer et al [13] they claimed that bacteria are integral components of soil, where their community structure and diversity have been found to be linked to many soil environmental characteristics, such as the physical and chemical properties of the soil.

5. Conclusion

In conclusion, in this study we found that Alphaproteobacteria, Bacilli, Acidobacteria, Actinobacteria and Nitrospira were the main phyla in the three sites. Our finding revealed that bacterial richness and diversity were relatively high in agricultural soil (MFILOU AND SNR) than in soil under ORSTOM. We found that sand and kind of nitrogen components, and carbon were the driver of bacterial communities respectively at SNR and ORSTOM while clay, pH and silt impacted the distribution of bacteria at MFILOU.

Acknowledgments

This research was funded by the corresponding author of this article. The National Institute for Research in Exact and Natural Sciences provided technical support.

Conflict of Interest

The authors declare no conflict of interest in this work.

References

- [1] Van der Heijden M. G. A., Bardgett R. D. and Van Straalen N. M., 2008. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems, *Ecol. Lett.*, vol. 11, pp. 296-310.
- [2] Zak D. R., Holmes W. E., White D. C., Peacock A. D. and Tilman D., 2003. Plant diversity, soil microbial communities, and ecosystem function: are there any links? *Ecology*, vol. 84, pp. 2042-2050.
- [3] Kuczynski J., Liu Z. Z., Lozupone C., McDonald D., Fierer N. and Knight R., 2010. Microbial community resemblance methods differ in their ability to detect biologically relevant patterns, *Nat. Methods*, vol. 7, pp. 813-817.

- [4] Girvan M. S., Bullimore J., Pretty J. N., Osborn A. M. and Ball A. S., 2003. Soil type is the primary determinant of the composition of the total and active bacterial communities in arable soils, *Appl. Environ. Microbiol.*, vol. 69, pp. 1800-1809.
- [5] Bft B., Prescott C. E. and Grayston S. J., 2012. Soil moisture is the major factor influencing microbial community structure and enzyme activities across seven biogeoclimatic zones in western Canada, *Soil Biology and Biochemistry*, vol. 44, pp. 9-20.
- [6] Liu J., He X. X., Lin X. R., Chen W. C., Zhou Q. X., Shu W. S. and Huan L. N., 2015. Ecological effects of combined pollution associated with e-waste recycling on the composition and diversity of soil microbial communities, *Environmental Science and Technology*, vol. 49, pp. 6438-6447.
- [7] Zhang K., Cheng X., Shu X., Liu Y. and Zhang Q., 2018. Linking soil bacterial and fungal communities to vegetation succession following agricultural abandonment, *Plant and Soil*, vol. 431, pp. 19-36.
- [8] Wang L., Wang J., Guo D. and Jiang A., 2020. Catabolic Activity and Structural Diversity of Bacterial Community in Soil Covered by Halophytic Vegetation., *Current Microbiology*, vol. 1 (2).
- [9] Fierer N. and Jackson R. B., 2006. The diversity and biogeography of soil bacterial communities, *Proc. Natl. Acad. Sci. U. S. A.*, vol. 103, pp. 626-631.
- [10] Wallenstein M. D., McMahon S. and Schimel J., 2007. Bacterial and fungal community structure in Arctic tundra tussock and shrub soils, *FEMS Microbiol. Ecol.*, vol. 59, pp. 428-435.
- [11] Hayat R., Ali S., Amara U., Khalid R. and Ahmed I., 2010. Soil beneficial bacteria and their role in plant growth promotion: a review, *Ann Microbiol*, vol. 60, pp. 579-598.
- [12] Lauber C. L., Hamady M., Knight R. and Fierer N., 2009. Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Appl Environ Microbiol.*, vol. 75, pp. 5111-5120.
- [13] Fierer N., Lauber C. L., Ramirez K. S., Zaneveld J., Bradford M. A. and R. Knight, 2012. Comparative metagenomic, phylogenetic and physiological analyses of soil microbial communities across nitrogen gradients., *ISME J.*, vol. 6, pp. 1007-1017.
- [14] Goma-Tchimbakala J. and Lebonguy A. A., 2020. Assessment of Bacterial Diversity of Sandy-Loam Soil Polluted by Hydrocarbons Using 454 Pyrosequencing, *American Journal of Microbiological Research*, vol. 8(11), pp. 34-41.
- [15] Nelson D. W. and Sommers L. E., 1982. Total carbon, organic carbon and organic matter, in *Methods of soil analysis*, Part 2 American Society of Agronomy, Madison, 1982, pp. 539-579.
- [16] Bremner J. M. and Mulvaney C. S., 1982. Nitrogen-Total,» chez *Methods of soil analysis, Chemical and Microbiological properties*, American Society of Agronomy, Soil Science Society of Americana, Madison, 1982, pp. 595-624.
- [17] Murphey J. and Riley PP., 1962. A modified single solution method for the determination of phosphate in natural waters. *Anal. Chimim. Acta.*, vol. 27, pp. 31-36.
- [18] Segalen PP., 1971. La détermination du fer libre dans les sols à sesquioxides,» *Cah. ORSTOM, sér. Pédol.*, vol. 9(11), pp. 3-27.
- [19] Zhao J., Zhang R., Xue C., Xun W., Sun L., Xu Y. and Shen Q., 2014. Pyrosequencing Reveals Contrasting Soil Bacterial Diversity and Community Structure of Two Main Winter Wheat Cropping Systems in China. *Microb. Ecol.*, vol. 67, pp. 443-453.
- [20] Alami M. M., Xue J., Ma Y., Zhu D., Gong Z., Shu S. and Wang X., 2020. Structure, Diversity, and Composition of Bacterial Communities in Rhizospheric Soil of *Coptis chinensis* Franch under Continuously Cropped Fields, *Diversity*, vol. 1(212), 57.
- [21] Sun Y., Luo C., Jiang L., Song M., Zhang D., Li J., Li Y., Ostle N. J. and G. Zhang, 2020. Land-use changes alter soil bacterial composition and diversity in tropical forest soil in China,» *Science of the Total Environment*, vol. 712. 136526.
- [22] Zhang M. M., Fan S. H., Guan F. Y., Yan X. R. and Yin Z. X., 2020. Soil bacterial community structure of mixed bamboo and broad-leaved, *Scientific reports, nature research*, vol. 1(210). 6522.
- [23] Wang Y., Osman J. R. and DuBow M. S., 2020. Bacterial Communities on the Surface of the Mineral Sandy Soil from the Desert of Maine (USA),» vol. 1(2).
- [24] Kennedy A. C. and Smith K. L., 1995. Soil microbial diversity and the sustainability of agriculture soils,» *Plant Soil*, vol. 170, pp. 75-86.
- [25] Hamarashid N. H., Othman M. A. and Hussain M. A. H., 2010. Effects of soil texture on chemical compositions, microbial populations and carbon mineralization in soil, *Egypt. J. Expp. Biol. (Bot.)*, vol. 6(11), pp. 59-64.
- [26] Klein E., Katan J. and Gamliel A., 2016. Soil suppressiveness by organic amendment to Fusarium disease in cucumber: Effect on pathogen and host., *Phytoparasitica*, vol. 44, pp. 239-249.
- [27] Qiu M., Zhang R., Xue C., Zhang S., Li S., Zhang N. and Shen Q., 2012. Application of bio-organic fertilizer can control Fusarium wilt of cucumber plants by regulating microbial community of rhizosphere soil, *Biol. Fertil. Soils*, vol. 48, pp. 807-816.
- [28] Sun B., Dong Z. X., Zhang X. X., Li Y., Cao H. and Cui Z. L., 2011. Rice to Vegetables: Short- Versus Long-Term Impact of Land-Use Change on the Indigenous Soil Microbial Community., *Microb. Ecol.*, vol. 1(262), pp. 474-485.
- [29] Xuan D. T., Guong V. T., Rosling A., Alström S., Chai B. and Högberg N., 2012. Different crop rotation systems as drivers of change in soil bacterial community structure and yield of rice, *Oryza sativa.*, *Biol. Fertil. Soils*, vol. 48, pp. 217-225.
- [30] Chen M., Chen B. and Marschner PP., 2008. Plant growth and soil microbial community structure of legumes and grasses grown in monoculture or mixture,» *Journal of Environmental Sciences*, vol. 20, pp. 1231-1237.
- [31] Zhang Y., Li Y., Wang L., Tang Y., Chen J., Hu Y., Fu X. and Le Y., 2013. Soil microbiological variability under different successional stages of the Chongming Dongtan wetland and its effect on soil organic carbon storage, *Ecological Engineering*, vol. 52, pp. 308-315.
- [32] Cheng J., Jing G., Wei L. and Jing Z., 2016. Long-term grazing exclusion effects on vegetation characteristics, soil properties and bacterial communities in the semi-arid grasslands of China, *Ecological Engineering*, vol. 97, pp. 170-178.
- [33] Xue L., Ren H., Li S., Leng X. and Yao X., 2017. Soil Bacterial Community Structure and Co-occurrence Pattern during Vegetation Restoration in Karst Rocky Desertification Area,» *Front Microbiol*, vol. 1(28), 2377.

