

Impact of COVID-19 on emergency department visits among palliative home care recipients: a retrospective population-based cohort study in the Piedmont region, Italy

Original

Impact of COVID-19 on emergency department visits among palliative home care recipients: a retrospective population-based cohort study in the Piedmont region, Italy / Scacchi, A., Conti, A., Politano, G., Dalmasso, M., Ostellino, S., Gianino, M.M.. - In: PALLIATIVE CARE AND SOCIAL PRACTICE. - ISSN 2632-3524. - 16:(2022), pp. 1-12. [10.1177/26323524221136880]

Availability:

This version is available at: 11583/2974358 since: 2023-01-05T11:36:19Z

Publisher:

SAGE Publications Ltd

Published

DOI:10.1177/26323524221136880

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Impact of COVID-19 on emergency department visits among palliative home care recipients: a retrospective population-based cohort study in the Piedmont region, Italy

Alessandro Scacchi, Alessio Conti^{ID}, Gianfranco Politano^{ID}, Marco Dalmasso, Sofia Ostellino^{ID} and Maria Michela Gianino^{ID}

Palliative Care & Social Practice

2022, Vol. 16: 1–12

DOI: 10.1177/

26323524221136880

© The Author(s), 2022.
Article reuse guidelines:
sagepub.com/journals-
permissions

Abstract

Background: Integrated palliative home care (IHPC) is delivered to patients with progressive end-stage diseases. During the COVID-19 pandemic, IHPC needed to provide high-quality home care services for patients who were treated at home, with the goal of avoiding unnecessary care, hospital admissions, and emergency department (ED) visits. This study aimed to compare the ED visits of IHPC recipients in a large Italian region before and during the first two waves of the COVID-19 pandemic and to find sociodemographic or clinical characteristics associated with changes in ED visits during the first two waves of COVID-19 pandemic, compared with the period before.

Methods: Administrative databases were used to identify sociodemographic and clinical variables of IHPC recipients admitted before and during the pandemic. The obtained data were balanced by applying a propensity score. The average number of ED visits before and during the pandemic was calculated by using the Welch's *t* test and stratified by all the variables.

Results: Before and during the pandemic, 5155 and 3177 recipients were admitted to IHPC, respectively. These individuals were primarily affected by neoplasms. ED visits of IHPC recipients reduced from 1346 to 467 before and during the pandemic, respectively. A reduced mortality among IHPC patients who had at least one ED visit during the pandemic (8% during the pandemic *versus* 15% before the pandemic) was found. The average number of ED visits decreased during the pandemic [0.143, confidence interval (CI) = (0.128–0.158) *versus* 0.264, CI = (0.242–0.286) before the pandemic; $p < 0.001$] for all ages and IHPC duration classes. The presence of a formal caregiver led to a significant decrease in ED use. Medium and high emergency ED admissions showed no difference, whereas a decrease in low-level emergency ED admissions during the pandemic [1.27, CI = (1.194–1.345) *versus* 1.439, CI = (1.3–1.579) before the pandemic; $p = 0.036$] was found.

Conclusion: ED visits among IHPC recipients were significantly decreased during the first two waves of the COVID-19 pandemic, especially in those individuals characterized by a low level of emergency. This did not result in an increase in mortality among IHPC recipients. These findings could inform the reorganization of home care services after the pandemic.

Keywords: COVID-19, emergency department visits, integrated home care, palliative care, retrospective cohort study

Received: 16 May 2022; revised manuscript accepted: 14 October 2022.

Correspondence to:
Alessio Conti
Department of Public Health and Paediatric Sciences, University of Turin, Via Santena, 5 bis, Turin 10126, Italy.
alessio.conti@unito.it
Alessandro Scacchi
Maria Michela Gianino
Department of Public Health and Paediatric Sciences, University of Turin, Turin, Italy
Gianfranco Politano
Sofia Ostellino
Department of Control and Computer Engineering, Politecnico di Torino, Torino, Italy
Marco Dalmasso
Epidemiology Unit, Local Health Unit T03, Grugliasco, Italy

Introduction

Palliative care has been developed in recent decades as a social movement and a medical specialty that addresses the physical, psychosocial, and spiritual needs of those individuals with life-threatening illnesses and their families.¹ In Italy, palliative care was officially recognized by the National Law 38/2010,² which guarantees the right of access to patients with progressive end-stage diseases, such as neoplasms, heart failure, chronic obstructive pulmonary disease, and dementia.³ This type of care is provided in each of the 20 regions of Italy in multiple settings, including hospices, hospitals, residential facilities, and at home, wherein it is referred to as integrated palliative home care (IHPC).

In Italy, IHPC is initiated following the request of the general practitioner, and it is delivered and administered by palliative care units, which create multiprofessional teams that ensure medical, nursing, rehabilitation, social and psychological support, pharmaceutical assistance, and diagnostic tests. The multiprofessional teams evaluate whether a recipient can receive IHPC based on multiprofessional assessment measures, and they then prepare an individual care plan. This plan must be shared with the recipient and their family and caregivers, as it constitutes a therapeutic care contract. The individual care plan aims to identify the goals of care and the most appropriate interventions in the case of any issue, as well as to guarantee a comprehensive, coordinated approach across different health care professionals, thus avoiding unnecessary care, hospital admissions, and emergency department (ED) visits.

Similar to any other health care system setting, the palliative home care program has also been influenced by the COVID-19 pandemic, albeit lacking evidence in the international literature. Indeed, an Italian study⁴ highlighted that some research focused on the crucial role of palliative care in managing individuals who were seriously infected during the COVID-19 pandemic,⁵⁻⁷ but few studies have investigated the impact of the pandemic on palliative care services.^{4,8,9} As it was clear from the early stages of the pandemic that hospital care services could quickly become a critical resource, it was also necessary to maintain high-quality home care services (whenever possible) for recipients who were treated at home during such period. For this reason, home care workflows were necessarily replanned in terms of

the readaptation of processes, flexibility, and precautions with the recipients.¹⁰

From this perspective, it could be valuable to evaluate whether IHPC recipients had reduced ED visits in 2020 during the first two waves of the COVID-19 pandemic, compared with 2019. This is due to the fact that Italy was the first European country to be seriously affected by COVID-19, with an exponential increase in infections and deaths. In February 2020, a few positive cases of COVID-19 were reported in northern Italy; by the beginning of March 2020, the virus had spread to all Italian regions, with particularly severe infection incidence rates in northern regions,¹¹ as Piedmont. The Piedmont region is the second largest of Italy's 20 regions, covering an area of 25,300 km² in the north-west of the country, and the seventh most populated (4.3 million inhabitants). IHPC recipients in Piedmont receive the same care provision in each of the 20 regions of Italy, as the service and its organization are designed on a national basis. As far as we know, ED visits among IHPC recipients have been understudied in Italy. It might be possible that, as reported for the general population, ED visits also have reduced for IHPC recipients during the early stages of the COVID-19 pandemic. Furthermore, sociodemographic and clinical characteristics of IHPC recipients could have been associated with a reduction in ED visits in this particular population. Therefore, the primary aim of this study was to compare the ED visits of IHPC recipients in a large northern region of Italy before and during the first two waves of the COVID-19 pandemic, analyzing if there has been a reduction of these visits during the first phase of the pandemic. Another objective of this study was to find any sociodemographic or clinical characteristics associated with a change in ED visits during the first two waves of the COVID-19 pandemic, compared with the period before.

Methods

We collected the data of all the individuals who were admitted to the IHPC program of Piedmont from 1 March 2020 to 31 October 2020 (during the first two waves of the pandemic), which encompassed the peak months of the COVID-19 pandemic in Italy. After the first outbreak and consequent decrease in the infection rate from May 2020 and during the summer season, Italy experienced a sudden increase in cases and hospitalizations during September and October 2020.¹²

Comparable data were also examined from the same time interval in 2019, from 1 March to 31 October (before the pandemic).

Data concerning the recipients and ED use were collected by merging two different sources: the Italian Official National Information System for Monitoring Palliative Home Care Services (the SIAD database) and the Italian National Information System for ED use database.

Data were collected for the following sociodemographic variables: sex, age, and presence of a formal caregiver. Age was stratified into four age groups (≤ 65 , 65–80, 81–90, and ≥ 91 years of age). The following clinical variables were collected: outcome (alive or dead), principal diagnosis at IHPC admission (as defined by the International Classification of Diseases, Ninth Revision), triage assessment, and principal diagnosis at the time of ED admission. Moreover, the IHPC duration in days (≤ 30 , 31–90, and > 90 days) and the arrival mode to the ED (autonomous, ambulance, or other modes) were collected.

The Italian triage assessment¹³ is based on a four-color code evaluation with the goal of a rapid determination of safe and comprehensible treatment priorities. Specifically, white defines non-critically ill patients who do not need treatment in a short time; green defines minimally critically ill patients whose interventions can be postponed; yellow defines moderately critically ill patients whose interventions cannot be postponed; and red defines highly critically ill patients who need immediate emergency interventions. White and green indicate a low level of emergency, whereas yellow and red indicate a medium and high level of emergency, respectively.

Moreover, the principal symptoms at the time of the ED visit were recorded during each triage examination per recipient. For the purposes of this study, eight different types of the most recurrent symptoms that are included on the Italian National Information System for ED use were categorized as follows: abdominal pain, dyspnea, traumatic problems, fever, cardiac rhythm alteration, urological symptoms, neurological symptoms, and undefined symptoms.

All the data originated from the Health Information System of the Piedmont region, which was designed and administered to comply with the regional, national, and European

regulations regarding the protection of personal data; in addition, it is used in support of the evaluation and monitoring of activities of the Piedmont region.

Data from these databases were merged by using the universal patient ID number, which is an anonymous, unique code that is centrally assigned to each patient. The data treatment phases have been exclusively performed by delegates of the regional epidemiology network (of which the Unit of Epidemiology-Regional Health Service-Local Health Unit TO3 is a node) who are in charge of the processing data for supporting regional evaluations stated in the regional regulation.¹⁴ A formal agreement regulates the cooperation between the units that were involved in this study. Therefore, ethics committee approval was unnecessary.

Statistical analysis

To guarantee that no confounding effect could arise because of age and sex imbalances between the two time periods, a propensity score (PS) balancing method was applied to preprocess the data. The PS method has been implemented in R by using the *WeighIt* package¹⁵ to create equivalent groups in the observations with entropy balancing,¹⁶ thus better balancing the covariates and minimizing the overdispersion of the weights that had been computed thus far.

After balancing the population, we resorted to an independent sample test for the two groups (the Welch's *t* test) to compare 2019 *versus* 2020 populations. The Welch's *t* test is a two-sample test that is used to test the hypothesis that two populations have equal means, and it is more reliable than the Student's *t* test when the two samples have unequal variances and/or unequal sample sizes.¹⁷

The test is also supported by the Bayes factor computation. Specifically, this method can make more robust assumptions by helping to index evidence in favor of the null hypothesis if it cannot be rejected with the standard null hypothesis significance test approach.

The average numbers of ED visits before and during the pandemic periods were calculated. Analyses were stratified by all the variables.

All the analyses were performed by using the R package version 3.6.1 (R Foundation for

Statistical Computing, Vienna, Austria). Statistical significance was set at $p < 0.05$ and 95% confidence intervals (CIs).

Results

IHPC recipients' characteristics and ED visits

A total of 5155 recipients were admitted to IHPC in 2019, and 3177 individuals were admitted in 2020. The characteristics of these two populations are shown in Table 1. The most noticeable differences among the groups were related to the IHPC duration, with 17.23% of the 2019 recipients compared with 11.61% of the 2020 recipients having a long (>90 days) IHPC duration, as well as the number of deaths (28.4% in 2019 *versus* 35.76% in 2020). For both groups, the principal diagnosis at IHPC admission was neoplasm (81.53% in 2019 and 84.95% in 2020).

The total numbers of ED visits of the IHPC recipients were 1346 in 2019 and 467 in 2020. A considerable number (20.4%; $n = 1054$) of IHPC recipients had at least one ED visit in 2019, whereas this number decreased to 12.1% ($n = 383$) in 2020. In particular, 15.39% of the IHPC patients who died in 2019 and 8.77% of those who died in 2020 had at least one ED visit.

ED visits before versus during the COVID-19 pandemic

The average number of ED visits among the IHPC recipients was reduced in 2020 compared with 2019 [0.264, CI = (0.242–0.286) in 2019 and 0.143, CI = (0.128–0.158) in 2020; $p < 0.001$].

The observed reduction in the average number of ED visits was affected by several variables (Table 2).

The average number of ED visits was reduced in 2020 for female and male recipients, for all age classes (except for those individuals aged >91 years), and for all IHPC duration classes. In addition, the presence of a formal caregiver led to a significant decrease in ED use [from 0.264, CI = (0.242–0.286) in 2019 to 0.143, CI = (0.128–0.158) in 2020; $p < 0.001$]. No significant differences were found in the arrival mode to the ED among the groups.

IHPC recipients with neoplasms and missing principal diagnoses at the time of IHPC

admission showed a significantly lower number of ED visits in 2020 than in 2019 [from 0.268, CI = (0.243–0.294) in 2019 to 0.142, CI = (0.126–0.158) in 2020 for neoplasms; from 0.356, CI = (0.281–0.432) in 2019 to 0.116, CI = (0.06–0.171) in 2020 for missing diagnoses].

No principal symptoms recorded at the time of the ED visits were significantly different among the IHPC recipient groups, except for a decrease in neurological symptoms [from 1.285, CI = (1.103–1.466) in 2019 to 1.063, CI = (0.945–1.18) in 2020; $p = 0.04$].

When accounting for the triage tags, which indicates the level of the emergency at the time of ED admission, the average number of ED visits with a low level of emergency significantly decreased in 2020 compared with 2019 [from 1.439, CI = (1.3–1.579) in 2019 to 1.27, CI = (1.194–1.345) in 2020; $p = 0.036$]. In contrast, the average number of medium and high emergency levels at the time of ED admission in 2020 showed no differences compared with 2019.

Discussion

This study aimed to describe the impact of the COVID-19 pandemic on ED visits among palliative care recipients in a large northern region of Italy. Our results highlighted the fact that during the early stages of the COVID-19 pandemic (when no vaccinations were available), there was a significant reduction in ED visits compared with 2019 among IHPC recipients. Our results confirm the general trend of a reduction in ED use during the pandemic compared with previous years, as has been observed in other studies.^{18–21} Furthermore, this reduction was mainly found for people with neoplasms, formal caregivers, and ED visits characterized by a low level of emergency.

The significant reduction that was observed in ED use for IHPC recipients with neoplasms during the COVID-19 pandemic could have different interpretations from both an individual and political perspective. Moreover, because of the fact that people with neoplasms represent a large majority of palliative care recipients,²² a minor ED use by this population could explain the overall amount of reduced ED admissions that occurred during 2020 in our sample. In this regard, this reduction could be more pronounced

Table 1. Characteristics of the 2019 and 2020 IHPC populations.

Sociodemographic and clinical characteristics	2019 (N = 5155)		2020 (N = 3177)	
	N	%	N	%
Sex				
Female	2438	47.29	1480	46.58
Male	2717	52.71	1697	53.42
Age (years)				
≤65	973	18.87	588	18.51
66–80	2110	40.93	1355	42.65
81–90	1750	33.95	1042	32.80
≥91	322	6.25	192	6.04
IHPC duration (days)				
≤30	2819	54.68	1896	59.68
31–90	1448	28.09	912	28.71
>90	888	17.23	369	11.61
Diagnosis at IHPC admission				
Other	64	1.24	30	0.94
Cardiovascular diseases	101	1.96	47	1.48
Digestive system disorder	57	1.11	29	0.91
Endocrine and metabolic diseases	32	0.62	23	0.72
Hematological disorder	31	0.60	11	0.35
Infectious disorder	14	0.27	9	0.28
Mental disorders	38	0.74	31	0.98
Missing	331	6.42	155	4.88
Musculoskeletal and connective	21	0.41	8	0.25
Neoplasms	4203	81.53	2699	84.95
Neurological disorders	153	2.97	70	2.20
Perinatal and congenital disorder	17	0.33	4	0.13
Respiratory diseases	47	0.91	28	0.88
Trauma and injury	17	0.33	8	0.25
Urogenital disorder	29	0.56	25	0.79
Outcomes				
Dead	3691	71.60	2041	64.24
Alive	1464	28.40	1136	35.76
IHPC, integrated palliative home care.				

Table 2. Comparison of the average number of ED visits in the 2019 and 2020 IHPC populations.

Variables	2019		2020		p value
	ED visits/Ave	CI	ED visits/Ave	CI	
Sex					
Female	0.265	[0.238 to 0.293]	0.139	[0.116 to 0.162]	<0.001*
Male	0.262	[0.228 to 0.296]	0.146	[0.127 to 0.166]	<0.001*
Age (years)					
≤65	0.257	[0.227 to 0.288]	0.152	[0.12 to 0.184]	<0.001*
66–80	0.27	[0.244 to 0.297]	0.135	[0.114 to 0.156]	<0.001*
81–90	0.257	[0.227 to 0.287]	0.152	[0.122 to 0.181]	<0.001*
≥91	0.276	[0.032 to 0.52]	0.119	[0.062 to 0.177]	0.218
Formal caregiver					
No	0.326	[−1.076 to 1.728]	NaN	[NaN-NaN]	= =
Yes	0.264	[0.242 to 0.286]	0.143	[0.128 to 0.158]	<0.001*
IHPC duration (days)					
≤30	0.154	[0.14 to 0.169]	0.092	[0.077 to 0.106]	<0.001*
31–90	0.304	[0.273 to 0.335]	0.199	[0.164 to 0.233]	<0.001*
>90	0.546	[0.44 to 0.652]	0.267	[0.209 to 0.324]	<0.001*
Diagnosis at IHPC admission					
Other	0.18	[0.079 to 0.28]	0.083	[−0.087 to 0.254]	0.327
Cardiovascular diseases	0.119	[0.045 to 0.193]	0.113	[0.015 to 0.211]	0.925
Digestive system disorder	0.248	[0.116 to 0.38]	0.25	[0.05 to 0.45]	0.986
Endocrine and metabolic diseases	0.125	[0.003 to 0.246]	0.216	[0.034 to 0.397]	0.394
Hematological disorder	0.57	[0.118 to 1.023]	0.089	[−0.109 to 0.287]	0.051
Infectious disorder	0.15	[−0.076 to 0.377]	0.584	[−0.09 to 1.257]	0.193
Mental disorders	0.054	[−0.023 to 0.132]	0	[0 to 0]	0.162
Missing	0.356	[0.281 to 0.432]	0.116	[0.06 to 0.171]	<0.001*
Musculoskeletal and connective	0.236	[−0.004 to 0.476]	0.16	[−0.219 to 0.54]	0.708
Neoplasms	0.268	[0.243 to 0.294]	0.142	[0.126 to 0.158]	<0.001*
Neurological disorders	0.177	[0.103 to 0.252]	0.136	[0.037 to 0.235]	0.509
Perinatal and congenital disorder	0.133	[−0.06 to 0.325]	0.526	[−0.446 to 1.498]	0.292
Respiratory diseases	0.174	[0.047 to 0.301]	0.164	[−0.172 to 0.5]	0.953
Trauma and injury	0.142	[−0.065 to 0.348]	0.16	[−0.219 to 0.54]	0.923

(Continued)

Table 2. (Continued)

Variables	2019		2020		p value
	ED visits/Ave	CI	ED visits/Ave	CI	
Urogenital disorder	0.18	[0.027 to 0.333]	0.25	[0.013 to 0.487]	0.613
ED triage codes					
Low level of emergency	1.439	[1.3 to 1.579]	1.27	[1.194 to 1.345]	0.036*
Medium and high level of emergency	1.09	[1.052 to 1.128]	1.054	[0.995 to 1.114]	0.323
Symptoms at ED visit					
Neurological symptoms	1.285	[1.103 to 1.466]	1.063	[0.945 to 1.18]	0.041*
Abdominal pain	1.337	[1.12 to 1.555]	1.198	[0.89 to 1.505]	0.453
Dyspnea	1.169	[1.074 to 1.265]	1.061	[0.988 to 1.134]	0.074
Traumatic problem	1.347	[1.162 to 1.531]	1.258	[1.089 to 1.428]	0.478
Fever	1.314	[0.884 to 1.744]	1.266	[0.75 to 1.781]	0.879
Cardiac rhythm alteration	1.566	[0.533 to 2.599]	1.013	[0.88 to 1.145]	0.247
Gynecological symptoms	1.387	[1.207 to 1.567]	1.377	[1.029 to 1.725]	0.958
Undefined symptoms	1.298	[1.165 to 1.431]	1.196	[1.127 to 1.265]	0.181
Arrival mode					
Ambulance	1.258	[1.143 to 1.373]	1.138	[1.084 to 1.192]	0.064
Autonomous	1.406	[1.292 to 1.52]	1.347	[1.174 to 1.521]	0.576
Others	1.193	[1.068 to 1.317]	1.182	[1.075 to 1.288]	0.892

CI, confidence interval; ED, emergency department; ED visits/Ave, average number of ED visits; IHPC, integrated palliative home care.
* $p < 0.05$.

in our sample, which almost entirely consisted of oncological care recipients. Nevertheless, this high ratio of people with neoplasms taken care of by IHPC is not surprising. In the Italian context, advanced nononcological diseases are usually enrolled in general home care services. Specifically, from a report from the Italian Ministry of Health, the rate of IHPC recipients with neoplasms was 80%, while those with other chronic conditions accounted for 20% in both 2016 and 2017.²³

People with neoplasms are frail and at higher risks for COVID-19 infection, which is frequently associated with complications, intensive care admission, and increased mortality.^{24,25} During the pandemic, more frail and more immunocompromised individuals could have felt anxious and

afraid of being infected in health care facilities, thus leading them to reduce the use of medical services, such as ED. In this regard, a review by O’Rielly *et al.*²⁶ reported that during the early stages of the COVID-19 pandemic, many patients canceled or delayed their routine visits or procedures because of concerns with safety and nosocomial infections. National health policies may also have enhanced this trend. Indeed, significant efforts have been made throughout the world to reduce the risk of contagion in these individuals and to maintain the treatment of frail patients outside of health care facilities and hospitals. The Italian Ministry of Health and National Health Service (NHS) has recommended that the population should avoid being directly admitted to hospitals; instead, they should dial the emergency number dedicated to evaluating patients at home

and decide on eventual transport to the hospital, thus giving priority to the access of EDs through emergency vehicles.²⁷ Even in the United Kingdom, during the first lockdown period, the public was instructed to ‘stay at home, save lives, and protect the NHS’. This resulted in increased anxiety in the population over visiting hospitals, and a higher use of emergency numbers and helplines occurred,²⁸ which resulted in a decline in the number of ED visits with minor health issues.^{19,28,29} Fear of contagion, the deterrence of accessing the ED, and the effect of travel restrictions likely led to seeking medical care for more severe conditions. Our finding supports the assumption that ED visits for medium and high emergency levels did not show a significant decrease during the COVID-19 pandemic. Although highlighted in a small population, the significant reduction in ED visits in oncologic IHPC recipients may be explained by the tendency to protect frail people during the first two waves of the COVID-19 pandemic.

In the midst of the COVID-19 pandemic, home palliative care teams may have played an essential role in providing care during a time period that was characterized by extensive changes in health care services while facing an emergency. In the home care setting, an infectious disease spread represents a severe problem from four points of view. The first view is the need to readjust care provision, adapt therapeutic processes, and guarantee flexibility. The second view is the risk of the patients being infected, which leads to the adoption of precautions with patients.¹⁰ The third view is the risk of health care workers being infected,¹⁰ whereas the fourth view is protecting the caregiver’s health, whose role was pivotal during the COVID-19 pandemic and whose contagion may limit the capacity to care for their patients.³⁰ In this context, ensuring continuity and quality of home care for patients, and reducing unnecessary hospital admissions and avoidable ED visits during a pandemic were essential and challenging aspects of health management. A common response by health care policies to the COVID-19 pandemic has been the digitalization of care via increased technological provisions and skill development by health care professionals, recipients, and their caregivers.³¹ IHCP teams coped with new work challenges by adapting their usual care provision routines, improving their communication with caregivers, and implementing frequent phone or video calls for both recipients and colleagues. During the COVID-19 pandemic, the

key messages from IHCP providers have been flexible, resilient, and able to rapidly implement new solutions in response to evolving care needs, such as by replacing traditional nurse visits with video consultations and improving compliance with medications and the education of the recipients.³² The increase in telemedicine use contributed to an increase in virtual primary care consultations, thus filling the gap in primary care due to the COVID-19 pandemic.^{19,29,33} In our setting, a faster symptom assessment and the ability to communicate in real time by adjusting the medication regimen, combined with a lower number of IHCP recipients, may have contributed to a more timely response by IHCP teams, with the result of a reduced use of ED, especially for less severe emergencies.

The findings from this study showed that the presence of formal caregivers was associated with a significant reduction in ED visits during the COVID-19 pandemic. Previous evidence has shown that the presence of formal caregivers could reduce the risk of ED visits, as they are often trained in patient monitoring and supervision techniques, as well as being completely committed to their care and assistance.^{34,35} In addition, patients who rely on available and skilled caregivers have been shown to be significantly more inclined to meet health care goals.³⁶ Hence, it is plausible that during the COVID-19 pandemic, having a formal caregiver could have represented a protective factor to deal with symptoms and struggles that are treated in the home setting, instead of accessing the ED. The COVID-19 pandemic may have played a positive role in leading IHCP teams, recipients, and their caregivers to enhance symptom management at home,^{18,37} thus resulting in the referral of those urgent cases to the ED and limiting the use of emergency services for less acute visits. This is because, especially during the early stages of the COVID-19 pandemic, ED visits through emergency vehicles were exclusively directed to those infected or in life-threatening situations.³⁸ In this respect, the reduction in the mortality rate in 2020 compared with 2019 would appear to be congruent with the timely management of IHCP recipients that could potentially incur any clinical deterioration and, especially in the palliative care setting, experience secondary conditions to hospitalization. Our results may suggest that the forced response of IHCP teams to the COVID-19 pandemic positively influenced the care that is provided to IHCP recipients and their families.

This may have been mediated by a more rapid response by IHCP teams to the needs of a smaller number of recipients and a better coordination with their caregivers.

Taken together, these results could prompt the conduction of studies aimed at testing the presence of a positive correlation between the pandemic and an improvement of the provision of IHPC at home, which could be suggested by the reduced low-acuity ED visits and reduced mortality. During the first wave of the COVID-19 pandemic, Italian mortality was high, exceeding 74,000 expected deaths in the Northern regions.¹² Because this increase was probably due to deaths related to COVID-19, it is conceivable that the opposite mortality trend we observed in IHPC recipients in the first two waves of the pandemic was due to better caring of this fragile population by home care services. This hypothesis could be explained by the forced resilience of home care services and the improved care delivery (which was not necessarily demonstrated in the presence of health care workers) that occurred during the most acute phase of the pandemic. On the other hand, there may have been unmet care needs by people necessitating IHPC, potentially leading to receiving such care, and deaths that were not detected.

Although supporting the hypotheses that during the early stages of the COVID-19 pandemic, ED visits among IHPC recipients decreased and that there were sociodemographic and clinical characteristics associated with this reduction, our study presented several limitations. The study population refers to a specific region of Northern Italy, and IHPC recipients were mostly people with neoplasms; therefore, our findings cannot be generalized to other regions or countries where the first two waves of the pandemic had a different spread or in which other models of palliative home care delivery were adopted. Moreover, our results are based on administrative databases and may correspondingly suffer in regard to coding accuracy because of clerical error and the limited precision of codes to describe conditions or procedures in detail. Finally, the examined administrative databases could not trace the number of home care visits made by the palliative care team. An in-depth study of the differences in providing such visits before and during the COVID-19 pandemic could comprehensively explain the reduced ED visits among IHPC recipients. While unable to provide a causal link between IHPC provision

and reduced ED visits, which can be obtained only through longitudinal studies, our study represents the first investigation of this phenomenon during the early stages of the pandemic. Indeed, our focus on a cohort of patients who were admitted to the IHPC program, as well as the availability of data individually linked to administrative databases, represents a strength of our study because they provide evidence that the first two waves of COVID-19 have also had a significant impact on health service delivery at home.

Conclusion

This study demonstrated a significant reduction in ED visits among IHPC recipients during the COVID-19 pandemic, especially those individuals with neoplasms who relied on formal caregivers. Correspondingly, the ED visits characterized by a low level of emergency were significantly reduced. The decline in ED visits did not result in an increase in mortality among IHPC recipients. The rationalization of resources by identifying an appropriate ratio between IHPC providers and implementations aimed at improving telemedicine in palliative care represents future research topics in this area. Moreover, it would be worth understanding whether, in the Italian context, the reduction in ED visits from IHPC during the COVID-19 pandemic is comparable to that experienced in general home care. Future studies should demonstrate the efficacy of IHPC longitudinally in reducing the use of emergency services, providing policymakers with a broader perspective to enhance the delivery of care to all patients by strengthening IHPC teams. The adaptation of existing home care services by implementing innovative solutions should encourage health policies to maintain good practices adopted during the COVID-19 pandemic, thus reducing the burden on emergency services and improving the care of recipients and their families.

Declarations

Ethics approval and consent to participate

The Italian National Information System databases are official, anonymized Ministerial Health information systems. All the patient information is centrally anonymized and available to authorized institutions to be used for epidemiological and/or health organization studies without any further authorizations. Hence, ethics committee approval was not needed. Personal data

treatments are conducted in compliance with the current rules set out in EU Regulation 2016/679 and the current legislation on the Protection of Personal Dataset out in Legislative Decree 101/2018, as well as Legislative Decree 196/2003, and subsequent amendments and additions. Our study used statistical and aggregate data that were shared according to the current deontological rules for the processing of National Statistical System data for scientific research purposes, in accordance with the provisions of Art. 5-ter of Legislative Decree 33/2013 as amended by Legislative Decree 97/2016 and Legislative Decree 101/2018.

Consent for publication

Not applicable.

Author contributions

Alessandro Scacchi: Conceptualization; Data curation; Methodology; Project administration; Writing – original draft; Writing – review & editing.

Alessio Conti: Data curation; Methodology; Writing – original draft; Writing – review & editing.

Gianfranco Politano: Conceptualization; Data curation; Formal analysis; Methodology; Writing – original draft; Writing – review & editing.

Marco Dalmasso: Conceptualization; Data curation; Investigation; Methodology; Writing – original draft; Writing – review & editing.

Sofia Ostellino: Data curation; Methodology; Writing – original draft; Writing – review & editing.

Maria Michela Gianino: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Writing – original draft; Writing – review & editing.

Acknowledgements

Not applicable.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.


Competing interests


The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Availability of data and materials

The datasets generated and/or analyzed during this study are not publicly available but are available from the corresponding author on reasonable request.

ORCID iDs

Alessio Conti  <https://orcid.org/0000-0003-4301-8924>

Gianfranco Politano  <https://orcid.org/0000-0001-5268-9899>

Sofia Ostellino  <https://orcid.org/0000-0002-6275-3214>

Maria Michela Gianino  <https://orcid.org/0000-0003-4239-7475>

References


1. Borasio GD. Translating the World Health Organization definition of palliative care into scientific practice. *Palliat Supp Care* 2011; 9: 1–2.
2. GU della Repubblica Italiana IMP. Disposizioni per garantire l'accesso alle cure palliative e alla terapia del dolore, 2010. <https://www.parlamento.it/parlam/leggi/10038l.htm> (accessed 8 November 2022).
3. Scaccabarozzi G, Lovaglio PG, Limonta F, et al. Monitoring the Italian home palliative care services. *Healthcare* 2019; 7: 4, <https://www.mdpi.com/2227-9032/7/1/4>
4. Franchini L, Varani S, Ostan R, et al. Home palliative care professionals perception of challenges during the Covid-19 outbreak: a qualitative study. *Palliat Med* 2021; 35: 862–874.
5. Mitchell S, Maynard V, Lyons V, et al. The role and response of primary healthcare services in the delivery of palliative care in epidemics and pandemics: a rapid review to inform practice and service delivery during the COVID-19 pandemic. *Palliat Med* 2020; 34: 1182–1192.
6. Tanzi S, Alquati S, Martucci G, et al. Learning a palliative care approach during the COVID-19 pandemic: a case study in an Infectious Diseases Unit. *Palliat Med* 2020; 34: 1220–1227.
7. Nouvet E, Sivaram M, Bezanson K, et al. Palliative care in humanitarian crises: a review of the literature. *J Int Humanit Action* 2018; 3: 5.
8. Rosenbaum L. Facing Covid-19 in Italy – ethics, logistics, and therapeutics on the epidemic's front line. *N Engl J Med* 2020; 382: 1873–1875.
9. Radbruch L, Knäul FM, de Lima L, et al. The key role of palliative care in response to the

- COVID-19 tsunamis of suffering. *Lancet (London, England)* 2020; 395: 1467–1469.
10. Porzio G, Cortellini A, Bruera E, *et al.* Home care for cancer patients during COVID-19 pandemic: the double triage protocol. *J Pain Symptom Manage* 2020; 60: e5–e7.
 11. Brogi F, Guardabascio B and Barcaroli G. Covid-19 in Italy: actual infected population, testing strategy and imperfect compliance, 2020. DOI: 10.13140/RG.2.2.18275.50729.
 12. Dorrucchi M, Minelli G, Boros S, *et al.* Excess mortality in Italy during the COVID-19 pandemic: assessing the differences between the first and the second wave, year 2020. *Front Public Health* 2021; 9: 669209.
 13. Ministero della Salute. DM 15 maggio 1992 – Gazzetta Ufficiale n. 121 del 25/5/92- Serie Generale Criteri e requisiti per la classificazione degli interventi di emergenza. *Gazz Uff* 1992; 121del, <https://www.fnopi.it/wp-content/uploads/DM150592.pdf>
 14. Regione Piemonte. Deliberazione della Giunta Regionale 10 gennaio 2012, n. 3-3259. *Boll Uff* 2015: 3–8.
 15. Greifer N. *WeightIt: weighting for covariate balance in observational studies*. R package version 0.12.0., 2021, <https://cran.r-project.org/package=WeightIt> (accessed 11 October 2021).
 16. Hainmueller J. Entropy balancing for causal effects: a multivariate reweighting method to produce balanced samples in observational studies. *Polit Anal* 2012; 20: 25–46.
 17. Tukey JW. Comparing individual means in the analysis of variance. *Biometrics* 1949; 5: 99–114. <http://www.jstor.org/stable/3001913>
 18. Vanni G, Legramante JM, Pellicciaro M, *et al.* Effect of lockdown in surgical emergency accesses: experience of a COVID-19 hospital. *In Vivo* 2020; 34: 3033–3038.
 19. Panovska-Griffiths J, Ross J, Elkhodair S, *et al.* Exploring overcrowding trends in an inner city emergency department in the UK before and during COVID-19 epidemic. *BMC Emerg Med* 2021; 21: 43.
 20. Uimonen M, Kuitunen I, Jämsen E, *et al.* Emergency visits by older adults decreased during COVID-19 but increased in the oldest old. *J Am Geriatr Soc* 2021; 69: 1738–1740.
 21. De Vito B, Castagno E, Garrone E, *et al.* Narrating care during the COVID-19 pandemic in a paediatric emergency department. *Reflective Pract* 2022; 23: 219–231.
 22. World Health Organization. Palliative care, 2020, <https://www.who.int/news-room/fact-sheets/detail/palliative-care>
 23. Ministero della Salute. Direzione generale della programmazione sanitaria: disposizioni per garantire l'accesso alle cure palliative e alla terapia del dolore. *Rapp Al Parlam Sullo Stato Di Attuazione Della Legge N 38 Del 15 Marzo* 2010; 2017: 1–81.
 24. Minichsdorfer C, Jeryczynski G, Krall C, *et al.* Impact of COVID-19 lockdown on routine oncology versus emergency care at a high volume cancer centre. *Eur J Clin Invest* 2021; 51: e13623.
 25. Tian Y, Qiu X, Wang C, *et al.* Cancer associates with risk and severe events of COVID-19: a systematic review and meta-analysis. *Int J Cancer* 2021; 148: 363–374.
 26. O'Rielly C, Ng-Kamstra J, Kania-Richmond A, *et al.* Surgery and COVID-19: a rapid scoping review of the impact of the first wave of COVID-19 on surgical services. *BMJ Open* 2021; 11: e043966.
 27. Fagoni N, Perone G, Villa GF, *et al.* The lombardy emergency medical system faced with COVID-19: the impact of out-of-hospital outbreak. *Prehosp Emerg Care* 2021; 25: 1–7.
 28. McConkey R and Wyatt S. Exploring the fall in A&E visits during the pandemic [Internet]. *The Health Foundation report*, 2020, <https://www.health.Org.uk/news-and-comment/charts-and-infographics/exploring-the-fall-in-a-e-visits-during-the-pandemic> (accessed 7 April 2022).
 29. The Royal College of Emergency Medicine. COVID-19: resetting emergency department care. *The Royal College of Emergency Medicine report*, <https://rcem.ac.uk/covid-19-resetting-emergency-care/> (accessed 8 April 2022).
 30. Buttiron Webber T, Giuliano S, Patrone C, *et al.* Home se-cure: a home care service for cancer patients during the COVID-19 pandemic. *Int J Environ Res Pub Heal* 2021; 18: 10913.
 31. Sutherland AE, Stickland J and Wee B. Can video consultations replace face-to-face interviews? Palliative medicine and the Covid-19 pandemic: rapid review. *BMJ Suppl Palliat Care* 2020; 10: 271–275, <https://pubmed.ncbi.nlm.nih.gov/32457086>
 32. Mojtahedi Z and Shen JJ. Home palliative care during the COVID-19 pandemic: a scoping review. *Am J Hosp Palliat Care*.

Epub ahead of print 3 May 2022. DOI:
10.1177/10499091221093314.

33. Warner I. Telemedicine applications for home health care. *J Telemed Telecare* 1997; 3(Suppl. 1): 65–66.
34. Borraccino A, Campagna S, Politano G, *et al.* Predictors and trajectories of ED visits among patients receiving palliative home care services: findings from a time series analysis (2013–2017). *BMC Palliat Care* 2020; 19: 126, <https://pubmed.ncbi.nlm.nih.gov/32799860>
35. Gagnon-Roy M, Hami B, Génèreux M, *et al.* Preventing emergency department (ED) visits and hospitalisations of older adults with cognitive impairment compared with the general senior population: what do we know about avoidable incidents? Results from a scoping review. *BMJ* *Open* 2018; 8: e019908, <https://pubmed.ncbi.nlm.nih.gov/29666129>
36. O’Sullivan MJ and Volicer B. Factors associated with achievement of goals for home health care. *Home Health Care Serv Q* 1997; 16: 21–34.
37. Pinarelli L, Colais P, Mataloni F, *et al.* Access to the emergency at times of COVID-19: an analysis of the first three months in the Lazio region. *Epidemiol Preven* 2021; 44: 359–366, https://epiprev.it/articoli_scientifici/laccesso-al-pronto-soccorso-ai-tempi-del-covid-19-unanalisi-dei-primi-tre-mesi-nella-regione-lazio
38. Perlini S, Canevari F, Cortesi S, *et al.* Emergency department and out-of-hospital emergency system (112-AREU 118) integrated response to coronavirus disease 2019 in a Northern Italy centre. *Intern Emerg Med* 2020; 15: 825–833.

Visit SAGE journals online
[journals.sagepub.com/
home/pcr](https://journals.sagepub.com/home/pcr)

 SAGE journals