

Microphysical time scales, supersaturation fluctuations and at a warm Cloud Top Boundary in temporal decay

Original

Microphysical time scales, supersaturation fluctuations and at a warm Cloud Top Boundary in temporal decay / Fossa', Ludovico; Abdunabiev, Shahbozbek; Golshan, Mina; Tordella, Daniela. - ELETTRONICO. - 18:(2023), pp. 1-1. (18th European Turbulence Conference Valencia, Spain September 4-6, 2023).

Availability:

This version is available at: 11583/2980162 since: 2023-09-08T14:32:26Z

Publisher:

EUROMECH - European Mechanics Society

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Microphysical time scales, supersaturation fluctuations and at a warm Cloud Top Boundary intemporal decay

L. Fossa^{*,*}, S. Abdunabiev[†], M. Golshan[†], D. Tordella[†]

Recent results have shown that there is an acceleration in the spread of the size distribution of droplet populations in the region bordering the cloud and undersaturated ambient¹. We have analyzed the supersaturation balance in this region, which is typically a highly intermittent shearless turbulent mixing layer, under a condition where there is no mean updraft. We have investigated the evolution of the cloud - clear air interface and of the droplets therein via direct numerical simulations. We have compared horizontal averages of the phase relaxation, evaporation, reaction and condensation times within the cloud-clear air interface for the size distributions of the initial monodisperse and polydisperse droplet populations². For the monodisperse population, a clustering of the values of the reaction, phase and evaporation times, that is around 20-30 seconds, is observed in the central area of the mixing layer, just before the location where the maximum value of the supersaturation turbulent flux occurs. This clustering of values is similar for the polydisperse population but also includes the condensation time. The mismatch between the time derivative of the supersaturation and the condensation term in the interfacial mixing layer is correlated with the planar covariance of the horizontal longitudinal velocity derivatives of the carrier air flow and the supersaturation field, thus suggesting that a quasi-linear relationship may exist between these quantities.




Figure 1: Vertical distribution of the evaporation τ_{evap} , phase relaxation τ_{phase} and reaction τ_{react} time scales computed inside each grid cell and then averaged on horizontal planes inside the cloud and mixing regions. Data are displayed for two different time steps at the beginning and the end of the transient. The planar average of supersaturation is also plotted for comparison purposes.

*The University of Sheffield, UK

[†]Dip. Scienza Applicata e Tecnologia (DISAT), Politecnico di Torino, Italy

¹ Golshan et al., *International Journal of Multiphase Flow* **140**, (2021).

² Fossà et al., *Physics of fluids* **34**, (2022).