

Diastereoselective Self-Assembly of Low-Symmetry Pd<sub>n</sub>L<sub>2n</sub> Nanocages through Coordination-Sphere Engineering

*Original*

Diastereoselective Self-Assembly of Low-Symmetry Pd<sub>n</sub>L<sub>2n</sub> Nanocages through Coordination-Sphere Engineering / Molinska, Paulina; Tarzia, Andrew; Male, Louise; Jelfs, Kim E; Lewis, James. - In: ANGEWANDTE CHEMIE. - ISSN 1521-3773. - 62:51(2023). [10.1002/anie.202315451]

*Availability:*

This version is available at: 11583/2983609 since: 2023-11-06T09:31:53Z

*Publisher:*

Wiley

*Published*

DOI:10.1002/anie.202315451

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

A Journal of the Gesellschaft Deutscher Chemiker

# Angewandte Chemie

GDCh

International Edition

www.angewandte.org

## Accepted Article

**Title:** Diastereoselective Self-Assembly of Low-Symmetry Pd<sub>n</sub>L<sub>2n</sub> Nanocages through Coordination-Sphere Engineering

**Authors:** Paulina Molinska, Andrew Tarzia, Louise Male, Kim E. Jelfs, and James Lewis

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

**To be cited as:** *Angew. Chem. Int. Ed.* **2023**, e202315451

**Link to VoR:** <https://doi.org/10.1002/anie.202315451>

# Diastereoselective Self-Assembly of Low-Symmetry Pd<sub>n</sub>L<sub>2n</sub> Nanocages through Coordination-Sphere Engineering

Paulina Molinska,<sup>a</sup> Andrew Tarzia,<sup>b</sup> Louise Male,<sup>a</sup> Kim E. Jelfs<sup>c</sup> and James E. M. Lewis<sup>a\*</sup>

<sup>a</sup>School of Chemistry, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK. Email: j.e.m.lewis@bham.ac.uk.

<sup>b</sup>Department of Applied Science and Technology, Politecnico di Torino, Corso Duca degli Abruzzi 24, Torino 10129, Italy.

<sup>c</sup>Department of Chemistry, Imperial College London, Molecular Sciences Research Hub, White City Campus, Wood Lane, London, W12 0BZ, UK.

Metal-organic cages (MOCs) are popular host architectures assembled from ligands and metal ions/nodes. Assembling structurally complex, low-symmetry MOCs with anisotropic cavities can be limited by the formation of statistical isomer libraries. We set out to investigate the use of primary coordination-sphere engineering (CSE) to bias isomer selectivity within homo- and heteroleptic Pd<sub>n</sub>L<sub>2n</sub> cages. Unexpected differences in selectivities between alternative donor groups led us to recognise the significant impact of the second coordination sphere on isomer stabilities. From this, molecular-level insight into the origins of selectivity between *cis* and *trans* diastereoisomers was gained, highlighting the importance of both host-guest and host-solvent interactions, in addition to ligand design. This detailed understanding allows precision engineering of low-symmetry MOC assemblies without wholesale redesign of the ligand framework, and fundamentally provides a theoretical scaffold for the development of stimuli-responsive, shape-shifting MOCs.


## Introduction

Metal-organic cages (MOCs) are discrete, porous supramolecular architectures assembled from metal ions/nodes and coordinating ligands.<sup>1</sup> The ability to encapsulate guest molecules within the cavities of MOCs has led to investigations for their use in catalysis,<sup>2</sup> sensing,<sup>3</sup> drug delivery<sup>4</sup> and stabilising reactive species.<sup>5</sup>

Detailed principles behind the self-assembly of high-symmetry MOCs have been elucidated over the last four decades. To generate more sophisticated systems<sup>6</sup> with advanced functionality,<sup>7</sup> attention has recently turned to the development of methodologies to access lower symmetry cages.<sup>8</sup> These include the design of mixed-ligand (heteroleptic)<sup>9</sup> (Fig. 1a) and mixed-metal (heteronuclear) MOCs (Fig. 1b),<sup>10</sup> as well as those assembled from low-symmetry ligands (Fig. 1c).<sup>11</sup> Using these approaches, low-symmetry MOCs have been realised that exhibit shape-<sup>12</sup> and orientation-selective<sup>13</sup> guest binding.

The inherent directionality of unsymmetrical ligands gives rise to multiple possible constitutional isomers of their metal-organic assemblies (Fig. 1c). Various strategies have been investigated towards the high-fidelity, isomer-selective self-assembly of low-symmetry ligands.<sup>11</sup> Aside from developing ligands with mixed-denticity donors,<sup>14</sup> these include geometric design parameters,<sup>15,16</sup> use of non-covalent interactions within the ligand backbone,<sup>17</sup> and coordination-sphere engineering (CSE; also known as side-chain directing).<sup>18</sup> CSE strategies can be subdivided into two further categories: those that

use attractive interactions, such as hydrogen-bonding,<sup>19</sup> and those that use repulsive interactions, such as steric hindrance.



**Fig. 1** Cartoons of different unsymmetrical MOCs: a) *cis*- and *trans*-M<sub>2</sub>L<sub>2</sub>L'<sub>2</sub> heteroleptic cages, b) MM'L<sub>4</sub> heteronuclear cage, and c) potential isomers of M<sub>2</sub>L<sub>4</sub> cages assembled from an unsymmetrical ditopic ligand.

The use of CSE approaches in the metal-organic self-assembly of low-symmetry ligands has been limited,<sup>20</sup> despite its success in directing the self-assembly of heteroleptic MOCs.<sup>21,22,23,24</sup> We have previously reported preliminary findings of two systems that use steric parameters, by themselves and in combination with geometric designs,<sup>15</sup> whilst Crowley and co-workers have used hydrogen-bonding interactions to direct formation of a *cis*-Pd<sub>2</sub>L<sub>4</sub> cage.<sup>19</sup> Aside from these individual examples, CSE strategies to direct the isomer-selective assembly of homoleptic MOCs remains a significant, and under-investigated, challenge.

It was envisaged that combinations of unsubstituted and sterically bulky coordinating groups in unsymmetric ditopic ligands would bias self-assembly with Pd(II) ions towards specific isomers of  $M_nL_{2n}$  cages. Given their prior utility in directing the self-assembly of heteroleptic MOCs, picoline<sup>22</sup> ( $L^P$ ) and quinoline<sup>24</sup> ( $L^Q$ ) coordinating groups were chosen for investigation.

In this work, the successful use of CSE in the selective synthesis of  $Pd_nL_{2n}$  MOC isomers is reported. For both quinoline and picoline ligands, biasing towards assemblies with a 2:2 stoichiometry of donors at the metal nodes was observed. Intriguingly, the different coordinating groups were selective for alternative donor arrangements around the metal ions,<sup>25</sup> namely *cis* ( $L^Q$ ) and *trans* ( $L^P$ ).

Through careful investigation, the diastereoselectivities between *cis* and *trans* cages were rationalised, and molecular origins for this effect identified, demonstrating the importance of considering the combined effects of both first and second coordination sphere interactions in the design of these supramolecular systems. This detailed understanding has ramifications for the future design of MOCs, particularly those of low-symmetry, and also stimuli-responsive systems using CSE approaches.


CSE strategies allow the targeted assembly of MOCs with different symmetries whilst maintaining the structural formulation resulting from the design of the core ligand scaffold. As such, this nuanced approach provides a route for precision engineering the shape of low-symmetry MOCs, and the cavity spaces within, towards the development of more sophisticated, functional supramolecular hosts.

## Results & Discussion

### $Pd_2L_4$ cages

Based on a dipyriddy ligand motif originally reported by Chand and co-workers,<sup>26</sup>  $L^Q$  and  $L^P$  (Fig. 2a) were synthesised by ester condensation between commercially available 3-(hydroxymethyl)pyridine and the appropriate carboxylic acid. Each ligand was then combined with  $Pd(NO_3)_2 \cdot 2H_2O$  in a 2:1 ratio in  $d_6$ -DMSO ( $[L1] = 40$  mM) and heated at 50 °C for 24 h; no further changes were observed by  $^1H$  NMR with prolonged heating.

Analysis by electrospray ionisation mass spectrometry (ESI-MS) indicated formation of  $Pd_2L_4$  assemblies ( $C1$ ) for both systems (Fig. S22-25 and S53-56). Diffusion-orientated spectroscopy (DOSY) further supported this, with each system displaying diffusion coefficients ( $D = 9.51 \times 10^{-11}$  and  $11.0 \times 10^{-11}$  m<sup>2</sup>s<sup>-1</sup> for  $C1^Q$  and  $C1^P$ , respectively) consistent with related systems.<sup>26</sup>



**Fig. 2** a) Ligands  $L1^Q$  and  $L1^P$ . Partial  $^1H$  NMR (500 MHz,  $d_6$ -DMSO, 298 K) of b)  $C1^Q$ , and c)  $C1^P$ . SCXRD structures of d) *cis*- $C1^Q \cdot Cl$  (only one crystallographically independent conformer shown), and e) *trans*- $C1^P \cdot Cl$ . Exohedral interactions of  $BF_4^-$  counteranions with the Pd(II) coordination sphere in f)  $C1^Q$  (F...H 2.31-2.45 Å, C-H...F 139-156°), and g)  $C1^P$ .

In both cases a mixture of isomers of  $C1$  was observed to form by  $^1H$  NMR (Fig. 2b and 2c), but with a major component arising at a compositional fraction greater than expected from a statistical library (i.e. 25%), demonstrating successful induction of isomer selectivity. The percentage composition of the predominant species was estimated through a comparison of integrals between

methylene signals ( $H_e$ ) and isolated signals in the aromatic region of the  $^1\text{H}$  NMR spectra, giving values of approximately 70% and 50% for  $\mathbf{C1}^Q$  and  $\mathbf{C1}^P$ , respectively (Fig. S21 and S52).

For both major species of  $\mathbf{C1}$ , nuclear Overhauser effect spectroscopy (NOESY) revealed through-space interactions between signals assigned to the different coordinating groups of  $\mathbf{L1}$  (Fig. S17 and S50), identifying these as either the *cis* or *trans* isomers (Fig. 1c). In the case of  $\mathbf{C1}^Q$ , apparent diastereotopic splitting of the  $\text{CH}_2$  signal ( $J \approx 14$  Hz; Fig. 2b) was consistent with formation of *cis*- $\mathbf{C1}^Q$  as the major species. The absence of diastereotopic splitting for the major isomer of  $\mathbf{C1}^P$  led to the conclusion that this was most likely *trans*- $\mathbf{C1}^P$ .

It was also possible to observe second, minor isomers for  $\mathbf{C1}^Q$  (~9%) and  $\mathbf{C1}^P$  (~14%) that were identified as the alternative diastereoisomers *trans*- $\mathbf{C1}^Q$  (ESI section S2.3) and *cis*- $\mathbf{C1}^P$  (ESI section S2.8), respectively.

Related Pd(II) cages are known to encapsulate an  $\text{NO}_3^-$  anion that can be exchanged for stronger binding halide anions.<sup>27</sup> Upon addition of 1 eq. of  $\text{Bu}_4\text{NCl}$  to  $\mathbf{C1}^Q$  and  $\mathbf{C1}^P$ , encapsulation of  $\text{Cl}^-$  was evidenced by notable downfield shifts of signals assigned to the endohedral protons of the cage (e.g.  $H_a$  and  $H_h$ ,  $\Delta\delta = 0.49$  and  $0.65$ , respectively, for  $\mathbf{C1}^P$ ; Fig. S30 and S60).

$\mathbf{C1} \supset \text{Cl}^-$  were subsequently prepared and isolated through self-assembly of  $\mathbf{L1}$  with  $[\text{Pd}(\text{CH}_3\text{CN})_4](\text{BF}_4)_2$  in the presence of 1 eq. of  $\text{Bu}_4\text{NCl}$ . Unexpectedly, although the switch to  $\text{BF}_4^-$  counterions and encapsulation of a  $\text{Cl}^-$  guest in place of  $\text{NO}_3^-$  anions did not change the identity of the major host isomer, the selectivity values were altered:<sup>28</sup> for  $\mathbf{C1}^Q$  the *cis* isomer fell to ~40% of the mixture (Fig. S32), whilst the *trans* isomer of  $\mathbf{C1}^P$  increased to ~70% (Fig. S68).

The solid-state structures of  $\mathbf{C1} \supset \text{Cl}^-$  were determined by single-crystal X-ray diffraction (SCXRD) analysis, which revealed the anticipated *cis*- and *trans*- $[\text{Pd}_2(\mathbf{L1})_4 \supset \text{Cl}]^{3+}$  assemblies for  $\mathbf{C1}^Q$  (Fig. 2d) and  $\mathbf{C1}^P$  (Fig. 2e), respectively.<sup>29</sup> For  $\mathbf{C1}^P$ , steric clash of the methyl groups was avoided through induction of a helical twist, with an azimuthal angle ( $\alpha$ ) of ~27°. Whilst  $\mathbf{C1}^Q$  displayed no significant helical twist ( $\alpha \approx 1-3^\circ$ ), resulting in a slightly larger Pd...Pd distance (6.94-7.05 Å compared to 6.81 Å for  $\mathbf{C1}^P$ ), the planes of pyridine and quinoline units *trans* to each other were rotated to reduce interactions ( $\Theta = 3-28^\circ$ ; Fig. S250 and Table S2).

### Molecular origins of diastereoselectivity

Interestingly, the relative energies of the *cis* and *trans* isomers of both  $\mathbf{C1}^Q$  and  $\mathbf{C1}^P$ , computed using density functional theory (DFT) calculations

(HSE06 functional and Def2-SVP basis set with implicit DMSO solvation), suggested the *trans* isomer should be most stable for both systems (ESI section S3). This implied that additional external influences, beyond inherent structural factors, were responsible for the observed speciation. Thus, the question arose: why did  $\mathbf{C1}^Q$  exhibit selectivity towards the *cis* isomer?

The SCXRD structures showed exohedral  $\text{BF}_4^-$  anions located in proximity to the Pd(II) ions<sup>30</sup> for both assemblies (Fig. 2f and 2g). For  $\mathbf{C1}^P$  the steric bulk of the methyl groups resulted in a greater F...Pd distance compared to  $\mathbf{C1}^Q$ . Indeed, for  $\mathbf{C1}^Q$ , interactions between the counterions and C-H of both pyridine and quinoline donors were observed (F...H 2.3-2.8 Å, C-H...F 139-166°, Fig. 2f). This initially led us to consider that the different diastereoselectivities resulted from differences in interactions between cage and counteranions. Specifically, non-covalent interactions between the  $\text{BF}_4^-$  anions and  $\mathbf{C1}^Q$  stabilised the *cis* isomer to such an extent as to make it lower in energy than the more sterically favourable *trans* isomer.


This hypothesis was probed on two fronts: through dilution ( $[\mathbf{C1}] 10 - 1$  mM) in  $d_6$ -DMSO (Fig. 3b, 3c, S75 and S76) to investigate the effect of reducing the anion concentration, and the synthesis of  $\mathbf{C1}^Q \supset \text{NO}_3^-$  in the presence of excess  $\text{BF}_4^-$  or  $\text{OTf}^-$  to monitor the effect of increased anion concentration and stoichiometry (Fig. S71). No impact on the diastereoselectivity was observed from either study (ESI section 2.11).

The thought occurred that, whilst the  $\text{BF}_4^-$  anions were located around the Pd(II) nodes in the solid-state, in solution the concentration of strongly hydrogen-bond accepting DMSO molecules would be orders of magnitude higher. The idea that solvent molecules, rather than anions, interacting with the coordination sphere of the cages were responsible for stabilisation of the *cis* assemblies was thus examined experimentally.

The dilution studies were revisited using  $\text{CD}_3\text{CN}$  – a weaker hydrogen bond acceptor – as titrant (DMSO and MeCN have hydrogen bond acceptor parameters,  $\beta$ , of 8.9 and 4.7, respectively<sup>31</sup>). In this instance, the proportion of the *trans* isomer of  $\mathbf{C1}^P$  increased slightly with dilution (Fig. S81 and S82). Meanwhile, for  $\mathbf{C1}^Q$ , the percentage of the *cis* cage decreased; as the proportion of  $\text{CD}_3\text{CN}$  increased, the minor *trans* isomer became more prominent (Fig. S77 and S78). At a solvent ratio of 9:1  $\text{CD}_3\text{CN}/d_6$ -DMSO, the percentage composition of *trans*- $\mathbf{C1}^Q$  actually superceded that of the *cis* isomer (Fig. 3d). Similarly, addition of  $\text{D}_2\text{O}$  ( $\beta = 4.5$ )<sup>31</sup> to a  $d_6$ -DMSO solution of

**C1<sup>Q</sup>** also resulted in enhancement of the *trans* isomer at the expense of the *cis* (Fig. S72 and S74).

It was concluded that, in the absence of additional effects, the *trans* isomer is favoured for both **C1<sup>P</sup>** and **C1<sup>Q</sup>** purely on grounds of steric hindrance. The *cis* isomers, however, provide a suitable site around the coordination sphere for interacting with hydrogen bond acceptors (Fig. 3a). Consequently, employing as solvent DMSO – a strong hydrogen bond acceptor – led to a reduction in the relative energy of the *cis* isomer. For **C1<sup>Q</sup>** this effect was more pronounced due to the four polarised aromatic C-H bonds (H<sub>b</sub> and H<sub>j</sub>) around each Pd(II) ion (compared to just two for **C1<sup>P</sup>** - H<sub>b</sub>).



**Fig. 3** a) Second coordination sphere effects with solvent molecules alter the equilibrium between *cis*- and *trans*-**C1<sup>Q</sup>**. Partial <sup>1</sup>H NMR (300 MHz, 298 K) of **C1<sup>Q</sup>**NO<sub>3</sub> b) 10 mM (*d*<sub>6</sub>-DMSO) with *cis* isomer peaks labelled, c) 1 mM (*d*<sub>6</sub>-DMSO), and d) 1 mM (9:1 CD<sub>3</sub>CN/*d*<sub>6</sub>-DMSO) with *trans* isomer peaks labelled.

The difference in isomer selectivity between **C1**NO<sub>3</sub><sup>-</sup> and **C1**Cl<sup>-</sup> is proposed to arise partly from a reduction in Pd...Pd distance, with the smaller guest inducing an increased helical pitch.<sup>32</sup> This increased the offset of bulky donor groups in the *trans* isomer, further reducing its relative energy, observed as an increase in diastereoselectivity towards *trans*-**C1**. For **C1<sup>Q</sup>**, the conformational changes upon guest exchange may also impact the complementarity between the external binding pocket and solvent molecules, reducing effective stabilisation of the *cis* isomer. This was suggested by a lack of shift in the quinoline resonance H<sub>j</sub> for *trans*-**C1<sup>Q</sup>** (Δδ = 0.01 ppm), in contrast to *cis*-**C1<sup>Q</sup>** (Δδ = 0.11 ppm), upon increasing CD<sub>3</sub>CN composition

(Fig. 3b-d), indicative of H<sub>j</sub> being incapable of significant interactions with solvent molecules in the *trans* cage isomer.

From the combined data, we have constructed a molecular-level picture of the multiple interactions that influence the observed diastereoselectivities between *cis* and *trans* cages. The ligand design, endohedral interactions between host and guest, and exohedral interactions with solvent molecules all contribute to the relative energies of the diastereomers. Thus, both first and second coordination spheres<sup>33</sup> play an important role in directing the self-assembly process. Modulation of these factors enables control over isomer selectivity, and opens up the possibility for stimuli-responsive switching of the equilibrium position within isomer libraries.<sup>34</sup> More detailed investigations into this effect are underway and will be reported in due course.


### Larger Pd<sub>2</sub>L<sub>4</sub> cages

To probe the utility of these designs with alternative ligand scaffolds, ligand **L2<sup>Q</sup>** (Fig. 4a) was investigated. We have previously reported the self-assembly of ligand **L2<sup>P</sup>**; in *d*<sub>6</sub>-DMSO a mixture of the *cis*- and *trans*-Pd<sub>2</sub>L<sub>4</sub> cage (**C2**) isomers formed, whilst in CD<sub>3</sub>CN *trans*-**C2<sup>P</sup>** formed essentially exclusively.<sup>15</sup> It had been suggested that this behaviour arose from the higher polarity DMSO solvent stabilising the more polar *cis* isomer,<sup>35</sup> without being able to provide a more detailed explanation. In light of the new investigations with **C1**, it is now proposed that the stronger hydrogen bond acceptor nature of DMSO, compared to CH<sub>3</sub>CN, leads to enhanced stabilisation of the *cis*-**C2<sup>P</sup>** isomer specifically through hydrogen bonding interactions between solvent molecules and the exohedral face of the Pd(II) coordination sphere. These interactions are less favoured with the *trans* isomer which, excluding other factors, provides the least sterically hindered primary coordination sphere.

Self-assembly of **L2<sup>Q</sup>** with Pd(II) (as the BF<sub>4</sub><sup>-</sup> salt) in *d*<sub>6</sub>-DMSO resulted in what, superficially, appeared to be near-quantitative (estimated at 70% by NMR integration; Fig. S114) formation of a single species, **C2<sup>Q</sup>** (Fig. 4b). In combination, the high-symmetry NMR spectra (Fig. 4b and S104-106), ESI-MS (Fig. S115-118) and NOESY (H<sub>a</sub>...H<sub>n</sub>, H<sub>b</sub>...H<sub>m</sub>; Fig. S112) data identified **C2<sup>Q</sup>** as either *cis*- or *trans*-Pd<sub>2</sub>(**L2<sup>Q</sup>**)<sub>4</sub>. The absence of prochiral units within the ligand structure prevented the use of diastereotopic splitting (or lack thereof) as a diagnostic tool to differentiate the two isomers in solution.

The solid-state structure of **C2<sup>Q</sup>** was determined by SCXRD and revealed, as expected, a *cis* arrangement of ligands within the assembly (Fig.

4c-e).<sup>29</sup> Consequently, it seemed the preference under these conditions for *cis* and *trans* coordination environments when pairing pyridine with quinoline and picoline, respectively, holds for different ligand scaffolds.




**Fig. 4** a) Ligands  $L2^Q$  and  $L2^P$ ; b) Partial  $^1H$  NMR (500 MHz,  $d_6$ -DMSO, 298 K) of  $C2^Q$ ; SCXRDR structure of *cis*- $C2^Q$  showing c) *cis*- $C2^Q$ →DMSO<sub>2</sub>, d) view down Pd...Pd axis, and e) *cis* coordination environment around a Pd(II) ion.

### $Pd_3L_6$ 'double-walled' triangles

To explore higher nuclearity systems,  $L3^Q$  and  $L3^P$  (Fig. 5a) were synthesised. Symmetric, dipyriddy analogues of these ditopic ligands<sup>26</sup> have been shown to assemble into 'double-walled'  $Pd_3L_6$  triangles (Fig. 5b).<sup>36,37</sup> For such structures assembled from an unsymmetrical ligand, one instance of which has been reported by Chand and co-workers,<sup>38</sup> there are 9 possible isomers. Two of these provide all-*cis* or all-*trans* arrangements of donors at the three metal nodes (Fig. 5b). From our understanding of isomer selectivities induced by CSE, it was hypothesised that self-assembly of  $L3^Q$  with Pd(II) in DMSO would favour formation of the *cis*- $Pd_3L_6$  isomer of  $C3$ , whilst  $L3^P$  would be biased towards the *trans* assembly.

$C3^Q$  exhibited a set of dominant signals in the  $^1H$  NMR spectrum (Fig. 5c). The diffusion coefficient of this major species, derived from DOSY, corresponded to a solvodynamic radius of 15 Å, whilst isotopic patterns observed by ESI-MS were

consistent with assemblies possessing the anticipated  $Pd_3L_6$  formulation (Fig. S139-146). This major species was estimated by integration of  $^1H$  NMR signals to constitute ~35% of the isomeric mixture (Fig. S138), over three-fold that of a statistical library (11%). As with the previous systems, the high symmetry of the NMR spectra and through-space interactions observed by NOESY ( $H_a \cdots H_n$  and  $H_b \cdots H_m$ ) were consistent with only the all-*cis* or all-*trans* isomers (Fig. 5b).



**Fig. 5** a) Ligands  $L3^Q$  and  $L3^P$ ; b) cartoons of  $Pd_3L_6$  'double-walled' triangle isomers with all-*cis* and all-*trans* donor arrangements; c)  $^1H$  NMR (500 MHz,  $d_6$ -DMSO) of  $C3^Q$ ; SCXRDR structure of *cis*- $C3^Q$  showing d) *cis*- $C3^Q$ →DMF<sub>2</sub>DMSO, and space filling representation e) from the top, and f) from the side; g)  $^1H$  NMR (500 MHz,  $CD_3CN$ ) and h)  $^{13}C$  NMR (126 MHz,  $CD_3CN$ ) of  $C3^P$ .

Weakly diffracting crystals were grown that required the use of synchrotron radiation to obtain

satisfactory SCXRD data.<sup>29</sup> The solid-state structure (Fig. 5d-f), however, unambiguously revealed each pair of ligands along the three “walls” to be aligned parallel, with a ‘head-to-tail’ arrangement of the ligands on each face of the triangle. This gave, at each of the three metal nodes, the anticipated *cis* arrangement of quinoline and pyridine donors.

In contrast to **L3<sup>Q</sup>**, the equilibrated self-assembly mixture of **L3<sup>P</sup>** and Pd(II) in *d*<sub>6</sub>-DMSO resulted in a <sup>1</sup>H NMR spectrum that appeared to show formation of a mixture of assemblies, with no clear dominant species. Given the established impact of solvent on the equilibrium between *cis* and *trans* isomers, self-assembly with **L3<sup>P</sup>** was re-examined in CD<sub>3</sub>CN. This yielded a much simpler <sup>1</sup>H NMR spectrum (Fig. 5g), with a clear dominant species (~25% of mixture, Fig. S175).

The major product was of high symmetry, as observed by both <sup>1</sup>H (Fig. 5g) and <sup>13</sup>C NMR (Fig. 5h) which, combined with NOE interactions between the pyridyl and picolyl donor groups (H<sub>b</sub>...H<sub>i</sub>; Fig. S174) and ESI-MS data (Fig. S176-186) again indicated formation of either the *cis*- or *trans*-Pd<sub>3</sub>L<sub>6</sub> assembly as the major species. Despite multiple attempts, single crystals suitable for X-ray diffraction could not be obtained in our hands and the lack of prochiral units prevented the use of NMR spectroscopy to distinguish between the two possible isomers. Consequently, we turned to DFT to investigate the relative energies of the *cis* and *trans* assemblies.

Unexpectedly, use of different functionals for the geometry optimisations resulted in a switching of the *cis* (HSE) or *trans* (PBE0) **C3<sup>P</sup>** isomer being lower in energy (ESI section S3). It has been demonstrated how environmental perturbations significantly impact the stability of the individual isomers. Thus, without the suitable inclusion of explicit encapsulated and exohedral solvent molecules and anions within these models, the balance of calculated energies between isomers can be easily swayed. Based on the experimental data obtained for **C1**, and the DFT calculations previously performed on the more rigid **C2** systems, it is tentatively suggested that the most likely identity of the major isomer of **C3<sup>P</sup>** is the *trans* assembly.

### Heteroleptic self-assembly

Preliminary investigations to extend these CSE designs to Pd<sub>4</sub>L<sub>8</sub> ‘double-walled’ tetrahedra<sup>37,39,40</sup> assembled from ligands **L4** (Fig. 6a) – synthesised through copper-free Sonogashira couplings<sup>41</sup> – proved prohibitively difficult, with complex NMR spectra obtained from equilibrated mixtures with Pd(II) (ESI Section S2.22 and S2.24). This is perhaps unsurprising. In a statistical library, each of the 35 possible isomers<sup>42</sup> would constitute <3% of

the mixture, with even the most symmetrical cages (i.e. *cis* and *trans*) possessing two ligand environments. Without near quantitative selectivity, identification of NMR signals for a particular isomer would likely be an insurmountable challenge. Consequently, further exploration of these systems was not attempted.

Inspired by a recent report of a heteroleptic *cis*-Pd<sub>2</sub>L<sub>2</sub>L'<sub>2</sub> MOC,<sup>40</sup> there was motivation to investigate the potential integrative self-assembly between **L4** and symmetric ligand **L2<sup>H</sup>**. From such a mixed-ligand assembly there would be two possible isomers: the *syn* and the (chiral) *anti* isomer, with the pair of **L4** ligands arranged in the same or opposite directions, respectively (Fig. 6c).


After equilibrating a stoichiometric mixture of **L2<sup>H</sup>** and **L4<sup>Q</sup>** with Pd(II) in CD<sub>3</sub>CN, DOSY (Fig. S224) and ESI-MS (Fig. S226-230) confirmed formation of heteroleptic assemblies with the anticipated Pd<sub>2</sub>(**L2<sup>H</sup>**)<sub>2</sub>(**L4<sup>Q</sup>**)<sub>2</sub> formulation. Meanwhile, NMR analysis demonstrated the presence of two spectroscopically similar species in an approximately 3:1 ratio (Fig. 6b and S225).

The lack of bilateral symmetry in **L4<sup>Q</sup>** induced a lowering of the symmetry of **L2<sup>H</sup>** in both cage isomers, resulting in distinct signals for all 12 protons, corroborated by the <sup>13</sup>C NMR spectrum (Fig. S211). Whilst analysis of the <sup>1</sup>H NMR spectrum was made challenging by significant signal overlap, 2D NMR techniques (COSY, HMBC and TOCSY) enabled assignment of all peaks for both isomers (Fig. S205).

NOESY and ROESY were employed to determine the identities of the two isomers (Fig. 6c). Observation of particular through-space interactions (H<sub>B</sub>...H<sub>B'</sub> and H<sub>B</sub>...H<sub>A</sub>...H<sub>B</sub>) for the minor species (Fig. 6d) led to the conclusion that this was the less sterically congested *anti* isomer. The more limited NOE interactions observed for the major assembly demonstrated that this was the *syn* isomer, with both **L4<sup>Q</sup>** ligands arranged parallel. This result further supported the conclusion that, rather than selectivity towards the *cis* isomers of **C1-C3** being purely driven by repulsive steric effects, additional stabilising interactions promoted this ligand arrangement. In this instance, presumably the two different second coordination sphere sites around the Pd(II) ions in the *syn* isomer provide more favourable interactions with solvent molecules compared to those of the *anti* cage.

The anticipated structure of *syn*-Pd<sub>2</sub>(**L2<sup>H</sup>**)<sub>2</sub>(**L4<sup>Q</sup>**)<sub>2</sub> was confirmed by SCXRD, with both **L4<sup>Q</sup>** ligands arranged parallel to each other (Fig. 6e-g).<sup>29</sup> The solid-state structure of the heteroleptic cage was found with one BF<sub>4</sub><sup>-</sup> anion encapsulated within the cavity, and external counterions interacting

with the external face of the coordination spheres around the two Pd(II) ions (Fig. 6e).



**Fig. 6** a) Ligands  $\mathbf{L4^Q}$ ,  $\mathbf{L4^P}$  and  $\mathbf{L2^H}$ ; b)  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{CN}$ ) of  $[\text{Pd}_2(\mathbf{L2^H})_2(\mathbf{L4^Q})_2](\text{BF}_4)_4$  with major *syn* isomer peaks labelled; c) through-space interactions observed by NOESY in the two heteroleptic isomers; d) NOESY (500 MHz,  $\text{CD}_3\text{CN}$ ) with minor *anti* isomer peaks labelled. SCXRD structure of *syn*- $\text{Pd}_2(\mathbf{L2^H})_2(\mathbf{L4^Q})_2$  e) with endo- and exohedral  $\text{BF}_4^-$  counteranions, f) viewed down the Pd...Pd axis, and g) second crystallographically independent molecule with alternative conformation of internal  $\text{BF}_4^-$  anion.

The integrative self-assembly between  $\mathbf{L2^H}$  and  $\mathbf{L4^P}$  was also attempted. Whilst ESI-MS (Fig. S233-241) demonstrated the presence of the heteroleptic assembly (alongside minor signals for the homoleptic  $\text{Pd}_2(\mathbf{L2^H})_4$ ), signal resolution in the NMR spectrum was insufficient to enable effective analysis (Fig. S232).

We have previously been able to arrange two different unsymmetrical ligand scaffolds in defined relative orientations through covalent tethering, forming *pseudo*-heteroleptic MOCs.<sup>43</sup> Using CSE, we now demonstrate the ability to assemble truly heteroleptic MOCs, derived through integrative self-assembly of ligands with the same denticity, incorporating unsymmetrical scaffolds in an orientationally-selective manner.<sup>44</sup>

## Conclusion

We have prepared a range of unsymmetrical ditopic ligands, with varying backbone scaffolds, incorporating a pyridine donor paired with a sterically bulky quinoline or picoline moiety, and investigated their self-assembly into  $\text{Pd}_n\text{L}_{2n}$  architectures. This coordination-sphere engineering approach was successful in biasing self-assembly towards specific isomers from a statistical library. Interestingly, quinoline and picoline units promoted *cis* and *trans* arrangements of donors at the metal centres, respectively, resulting in diastereoselectivity of the self-assembly process towards different isomers of particular architectures. The ability to use this relatively subtle difference to target the formation of specific metal-organic assembly isomers provides a nuanced approach towards directing the self-assembly of unsymmetrical ligand scaffolds.

After probing the source of this difference in selectivity, it was concluded that interactions between solvent molecules and the exterior of the cage around the metal nodes play a crucial role in determining the relative stabilities of isomers. Thus, not only is the first coordination sphere important in directing the self-assembly, but second coordination sphere effects play a critical role and can, in fact, supersede the directing effects of the primary structure. This insight further opens up the potential for designing stimuli-responsive,<sup>45</sup> shape-shifting systems<sup>46</sup> that respond to changes in their environment.

The orientationally selective incorporation of an unsymmetrical ligand scaffold into a heteroleptic  $\text{M}_2\text{L}_2\text{L}'_2$  MOC was also demonstrated. Consistent with the homoleptic assemblies, the major isomer that formed had both bulky quinoline donor units coordinating to the same Pd(II) ion. This result

further supported the hypothesis that isomer selectivity can be affected by interactions beyond those simply between components within the covalent structure.

This flexible strategy adds a new approach to preparing metal-organic hosts with increased anisotropy to the metallosupramolecular chemist's toolbox. The continued development of methods to access more structurally sophisticated metal-organic cages<sup>6</sup> will lead to supramolecular hosts exhibiting higher-level behaviours,<sup>7</sup> reminiscent of the impressive properties of natural architectures, like enzymes, that have long provided a source of inspiration for chemists.

## Author contributions

PM – synthesis and characterisation, analysis; LM – SCXRD analysis; AT and KEJ – computational modelling; JEML – synthesis and characterisation, analysis, conceptualisation, supervision, writing – original draft. All authors contributed to reviewing of the final manuscript.

## Acknowledgements

Special thanks to Dr Georgia Orton for assistance with the collection of synchrotron SCXRD data, Dr Cécile Le Duff for assistance with the collection of NMR data, Dr Christopher Williams for the collection of MS data, and Prof. Steve Goldup for useful discussions. We acknowledge Diamond Light Source for time on Beamline I19 under Proposal CY28766. AT acknowledges funding received from the European Union under the NextGenerationEU program (grant CAGEX, SOE\_0000033). KEJ acknowledges the Royal Society for a University Research Fellowship and Enhancement Award and the ERC through Agreement No. 758370 (ERC-StG-PE5-CoMMaD). This work used the ARCHER2 UK National Supercomputing Service (<https://www.archer2.ac.uk>) via our membership of the UK's Materials Chemistry HEC Consortium, which is funded by EPSRC (EP/X035859/1), and Imperial College London's Research Computing Service, DOI: 10.14469/hpc/2232. This work was supported by the Royal Society (URFAR1\221740 and R\FERE\221016) and the University of Birmingham. JEML is a Royal Society University Research Fellow.


**Keywords:** low-symmetry • coordination-sphere • coordination cages • self-assembly • metallosupramolecular

## References

- D. L. Caulder, K. N. Raymond, *Acc. Chem. Res.* **1999**, *32*, 875-982; R. Chakrabarty, P. S. Mukherjee, P. J. Stang, *Chem. Rev.* **2011**, *111*, 6810-6918; K. Harris, D. Fujita, M. Fujita, *Chem. Commun.* **2013**, *49*, 6703-6712; M. Han, D. M. Engelhard, G. H. Clever, *Chem. Soc. Rev.* **2014**, *43*, 1848-1860; M. Yoshizawa, M. Yamashina, *Chem. Lett.* **2017**, *46*, 163-171; R. A. S. Vasdev, D. Preston, J. D. Crowley, *Chem. Asian J.* **2017**, *12*, 2513-2523; D. Zhang, T. K. Ronson, J. R. Nitschke, *Acc. Chem. Res.* **2018**, *51*, 2423-2436; P. M. Bogie, T. F. Miller, R. J. Hooley, *Isr. J. Chem.*, **2019**, *59*, 130-139; B. S. Pilgrim, N. R. Champness, *ChemPlusChem*, **2020**, *85*, 1842-1856; J. E. M. Lewis, *Chem. Commun.* **2022**, *58*, 13873-13886; T. Tateishi, M. Yoshimura, S. Tokuda, F. Matsuda, D. Fujita, S. Furukawa, *Coord. Chem. Rev.* **2022**, *467*, 414612; A. J. McConnell, *Chem. Soc. Rev.* **2022**, *51*, 2957-2971; R. Banerjee, D. Chakraborty, P. S. Mukherjee, *J. Am. Chem. Soc.* **2023**, *145*, 7692-7711.
- C. J. Brown, F. D. Toste, R. G. Bergman, K. N. Raymond, *Chem. Rev.* **2015**, *115*, 3012-3035; C. Tan, D. Chu, X. Tang, Y. Liu, W. Xuan, Y. Cui, *Chem. Eur. J.*, **2019**, *25*, 662-672; Y. Xue, X. Hang, J. Ding, B. Li, R. Zhu, H. Pang, Q. Xu, *Coord. Chem. Rev.* **2021**, *430*, 213656; R. Ham, C. J. Nielsen, S. Pullen, J. N. H. Reek, *Chem. Rev.* **2023**, *123*, 5225-5261; T. K. Piskorz, V. Marti-Centelles, R. L. Spicer, F. Duarte, P. J. Lusby, *Chem. Sci.* **2023**, DOI: 10.1039/D3SC02586A. For recent examples, see: J. Wang, T. A. Young, F. Duarte, P. J. Lusby, *J. Am. Chem. Soc.* **2020**, *142*, 17743-17750; M. D. Ludden, C. G. P. Taylor, M. B. Tipping, J. S. Train, N. H. Williams, J. C. Dorrat, K. L. Tuck, M. D. Ward, *Chem. Sci.* **2021**, *12*, 14781-14791; A. C. Ghosh, A. Legrand, R. Rajapaksha, G. A. Craig, C. Sassoey, G. Balázs, D. Farrusseng, S. Furukawa, J. Canivet, F. M. Wissner, *J. Am. Chem. Soc.* **2022**, *144*, 3626-3636; D. O. Bobylev, J. Ruijter, D. A. Poole III, S. Matthew, B. de Bruin, J. N. H. Reek, *Angew. Chem. Int. Ed.* **2023**, *62*, e202218162; Y. Yang, X. Jing, Y. Shi, Y. Wu, C. Duan, *J. Am. Chem. Soc.* **2023**, *145*, 10136-10148; G. R. Genov, H. Takezawa, H. Hayakawa, M. Fujita, *J. Am. Chem. Soc.* **2023**, *145*, 17013-17017.
- N. Dey, C. J. E. Haynes, *ChemPlusChem* **2021**, *86*, 418-433 A. Brzechwa-Chodzyńska, W. Drożdż, J. Harrowfield, A. R. Stefankiewicz, *Coord. Chem. Rev.* **2021**, *434*, 213820. For recent examples, see: Z. Zhang, Z. Zhao, L. Wu, S. Lu, S. Ling, G. Li, L. Xu, L. Ma, Y. Hou, X. Wang, X. Li, G. He, K. Wang, B. Zhou, M. Zhang, *J. Am. Chem. Soc.* **2020**, *142*, 2592-2600; Z. Meng, F. Yang, X. Wang, W.-L. Shan, D. Liu, L. Zhang, G. Yuan, *Inorg. Chem.* **2023**, *62*, 1297-1305; Y. Wang, J. Liu, Y. Niu, L. He, Y. Wang, Y. Ma, Y. Yao, J. Ye, *Adv. Optical Mater.* **2023**, *11*, 2202940; T.-P. Sheng, C.-Z. Sun, F.-R. Dai, *ACS Appl. Mater. Interface* **2023**, *15*, 29252-29258; Y. He, D. Luo, V. M. Lynch, M. Ahmed, J. L. Sessler, X. Chi, *Chem* **2023**, *9*, 93-101; O. Calvo-Lozano, L. Hernández-López, L. Gomez, A. Carné-Sánchez, C. von Baeckmann, L. M. Lechuga, D. Maspoch, *ACS Appl. Mater. Interfaces* **2023**, *15*, 39523-39529.
- A. Casini, B. Woods, M. Wenzel, *Inorg. Chem.* **2017**, *56*, 14715-14729; S. K. Samanta, L. Isaacs, *Coord. Chem. Rev.* **2020**, *410*, 213181; W.-T. Dou, C.-Y. Yang, L.-R. Hu, B. Song, T. Jin, P.-P. Jia, X. Ji, F. Zheng, H.-B. Yang, L. Xu, *ACS Materials Lett.* **2023**, *5*, 1061-1082; B. Therrien, G. Süß-Fink, P. Govindaswamy, A. K. Renfrew, P. J. Dyson, *Angew. Chem. Int. Ed.* **2008**, *47*, 3773-3776; J. E. M. Lewis, E. L. Gavey, S. A. Cameron, J. D. Crowley, *Chem. Sci.* **2012**, *3*, 778-784; E. O. Bobylev, Y. Zeng, K. Weijertse, E. Koelman, E. M. Meijer, B. de Bruin, A. Kros, J. N. H. Reek, *Chem* **2023**, *9*, 1578-1593.
- A. Galan, P. Ballester, *Chem. Soc. Rev.* **2016**, *45*, 1720-1737; P. Mal, B. Breiner, K. Rissanen, J. R. Nitschke, *Science* **2009**, *324*, 1697-1699; M. Yamashina, Y. Sei, M. Akita, M. Yoshizawa, *Nat. Commun.* **2014**, *5*, 4662; M. Canton, A. B. Grommet, L. Pesce, J. Gemen, S. Li, Y. Diskin-Posner, A. Credi, G. M. Pavan, K.

- Andréasson, R. Klajn, *J. Am. Chem. Soc.* **2020**, *142*, 14557-14565; R. Sumida, Y. Tanaka, K. Niki, Y. Sei, S. Toyota, M. Yoshizawa, *Chem. Sci.* **2021**, *12*, 9946-9951; S. Hasegawa, S. L. Meichsner, J. J. Holstein, A. Baksi, M. Kasanmascheff, G. H. Clever, *J. Am. Chem. Soc.* **2021**, *143*, 9718-9723; M. Yuasa, R. Sumida, Y. Tanaka, M. Yoshizawa, *Chem. Eur. J.* **2022**, *28*, e202104101; J. C. Dorrat, R. J. Young, C. G. P. Taylor, M. B. Tipping, A. J. Blok, D. R. Turner, A. I. McKay, S. Ovenden, M. D. Ward, G. H. Dennison, K. L. Tuck, *Dalton Trans.* **2023**, *52*, 11802-11814.
- <sup>6</sup> S. Pullen, J. Tessarolo, G. H. Clever, *Chem. Sci.* **2021**, *12*, 7269-7293.
- <sup>7</sup> J. E. M. Lewis, *Trends Chem.* **2023**, *5*, 717-719; A. C. Percy, J. D. Crowley, *Chem. Eur. J.* **2023**, *29*, e202203752.
- <sup>8</sup> C. T. McTernan, J. A. Davies, J. R. Nitschke, *Chem. Rev.* **2022**, *122*, 10393-10437; J. L. Algar, D. Preston, *Chem. Commun.* **2022**, *58*, 11637-11648.
- <sup>9</sup> W. M. Bloch, G. H. Clever, *Chem. Commun.* **2017**, *53*, 8506-8516; L. R. Holloway, P. M. Bogie, R. J. Hooley, *Dalton Trans.* **2017**, *46*, 14719-14723; D. Bardhan, D. K. Chand, *Chem. Eur. J.* **2019**, *25*, 12241-12269; C.-B. Tian, Q.-F. Sun, *Chem. Eur. J.* **2023**, *29*, e202300195. For recent examples, see: D. Zhang, T. K. Ronson, L. Xu, J. R. Nitschke, *J. Am. Chem. Soc.* **2020**, *142*, 9152-9157; L. He, S.-C. Wang, L.-T. Lin, J.-Y. Cai, L. Li, T.-H. Tu, Y.-T. Chan, *J. Am. Chem. Soc.* **2020**, *142*, 7134-7144; R.-J. Li, F. Fadaei-Tirani, R. Scopelliti, K. Severin, *Chem. Eur. J.* **2021**, *27*, 9439-9445; Y. Liu, S.-H. Liao, W.-T. Dai, Q. Bai, S. Lu, H. Wang, X. Li, Z. Zhang, P. Wang, W. Lu, Q. Zhang, *Angew. Chem. Int. Ed.* **2023**, *62*, e202217215; T. Abe, N. Sanada, K. Takeuchi, A. Okazawa, S. Hiraoka, *ChemRxiv* **2023**, 10.26434/chemrxiv-2023-hht0b; K. Wu, E. Benchimol, A. Baksi, G. H. Clever, *ChemRxiv* **2023**, 10.26434/chemrxiv-2023-5gb4q.
- <sup>10</sup> F. Li, L. F. Lindoy, *Aus. J. Chem.* **2019**, *72*, 731-741; M. Hardy, A. Lützen, *Chem. Eur. J.* **2020**, *26*, 13332-13346. For recent examples, see: S. M. Jansze, M. D. Wise, A. V. Vologzhanina, R. Scopelliti, K. Severin, *Chem. Sci.* **2017**, *8*, 1901-1908; C. Shen, A. D. W. Kennedy, W. A. Donald, A. M. Torres, W. S. Price, J. E. Beves, *Inorg. Chim. Acta* **2017**, *458*, 122-128; D. Preston, J. J. Sutton, K. C. Gordon, J. D. Crowley, *Angew. Chem. Int. Ed.* **2018**, *57*, 8659-8663; D. Yang, J. L. Greenfield, T. K. Ronson, L. K. S. von Krbek, L. Yu, J. R. Nitschke, *J. Am. Chem. Soc.* **2020**, *142*, 19856-19861; L. S. Lisboa, J. A. Findlay, L. J. Wright, C. G. Hartinger, J. D. Crowley, *Angew. Chem. Int. Ed.* **2020**, *59*, 11101-11107; M. T. Yong, O. M. Linder-Patton, W. M. Bloch, *Inorg. Chem.* **2022**, *61*, 12863-12869; L. S. Lisboa, D. Preston, C. J. McAdam, L. J. Wright, C. G. Hartinger, J. D. Crowley, *Angew. Chem. Int. Ed.* **2022**, *61*, e202201700; Q.-Y. Zhu, L.-P. Zhou, L.-X. Cai, S.-J. Hu, X.-Z. Li, Q.-F. Sun, *Inorg. Chem.* **2022**, *61*, 16814-16821; L. S. Lisboa, M. Riisom, H. J. Dunne, D. Preston, S. M. F. Jamieson, L. J. Wright, C. G. Hartinger, J. D. Crowley, *Dalton Trans.* **2022**, *51*, 18438-18445.
- <sup>11</sup> J. E. M. Lewis, J. D. Crowley, *ChemPlusChem* **2020**, *85*, 815-827; D. Tripathy, N. B. Debata, K. C. Naik, H. S. Sahoo, *Coord. Chem. Rev.* **2022**, *456*, 214396.
- <sup>12</sup> W. M. Bloch, Y. Abe, J. J. Holstein, C. M. Wandtke, B. Dittrich, G. H. Clever, *J. Am. Chem. Soc.* **2016**, *138*, 13750-13755
- <sup>13</sup> A. C. Percy, L. S. Lisboa, D. Preston, N. B. Page, T. Lawrence, L. J. Wright, C. Hartinger, J. D. Crowley, *Chem. Sci.* **2023**, *14*, 8615-8623.
- <sup>14</sup> D. Preston, P. E. Kruger, *ChemPlusChem* **2020**, *85*, 454-465.
- <sup>15</sup> J. E. M. Lewis, A. Tarzia, A. J. P. White, K. E. Jelfs, *Chem. Sci.* **2020**, *11*, 677-683.
- <sup>16</sup> D. Ogata, J. Yuasa, *Angew. Chem. Int. Ed.* **2019**, *58*, 18424-18428; S. S. Mishra, S. V. K. Kompella, S. Krishnaswamy, S. Balasubramanian, D. K. Chand, *Inorg. Chem.* **2020**, *59*, 12884-12894; J. E. M. Lewis, *Chem. Eur. J.* **2021**, *27*, 4454-4460; A. Tarzia, J. E. M. Lewis, K. E. Jelfs, *Angew. Chem. Int. Ed.* **2021**, *60*, 20879-20887; H. Yu, J. Li, C. Shan, T. Lu, X. Jiang, J. Shi, L. Wojitas, H. Zhang, M. Wang, *Angew. Chem. Int. Ed.* **2021**, *60*, 26523-26527; J. Shi, Y. Li, X. Jiang, H. Yu, J. Li, H. Zhang, D. J. Trainer, S. W. Hla, H. Wang, M. Wang, X. Li, *J. Am. Chem. Soc.* **2021**, *143*, 1224-1234; R.-J. Li, A. Tarzia, V. Posligua, K. E. Jelfs, N. Sanchez, A. Marcus, A. Baksi, G. H. Clever, F. Fadaei-Tirani, K. Severin, *Chem. Sci.* **2022**, *13*, 11912-11917; S. S. Mishra, D. K. Chand, *Dalton Trans.* **2022**, *51*, 11650-11657.
- <sup>17</sup> A. M. Johnson, R. J. Hooley, *Inorg. Chem.* **2011**, *50*, 4671-4673; S. K. Sen, R. Natarajan, *Inorg. Chem.*, **2019**, *58*, 7180-7188.
- <sup>18</sup> M. Yoshizawa, M. Nagao, K. Kumazawa, M. Fujita, *J. Organomet. Chem.* **2005**, *690*, 5383-5388.
- <sup>19</sup> R. A. S. Vasdev, D. Preston, C. A. Casey-Stevens, V. Martí-Centelles, P. J. Lusby, A. L. Garden, J. D. Crowley, *Inorg. Chem.* **2023**, *62*, 1833-1844.
- <sup>20</sup> L. Zhao, B. H. Northrop, Y. R. Zheng, H. B. Yang, J. L. Hyo, M. L. Young, Y. P. Joo, K. W. Chi, P. J. Stang, *J. Org. Chem.* **2008**, *73*, 6580-6586; M. L. Saha, K. Mahata, D. Samanta, V. Kalsani, J. Fan, J. W. Bats, M. Schmittel, *Dalton Trans.* **2013**, *42*, 12840-12843.
- <sup>21</sup> M. Yoshizawa, J. Nakagawa, K. Kumazawa, M. Nagao, M. Kawano, T. Ozeki, M. Fujita, *Angew. Chem. Int. Ed.* **2005**, *44*, 1810-1813; S. E. Walker, S. A. Boer, T. Malcomson, M. J. Paterson, K. L. Tuck, D. A. Turner, *Chem. Commun.* **2021**, *57*, 12456-12459.
- <sup>22</sup> R. Zhu, W. M. Bloch, J. J. Holstein, S. Mandal, L. V. Schäfer, G. H. Clever, *Chem. Eur. J.* **2018**, *24*, 12976-12982.
- <sup>23</sup> D. Preston, J. E. Barnsley, K. C. Gordon, J. D. Crowley, *J. Am. Chem. Soc.* **2016**, *138*, 10578-10585.
- <sup>24</sup> R.-J. Li, J. Tessarolo, H. Lee, G. H. Clever, *J. Am. Chem. Soc.* **2021**, *143*, 3865-3873; B. Chen, J. J. Holstein, A. Platzek, L. Schneider, K. Wu, G. H. Clever, *Chem. Sci.* **2022**, *13*, 1829-1834.
- <sup>25</sup> Recently, an  $M_4L_4$  tetrahedron has been shown to undergo guest-induced rearrangement from an all-*mer* configuration with  $S_4$  symmetry to all-*fac* with  $T$  symmetry: W. Xue, K. Wu, N. Ouyang, T. Brotin, J. R. Nitschke, *Angew. Chem. Int. Ed.* **2023**, *62*, e202301319.
- <sup>26</sup> S. Samantray, S. Krishnaswamy, D. K. Chand, *Nat. Commun.* **2020**, *11*, 880.
- <sup>27</sup> S. Bandi, A. K. Pal, G. S. Hanan, D. K. Chand, *Chem. Eur. J.* **2014**, *20*, 13122-13126; S. Bandi, S. Samantray, R. D. Chakravarthy, A. K. Pal, G. S. Hanan, D. K. Chand, *Eur. J. Inorg. Chem.* **2016**, 2816-2827.
- <sup>28</sup> Encapsulation of different anions has previously been shown to alter the diastereoselectivity within a tetrahedral MOC: J. K. Clegg, J. Cremers, A. J. Hogben, B. Breiner, M. M. J. Smulders, J. D. Thoburn, J. R. Nitschke, *Chem. Sci.* **2013**, *4*, 68-76.

- <sup>29</sup> Deposition numbers 2300291 (C1<sup>Q</sup>), 2300288 (C1<sup>P</sup>), 2300289 (C2<sup>Q</sup>), 2300292 (C3<sup>Q</sup>), and 2300290 ([Pd<sub>2</sub>(L2<sup>H</sup>)<sub>2</sub>(L4<sup>Q</sup>)<sub>2</sub>](BF<sub>4</sub>)<sub>4</sub>) contain the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structure service.
- <sup>30</sup> J. E. M. Lewis, J. D. Crowley, *Supramol. Chem.* **2014**, *26*, 173-181.
- <sup>31</sup> C. A. Hunter, *Angew. Chem. Int. Ed.* **2004**, *43*, 5310-5324.
- <sup>32</sup> Encapsulation of different anions has previously been shown to alter the helical pitch of M<sub>2</sub>L<sub>4</sub>-type MOCs: L.-P. Zhou, Q.-F. Sun, *Chem. Commun.* **2015**, *51*, 16767-16770; J.-Y. Wu, M.-S. Zhong, M.-H. Chiang, *Chem. Eur. J.* **2017**, *23*, 15957-15965; P. J. Steel, D. M. McMorran, *Chem. Asian J.* **2019**, *14*, 1098-1101; W. M. Bloch, S. Horiuchi, J. J. Holstein, C. Drechsler, A. Wuttke, W. Hiller, R. A. Mata, G. H. Clever, *Chem. Sci.* **2023**, *14*, 1524-1531.
- <sup>33</sup> W. Liu, P. J. Das, H. M. Colquhoun, J. F. Stoddart, *CCS Chem.* **2022**, *4*, 755-784; M. W. Drover, *Chem. Soc. Rev.* **2022**, *51*, 1861-1880.
- <sup>34</sup> D. Zhang, T. K. Ronson, S. Güryel, J. D. Thoburn, D. J. Wales, J. R. Nitschke, *J. Am. Chem. Soc.* **2019**, *141*, 14534-15438; T. Tsutsui, L. Catti, K. Yoza, M. Yoshizawa, *Chem. Sci.* **2020**, *11*, 8145-8150; J. L. Algar, J. A. Findley, J. D. Evans, D. Preston, *Angew. Chem. Int. Ed.* **2022**, *61*, e202210476.
- <sup>35</sup> M. Hutin, G. Bernardinelli, J. R. Nitschke, *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 17655-17660.
- <sup>36</sup> K. Suzuki, M. Kawano, M. Fujita, *Angew. Chem. Int. Ed.* **2007**, *46*, 2819-2811; R. A. S. Vasdev, J. A. Findlay, D. R. Turner, J. D. Crowley, *Chem. Asian J.* **2021**, *16*, 39-43.
- <sup>37</sup> D. K. Chand, K. Biradha, M. Kawano, S. Sakamoto, K. Yamaguchi, M. Fujita, *Chem. Asian J.* **2006**, *1*, 82-90; J. Tessarolo, H. Lee, E. Sakuda, K. Umakoshi, G. H. Clever, *J. Am. Chem. Soc.* **2021**, *143*, 6339-6344; A. Kumar, R. Banerjee, E. Zangrando, P. S. Mukherjee, *Inorg. Chem.* **2022**, *61*, 2368-2377; A. D. W. Kennedy, R. G. DiNardi, L. L. Fillbrook, W. A. Donald, J. E. Beves, *Chem. Eur. J.* **2022**, *28*, e202104461.
- <sup>38</sup> S. S. Mishra, D. K. Chand, *Dalton Trans.* **2022**, *51*, 11650-11657.
- <sup>39</sup> C. Klein, C. Gütz, M. Bogner, F. Topić, K. Rissanen, A. Lützen, *Angew. Chem. Int. Ed.* **2014**, *53*, 3739-3742; J. A. Findlay, K. M. Patil, M. G. Gardiner, H. I. MacDermott-Opeskin, M. L. O'Mara, P. E. Kruger, D. Preston, *Chem. Asian J.* **2022**, *17*, e202200093.
- <sup>40</sup> R.-J. Li, J. de Montmollin, F. Fadaei-Tirani, R. Scopelliti, K. Severin, *Dalton Trans.* **2023**, *52*, 6451-6456.
- <sup>41</sup> A. Mori, J. Kawashima, T. Shimada, M. Suguro, K. Hirabayashi, Y. Nishihara, *Org. Lett.* **2000**, *2*, 2935-2937; Y. Liang, Y.-X. Xie, J.-H. Li, *J. Org. Chem.* **2006**, *71*, 379-381; M. Lehr, T. Paschelke, V. Bendt, A. Petersen, L. Pietsch, P. Harders, A. J. McConnell, *Eur. J. Org. Chem.* **2021**, 2728-2735.
- <sup>42</sup> R.-J. Li, A. Marcus, F. Fadaei-Tirani, K. Severin, *Chem. Commun.* **2021**, *57*, 10023-10026.
- <sup>43</sup> J. E. M. Lewis, *Angew. Chem. Int. Ed.* **2022**, *61*, e202212392.
- <sup>44</sup> The incorporation of chiral ligands into heteroleptic cages has been recently reported: K. Wu, J. Tessarolo, A. Baksi, G. H. Clever, *Angew. Chem. Int. Ed.* **2022**, *61*, e2022057; J. Tessarolo, E. Benchimol, A. Jouaiti, M. W. Hosseini, G. H. Clever, *Chem. Commun.* **2023**, *59*, 3467-4370.
- <sup>45</sup> A. J. McConnell, C. S. Wood, P. P. Neelakandan, J. R. Nitschke, *Chem. Rev.* **2015**, *115*, 7729-7793.
- <sup>46</sup> A. E. Martín Díaz, J. E. M. Lewis, *Front. Chem.* **2021**, *9*, 706462.



Often overlooked second coordination sphere effects are shown to be of significant importance in isomer selectivity with self-assembled, low-symmetry metal-organic cages. In some instances, these effects override primary structural factors that are often the only consideration when designing such systems.

@LewisChemistry  
@andrew\_tarzia  
@JelfsChem  
@UoBChemistry  
@unibirmingham