

Short-Term and Long-Term Effects of Cement Kiln Dust Stabilization of Construction and Demolition Waste

Original

Short-Term and Long-Term Effects of Cement Kiln Dust Stabilization of Construction and Demolition Waste / Bassani, M., Riviera, P.P., Tefa, L.. - In: JOURNAL OF MATERIALS IN CIVIL ENGINEERING. - ISSN 0899-1561. - STAMPA. - 29:5(2017), pp. 1-11. [10.1061/(ASCE)MT.1943-5533.0001797]

Availability:

This version is available at: 11583/2657615 since: 2016-11-25T09:02:46Z

Publisher:

American Society of Civil Engineering

Published

DOI:10.1061/(ASCE)MT.1943-5533.0001797

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

ASCE postprint/Author's Accepted Manuscript

This material may be downloaded for personal use only. Any other use requires prior permission of the American Society of Civil Engineers. This material may be found at [http://dx.doi.org/10.1061/\(ASCE\)MT.1943-5533.0001797](http://dx.doi.org/10.1061/(ASCE)MT.1943-5533.0001797).

(Article begins on next page)

Biocorrosion of speleothems in show caves: analysis of young speleothems exposed to two years of new LED lights

Valentina Balestra*^{†1,2,3}, Aurora Peruzzi⁴, Paola Marini¹, and Rossana Bellopede¹

¹Department of Environment, Land and Infrastructure Engineering, Politecnico di Torino – Italy

²Biologia Sotterranea Piemonte - Gruppo di Ricerca – Italy

³Underground Karst Laboratory of Bossea Cave - S.O. Bossea C.A.I. – Italy

⁴Politecnico di Torino – Italy

Abstract

Speleothems in show caves are the most important attractions, however, they are often subjected to tourism-driven alterations, including corrosion due to CO₂ increase, undersaturated water and photosynthetic biofilms. The so-called "lampenflora" is a community of photosynthetic microorganisms, mainly cyanobacteria, diatoms and green algae, but may also include mosses, ferns and higher plants. It develops on speleothems in show caves, following the installation of artificial lights, which allow the photosynthetic activity in an otherwise dark environment. The growth of the lampenflora causes physical, chemical and aesthetic damage to speleothems and changes the underground ecosystem, as it modifies the food chain and can therefore damage the organisms that naturally inhabit the cave itself. Moreover, the biochemical reactions given by the metabolic activity of microorganisms can release substances that lead to the deterioration of the rock, resulting in erosion and damage to speleological formations.

Bossea cave, Piedmont, Italy, was opened to the public in 1874, and receives about 12,000 tourists/year. The cavity develops for about 2800 m in a tectonic contact between dolostone, carbonate rocks and metavolcanics and it is crossed by a subterranean river. Different underground karst laboratories to study hydrogeology, climatology, radon activity and subterranean biology are located in the cave, managed by S.O. Bossea CAI, the DIATI of the Politecnico di Torino, and Biologia Sotterranea Piemonte – Gruppo di Ricerca. In the Bossea cave, the presence of lampenflora and biocorrosion on speleothems along the tourist path was previously documented. A new LED lights system was installed in 2021, which should allow a slowdown of the phenomenon of lampenflora.

This study aims at observing the growth of lampenflora with the new LED lamps, evaluating changes in calcite crystal habits over time. A homogenous slab of speleothem recently formed as a patina on a cover of the central underground laboratory instrumentation was collected, divided into several parts and observed with a Scanning Electron Microscopy (SEM). After, 17 small slab parts were placed along the tourist path in six different areas, from one to seven parts per point, positioned with different inclinations, where they remained for two years. These areas were chosen taking into account the presence of lampenflora due to the previous halogenic lamps. Then, the samples were collected to verify the degree of biocorrosion. For each sample, different points were analysed with SEM and classified according to a predetermined index.

*Speaker

†Corresponding author: valentina.balestra@polito.it

Keywords: show caves, lampenflora, LED lights, biocorrosion, speleothems