

Groundwater: An Endangered Invisible Resource. Microplastic Pollution in Underground Karst Systems, from Surface Watercourses to Cave Waters

*Original*

Groundwater: An Endangered Invisible Resource. Microplastic Pollution in Underground Karst Systems, from Surface Watercourses to Cave Waters / Balestra, Valentina; Vigna, Bartolomeo; Bellopede, Rossana. - ELETTRONICO. - (2023), pp. 89-95. ( 3rd International Conference on Microplastic Pollution in the Mediterranean Sea Naples 25-28 September 2022) [10.1007/978-3-031-34455-8\_13].

*Availability:*

This version is available at: 11583/2983691 since: 2024-06-18T14:50:09Z

*Publisher:*

Springer

*Published*

DOI:10.1007/978-3-031-34455-8\_13

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

Springer postprint/Author's Accepted Manuscript

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: [http://dx.doi.org/10.1007/978-3-031-34455-8\\_13](http://dx.doi.org/10.1007/978-3-031-34455-8_13)

(Article begins on next page)

# Metadata of the chapter that will be visualized in SpringerLink

Book Title	Proceedings of the 3rd International Conference on Microplastic Pollution in the Mediterranean Sea	
Series Title		
Chapter Title	Groundwater: An Endangered Invisible Resource. Microplastic Pollution in Underground Karst Systems, from Surface Watercourses to Cave Waters	
Copyright Year	2023	
Copyright HolderName	The Author(s), under exclusive license to Springer Nature Switzerland AG	
Corresponding Author	Family Name	<b>Balestra</b>
	Particle	
	Given Name	<b>Valentina</b>
	Prefix	
	Suffix	
	Role	
	Division	
	Organization	Politecnico Di Torino
	Address	Corso Duca Degli Abruzzi 24, Torino, Italy
	Email	valentina.balestra@polito.it
	ORCID	<a href="http://orcid.org/0000-0002-8529-468X">http://orcid.org/0000-0002-8529-468X</a>
Author	Family Name	<b>Vigna</b>
	Particle	
	Given Name	<b>Bartolomeo</b>
	Prefix	
	Suffix	
	Role	
	Division	
	Organization	Bartolomeo Vigna, Politecnico Di Torino
	Address	Rossana BellopedeCorso Duca Degli Abruzzi 24, Torino, Italy
	Email	
	ORCID	<a href="http://orcid.org/0000-0003-2272-8740">http://orcid.org/0000-0003-2272-8740</a>
Author	Family Name	<b>Bellopede</b>
	Particle	
	Given Name	<b>Rossana</b>
	Prefix	
	Suffix	
	Role	
	Division	
	Organization	Bartolomeo Vigna, Politecnico Di Torino
	Address	Rossana BellopedeCorso Duca Degli Abruzzi 24, Torino, Italy
	Email	
	ORCID	<a href="http://orcid.org/0000-0002-9569-0678">http://orcid.org/0000-0002-9569-0678</a>

## Abstract

Microplastic (MP) pollution has been found in environments worldwide, including remote areas [e.g. Zhang in *Sci Total Environ* 758, 2021, Ambrosini in *Environ Pollut* 253:297–301, 2019, Cincinelli in *Chemosphere* 175:391–400, 2017]. MPs have a dimension less than 5 mm and are a global problem because they are easily transportable even at long distances, tend to adsorb and transport hazardous substances such as heavy metals, pesticides, persistent organic pollutants (POPs) and antibiotics [e.g. Li in *Water Res* 157:228–237, 2019, Li et al. in *Environ Pollut* 237:460–467, 2018; Rochman in *Sci Rep* 3:1–7, 2013; Selvam in *J Hazard Mater* 402, 2021; Zhou et al. in *Sci Total Environ* 694, 2019;], and pose a serious threat to living beings, ecosystem, and human health [e.g. Devereux in *Mar Pollut Bull* 173, 2021, Assas in *Mar Pollut Bull* 158, 2020].

---

# Chapter 13

## Groundwater: An Endangered Invisible Resource. Microplastic Pollution in Underground Karst Systems, from Surface Watercourses to Cave Waters



Valentina Balestra , Bartolomeo Vigna , and Rossana Bellopede 

### 13.1 Introduction

Microplastic (MP) pollution has been found in environments worldwide, including remote areas [e.g. 1, 2, 3]. MPs have a dimension less than 5 mm and are a global problem because they are easily transportable even at long distances, tend to adsorb and transport hazardous substances such as heavy metals, pesticides, persistent organic pollutants (POPs) and antibiotics [e.g. 4, 5–8], and pose a serious threat to living beings, ecosystem, and human health [e.g. 9, 10].

Despite groundwater in karst aquifers constitutes a quarter of the global drinking water sources [11], MP pollution in karst areas and underground environments is still poorly known. This fact can be related to the invisibility of subterranean karst systems and their hard access, making them difficult to study. Being open systems, karst aquifers are vulnerable to contamination by surface pollution: MP presence in groundwater can be due to human activities, wastewater, surface waters and soil pollution, atmospheric deposition and litter. The subsoil characteristics play a fundamental role in the MP transport to groundwater: MPs deposited on the ground can be transported vertically in the subsoil throughout the soil pores and rock fractures [12, 13]. Precipitations facilitate the MP infiltration into the subsoil [14], as well as

---

V. Balestra (✉)  
Politecnico Di Torino, Corso Duca Degli Abruzzi 24, Torino, Italy  
e-mail: [valentina.balestra@polito.it](mailto:valentina.balestra@polito.it)

B. Vigna · R. Bellopede  
Bartolomeo Vigna, Politecnico Di Torino, Rossana Bellopede Corso Duca Degli Abruzzi 24,  
Torino, Italy

© The Author(s), under exclusive license to Springer Nature Switzerland AG 2023  
M. Cocca et al. (eds.), *Proceedings of the 3rd International Conference on Microplastic Pollution in the Mediterranean Sea*, Springer Water,  
[https://doi.org/10.1007/978-3-031-34455-8\\_13](https://doi.org/10.1007/978-3-031-34455-8_13)

1

17 transport them over long distances [15]. The hydrogeology of the study area, the flow  
18 regime of streams and their seasonal variation must be taken into account too.

19 Karst environments have an important role in natural ecosystems and karst aquifer  
20 are a fundamental drinking water reserve, therefore, investigations and monitoring  
21 are required for understanding the system dynamics and identify potential sources  
22 of contamination. In this work, we conducted preliminary investigations on MP  
23 pollution in different water samples from a karst area of NW Italy, considering  
24 connected surface and underground (cave) waters.

## 25 13.2 Experimental

### 26 13.2.1 Materials

27 Bossea cave (Frabosa Soprana, Piedmont, Italy, 836 m a.s.l.), the oldest Italian show  
28 cave, was firstly explored in the first half of the nineteenth century and opened to  
29 the public in 1874. Today, it receives about 13,000 tourist/year (2021). The cavity  
30 has a single entrance and an ascending structure, developing for about 2800 m,  
31 with a vertical drop of about 200 m. It represents the final part of the karst system  
32 developing in the Maudagna-Corsaglia watershed. The cave is fed by a system with  
33 primary (precipitations infiltrating directly into the aquifer) and secondary supply  
34 (superficial water streams losses), such as Rio Roccia bianca, the main secondary  
35 supply contributing to the Bossea karst system recharge [16]. Infiltration waters feed  
36 an important collector which cross the Bossea cave for more than 1.5 km and directly  
37 gushes out in the Corsaglia River with a set of springs [17]. A series of tests using two  
38 different dyes, previously carried out in Rio Roccia bianca, highlighted that water  
39 coming from Rio Roccia bianca arrive in the collector of Bossea cave in short time  
40 and with high concentration [16]. Different discontinuities are present in the cave,  
41 characterized by low flow circulation [17, 18].

42 Spot sampling were carried out in Rio Roccia bianca superficial tributary (one  
43 sample), in the Bossea cave waters (four samples), and in the Corsaglia stream, where  
44 the waters of the underground collector emerge (one sample). In the Bossea cave, two  
45 samples were collected in the speleological traits (Sifone and Polla delle anatre), as  
46 reference samples of unpolluted waters, and two in the touristic area (Uovo and Sala  
47 frane). In Sifone, in the final part of the cave, the water comes out from a siphoning  
48 tract and are collected in a lake. Polla delle anatre is a small pool of water flowing out  
49 of a fracture in the rock, entering in the inner collector [18]. Along the tourist path,  
50 in Uovo, water flows between different gours, slowing down, while, in the initial part  
51 of the cave, in Sala frane, the collector waters flow quickly before leaving the cave.  
52 Groundwater and superficial water samples were collected into 1 L glass vessels, in  
53 September 2021, under low-flow conditions, transported to the laboratory in coolers,  
54 and kept refrigerated at 6 °C until analysis.

### 13.2.2 Methods

MP samples were analysed according to the previously published method for cave sediments [see 19 for details], adapted to the liquid matrix. Glass filters were replaced to silver ones to determine MP typology using the infrared (IR) spectroscopy. MPs were detected by exploiting their fluorescence under ultraviolet (UV) light, given by Fluorescent Whitening Agents (FWAs), commonly used additives in plastic production [20, 21] (Fig. 13.1). All researchers used nitrile gloves and cotton coats. For every step, all surfaces and materials were cleaned with ethanol and distilled water to avoid external contaminations. Plastic equipment was replaced with glass or metal one.

#### 13.2.2.1 Preparation of Samples

Water samples were filtered through a 0.8  $\mu\text{m}$  pore size silver filter (GVS Life Sciences, Membrane Disk 47 mm). Filters, positioned on glass petri dishes and covered with aluminium foil, were dried in an oven at 40 °C for 2 h. The organic matter removal was carried out on dried filters through the application of 2 ml of 15% hydrogen peroxide solution, left to react for 30 min at room temperature and, finally, dried again for 1 h at 50 °C.




Fig. 13.1 Microplastic fibre under UV light

### 13.2.2.2 Analytical Techniques

Filters were observed with and without an UV flashlight (Alonefire SV10 365 nm UV flashlight 5W) under a Leitz ORTHOLUX II POL-MK microscope equipped with a DeltaPix Invenio 12EIII 12 Mpx Camera, with a 2.5x, 4x, 10 × or higher magnifications. Images of natural, artificial and synthetic fibres under microscope [e.g. 22, 23, 24] were used to compare particles observed under microscope, especially transparent fibres. The UV flashlight was positioned with an inclination of 45 °C. Visual identification was used to count MPs, according to strict selection criteria described in previous research [25–27]. A cut off of the particles smaller than 0.1 mm was made as suggested in European Commission [28]. Particles that could not be identified as MPs were not take into consideration. The counted MPs were described using the Standardised size and colour sorting system (SCS) [25].

Finally, a 10% of MPs of each filter were analysed using a micro-Fourier Transform Infrared Spectroscopy (micro-FTIR) Shimadzu AIM-9000 microscope equipped with a Shimadzu IRTracer-100 spectrophotometer. Spectra were compared with the Shimadzu standard library.

## 13.3 Results and Discussion


### 13.3.1 Results

MPs were found in all water samples: 54 items/L in Sifone, 16 items/L in Sala frane, 12 items/L in Polla delle anatre, 30 items/L in Uovo, 23 items/L in Rio Rocchia bianca and 29 items/L in Corsaglia River. Fibres represented the majority of the MPs present in the karst system (95.1%), followed by fragments (4.3%) and beads (0.6%). Mini-MPs accounted for 82.9% of the total MPs found in water karst system. The highest MP abundance was fluorescent under UV light (84.8%). Of the fluorescent particles, 46% were transparent, 16.5% red, 10.8% blue, 7.2% beige, and 26.7% other colours. Non-fluorescent MPs were mainly black (68%) or blue (20%). Polyethylene (51.4%) and polyvinyl alcohol (31.4%) are the main types of MPs found in the karst system waters, followed by polyester (8.9%), ethylene vinyl alcohol (2.9%), polyvinyl chloride (2.9%) and acrylic adhesive (2.9%).


### 13.3.2 Discussion

At the moment, a standardized method for assessing MPs do not exist. Using different methodology makes it difficult to compare the results. However, some considerations on MP amount in karst environment can be equally done. The only work detecting MP pollution in springs and wells in a karst water area has documented a maximum

value of 15.2 MPs/L, under low-flow conditions [11], a fairly low value compared to those found in the Bossea karst system. Instead, concerning MP characteristics, the data about MPs in cave waters are similar to those described in Balestra and Bellopede [19] for Bossea cave sediments. Five sediment samples were collected along different part of the tourist path and one in the not-tourist area of the Bossea cave. Comparisons between the mean values obtained from sediment and water samples analysis inside Bossea cave are reported in Figs. 13.2, 13.3. The major part of MPs in Bossea cave waters are fluorescent, fibre-shape, and have a dimension between 0.99 and 0.1 mm, as in cave sediments (Fig. 13.2). The most of the fluorescent particles in Bossea cave waters are transparent, as in cave sediments, and the other main colours are similar too, with except of red particles, greater in water (Fig. 13.3). The non-fluorescent MPs were mainly black and blue, confirming the data found in Bossea cave sediments (Fig. 13.3). However, percentages of these data are different, as the colour of the less abundant MPs (Fig. 13.3). It could be related to different kind of biodeterioration, or to the diverse density of certainly MPs, more easily transportable in liquid medium. MPs in groundwater of Bossea karst system could be related to outside pollution, linked to the air transport of particles, soil contamination, winter activities related to the ski resorts and pollution from the near countries and could be transported especially by the main secondary supplies such as Rio Rocchia bianca. The waters in the touristic part of the cave could be enriched with pollutants too, brought inside by tourists [19], which could be transported in different part of the cave and outside, in the Corsaglia river.



**Fig. 13.2** Comparison of size, shape and fluorescence between microplastics found in Bossea cave waters and sediments



**Fig. 13.3** Comparison of colours between fluorescent and non-fluorescent microplastics found in Bossea cave waters and sediments

## 13.4 Conclusions

This study documented the presence of microplastics in karst systems, from surface watercourses to water in cave. Microplastics with a dimension less than 1 mm and fibre shape dominate the samples and polyethylene is the main types of microplastics found in the karst system waters. Fractured aquifers may be vulnerable to microplastic pollution, contaminating waters and nearby environments, therefore, MP detection is essential to understand the pollution amount and define strategies for conservation. Moreover, karst aquifers are open systems, susceptible to pollutant contamination from surface, therefore, the above-ground areas should be monitored and protected too. Greater endeavours should be done to karst areas protection, implementing pollutants monitoring and sustainably managing important resources as water.

## References

- Zhang Y et al (2021) Microplastics in glaciers of the Tibetan Plateau: Evidence for the long-range transport of microplastics. *Sci Total Environ* 758:143634
- Ambrosini R et al (2019) First evidence of microplastic contamination in the supraglacial debris of an alpine glacier. *Environ Pollut* 253:297–301
- Cincinelli A et al (2017) Microplastic in the surface waters of the Ross Sea (Antarctica): occurrence, distribution and characterization by FTIR. *Chemosphere* 175:391–400
- Li X et al (2019) Enhancement in adsorption potential of microplastics in sewage sludge for metal pollutants after the wastewater treatment process. *Water Res* 157:228–237
- Li J, Zhang K, Zhang H (2018) Adsorption of antibiotics on microplastics. *Environ Pollut* 237:460–467
- Rochman CM et al (2013) Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. *Sci Rep* 3(1):1–7

- 152 7. Selvam S et al (2021) Hazardous microplastic characteristics and its role as a vector of heavy  
153 metal in groundwater and surface water of coastal south India. *J Hazard Mater* 402:123786
- 154 8. Zhou Y, Liu X, Wang J (2019) Characterization of microplastics and the association of heavy  
155 metals with microplastics in suburban soil of central China. *Sci Total Environ* 694:133798
- 156 9. Devereux R et al (2021) The abundance of microplastics in cnidaria and ctenophora in the  
157 North Sea. *Mar Pollut Bull* 173:112992
- 158 10. Assas M et al (2020) Bioaccumulation and reproductive effects of fluorescent microplastics in  
159 medaka fish. *Mar Pollut Bull* 158:111446
- 160 11. Panno SV et al (2019) Microplastic contamination in karst groundwater systems. *Groundwater*  
161 57(2):189–196
- 162 12. Chia RW, et al. (2021) Microplastic pollution in soil and groundwater: a review. *Environmental*  
163 *Chemistry Letters*, 1–14
- 164 13. Wanner P (2021) Plastic in agricultural soils—a global risk for groundwater systems and drinking  
165 water supplies?—a review. *Chemosphere* 264:128453
- 166 14. Zhou Y et al (2021) Microplastic contamination is ubiquitous in riparian soils and strongly  
167 related to elevation, precipitation and population density. *J Hazard Mater* 411:125178
- 168 15. Allen S et al (2019) Atmospheric transport and deposition of microplastics in a remote mountain  
169 catchment. *Nat Geosci* 12(5):339–344
- 170 16. Vigna, B (2020) Assetto geologico ed idrogeologico del Sistema carsico di Bossea (SW  
171 Piemonte, Italy), in *Atti del Convegno Nazionale “L’uomo domanda, la grotta risponde”*.  
172 Cinquantésimo anniversario del Laboratorio Carsologico Sotterraneo di Bossea, Frabosa  
173 Soprana (CN), Italy. p 283–300
- 174 17. Vigna B, (2021) Vadose Zone Hydrogeology In The Bossea Cave System (Southern Piedmont,  
175 Northern Italy), in *Proceedings of the 17th International Congress of Speleology*, Sidney. p  
176 222–225
- 177 18. Balestra V, Fiorucci A, Vigna B (2022) Study of the trends of chemical-physical parameters in  
178 different karst aquifers: some examples from italian alps. *Water* 14(3):441
- 179 19. Balestra V, Bellopede R (2022) Microplastic pollution in show cave sediments: First evidence  
180 and detection technique. *Environ Pollut* 292:118261
- 181 20. Ehlers SM, Maxein J, Koop JH (2020) Low-cost microplastic visualization in feeding  
182 experiments using an ultraviolet light-emitting flashlight. *Ecol Res* 35(1):265–273
- 183 21. Klein M, Fischer EK (2019) Microplastic abundance in atmospheric deposition within the  
184 Metropolitan area of Hamburg. Germany. *Sci. Total Environ.* 685:96–103
- 185 22. Houck MM (2009) Identification of textile fibers
- 186 23. Khan A, et al. (2017) A review paper on Textile Fiber Identification. *IOSR Journal of Polymer*  
187 *and Textile Engineering (IOSR-JPTE)* 4, 14–20
- 188 24. Zhang, X (2014) Fundamentals of fiber science
- 189 25. Crawford CB, Quinn B (2016) Microplastic pollutants, Amsterdam
- 190 26. Noren F (2007) Small plastic particles in coastal swedish waters, in n-research report,  
191 commissioned by KIMO, Sweden
- 192 27. Hidalgo-Ruz V et al (2012) Microplastics in the marine environment: a review of the methods  
193 used for identification and quantification. *Environ Sci Technol* 46(6):3060–3075
- 194 28. European Commission: Guidance on monitoring of marine litter in European seas. A guidance  
195 document within the common implementation strategy for the Marine Strategy Framework  
196 Directive. Ispra: European Commission, Joint Research Centre, MSFD Technical Subgroup on  
197 Marine Litter. p 126 (2013)