

Diaphragmatic ultrasound and esophageal pressure in COVID-19 pneumonia during helmet CPAP

*Original*

Diaphragmatic ultrasound and esophageal pressure in COVID-19 pneumonia during helmet CPAP / Steinberg, I., Chiodaroli, E., Gattarello, S., Cappio Borlino, S., Chiumello, D.. - In: INTENSIVE CARE MEDICINE. - ISSN 0342-4642. - 48:8(2022), pp. 1095-1096. [10.1007/s00134-022-06785-z]

*Availability:*

This version is available at: 11583/2993700 since: 2024-10-25T13:43:40Z

*Publisher:*

Springer Science and Business Media Deutschland

*Published*

DOI:10.1007/s00134-022-06785-z

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

LETTER



# Diaphragmatic ultrasound and esophageal pressure in COVID-19 pneumonia during helmet CPAP

Irene Steinberg<sup>1\*</sup> , Elena Chiodaroli<sup>2</sup>, Simone Gattarello<sup>3</sup>, Simone Cappio Borlino<sup>2</sup> and Davide Chiumello<sup>2</sup>

© 2022 The Author(s)

Dear Editor,

Diaphragmatic thickening fraction was reported to correlate with inspiratory effort [1, 2]. In this prospective study, we investigated whether ultrasound-measured diaphragmatic excursion and thickening may surrogate esophageal pressure swing during Continuous Positive Airway Pressure (CPAP).

From December 2020 to August 2021, 46 consecutive patients admitted to the High-Dependency Unit (HDU) of San Carlo University Hospital (Milan, Italy) for pneumonia related to coronavirus disease 2019 (COVID-19) (confirmed by acute respiratory failure, positive test and infiltrates at lung computed tomography [CT]) were prospectively enrolled. The study was approved by the ethical committee (Comitato Etico Milano Area I; 17263/2020-2020/ST/095) and written informed consent was obtained. At HDU admission, according to the local protocol of clinical monitoring in COVID-19 patients, an esophageal balloon catheter was inserted. Esophageal pressure, ultrasound and blood gas analysis were simultaneously obtained during helmet CPAP. Measurements were repeated at zero end-expiratory pressure (ZEEP). The ultrasounds were performed by ultrasound-experienced physicians with good reproducibility of measures.

The study population had a median [interquartile range (IQR)] age of 58 [54;65] years and 61% were male, body mass index (BMI) was 27 [25;28]. PaO<sub>2</sub>/FiO<sub>2</sub> ratio and

PaCO<sub>2</sub> at admission were 184 [136;243] mmHg and 39 [36;42] mmHg. Charlson Comorbidity Index was 1 [1;2] and Sequential Organ Failure Assessment (SOFA) score was 3 [2;3]. Thirty-nine patients had a positive outcome, 7 underwent mechanical ventilation and 3 of them ultimately died.

As shown in Fig. 1, no association was observed between esophageal pressure swings and: (1) thickening fractions (Panel A) and (2) diaphragmatic excursions (Panel B). The correlation between changes in esophageal pressure swing and thickening fraction between PEEP and ZEEP was significantly positive but moderate ( $p = 0.038$ , Spearman's rho 0.35) (Panel C).

The main finding of the present study is that neither thickening fraction nor diaphragmatic excursion were able to estimate esophageal pressure swing in patients with COVID-19 pneumonia. Despite previous positive studies [1, 2], also Poulard et al. [3] observed a poor correlation between thickening fraction and change in transdiaphragmatic pressure. The two variables were repeatedly tested in a single subject and a correlation was found only in few of them. This suggests a non-fixed relationship between diaphragm contraction and change in transdiaphragmatic pressure and agrees with the dispersion we observed in the relationship between changes in esophageal pressure swing and thickening fraction between PEEP and ZEEP.


The change in transpulmonary pressure results from the applied pressure and the muscular inspiratory effort according to the following equation [4]:

$$\Delta P_{es} = \Delta P_{aw} \cdot \frac{E_w}{E_{TOT}} - \Delta P_{musc} \cdot \frac{E_L}{E_{TOT}},$$

\*Correspondence: irene@steinberg.it

<sup>1</sup> Department of Anesthesiology, Medical University of Göttingen, Robert Koch Straße 40, 37075 Göttingen, Germany

Full author information is available at the end of the article



where  $\Delta P_{es}$  is the esophageal pressure swing,  $\Delta P_{aw}$  is the applied airway pressure,  $\Delta P_{musc}$  is the pressure generated by the patient's muscles,  $E_W$  is the chest wall elastance,  $E_L$  is the lung elastance and  $E_{TOT}$  is the total elastance. Therefore, a different ratio between chest wall and lung elastance, at the same  $\Delta P_{musc}$ , causes different changes in esophageal pressure and the variability of this ratio may account for the lack of correlation between diaphragmatic contraction, a manifestation of diaphragmatic  $\Delta P_{musc}$ , and esophageal pressure. Moreover,  $\Delta P_{musc}$  is the sum of the pressure generated by the diaphragm (as revealed by the thickening fraction) and the pressure generated by the intercostal muscles and during increasing inspiratory effort the contribution of the intercostal muscles increases progressively more in comparison to the activity of the diaphragm [5]. Therefore, we may hypothesize that, at least in COVID-19 patients, an increased inspiratory effort, as measured by esophageal pressure swings, is poorly related to the pressure generated by the diaphragm and inferred by the thickening fraction.

#### Author details

<sup>1</sup> Department of Anesthesiology, Medical University of Göttingen, Robert Koch Straße 40, 37075 Göttingen, Germany. <sup>2</sup> Department of Anesthesiology and Intensive Care, ASST Santi e Paolo Hospital, University of Milan, Milan, Italy. <sup>3</sup> Department of Anesthesiology and Intensive Care, IRCCS San Raffaele Scientific Institute, Milan, Italy.

#### Author contributions

Conception and design: IS, EC, and DC. Analysis and interpretation: IS, EC, SG, SCB, and DC. Drafting the manuscript for important intellectual content: IS and DC. Guarantor of this paper and responsible for the integrity of the work: IS and DC.

#### Funding

Open Access funding enabled and organized by Projekt DEAL. The support for this work was provided from departmental sources and from two unrestricted grants for research in respiratory medicine, apportioned by Sartorius AG (Otto-Brenner-Straße 20, 37079, Göttingen, Germany).

#### Declarations

#### Conflicts of interest

The authors report no conflicts of interest.

#### Open Access

This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Accepted: 8 June 2022

Published: 21 June 2022

#### References

- Vivier E, Mekontso Dessap A, Dimassi S et al (2012) Diaphragm ultrasonography to estimate the work of breathing during non-invasive ventilation. *Intensive Care Med* 38:796–803. <https://doi.org/10.1007/s00134-012-2547-7>
- Umbrello M, Formenti P, Lusardi AC et al (2020) Oesophageal pressure and respiratory muscle ultrasonographic measurements indicate inspiratory effort during pressure support ventilation. *Br J Anaesth* 125:e148–e157. <https://doi.org/10.1016/j.bja.2020.02.026>
- Poulard T, Bachasson D, Fossé Q et al (2021) Poor correlation between diaphragm thickening fraction and transdiaphragmatic pressure in mechanically ventilated patients and healthy subjects. *Anesthesiology*. <https://doi.org/10.1097/ALN.0000000000004042>
- Gattinoni L, Marini JJ, Collino F et al (2017) The future of mechanical ventilation: lessons from the present and the past. *Crit Care* 21:183. <https://doi.org/10.1186/s13054-017-1750-x>
- Mengeot PM, Bates JH, Martin JG (1985) Effect of mechanical loading on displacements of chest wall during breathing in humans. *J Appl Physiol* 58:477–484. <https://doi.org/10.1152/jappl.1985.58.2.477>