

Muscle pre-activation prior to landing in athletes with anterior cruciate ligament reconstruction: detection of EMG onset using artificial intelligence

Original

Muscle pre-activation prior to landing in athletes with anterior cruciate ligament reconstruction: detection of EMG onset using artificial intelligence / Russo, F., Ghislieri, M., Baldazzi, A., Rum, L., Bergamini, E., Agostini, V.. - In: GAIT & POSTURE. - ISSN 0966-6362. - ELETTRONICO. - 114:(2024), pp. 38-39. (24th National Congress of SIAMOC Stresa (Italy) 2-5 October 2024) [10.1016/j.gaitpost.2024.08.064].

Availability:

This version is available at: 11583/2994261 since: 2024-11-08T07:47:32Z

Publisher:

Elsevier

Published

DOI:10.1016/j.gaitpost.2024.08.064

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Elsevier postprint/Author's Accepted Manuscript

© 2024. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:
<http://dx.doi.org/10.1016/j.gaitpost.2024.08.064>

(Article begins on next page)

Authors:

Federica Russo¹, Marco Ghislieri¹, Andrea Baldazzi², Lorenzo Rum³, Elena Bergamini⁴, Valentina Agostini¹

Affiliations:

¹ Polito^{BIO}Med Lab, Department of Electronics and Telecommunications, Politecnico di Torino, Turin, Italy

² Dipartimento di Scienze Motorie, Umane e della Salute, Università degli Studi di Roma “Foro Italico”

³ Dipartimento di Scienze Biomediche, Università degli Studi di Sassari

⁴ Dipartimento di Ingegneria Gestionale, dell'Informazione e della Produzione, Università degli Studi di Bergamo

Title

Muscle pre-activation prior to landing in athletes with anterior cruciate ligament reconstruction: detection of EMG onset using artificial intelligence

Keywords:

ACL-R; asymmetry; EMG onset; deep learning; motion analysis

ABSTRACT***Introduction***

In Return-to-Sport evaluation of athletes after Anterior Cruciate Ligament Reconstruction (ACL-R), motion analysis of specific tasks, such as Single-Leg Hop (SLH), can provide unique measurements. Muscle pre-activation prior to SLH landing is adopted as a discriminant parameter between ACL-R and control athletes [1]. To detect muscle pre-activation from electromyography (EMG) signals, algorithms for activation-interval detection can overcome the limitations of manual segmentation. Long Short-Term Memory recurrent neural network for Muscle Activity Detection (LSTM-MAD) [2] proved successful, in gait analysis, as a tool for EMG-onset detection based on Artificial Intelligence (AI). LSTM-MAD directly works on raw EMG signals without requiring additional pre-processing or input parameters (e.g., background-noise power and Signal-to-Noise Ratio). The aim of this work is to quantify muscle pre-activation and asymmetry in ACL-R and control athletes, adapting LSTM-MAD to the evaluation of SLH task.

Methods

The EMG activity of 4 lower-limb muscles - Vastus Lateralis (VL), Vastus Medialis (VM), Semitendinosus (ST), and Biceps Femoris (BF) – was acquired bilaterally from 12 ACL-R and 17 control athletes. LSTM-MAD was applied to evaluate muscle pre-activation before initial contact while landing. The median value of the EMG pre-activation across 3 trials was considered for the analysis. A derived parameter - named “pre-

activation asymmetry” - was obtained for each subject, defined as $|EMG\ preactivation_{left} - EMG\ preactivation_{right}|$. In other words, the asymmetry between the reconstructed and contralateral side of patients was calculated and compared against the “physiological” asymmetry between the dominant and non-dominant side of controls. Two 2-way ANOVAs with Bonferroni adjustment for multiple comparisons were performed to test differences in muscle pre-activation and pre-activation asymmetry between populations (ACL-R athletes vs. controls) and knee flexor/extensor muscle groups (knee flexors: ST and BF; knee extensors: VL and VM).

Results

Figure 1 shows the muscle activation intervals of a representative ACL-R athlete, and results on ACL-R and control populations.

Statistically significant differences in muscle pre-activation were found between populations (ACL-R: 167 ± 6 ms (mean $\pm$ SE); controls: 137 ± 5 ms; $p < 0.001$) and muscle groups (knee flexors: 169 ± 5 ms; knee extensors: 135 ± 5 ms; $p < 0.001$). Moreover, pre-activation asymmetry was higher in ACLR athletes compared to controls (ACL-R: 55 ± 6 ms; controls: 29 ± 4 ms; $p < 0.001$) and in knee extensors compared to knee flexors (knee flexors: 29 ± 5 ms; knee extensors: 55 ± 5 ms; $p < 0.001$).




Figure 1. Example of LSTM-based detection of EMG activation intervals and boxplots of muscle pre-activation and asymmetry in athletes after ACL-R and controls.

Fig. 1A shows activation intervals of the VL, VM, BF, and ST muscles, detected by LSTM-MAD, of a representative ACL-R athlete during SLH task. Boxplots showing differences in muscle pre-activation and asymmetry between the ACL-R group (affected and sound side) and control group (dominant and non-dominant side) are represented in Fig. 1B and Fig. 1C, respectively.

Discussion

AI proved useful in the automatic detection of muscle activity, allowing for the extraction of muscle pre-activation and derived parameters such as pre-activation asymmetry during SLH landing in ACL-R and control athletes.

References

- [1] A. Baldazzi, C. Bosio, L. Rum, H. Pillet, F. Margheritini, A. Macaluso, E. Bergamini, “Muscle activation timing and knee biomechanics during single leg hop in subjects with and without anterior cruciate ligament reconstruction: a case study”, *Gait & Posture* 97, 2 (2022), Doi: 10.1016/j.gaitpost.2022.09.054.
- [2] M. Ghislieri, G.L. Cerone, M. Knaflitz, V. Agostini, “Long short-term memory (LSTM) recurrent neural network for muscle activity detection”, *J NeuroEngineering Rehabil* 18, 153 (2021), Doi: 10.1186/s12984-021-00945-w.