

Erosion and deposition in the JET divertor during the second ITER-like wall campaign

Original

Erosion and deposition in the JET divertor during the second ITER-like wall campaign / Mayer, M; Krat, S; Baron-Wiechec, A; Gasparyan, Yu; Heinola, K; Koivuranta, S; Likonen, J; Ruset, C; de Saint-Aubin, G; Widdowson, A; Subba, F. - In: PHYSICA SCRIPTA. - ISSN 0031-8949. - T170:T170(2017). [10.1088/1402-4896/aa8ff9]

Availability:

This version is available at: 11583/2986867 since: 2024-03-12T14:51:13Z

Publisher:

IOP Publishing Ltd

Published

DOI:10.1088/1402-4896/aa8ff9

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

IOP preprint/submitted version

This is the version of the article before peer review or editing, as submitted by an author to PHYSICA SCRIPTA. IOP Publishing Ltd is not responsible for any errors or omissions in this version of the manuscript or any version derived from it. The Version of Record is available online at <https://dx.doi.org/10.1088/1402-4896/aa8ff9>.

(Article begins on next page)

A 3D multimodal and multiscale approach for the study of Upper Palaeolithic ground stone tools

G. Sorrentino * ^(a,b), L. Vigorelli ^(c,b), A. Re ^(a,b), L. Longo ^(d), M. Paggi ^(e), F. Menna ^(f), A. Karami ^(f), L. Mancini ^(g) & A. Lo Giudice ^(a,b)

^(a)Department of Physics, University of Turin, Italy

^(b) INFN, Turin Division, Italy

^(c) Department of Electronics and Telecommunications, Polytechnic of Turin, Italy

^(d) DAIS, Ca' Foscari University of Venice, Italy

^(e) IMT School for Advanced Studies Lucca, Italy

^(f) Bruno Kessler Foundation, Trento, Italy

^(g) Elettra-Sincrotrone Trieste S.C.p.A., Basovizza, Trieste, Italy

* Presenting author

giusi.sorrentino@unito.it

The term “ground stone” refers to non-flaked industry, and includes a multitude of different instruments that have not undergone an intensive stone shaping. This natural and non-modified appearance of the stones lends to them the high potentiality of being utilised in the most varied uses as the processing of different materials such as bone, mineral and plant [1]. Among them we are here referring to the elaboration of vegetal resources that leave evanescent traces and perishable residues on the stone surfaces, which call for a tailored approach. Of great support for the comprehension of these stone tools’ function(s) is the use of analytical techniques that allow a 3dimensional characterisation of the item and of wear affecting the surfaces as well as of morphotextural properties within the sample volume [2]. Our approach is based on multimodal and multidimensional morphological approach, by applying imaging techniques ranging from macro to sub-micro scales aiming at elaborate a digital model of the tools and at performing a wide range of analysis. In particular, the procedure exploits the advantage of (i) photogrammetric acquisition of the ground stones and 3D model elaborations to inspect their geometry and macroscale investigation of their surface, (ii) profilometry measurements of selected areas of the stone surface to characterise the microtopography and highlight the presence of use-wear traces, (iii) high-resolution synchrotron X-ray tomographic imaging to investigate the volumetric microstructure by a non-destructive approach and to verify the presence of crevices that served as putative traps for biogenic residues deriving from the plants grinding and pounding.

We present the pipeline developed for the analysis of the artefacts from several Marine Isotopic Stage 3 (MIS 3, 60-25 ka) sites, and in particular, tailored for the ground stones retrieved in the Aurignacian cultural level III of the Brînzei I cave (north-west Moldova) [3]. The proposed analysis is tested on experimental replicas used to treat plants starch-rich organs compatible with the biome of the Pontic steppe during the MIS 3 [4].

References

- [1] Longo L., Skakun N., Pantyukhina I., Terekhina V., Sorrentino G., Aurignacian grinding stone from Surein I (Crimea): “trace-ing” the roots of starch-based diet. *J. Archaeol. Sci. Rep.*, 38 (2021). [2] Zambrano M., Volatili T., Mancini L., Pitts A., Giorgioni M., Tondi E., Pore-scale dual-porosity and dual-permeability modeling in an exposed multi-facies porous carbonate reservoir. *Mar. Pet. Geol.*, 128 (2021). [3] Allsworth-Jones P., Borziac I.A., Chetaru N.A., French C.A.I., Medyanik S.I., Brînzei: a multidisciplinary study of an upper palaeolithic site in Moldova. *Proc. Prehist. Soc.*, 84 (2018). [4] Sorrentino G., Lo Giudice A., Re A., Borghi A., Longo L., Più di un semplice ciottolo: Un protocollo sperimentale per la comprensione del ruolo di strumenti macrolitici nella trasformazione delle risorse vegetali nel Paleolitico Superiore in Eurasia. *Archeologie sperimentali*, 2 (2022).