

Particle toxicity and air quality standards. A focus on the Po Valley, Italy

Original

Particle toxicity and air quality standards. A focus on the Po Valley, Italy / Ravina, M., De Maria, R., Maringo, M., Sacco, M., Bardi, L., Bruno, A., Panepinto, D., Zanetti, M., Brizio, E., Barbero, S.. - (2024). (XI Convegno Nazionale sul Particolato Atmosferico (PM 2024)).

Availability:

This version is available at: 11583/2995012 since: 2024-12-04T16:31:40Z

Publisher:

IAS Italian Aerosol Society

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

XI Convegno Nazionale sul Particolato Atmosferico



TORINO PM 2024

ATTI

Un evento a cura di



Con il patrocinio di



Con il supporto organizzativo di



UNIVERSITÀ
DI TORINO

TORINO 28 – 31 Maggio 2024

Particle toxicity and air quality standards. A focus on the Po Valley, Italy.

Marco Ravina^{1*}, Roberta De Maria², Marilena Maringo², Milena Sacco², Luisella Bardi², Annalisa Bruno², Deborah Panepinto¹, Mariachiara Zanetti¹, Enrico Brizio², Secondo Barbero².

¹ Department of Environment, Land and Infrastructure Engineering, Politecnico di Torino;

² Environmental Protection and Research Agency of Piedmont (ARPA Piemonte)

* Corresponding author. Tel: +39 0110907699, E-mail: marco.ravina@polito.it

Keywords: atmospheric aerosols, particle toxicity, air quality planning, Po Valley

Introduction: Northern Italy still has difficulties complying with EU air quality standards, despite strong emission reductions. In the colder seasons, wind speed, PBL height, and atmospheric pressure occurring in the Po basin are three to five times less efficient in dispersing pollutants if compared to other European regions [1]. Different regions also show different aerosols composition, resulting in a changing toxic impact on human health. The characterization of aerosols toxicity is gaining importance, as it could act as a key point for the development of air quality planning policies. To this end, toxicity ranking methods have been recently proposed by researchers.

Methodology: In this study, the toxicity of PM₁₀ in different sites worldwide was calculated and compared. The PM₁₀ source apportionment and concentration trend were analysed based on data reported in the literature. The results of the LIFE PREPAIR project (<https://www.lifeprepare.eu/>) were also considered in this study. To evaluate particle toxicity, the methodology proposed by Park et al. [2] was applied. This method is based on the analysis of multiple biological and chemical endpoints, that were integrated for various source-specific aerosols to derive toxicity scores for particles originating from different sources. Average PM₁₀ toxicity of rural and urban sites was thus calculated based on the average chemical composition.

Results: the results showed that, on average, toxicity ranking of rural sites was lower than that of urban areas. In addition, the average contribution of secondary inorganic aerosols to the overall PM toxicity was analysed. A significant variability was observed among different sites. In the second phase of the study, the temporal variability of PM composition and toxicity of a rural and a urban area in the Po Valley was analysed and compared. Time series of toxicity-weighted concentration were defined.

Conclusion: The results reported show that, despite toxicity ranking methods are still not consolidated, if coupled with advanced PM source apportionment, reliable toxicity and epidemiological analyses, they could represent important tools to build a new consistent health metric for ambient PM.

Acknowledgement: Part of the data used in this study were elaborated in the PREPAIR - LIFE 15 IPE IT013 project – Action D6.

References

- [1] Robotto et al. (2022), <https://doi.org/10.3390/atmos13050642>
- [2] Park, M. et al. (2018), <https://doi.org/10.1038/s41598-018-35398-0>