

Non-Lorentzian supergravity and kinematical superalgebras

Original

Non-Lorentzian supergravity and kinematical superalgebras / Concha, P., Ravera, L.. - In: JOURNAL OF HIGH ENERGY PHYSICS. - ISSN 1029-8479. - ELETTRONICO. - 2025:3(2025), pp. 1-47. [10.1007/JHEP03(2025)032]

Availability:

This version is available at: 11583/3012399 since: 2026-06-24T14:13:04Z

Publisher:

Springer

Published

DOI:10.1007/JHEP03(2025)032

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

RECEIVED: December 13, 2024

REVISED: January 22, 2025

ACCEPTED: January 27, 2025

PUBLISHED: March 6, 2025

Non-Lorentzian supergravity and kinematical superalgebras

Patrick Concha ^{a,b} and Lucrezia Ravera ^{c,d,b}

^a*Departamento de Matemática y Física Aplicadas, Universidad Católica de la Santísima Concepción, Alonso de Ribera 2850, Concepción, Chile*

^b*Grupo de Investigación en Física Teórica, GIFT, Universidad Católica de la Santísima Concepción, Alonso de Ribera 2850, Concepción, Chile*

^c*DISAT, Politecnico di Torino - PoliTo, Corso Duca degli Abruzzi 24, 10129 Torino, Italy*

^d*Istituto Nazionale di Fisica Nucleare, Section of Torino - INFN, Via P. Giuria 1, 10125 Torino, Italy*

E-mail: patrick.concha@ucsc.cl, lucrezia.ravera@polito.it

ABSTRACT: In this paper, we present and classify the supersymmetric extensions of extended kinematical algebras, at the basis of non-Lorentzian physics theories. The diverse kinematical superalgebras are here derived by applying non- and ultra-relativistic expansion procedures involving different semigroups. We then build three-dimensional Chern-Simons non-Lorentzian supergravity theories based on such (extended) kinematical superalgebras, providing the supersymmetry transformation laws of the fields and the field equations of the models, which correspond to the vanishing of the curvature two-forms. In fact, the expansion procedure adopted allows to automatically end up with a non-degenerate bilinear invariant trace for the (extended) kinematical superalgebras. The latter is a crucial ingredient of the Chern-Simons field-theoretical formulation, as it allows to include a kinetic term for each gauge field of the theory, implying the vanishing of the curvature two-forms as field equations.

KEYWORDS: Supergravity Models, Chern-Simons Theories, Gauge Symmetry, Classical Theories of Gravity

ARXIV EPRINT: [2412.07665](https://arxiv.org/abs/2412.07665)

Contents

1	Introduction	1
2	Extended kinematical Lie algebras	3
3	Supersymmetric extensions of extended kinematical algebras	6
3.1	Ultra-relativistic expansions	8
3.2	Non-relativistic expansions	13
4	Three-dimensional non-Lorentzian supergravity theories based on extended kinematical superalgebras	21
4.1	$\mathcal{N} = 2$ AdS-Carroll Chern-Simons supergravity and beyond	22
4.2	$\mathcal{N} = 2$ extended Newton-Hooke Chern-Simons supergravity and its generalization	27
4.3	$\mathcal{N} = 2$ (AdS-)static Chern-Simons supergravity and its generalization	32
5	Discussion	36

1 Introduction

The gravitational interaction can be effectively described by Newtonian gravity for regimes of low velocities relative to the speed of light c and weak gravitational fields. Notably, a non-relativistic (NR) formulation of General Relativity (GR) was introduced by Cartan in [1, 2], commonly referred to Newton-Cartan (NC) gravity. The latter employs a NC geometry that can be seen as the NR analogue of Riemannian geometry for GR, providing a geometric interpretation of the Poisson equation. NR theories, also known as Galilean, are commonly derived as the $c \rightarrow \infty$ limit of relativistic ones. Conversely, the $c \rightarrow 0$ limit defines the ultra-relativistic (UR) regime, also referred to as the Carrollian limit [3, 4], in homage to Lewis Carroll, the author of *Alice’s Adventures in Wonderland*. Both the Galilean and the Carrollian algebras, also referred as non-Lorentzian (NL), can be obtained through specific contraction of the Poincaré one, and they form part of the broader classification of kinematical Lie algebras introduced by Bacry and Lévy-Leblond in [5].

NL symmetries have garnered increasing attention due to their emergence in a wide range of physical phenomena. On the one hand, NR theories and their corresponding geometries have been useful in the study of holography [6–20], Hořava-Lifshitz gravity [21–26], Quantum Hall effect [27–31], asymptotic symmetries [32–34], post-Newtonian corrections [35–37], among others. On the other hand, UR symmetries have received a renewed interest since they manifest within the context of tachyon condensation [38], tensionless strings [39–43], asymptotic structures [44–46], holography in asymptotically flat spacetimes [47–54], and black hole horizon [55–59].

In particular, NL gravity theories in three spacetime dimensions can be explored using the Chern-Simons (CS) formalism. The three-dimensional CS approach serves as a laboratory model for studying the properties of higher-dimensional gravity theories, as it shares with

them intriguing features, e.g. concerning black hole solutions and their associated thermodynamics [60]. Notably, three-dimensional GR with negative cosmological constant can be formulated as a CS action for the $\mathfrak{so}(2,2)$ algebra, commonly referred to as the AdS algebra [61–63]. In the limit of vanishing cosmological constant, the underlying symmetry is given by the $\mathfrak{iso}(2,1)$, namely Poincaré, algebra. While the Carrollian regime of relativistic CS gravity can be consistently derived as a UR limit [64–66], the NR counterpart presents significant challenges. Specifically, both the Newton-Hooke [5, 67–69] and the Galilean algebras, which emerge as NR limits of the AdS and Poincaré algebras, respectively, suffer from degeneracy issues of their invariant tensor, key ingredients of field-theoretic CS formulations. In fact, a CS action with a non-degenerate invariant bilinear trace admits a kinetic term for each gauge field which, in three spacetime dimensions, implies the vanishing of the curvature two-forms as field equations in the CS field-theoretical framework. One way to avoid degeneracy in the NR realm is to add two central charges to the corresponding NR algebras, reproducing the extended Newton-Hooke [70–76] and the extended Bargmann algebra [77, 78]. Both NR algebras can be obtained through a contraction process from the $\text{AdS} \oplus \mathfrak{u}(1)^2$ and the $\text{Poincaré} \oplus \mathfrak{u}(1)^2$ algebra, respectively [64].

More recently, an alternative approach to obtain NR symmetries, based on Lie algebra expansion procedures [79–82], has been extensively discussed in [66, 83–90]. Notably, the original “cube” of Bacry and Lévy-Leblond, exhibiting in a compact way the links among algebraic structures, was extended in [90] to include non-degenerate kinematical Lie algebras through the semigroup expansion (S -expansion) method [81]. This approach allows to recover centrally extended NR algebras by expanding relativistic algebras, without introducing $\mathfrak{u}(1)$ generators as required in the contraction process. The S -expansion method has been useful not only for deriving the commutation relations of an expanded algebra but also for obtaining the non-vanishing components of their invariant tensor in terms of the original ones [81]. Furthermore, the CS action for a given expanded Lie algebra can be recovered directly from the CS action of the original algebra by appropriately expanding the corresponding gauge fields.

Although supersymmetric extensions of gravity theories have been largely studied, their NR regime has only recently been approached, mainly due to the lack of clarity about possible NR supermultiplets [91]. Furthermore, the proper derivation of a NR superalgebra cannot be reproduced as a straightforward limit of a relativistic superalgebra as in the bosonic case, and instead requires alternative strategies. The first successful approach to NR supergravity was based on NC geometry in three spacetime dimensions [92, 93]. Later, a non-vanishing torsion was introduced in NC supergravity through a NR superconformal tensor calculus [94]. A third attempt for constructing NR supergravity was possible via the CS formalism in three spacetime dimensions. A CS supergravity action based on a supersymmetric extension of the Bargmann algebra was first presented in [78]. This approach was subsequently applied to develop extended Newtonian supergravity [95], Maxwellian extended Bargmann supergravity [96], extended Newton-Hooke, extended Lifshitz and extended Schrödinger supergravity [97]. The CS formalism has also been useful to construct UR supergravity models based on the Carroll superalgebra [98] and its $\mathcal{N}$ -extensions [99]. More recently, the Lie algebra expansion procedure [80, 81] has been used to derive different NR regimes of known three-dimensional relativistic CS supergravity models [69, 100–105]. Despite differences in the methods employed to construct NR supergravity theories, they are all defined in three spacetime dimensions

and require $\mathcal{N} = 2$ supersymmetry to express the time translation generators as bilinear combinations of supersymmetry generators. To our knowledge, aside from recent developments in ten and eleven spacetime dimensions,¹ no consistent NL supergravity models exist in higher spacetime dimensions, despite extensive investigation into relativistic supergravity theories.

In this work, motivated by the numerous applications of NL symmetries and by the absence of a general framework for exploring NR (super)gravity in spacetime dimensions higher than three, we present a systematic approach to derive NL regimes of relativistic kinematical superalgebras. To this aim, we combine the S -expansion method [81] with the CS formalism to construct three-dimensional NL supergravity actions. The derivation of consistent NL superalgebras is constrained by the non-degeneracy criterion, which allows us to identify “good” NL supersymmetries for constructing CS actions. As a result, we extend the original cube of Bacry and Lévy-Leblond [5] into a supersymmetric one, where the (extended) kinematical superalgebras are characterized by a non-degenerate bilinear invariant trace.

In the UR regime, we show that the S -expansion method used in [90] can be extended to the supersymmetric realm, first recovering the $\mathcal{N} = 2$ (AdS-)Carroll introduced in [99] and then generalizing the results of [99] to a family of generalized (AdS-)Carroll superalgebras. In the NR counterpart, we recover the $\mathcal{N} = 2$ extended Newton-Hooke superalgebra [85, 90] and the extended Bargmann superalgebra [78]. Notably, the S -expansion method allows us to classify these superalgebras within a broader framework inspired by the original cube of Bacry and Lévy-Leblond [5]. In this context, we present novel non-Lorentzian superalgebras, including the $\mathcal{N} = 2$ extended (AdS-)static, generalized Newton-Hooke and generalized Galilean superalgebras. Both the new results and those that were already known correspond to supersymmetric extensions of the general kinematical algebras discussed in [90]. Furthermore, we construct the CS supergravity action based on the kinematical superalgebras obtained here. It is worth emphasizing that the classification of these general kinematical superalgebras could serve for approaching higher-dimensional non-Lorentzian supergravity theories, which remain largely unexplored. To this end, the non-degeneracy criterion will be an essential starting point. Additional details on this issue, as well as potential extensions of our results, are discussed in the concluding section.

This paper is organized as follows: in section 2, we briefly review the extended kinematical Lie algebras which admit a non-degenerate bilinear invariant trace [90]. Sections 3 and 4 contain our main results. In section 3, we classify the supersymmetric extensions of the extended kinematical algebras by applying non- and ultra-relativistic expansions through appropriate semigroups. Section 4 is devoted to the construction of NL supergravity theories based on the aforementioned extended kinematical superalgebras. Section 5 concludes our work with some comments about our results and possible future developments.

2 Extended kinematical Lie algebras

Diverse strategies have been implemented to address the degeneracy issue present in the NR regime of the original kinematical Lie algebra [5]. Notably, the Lie algebra expansion method based on semigroups [81] has demonstrated effectiveness not only to derive NR symmetries

¹Recent results have been obtained in the study of NL version of ten-dimensional minimal supergravity [106] and 11-dimensional supergravity [107, 108].




Figure 1. This cube summarizes the different regimes starting from the AdS algebra. Both relativistic and non-Lorentzian algebras admit a non-degenerate bilinear invariant trace [90].

without degeneracy but also to construct the corresponding NR CS actions [90]. In this section, following [90], we briefly review the different expansions applied, starting from the $\mathfrak{so}(2, 2)$ algebra, which reproduce both the UR limit and the NR expansion (see figure 1).

The commutation relations of $\mathfrak{so}(2, 2)$ are given by

$$[\tilde{J}_A, \tilde{J}_B] = \epsilon_{ABC} \tilde{J}^C, \quad [\tilde{J}_A, \tilde{P}_B] = \epsilon_{ABC} \tilde{P}^C, \quad [\tilde{P}_A, \tilde{P}_B] = \frac{1}{\ell^2} \epsilon_{ABC} \tilde{J}^C, \quad (2.1)$$

where $\tilde{J}_A$ correspond to the Lorentz generators and $\tilde{P}_A$ are the spacetime translations. Here, the AdS radius (length parameter) ℓ is related to the cosmological constant through $\Lambda = -\frac{1}{\ell^2}$. In the vanishing cosmological constant limit $\ell \rightarrow \infty$, the AdS algebra reproduces the Poincaré one. Before applying the expansion, a decomposition of the relativistic rigid (Lorentz) indices $A = 0, 1, 2$ is required, by considering a time-space splitting such that $A = \{0, a\}$ with $a = 1, 2$. Then, implementing this decomposition, the AdS algebra reads

$$\begin{aligned} [J, G_a] &= \epsilon_{ab} G_b, & [J, P_a] &= \epsilon_{ab} P_b, & [G_a, G_b] &= -\epsilon_{ab} J, \\ [H, G_a] &= \epsilon_{ab} P_b, & [G_a, P_b] &= -\epsilon_{ab} H, & [H, P_a] &= \frac{1}{\ell^2} \epsilon_{ab} G_b, \\ [P_a, P_b] &= -\frac{1}{\ell^2} \epsilon_{ab} J, \end{aligned} \quad (2.2)$$

with $\epsilon_{ab} \equiv \epsilon_{0ab}$ and $\epsilon^{ab} \equiv \epsilon^{0ab}$. Here we have relabeled the AdS generators as follows:

$$J = \tilde{J}_0, \quad G_a = \tilde{J}_a, \quad H = \tilde{P}_0, \quad P_a = \tilde{P}_a. \quad (2.3)$$

Let $\mathfrak{so}(2, 2) = V_0 + V_1$ be a subspace decomposition of the $\mathfrak{so}(2, 2)$ algebra whose content depends on the desired expansion as shown in table 1. The subspaces V_0 and V_1 satisfy

Subspaces	NR expansion	UR limit
V_0	J, H	J, P_a
V_1	G_a, P_a	H, G_a

Table 1. Subspace decomposition of the AdS algebra.

a $\mathbb{Z}_2$ -graded Lie algebra,

$$[V_0, V_0] \subset V_0, \quad [V_0, V_1] \subset V_1, \quad [V_1, V_1] \subset V_0. \quad (2.4)$$

The UR limit can be reproduced considering the $S_E^{(1)}$ semigroup, whose elements fulfill the following multiplication rules:

$$\lambda_i \lambda_j = \begin{cases} \lambda_{i+j} & \text{if } i+j \leq 2, \\ \lambda_2 & \text{if } i+j > 2, \end{cases} \quad (2.5)$$

where $\lambda_2 = 0_S$ is the zero element of the semigroup which satisfies $0_S \lambda_i = \lambda_i 0_S = 0_S$. A subset decomposition of the semigroup having the same algebraic structure as the subspace decompositions provided in table 1 is given by

$$S_0 = \{\lambda_0, \lambda_2\}, \quad S_1 = \{\lambda_1, \lambda_2\}. \quad (2.6)$$

Such subset decomposition is said to be in resonance with (2.4), since it satisfies

$$S_0 \cdot S_0 \subset S_0, \quad S_0 \cdot S_1 \subset S_1, \quad S_1 \cdot S_1 \subset S_0. \quad (2.7)$$

On the other hand, the NR expansion requires to consider $S_E^{(2)}$ as the relevant semigroup [90], whose elements satisfy:

$$\lambda_i \lambda_j = \begin{cases} \lambda_{i+j} & \text{if } i+j \leq 3, \\ \lambda_3 & \text{if } i+j > 3, \end{cases} \quad (2.8)$$

with $\lambda_3 = 0_S$ being the zero element of the semigroup. A resonant subset decomposition of the semigroup reads

$$S_0 = \{\lambda_0, \lambda_2, \lambda_3\}, \quad S_1 = \{\lambda_1, \lambda_3\}. \quad (2.9)$$

Each NL regime appearing in the cube of figure 1 can be obtained after applying an S_E -resonant expansion of the relativistic $\mathfrak{so}(2, 2)$ algebra:

$$\mathfrak{G} = (S_0 \times V_0) \oplus (S_1 \times V_1), \quad (2.10)$$

and performing a 0_S -reduction, namely $0_S T_A = 0$. The S -expanded generators are written in terms of the $\mathfrak{so}(2, 2)$ ones and the semigroup elements as in table 2.

The explicit commutation relations of the expanded algebra are listed in table 3 and are derived by combining the multiplication rule of the semigroup S_E and the commutators of the starting algebra. Let us note that the AdS-Carroll algebra can be alternatively recovered

	NR expansion		UR limit	
λ_3				
λ_2	S, M			
λ_1		G_a, P_a		H, G_a
λ_0	J, H		J, P_a	
	J, H	G_a, P_a	J, P_a	H, G_a

Table 2. Expanded generators in terms of the relativistic ones and the semigroup elements. The relativistic generators have been organized according to the subspace decomposition of table 1.

	AdS-Carroll	Extended Newton-Hooke	Extended AdS-static
$[J, G_a]$	$\epsilon_{ab} G_b$	$\epsilon_{ab} G_b$	$\epsilon_{ab} G_b$
$[J, P_a]$	$\epsilon_{ab} P_b$	$\epsilon_{ab} P_b$	$\epsilon_{ab} P_b$
$[G_a, G_b]$	0	$-\epsilon_{ab} S$	0
$[H, G_a]$	0	$\epsilon_{ab} P_b$	0
$[G_a, P_b]$	$-\epsilon_{ab} H$	$-\epsilon_{ab} M$	$-\epsilon_{ab} M$
$[H, P_a]$	$\frac{1}{\ell^2} \epsilon_{ab} G_b$	$\frac{1}{\ell^2} \epsilon_{ab} G_b$	$\frac{1}{\ell^2} \epsilon_{ab} G_b$
$[P_a, P_b]$	$-\frac{1}{\ell^2} \epsilon_{ab} J$	$-\frac{1}{\ell^2} \epsilon_{ab} S$	$-\frac{1}{\ell^2} \epsilon_{ab} S$

Table 3. Commutation relations of the (extended) kinematical algebras. The vanishing cosmological constant limit $\ell \rightarrow \infty$ reproduces the Carroll algebra, the extended Bargmann algebra, and the extended static algebra, respectively.

as a resonant $S_E^{(1)}$ -expansion of $\mathfrak{so}(2, 2)$. Despite the isomorphism existing between the AdS-Carroll and the Poincaré algebra, exchanging the generators G_a and P_a leads to quite distinct physical implications [5]. On the other hand, from the cube of figure 1, the Carroll algebra can either be recovered as an $S_E^{(1)}$ -expansion of the Poincaré algebra or as a flat limit $\ell \rightarrow \infty$ applied to the AdS-Carroll one.

In the NR counterpart, we obtain extended kinematical algebras which are characterized by the presence of central charges. Such central extensions solve the degeneracy present in the NR regime of the original kinematical algebra [5]. Here, the extended Newton-Hooke [70–76] and extended Bargmann [77, 78] algebras appear after performing the $S_E^{(2)}$ -expansion of $\mathfrak{so}(2, 2)$ and the Poincaré algebra, respectively. Extensions of the static algebra, which have been related to infinitely massive systems that cannot move [5], can also be derived as expansions. Here, the extended AdS-static algebra and its flat limit can be obtained either as a resonant $S_E^{(2)}$ -expansion of the Carrollian algebras or as a resonant $S_E^{(1)}$ -expansion of the extended NR ones. However, as one can see from table 4, different subspace decompositions of the starting algebra have to be considered.

3 Supersymmetric extensions of extended kinematical algebras

In this section, we explore the supersymmetric extension of the kinematical Lie algebras [5] in three-dimensional spacetime, using the semigroup expansion method [81], starting from an

	NR expansion		UR limit	
λ_3				
λ_2	S, M			
λ_1		G_a, P_a		H, M, G_a
λ_0	J, H		J, S, P_a	
	J, H	G_a, P_a	J, S, P_a	H, M, G_a

Table 4. Extended (AdS-)static generators in terms of the non-Lorentzian ones and the semigroup elements. The UR limit column contains expanded kinematical generators in terms of the extended (AdS-)Carroll ones. On the other hand, the extended (AdS-)static generators are obtained from the extended Bargmann (or extended Newton-Hooke) ones in the NR expansion column.

$\mathcal{N} = 2$ relativistic superalgebra. The use of $\mathcal{N} = 2$ ensures that the expanded non-relativistic superalgebras yield genuine supergravity actions, in which the time translational generators can be expressed as bilinear combinations of supersymmetry generators. Additionally, in the NR regime we will derive supersymmetric extensions of the extended kinematic algebras discussed in [90], which are distinguished by a non-degenerate invariant bilinear form. The non-degeneracy of the three-dimensional CS action guarantees a kinetic term for each gauge field leading to the vanishing of the curvatures as equations of motion.

In three spacetime dimensions, the $\mathcal{N} = 2$ supersymmetric extension of the AdS algebra with a well-defined Poincaré limit $\ell \rightarrow \infty$ is given by an $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra whose (anti-)commutation relations are given by [109]:

$$\begin{aligned}
 [\tilde{J}_A, \tilde{J}_B] &= \epsilon_{ABC} \tilde{J}^C, \\
 [\tilde{J}_A, \tilde{P}_B] &= \epsilon_{ABC} \tilde{P}^C, \\
 [\tilde{P}_A, \tilde{P}_B] &= \frac{1}{\ell^2} \epsilon_{ABC} \tilde{J}^C, \\
 [\tilde{J}_A, \tilde{Q}_\alpha^i] &= -\frac{1}{2} (\gamma_A)_\alpha^\beta \tilde{Q}_\beta^i, \\
 [\tilde{P}_A, \tilde{Q}_\alpha^i] &= -\frac{1}{2\ell} (\gamma_A)_\alpha^\beta \tilde{Q}_\beta^i, \\
 [\tilde{\mathcal{T}}, \tilde{Q}_\alpha^i] &= \frac{1}{2} \epsilon^{ij} \tilde{Q}_\beta^j, \\
 \{\tilde{Q}_\alpha^i, \tilde{Q}_\beta^j\} &= -\frac{1}{\ell} \delta_{ij} (\gamma^A C)_{\alpha\beta} \tilde{J}_A - \delta_{ij} (\gamma^A C)_{\alpha\beta} \tilde{P}_A - C_{\alpha\beta} \epsilon^{ij} \left(\tilde{\mathcal{U}} + \frac{1}{\ell} \tilde{\mathcal{T}} \right), \quad (3.1)
 \end{aligned}$$

where $\tilde{Q}_\alpha$ is a Majorana spinor charge. On the other hand, the presence of the generators $\{\tilde{\mathcal{T}}, \tilde{\mathcal{U}}\}$ ensures the non-degeneracy of the bilinear invariant form in the Poincaré limit in which $\tilde{\mathcal{U}}$ becomes the central charge. Here, $A, B, \dots = 0, 1, 2$ denote the Lorentz indices which are raised and lowered with the Minkowski metric $\eta_{AB} = (-1, 1, 1)$ and ϵ_{ABC} is the Levi-Civita tensor, while $\alpha, \beta = 1, 2$, $i, j = 1, 2$ denote the number of supercharges, the gamma matrices in three dimensions are denoted by γ_A , and C is the charge conjugation matrix, satisfying $C^T = -C$ and $C\gamma^A = (C\gamma^A)^T$. The $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra admits

the following non-vanishing components of a non-degenerate invariant tensor:

$$\begin{aligned}
 \langle \tilde{J}_A \tilde{J}_B \rangle &= \tilde{\alpha}_0 \eta_{AB}, & \langle \tilde{J}_A \tilde{P}_B \rangle &= \tilde{\alpha}_1 \eta_{AB}, \\
 \langle \tilde{P}_A \tilde{P}_B \rangle &= \frac{\tilde{\alpha}_0}{\ell^2} \eta_{AB}, & \langle \tilde{T} \tilde{T} \rangle &= \tilde{\alpha}_0, \\
 \langle \tilde{T} \tilde{U} \rangle &= \tilde{\alpha}_1, & \langle \tilde{U} \tilde{U} \rangle &= -\frac{\tilde{\alpha}_1}{\ell}, \\
 \langle \tilde{Q}_\alpha^i \tilde{Q}_\beta^j \rangle &= 2 \left(\tilde{\alpha}_1 + \frac{\tilde{\alpha}_0}{\ell} \right) C_{\alpha\beta} \delta^{ij}.
 \end{aligned} \tag{3.2}$$

Before applying the expansion, let us consider a time-space splitting of the relativistic Lorentz index $A = \{0, a\}$ with $a = 1, 2$. Then, the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra is spanned by $\{J, G_a, H, P_a, T, U, Q_\alpha^\pm\}$, where we have considered the redefinitions (2.3) together with the following ones:

$$T = \tilde{T}, \quad U = \tilde{U}, \quad Q_\alpha^\pm = \frac{1}{\sqrt{2}} \left(\tilde{Q}_\alpha^1 \pm (\gamma^0)_{\alpha\beta} \tilde{Q}_\beta^2 \right). \tag{3.3}$$

Then, the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra can be written as

$$\begin{aligned}
 [J, G_a] &= \epsilon_{ab} G_b, & [J, P_a] &= \epsilon_{ab} P_b, & [G_a, G_b] &= -\epsilon_{ab} J, \\
 [H, G_a] &= \epsilon_{ab} P_b, & [G_a, P_b] &= -\epsilon_{ab} H, & [H, P_a] &= \frac{1}{\ell^2} \epsilon_{ab} G_b, \\
 [P_a, P_b] &= -\frac{1}{\ell^2} \epsilon_{ab} J, & [J, Q_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \tilde{Q}_\beta^\pm, & [H, Q_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta \tilde{Q}_\beta^\pm, \\
 [G_a, Q_\alpha^\pm] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta \tilde{Q}_\beta^\mp, & [P_a, Q_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta \tilde{Q}_\beta^\mp, & [T, Q_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_{\alpha\beta} \tilde{Q}_\beta^\pm, \\
 \{Q_\alpha^+, Q_\beta^+\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} J + H \right) - (\gamma^0 C)_{\alpha\beta} \left(U + \frac{1}{\ell} T \right), \\
 \{Q_\alpha^+, Q_\beta^-\} &= -(\gamma^a C)_{\alpha\beta} \left(\frac{1}{\ell} G_a + P_a \right), \\
 \{Q_\alpha^-, Q_\beta^-\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} J + H \right) + (\gamma^0 C)_{\alpha\beta} \left(U + \frac{1}{\ell} T \right).
 \end{aligned} \tag{3.4}$$

3.1 Ultra-relativistic expansions

The Carrollian regime of the kinematical superalgebras can be obtained by expanding the relativistic ones through a pertinent semigroup and by implementing an explicit reduction procedure. Interestingly, a family of UR superalgebras can be derived by considering larger semigroups. Before applying the UR expansion, we require to consider a particular subspace decomposition of the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ given by

$$V_0 = \{J, P_a, T\}, \quad V_1 = \{Q_\alpha^+, Q_\alpha^-\}, \quad V_2 = \{H, G_a, U\}. \tag{3.5}$$

Here, the subspaces V_0 , V_1 , and V_2 satisfy the following algebraic structure:

$$\begin{aligned}
 [V_0, V_0] &\subset V_0, & [V_1, V_1] &\subset V_0 \oplus V_2, \\
 [V_0, V_1] &\subset V_1, & [V_1, V_2] &\subset V_1, \\
 [V_0, V_2] &\subset V_2, & [V_2, V_2] &\subset V_0.
 \end{aligned} \tag{3.6}$$

3.1.1 The $\mathcal{N} = 2$ (AdS-)Carroll superalgebra

Let $S_E^{(2)} = \{\lambda_0, \lambda_1, \lambda_2, \lambda_3\}$ be the finite Abelian semigroup whose elements obey the multiplication law (2.8) and whose zero element is given by λ_3 . A subset decomposition of the semigroup $S_E^{(2)} = S_0 \cup S_1 \cup S_2$, with

$$S_0 = \{\lambda_0, \lambda_2, \lambda_3\}, \quad S_1 = \{\lambda_1, \lambda_3\}, \quad S_2 = \{\lambda_2, \lambda_3\}, \quad (3.7)$$

is said to be resonant since it satisfies the same algebraic structure (3.6) as the subspaces V_0 , V_1 , and V_2 , namely

$$\begin{aligned} S_0 \cdot S_0 &\subset S_0, & S_1 \cdot S_1 &\subset S_0 \cap S_2, \\ S_0 \cdot S_1 &\subset S_1, & S_1 \cdot S_2 &\subset S_1, \\ S_0 \cdot S_2 &\subset S_2, & S_2 \cdot S_2 &\subset S_0. \end{aligned} \quad (3.8)$$

According to [81], the superalgebra $\mathfrak{G}_R = (S_0 \times V_0) \oplus (S_1 \times V_1) \oplus (S_2 \times V_2)$ corresponds to a resonant super subalgebra of the $S_E^{(2)}$ -expanded one. A smaller superalgebra can be extracted from the resonant one through a reduction procedure [81]. Let us consider a partition of the subsets $S_p = \hat{S}_p \cup \check{S}_p$ with

$$\begin{aligned} \check{S}_0 &= \{\lambda_0\}, & \hat{S}_0 &= \{\lambda_2, \lambda_3\}, \\ \check{S}_1 &= \{\lambda_1\}, & \hat{S}_1 &= \{\lambda_3\}, \\ \check{S}_2 &= \{\lambda_2\}, & \hat{S}_2 &= \{\lambda_3\}. \end{aligned} \quad (3.9)$$

Such partition satisfies $\hat{S}_p \cap \check{S}_p = \emptyset$ together with the following algebraic structure:

$$\begin{aligned} \check{S}_0 \cdot \hat{S}_0 &\subset \hat{S}_0, & \check{S}_1 \cdot \hat{S}_1 &\subset \hat{S}_0 \cap \hat{S}_2, \\ \check{S}_0 \cdot \hat{S}_1 &\subset \hat{S}_1, & \check{S}_1 \cdot \hat{S}_2 &\subset \hat{S}_1, \\ \check{S}_0 \cdot \hat{S}_2 &\subset \hat{S}_2, & \check{S}_2 \cdot \hat{S}_2 &\subset \hat{S}_0. \end{aligned} \quad (3.10)$$

Then, we have

$$\begin{aligned} \check{\mathfrak{G}}_R &= (\check{S}_0 \times V_0) \oplus (\check{S}_1 \times V_1) \oplus (\check{S}_2 \times V_2), \\ \hat{\mathfrak{G}}_R &= (\hat{S}_0 \times V_0) \oplus (\hat{S}_1 \times V_1) \oplus (\hat{S}_2 \times V_2), \end{aligned} \quad (3.11)$$

where $\check{\mathfrak{G}}_R$ corresponds to a *reduced* superalgebra of $\mathfrak{G}_R$ which satisfies

$$[\check{\mathfrak{G}}_R, \hat{\mathfrak{G}}_R] \subset \hat{\mathfrak{G}}_R. \quad (3.12)$$

The expanded generators of the reduced superalgebra $\check{\mathfrak{G}}_R$ are written in terms of the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ ones and the semigroup elements as in table 5.

The (anti-)commutation relations of the reduced superalgebra $\check{\mathfrak{G}}_R$ are obtained by combining the multiplication law of the $S_E^{(2)}$ semigroup and the (anti-)commutators of the

λ_3			
λ_2			H, G_a , U
λ_1		Q_α^+ , Q_α^-	
λ_0	J, P_a , T		
	J , P_a , T	Q_α^+ , Q_α^-	H , G_a , U

Table 5. Expanded generators of the reduced superalgebra $\check{\mathfrak{G}}_R$ in terms of the relativistic ones and the semigroup elements.

$\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2,2) \otimes \mathfrak{sp}(2)$ superalgebra (3.4),

$$\begin{aligned}
 [J, G_a] &= \epsilon_{ab} G_b, & [J, P_a] &= \epsilon_{ab} P_b, & [P_a, P_b] &= -\frac{1}{\ell^2} \epsilon_{ab} J, \\
 [H, P_a] &= \frac{1}{\ell^2} \epsilon_{ab} G_b, & [G_a, P_a] &= -\epsilon_{ab} H, & [J, Q_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta Q_\beta^\pm, \\
 [P_a, Q_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta Q_\beta^\mp, & [T, Q_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta Q_\beta^\pm, \\
 \{Q_\alpha^+, Q_\beta^+\} &= -(\gamma^0 C)_{\alpha\beta} (H + U), \\
 \{Q_\alpha^+, Q_\beta^-\} &= -\frac{1}{\ell} (\gamma^a C)_{\alpha\beta} G_a, \\
 \{Q_\alpha^-, Q_\beta^-\} &= -(\gamma^0 C)_{\alpha\beta} (H - U).
 \end{aligned} \tag{3.13}$$

The reduced superalgebra $\check{\mathfrak{G}}_R$ corresponds to the $\mathcal{N} = 2$ AdS-Carroll superalgebra² introduced in [99] and differs from the one presented in [110] due to the presence of the U generator. In the ultra-relativistic regime, the U generator becomes a central charge as in the Poincaré superalgebra. In particular, the $\mathcal{N} = 2$ AdS-Carroll superalgebra (3.13) is isomorphic to the $\mathcal{N} = 2$ Poincaré superalgebra under exchange of the ℓP_a and G_a generators. In the vanishing cosmological constant limit $\ell \rightarrow \infty$, the superalgebra reproduces the $\mathcal{N} = 2$ Carroll superalgebra. Interestingly, the $\mathcal{N} = 2$ Carroll superalgebra can alternatively be recovered by S -expanding the $\mathcal{N} = 2$ Poincaré one following the same procedure used to obtain the $\mathcal{N} = 2$ AdS-Carroll superalgebra (see figure 2).

3.1.2 Generalized $\mathcal{N} = 2$ AdS-Carroll superalgebras

A large semigroup $S_E^{(2N)}$ can be considered to obtain generalized $\mathcal{N} = 2$ AdS-Carroll superalgebras from the $\mathfrak{osp}(2,2) \otimes \mathfrak{sp}(2)$ superalgebra. Let us consider first $S_E^{(2N)} = \{\lambda_0, \lambda_1, \lambda_2, \dots, \lambda_{2N+1}\}$ as the finite semigroup whose elements satisfy

$$\lambda_\alpha \lambda_\beta = \begin{cases} \lambda_{\alpha+\beta} & \text{if } \alpha + \beta \leq 2N + 1, \\ \lambda_{2N+1} & \text{if } \alpha + \beta > 2N + 1, \end{cases} \tag{3.14}$$

²There are two inequivalent $\mathcal{N} = 2$ AdS-Carroll superalgebras: the so-called $\mathcal{N} = (1,1)$ and the $\mathcal{N} = (2,0)$. Here, $\mathcal{N} = 2$ refers to $\mathcal{N} = (2,0)$, since we started from the $\mathcal{N} = (2,0)$ AdS superalgebra. An $\mathcal{N} = (1,1)$ AdS-Carroll superalgebra can also be obtained from the $\mathcal{N} = (1,1)$ AdS one following the same procedure considered here.




Figure 2. Diagram summarizing the (reduced) UR expansion and flat limit starting from the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2,2) \otimes \mathfrak{sp}(2)$ superalgebra.

with $\lambda_{2N+1} = 0_S$ being the zero element of the semigroup. A subset decomposition of the semigroup $S_E^{(2N)} = S_0 \cup S_1 \cup S_2$ is given by

$$\begin{aligned} S_0 &= \{\lambda_{2m}, \text{ with } m = 0, \dots, N\} \cup \{\lambda_{2N+1}\}, \\ S_1 &= \{\lambda_{2m+1}, \text{ with } m = 0, \dots, N-1\} \cup \{\lambda_{2N+1}\}, \\ S_2 &= \{\lambda_{2m+2}, \text{ with } m = 0, \dots, N-1\} \cup \{\lambda_{2N+1}\}. \end{aligned} \quad (3.15)$$

The decomposition (3.15) is resonant since it satisfies the same algebraic structure as the subspace decomposition (3.6) of the original $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2,2) \otimes \mathfrak{sp}(2)$ superalgebra. The resonant expanded superalgebra is given by

$$\mathfrak{G}_R = (S_0 \times V_0) \oplus (S_1 \times V_1) \oplus (S_2 \times V_2), \quad (3.16)$$

where V_0 , V_1 , and V_2 are the subspaces of the starting relativistic superalgebra given in (3.5). A smaller superalgebra can be extracted from $\mathfrak{G}_R$ by considering a partition of the subsets $S_p = \hat{S}_p \cup \check{S}_p$ with

$$\begin{aligned} \check{S}_0 &= \{\lambda_{2m}, \text{ with } m = 0, \dots, N-1\}, & \hat{S}_0 &= \{\lambda_{2N}, \lambda_{2N+1}\}, \\ \check{S}_1 &= \{\lambda_{2m+1}, \text{ with } m = 0, \dots, N-1\}, & \hat{S}_1 &= \{\lambda_{2N+1}\}, \\ \check{S}_2 &= \{\lambda_{2m+2}, \text{ with } m = 0, \dots, N-1\}, & \hat{S}_2 &= \{\lambda_{2N+1}\}. \end{aligned} \quad (3.17)$$

One can check that this subset partition satisfies the conditions (3.10) and $\hat{S}_p \cap \check{S}_p = \emptyset$. Both conditions ensure that

$$[\check{\mathfrak{G}}_R, \hat{\mathfrak{G}}_R] \subset \hat{\mathfrak{G}}_R, \quad (3.18)$$

where

$$\begin{aligned} \check{\mathfrak{G}}_R &= (\check{S}_0 \times V_0) \oplus (\check{S}_1 \times V_1) \oplus (\check{S}_2 \times V_2), \\ \hat{\mathfrak{G}}_R &= (\hat{S}_0 \times V_0) \oplus (\hat{S}_1 \times V_1) \oplus (\hat{S}_2 \times V_2). \end{aligned} \quad (3.19)$$

Then, $\check{G}_R$ is a reduced superalgebra of the resonant superalgebra $\mathfrak{G}_R$ [81]. The generators of the reduced superalgebra $\check{G}_R$ are related to the relativistic ones through the semigroup elements as follows:

$$\begin{aligned} J^{(m)} &= \lambda_{2m} J, & P_a^{(m)} &= \lambda_{2m} P_a, & T^{(m)} &= \lambda_{2m} T, \\ H^{(m)} &= \lambda_{2m+2} H, & G_a^{(m)} &= \lambda_{2m+2} G_a, & U^{(m)} &= \lambda_{2m+2} U, \\ Q_\alpha^{+(m)} &= \lambda_{2m+1} Q_\alpha^+, & Q_\alpha^{-(m)} &= \lambda_{2m+1} Q_\alpha^-, \end{aligned} \quad (3.20)$$

where $m = 0, \dots, N-1$. The $\check{G}_R$ generators satisfy the following commutation relations:

$$\begin{aligned} [J^{(m)}, G_a^{(n)}] &= \epsilon_{ab} G_b^{(m+n)}, & [G_a^{(m)}, G_b^{(n)}] &= -\epsilon_{ab} J^{(m+n+2)}, \\ [J^{(m)}, P_a^{(n)}] &= \epsilon_{ab} P_b^{(m+n)}, & [G_a^{(m)}, P_a^{(n)}] &= -\epsilon_{ab} H^{(m+n)}, \\ [H^{(m)}, P_a^{(n)}] &= \frac{1}{\ell^2} \epsilon_{ab} G_b^{(m+n)}, & [P_a^{(m)}, P_b^{(n)}] &= -\frac{1}{\ell^2} \epsilon_{ab} J^{(m+n)}, \\ [H^{(m)}, G_a^{(n)}] &= \epsilon_{ab} P_b^{(m+n+2)}, & [J^{(m)}, Q_\alpha^\pm(n)] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta Q_\beta^\pm(m+n), \\ [H^{(m)}, Q_\alpha^\pm(n)] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta Q_\beta^\pm(m+n+1), & [P_a^{(m)}, Q_\alpha^\pm(n)] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta Q_\beta^\mp(m+n), \\ [G_a^{(m)}, Q_\alpha^\pm(n)] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta Q_\beta^\mp(m+n+1), & [T^{(m)}, Q_\alpha^\pm(n)] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta Q_\beta^\pm(m+n), \end{aligned} \quad (3.21)$$

together with the following anti-commutators:

$$\begin{aligned} \{Q_\alpha^{+(m)}, Q_\beta^{+(n)}\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} J^{(m+n+1)} + H^{(m+n)} \right) - (\gamma^0 C)_{\alpha\beta} \left(U^{(m+n)} + \frac{1}{\ell} T^{(m+n+1)} \right), \\ \{Q_\alpha^{+(m)}, Q_\beta^{-(n)}\} &= -(\gamma^a C)_{\alpha\beta} \left(\frac{1}{\ell} G_a^{(m+n)} + P_a^{(m+n+1)} \right), \\ \{Q_\alpha^{-(m)}, Q_\beta^{-(n)}\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} J^{(m+n+1)} + H^{(m+n)} \right) + (\gamma^0 C)_{\alpha\beta} \left(U^{(m+n)} + \frac{1}{\ell} T^{(m+n+1)} \right). \end{aligned} \quad (3.22)$$

Here, we considered the multiplication laws of the $S_E^{(2N)}$ semigroup (3.14) and the (anti-)commutation relations of the $\mathfrak{so}(2)$ -extension of $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ (3.4). In particular, for $m+n+1 > N-1$ we have $T_A^{(m+n+1)} = 0$. The obtained $\mathcal{N} = 2$ superalgebra corresponds to a generalization of the $\mathcal{N} = 2$ AdS-Carroll superalgebra which we have denoted as the $\mathcal{N} = 2$ **ads-car**^(N) superalgebra.³ In the flat limit $\ell \rightarrow \infty$, we find a generalized $\mathcal{N} = 2$ Carroll superalgebra denoted as $\mathcal{N} = 2$ **car**^(N), which can be seen as the higher-dimensional Carrollian analogue of supersymmetric extensions of the (post-)Newtonian symmetries discussed in [95, 111, 112]. Alternatively, the $\mathcal{N} = 2$ **car**^(N) superalgebra can also be derived from the $\mathcal{N} = 2$ Poincaré superalgebra considering the same semigroup and series of steps employed to obtain the **ads-car**^(N) superalgebra (see figure 3).

One can check that, for $N = 1$ (i.e., considering $S_E^{(2)}$ in the expansion procedure), the $\mathcal{N} = 2$ super **ads-car**⁽¹⁾ reproduces the $\mathcal{N} = 2$ AdS-Carroll superalgebra (3.13). A novel $\mathcal{N} = 2$ AdS-Carroll superalgebra appears for $N = 2$ which involves the inclusion of two additional

³The **ads-car**^(N) superalgebra can also be seen as a $\mathcal{N} = 2$ supersymmetric extension of the generalized para-Poincaré algebra introduced in [90].

Majorana fermionic charges $\mathbf{F}_\alpha^\pm$. The generators of the $\mathcal{N} = 2$ $\mathfrak{ad}\mathfrak{s}\text{-car}^{(2)}$ superalgebra satisfy the following commutators:

$$\begin{aligned}
 [\mathbf{J}, \mathbf{G}_a] &= \epsilon_{ab} \mathbf{G}_b, & [\mathbf{G}_a, \mathbf{P}_a] &= -\epsilon_{ab} \mathbf{H}, & [\mathbf{J}, \mathbf{P}_a] &= \epsilon_{ab} \mathbf{P}_b, \\
 [\mathbf{H}, \mathbf{P}_a] &= \frac{1}{\ell^2} \epsilon_{ab} \mathbf{G}_b, & [\mathbf{P}_a, \mathbf{P}_b] &= -\frac{1}{\ell^2} \epsilon_{ab} \mathbf{J}, & [\mathbf{J}, \mathbf{M}_a] &= \epsilon_{ab} \mathbf{M}_b, \\
 [\mathbf{J}, \mathbf{Z}_a] &= \epsilon_{ab} \mathbf{Z}_b, & [\mathbf{P}_a, \mathbf{M}_b] &= -\frac{1}{\ell^2} \epsilon_{ab} \mathbf{K}, & [\mathbf{K}, \mathbf{G}_a] &= \epsilon_{ab} \mathbf{Z}_b, \\
 [\mathbf{Z}, \mathbf{P}_a] &= \frac{1}{\ell^2} \epsilon_{ab} \mathbf{Z}_b, & [\mathbf{P}_a, \mathbf{Z}_b] &= -\epsilon_{ab} \mathbf{Z}, & [\mathbf{K}, \mathbf{P}_a] &= \epsilon_{ab} \mathbf{M}_b, \\
 [\mathbf{H}, \mathbf{M}_a] &= \frac{1}{\ell^2} \epsilon_{ab} \mathbf{Z}_b, & [\mathbf{G}_a, \mathbf{M}_b] &= -\epsilon_{ab} \mathbf{Z}, & [\mathbf{J}, \mathbf{Q}_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{Q}_\beta^\pm, \\
 [\mathbf{P}_a, \mathbf{Q}_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta \mathbf{Q}_\beta^\mp, & [\mathbf{H}, \mathbf{Q}_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta \mathbf{F}_\beta^\pm, & [\mathbf{G}_a, \mathbf{Q}_\alpha^\pm] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta \mathbf{F}_\beta^\mp, \\
 [\mathbf{J}, \mathbf{F}_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{F}_\beta^\pm, & [\mathbf{P}_a, \mathbf{F}_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta \mathbf{F}_\beta^\mp, & [\mathbf{K}, \mathbf{Q}_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{F}_\beta^\pm, \\
 [\mathbf{M}_a, \mathbf{Q}_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta \mathbf{F}_\beta^\mp, & [\mathbf{T}_1, \mathbf{Q}_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{Q}_\beta^\pm, & [\mathbf{T}_1, \mathbf{F}_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{F}_\beta^\pm, \\
 [\mathbf{T}_2, \mathbf{Q}_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{F}_\beta^\pm, & & & &
 \end{aligned} \tag{3.23}$$

and anti-commutators:

$$\begin{aligned}
 \{\mathbf{Q}_\alpha^\pm, \mathbf{Q}_\beta^\pm\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{K} + \mathbf{H} \right) \mp (\gamma^0 C)_{\alpha\beta} \left(\mathbf{U}_1 + \frac{1}{\ell} \mathbf{T}_2 \right), \\
 \{\mathbf{Q}_\alpha^+, \mathbf{Q}_\beta^-\} &= -(\gamma^a C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{G}_a + \mathbf{M}_a \right), \\
 \{\mathbf{Q}_\alpha^\pm, \mathbf{F}_\beta^\pm\} &= -(\gamma^0 C)_{\alpha\beta} (\mathbf{Z} \pm \mathbf{U}_2), \\
 \{\mathbf{Q}_\alpha^\pm, \mathbf{F}_\beta^\mp\} &= -\frac{1}{\ell} (\gamma^a C)_{\alpha\beta} \mathbf{Z}_a,
 \end{aligned} \tag{3.24}$$

where we have defined

$$\begin{aligned}
 \mathbf{J}^{(0)} &= \mathbf{J}, & \mathbf{J}^{(1)} &= \mathbf{K}, & \mathbf{H}^{(0)} &= \mathbf{H}, & \mathbf{H}^{(1)} &= \mathbf{Z}, \\
 \mathbf{G}_a^{(0)} &= \mathbf{G}_a, & \mathbf{G}_a^{(1)} &= \mathbf{Z}_a, & \mathbf{P}_a^{(0)} &= \mathbf{P}_a, & \mathbf{P}_a^{(1)} &= \mathbf{M}_a, \\
 \mathbf{T}^{(0)} &= \mathbf{T}_1, & \mathbf{T}^{(1)} &= \mathbf{T}_2, & \mathbf{U}^{(0)} &= \mathbf{U}_1, & \mathbf{U}^{(1)} &= \mathbf{U}_2, \\
 \mathbf{Q}_\alpha^{\pm(0)} &= \mathbf{Q}_\alpha^\pm, & \mathbf{Q}_\alpha^{\pm(1)} &= \mathbf{F}_\alpha^\pm.
 \end{aligned} \tag{3.25}$$

In the flat limit $\ell \rightarrow \infty$, the (anti-)commutation relations (3.23) and (3.24) reproduces an $\mathcal{N} = 2$ $\mathfrak{car}^{(2)}$ superalgebra which can alternatively be obtained as a resonant $\check{S}_E^{(4)}$ -expansion of the $\mathcal{N} = 2$ Poincaré superalgebra. As an ending remark, one could consider $S_E^{(2N+1)}$ as the relevant semigroup applied to the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra. Nonetheless, as we shall discuss in next section, the non-degeneracy of the invariant bilinear trace requires $\langle \mathbf{T}^{(m)} \mathbf{U}^{(m)} \rangle \neq 0$, which is fulfilled only for superalgebras obtained with $S_E^{(2N)}$.

3.2 Non-relativistic expansions

The NR sector of the kinematical superalgebra without degeneracy can be derived by expanding the $\mathcal{N} = 2$ relativistic superalgebras with $S_E^{(2)}$ as the relevant semigroup. As we




Figure 3. Diagram summarizing the generalized (reduced) UR expansion and flat limit starting from the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2,2) \otimes \mathfrak{sp}(2)$ superalgebra.




Figure 4. This cube summarizes the different expansion relations starting from the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2,2) \otimes \mathfrak{sp}(2)$ superalgebra.

shall see, the $S_E^{(2)}$ -expansion can also be applied to the Carrollian superalgebra obtained previously, which reproduces an $\mathcal{N}=2$ extended AdS-static superalgebra and its flat limit (see figure 4). The static sector can alternatively be recovered from the $\mathcal{N}=2$ NR superalgebras by applying a reduced S -expansion following the same procedure used to derive the $\mathcal{N}=2$ $(\mathfrak{ads})\ \mathfrak{car}^{(2)}$ superalgebra. Then, the expansion made with the $S_E^{(2)}$ semigroup along its reduced subset $\check{S}_E^{(2)}$ allows us to define the supersymmetric extension of the extended kinematical algebras [90]. Therefore, the NR expansions and UR ones not only extend the cube of Bacry and Lévy-Leblond [5] to supersymmetry but also solve the degeneracy issue presented in the original cube.

λ_3		
λ_2	S, M, T ₂ , U ₂ , R _α	
λ_1		G _a , P _a , Q _α ⁻
λ_0	J, H, T ₁ , U ₁ , Q _α ⁺	
	J, H, T, U, Q _α ⁺	G _a , P _a , Q _α ⁻

Table 6. Expanded generators in terms of the relativistic $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ ones and the semigroup elements.

3.2.1 The $\mathcal{N} = 2$ extended Newton-Hooke and extended Bargmann superalgebras

Before applying the NR expansion, we first consider a subspace decomposition of the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra different from the one considered in the UR expansion:

$$V_0 = \{J, H, T, U, Q_\alpha^+\}, \quad V_1 = \{G_a, P_a, Q_\alpha^-\}, \quad (3.26)$$

which satisfy a $\mathbb{Z}_2$ -graded Lie algebra (2.4). Let $S_E^{(2)}$ be the finite semigroup whose elements satisfy the multiplication rule (2.8) and with λ_3 being the zero element of the semigroup. A resonant decomposition of the semigroup $S_E^{(2)}$ is given by

$$S_0 = \{\lambda_0, \lambda_2, \lambda_3\}, \quad S_1 = \{\lambda_1, \lambda_3\}, \quad (3.27)$$

which satisfy the same algebraic structure as the subspaces of the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra. Then, an NR superalgebra is obtained after performing a resonant $S_E^{(2)}$ -expansion of the relativistic $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$,

$$\mathfrak{G} = (S_0 \times V_0) \oplus (S_1 \times V_1), \quad (3.28)$$

and considering the 0_S -reduction $\lambda_3 T_A = 0$. The expanded generators are related to the relativistic ones through the corresponding semigroup elements as in table 6.

The (anti-)commutators of the resonant $S_E^{(2)}$ -expansion of the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra are obtained by combining the relativistic (anti-)commutation relations (3.4) and the multiplication rule of the $S_E^{(2)}$ semigroup (2.8). In particular, the expanded generators satisfy the following commutation relations:

$$\begin{aligned}
 [G_a, G_b] &= -\epsilon_{ab} S, & [J, G_a] &= \epsilon_{ab} G_b, & [J, P_a] &= \epsilon_{ab} P_b, \\
 [G_a, P_b] &= -\epsilon_{ab} M, & [H, G_a] &= \epsilon_{ab} P_b, & [H, P_a] &= \frac{1}{\ell^2} \epsilon_{ab} G_b, \\
 [P_a, P_b] &= -\frac{1}{\ell^2} \epsilon_{ab} S, & [J, Q_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta Q_\beta^\pm, & [J, R_\alpha] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta R_\beta, \\
 [H, Q_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta Q_\beta^\pm, & [H, R_\alpha] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta R_\beta, & [S, Q_\alpha^+] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta R_\beta, \\
 [M, Q_\alpha^+] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta R_\beta, & [G_a, Q_\alpha^+] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta Q_\beta^-, & [G_a, Q_\alpha^-] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta R_\beta, \\
 [P_a, Q_\alpha^+] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta Q_\beta^-, & [P_a, Q_\alpha^-] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta R_\beta, & [T_1, Q_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta Q_\beta^\pm, \\
 [T_1, R_\alpha] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta R_\beta, & [T_2, Q_\alpha^+] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta R_\beta, & &
 \end{aligned} \quad (3.29)$$

together with the following anti-commutation relations:

$$\begin{aligned}
 \{\mathbf{Q}_\alpha^+, \mathbf{Q}_\beta^+\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{J} + \mathbf{H} \right) - (\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{T}_1 + \mathbf{U}_1 \right), \\
 \{\mathbf{Q}_\alpha^+, \mathbf{Q}_\beta^-\} &= -(\gamma^a C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{G}_a + \mathbf{P}_a \right), \\
 \{\mathbf{Q}_\alpha^-, \mathbf{Q}_\beta^-\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{S} + \mathbf{M} \right) + (\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{T}_2 + \mathbf{U}_2 \right), \\
 \{\mathbf{Q}_\alpha^+, \mathbf{R}_\beta\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{S} + \mathbf{M} \right) - (\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{T}_2 + \mathbf{U}_2 \right). \tag{3.30}
 \end{aligned}$$

The obtained superalgebra corresponds to the $\mathcal{N} = 2$ extended Newton-Hooke superalgebra [97, 102]. The extended Newton-Hooke algebra [70–76] appears as a subalgebra and is characterized by the central charges $\mathbf{S}$ and $\mathbf{M}$ which are crucial for avoiding degeneracy. In the presence of supersymmetry, the additional bosonic content $\{\mathbf{T}_1, \mathbf{T}_2, \mathbf{U}_1, \mathbf{U}_2\}$ ensures the non-degeneracy of the extended Bargmann superalgebra [78] obtained in the vanishing cosmological constant limit $\ell \rightarrow \infty$. In particular, both $\mathbf{T}_1$ and $\mathbf{T}_2$ appear as expansion of the R -symmetry generator T and act non-trivially on the fermionic charges. Let us note that the extended Bargmann superalgebra can also be derived by expanding the $\mathcal{N} = 2$ Poincaré superalgebra with the $S_E^{(2)}$ semigroup, following the same procedure considered to obtain the extended Newton-Hooke superalgebra (see figure 4).

3.2.2 The $\mathcal{N} = 2$ extended (AdS-)static superalgebra

The $\mathcal{N} = 2$ extended AdS-static superalgebra and its vanishing cosmological constant limit can be derived from the NR expansion of the $\mathcal{N} = 2$ AdS-Carroll superalgebra (3.13). To this end, let us consider the following subspace decomposition of the $\mathcal{N} = 2$ AdS-Carroll superalgebra:

$$V_0 = \{\mathbf{J}, \mathbf{H}, \mathbf{T}, \mathbf{U}, \mathbf{Q}_\alpha^+\}, \quad V_1 = \{\mathbf{G}_a, \mathbf{P}_a, \mathbf{Q}_\alpha^-\}, \tag{3.31}$$

which satisfies (2.4). Let us consider the same semigroup used to obtain the extended Newton-Hooke superalgebra, namely $S_E^{(2)} = \{\lambda_0, \lambda_1, \lambda_2, \lambda_3\}$. The elements of $S_E^{(2)}$ obey the multiplication laws (2.8) and λ_3 corresponds to the zero element of the semigroup. A resonant decomposition of the semigroup $S_E^{(2)}$ is given by (3.27), which satisfies the same algebraic structure as the subspaces of the $\mathcal{N} = 2$ AdS-Carroll superalgebra. Then, a supersymmetric extension of the extended AdS-static algebra⁴ is obtained after performing a resonant $S_E^{(2)}$ -expansion of the $\mathcal{N} = 2$ AdS-Carroll superalgebra and imposing the 0_S -reduction $\lambda_3 T_A = 0$. The expanded generators are related to the Carrollian ones through the semigroup elements as in table 7. Then, the expanded superalgebra is given by

$$\begin{aligned}
 [\mathbf{J}, \mathbf{G}_a] &= \epsilon_{ab} \mathbf{G}_b, & [\mathbf{J}, \mathbf{P}_a] &= \epsilon_{ab} \mathbf{P}_b, & [\mathbf{P}_a, \mathbf{P}_b] &= -\frac{1}{\ell^2} \epsilon_{ab} \mathbf{S}, \\
 [\mathbf{H}, \mathbf{P}_a] &= \frac{1}{\ell^2} \epsilon_{ab} \mathbf{G}_b, & [\mathbf{G}_a, \mathbf{P}_a] &= -\epsilon_{ab} \mathbf{M}, & [\mathbf{J}, \mathbf{Q}_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{Q}_\beta^\pm, \\
 [\mathbf{J}, \mathbf{R}_\alpha] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{R}_\beta, & [\mathbf{S}, \mathbf{Q}_\alpha^+] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{R}_\beta, & [\mathbf{P}_a, \mathbf{Q}_\alpha^+] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta \mathbf{Q}_\beta^-, \\
 [\mathbf{P}_a, \mathbf{Q}_\alpha^-] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta \mathbf{R}_\beta, & [\mathbf{T}_1, \mathbf{Q}_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{Q}_\beta^\pm, & [\mathbf{T}_1, \mathbf{R}_\alpha] &= \frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{R}_\beta, \\
 [\mathbf{T}_2, \mathbf{Q}_\alpha^+] &= \frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{R}_\beta, & & & &
 \end{aligned} \tag{3.32}$$

⁴Originally denoted as the extended para-Bargmann in [90], due to its isomorphism with the extended Bargmann algebra.

λ_3		
λ_2	S, M, T ₂ , U ₂ , R _{α}	
λ_1		G _{a} , P _{a} , Q _{α} ⁻
λ_0	J, H, T ₁ , U ₁ , Q _{α} ⁺	
	J, H, T, U, Q _{α} ⁺	G _{a} , P _{a} , Q _{α} ⁻

Table 7. Expanded generators in terms of the $\mathcal{N} = 2$ super AdS-Carroll ones and the semigroup elements.

λ_3			
λ_2			H, M, G _{a} , U ₁ , U ₂
λ_1		Q _{α} ⁺ , Q _{α} ⁻ , R _{α}	
λ_0	J, S, P _{a} , T ₁ , T ₂		
	J, S, P _{a} , T ₁ , T ₂	Q _{α} ⁺ , Q _{α} ⁻ , R _{α}	H, M, G _{a} , U ₁ , U ₂

Table 8. $\mathcal{N} = 2$ super extended AdS-static generators in terms of the extended Newton-Hooke ones and the semigroup elements.

and

$$\begin{aligned}
 \{Q_\alpha^+, Q_\beta^+\} &= -(\gamma^0 C)_{\alpha\beta} (H + U_1), & \{Q_\alpha^+, Q_\beta^-\} &= -\frac{1}{\ell} (\gamma^a C)_{\alpha\beta} G_a, \\
 \{Q_\alpha^-, Q_\beta^-\} &= -(\gamma^0 C)_{\alpha\beta} (M - U_2), & \{Q_\alpha^+, R_\beta\} &= -(\gamma^0 C)_{\alpha\beta} (M + U_2),
 \end{aligned} \tag{3.33}$$

where we have considered the multiplication laws of $S_E^{(2)}$ along the starting (anti-)commutation relations of the $\mathcal{N} = 2$ super AdS-Carroll superalgebra given by (3.13). The obtained superalgebra corresponds to a $\mathcal{N} = 2$ supersymmetric extension of the extended para-Bargmann algebra defined in [90]. As its bosonic sector, the $\mathcal{N} = 2$ extended AdS-static superalgebra (3.32) and (3.33) is isomorphic to the $\mathcal{N} = 2$ extended Bargmann superalgebra [78] by interchanging G_a and ℓP_a . In the vanishing cosmological constant limit $\ell \rightarrow \infty$, we get the $\mathcal{N} = 2$ supersymmetric extension of the extended static algebra introduced in [90].

An alternative procedure to recover the $\mathcal{N} = 2$ extended AdS-static superalgebra is starting from the $\mathcal{N} = 2$ extended Newton-Hooke and apply a resonant $\check{S}_E^{(2)}$ -expansion (see figure 4). In such case, the subspace decomposition of the extended Newton-Hooke superalgebra that one has to consider is the following:

$$V_0 = \{J, S, P_a, T_1, T_2\}, \quad V_1 = \{Q_\alpha^+, Q_\alpha^-, R_\alpha\}, \quad V_2 = \{H, M, G_a, U_1, U_2\}. \tag{3.34}$$

After performing a resonant $S_E^{(2)}$ -expansion of the extended Newton-Hooke superalgebra and extracting a $\check{S}_E^{(2)}$ reduced subalgebra, we can express the $\mathcal{N} = 2$ super extended AdS-static generators in terms of the extended Newton-Hooke ones as in table 8.

3.2.3 Generalized Newton-Hooke and Galilean superalgebras

Generalized kinematical superalgebras can be obtained by applying the NR expansion with a bigger semigroup $S_E^{(2N)} = \{\lambda_0, \lambda_1, \lambda_2, \dots, \lambda_{2N+1}\}$. Indeed, $\mathcal{N} = 2$ supersymmetric extension

of the generalized Newton-Hooke algebra [85, 90] can be derived from the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra. To this end, let us consider the subspace decomposition of the relativistic superalgebra given in (3.26), which satisfies a $\mathbb{Z}_2$ -graded Lie algebra (2.4). Let $S_E^{(2N)} = \{\lambda_0, \lambda_1, \lambda_2, \dots, \lambda_{2N+1}\}$ be the finite semigroup whose elements obey (3.14), and $\lambda_{2N+1} = 0_S$ corresponds to the zero element of the semigroup. A resonant subset decomposition of the semigroup $S_E^{(2N)} = S_0 \cup S_1$ reads

$$\begin{aligned} S_0 &= \{\lambda_{2m}, \text{ with } m = 0, \dots, N\} \cup \{\lambda_{2N+1}\}, \\ S_1 &= \{\lambda_{2m+1}, \text{ with } m = 0, \dots, N-1\} \cup \{\lambda_{2N+1}\}, \end{aligned} \quad (3.35)$$

which satisfy the same algebraic structure (2.4) as the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ subspaces. A resonant expanded superalgebra is given by

$$\mathfrak{G}_R = (S_0 \times V_0) \oplus (S_1 \times V_1), \quad (3.36)$$

where V_0 and V_1 are the relativistic subspaces given in (3.26). Then, generalized Newton-Hooke superalgebras are obtained after performing a 0_S -reduction, namely $0_S T_A = 0$. The generalized Newton-Hooke generators are related to the relativistic ones through the semigroup elements as follows

$$\begin{aligned} J^{(m)} &= \lambda_{2m} J, & P_a^{(m)} &= \lambda_{2m+1} P_a, & T^{(m)} &= \lambda_{2m} T, \\ H^{(m)} &= \lambda_{2m} H, & G_a^{(m)} &= \lambda_{2m+1} G_a, & U^{(m)} &= \lambda_{2m} U, \\ Q_\alpha^{+(m)} &= \lambda_{2m} Q_\alpha^+, & Q_\alpha^{-(m)} &= \lambda_{2m+1} Q_\alpha^-. \end{aligned} \quad (3.37)$$

Considering the (anti-)commutation relations of the starting $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra and the multiplication laws of the $S_E^{(2N)}$ semigroup, the generalized Newton-Hooke superalgebra is expressed as

$$\begin{aligned} [J^{(m)}, G_a^{(n)}] &= \epsilon_{ab} G_b^{(m+n)}, & [G_a^{(m)}, G_b^{(n)}] &= -\epsilon_{ab} J^{(m+n+1)}, \\ [J^{(m)}, P_a^{(n)}] &= \epsilon_{ab} P_b^{(m+n)}, & [G_a^{(m)}, P_a^{(n)}] &= -\epsilon_{ab} H^{(m+n+1)}, \\ [H^{(m)}, P_a^{(n)}] &= \frac{1}{\ell^2} \epsilon_{ab} G_b^{(m+n)}, & [P_a^{(m)}, P_b^{(n)}] &= -\frac{1}{\ell^2} \epsilon_{ab} J^{(m+n+1)}, \\ [H^{(m)}, G_a^{(n)}] &= \epsilon_{ab} P_b^{(m+n)}, & [J^{(m)}, Q_\alpha^\pm(n)] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta Q_\beta^\pm(m+n), \\ [H^{(m)}, Q_\alpha^\pm(n)] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta Q_\beta^\pm(m+n), & [P_a^{(m)}, Q_\alpha^\pm(n)] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta Q_\beta^\pm(m+n), \\ [P_a^{(m)}, Q_\alpha^-(n)] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta Q_\beta^{+(m+n+1)}, & [G_a^{(m)}, Q_\alpha^\pm(n)] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta Q_\beta^\pm(m+n), \\ [G_a^{(m)}, Q_\alpha^-(n)] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta Q_\beta^{+(m+n+1)}, & [T^{(m)}, Q_\alpha^\pm(n)] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta Q_\beta^\pm(m+n), \end{aligned} \quad (3.38)$$

together with the following anti-commutators:

$$\begin{aligned} \{Q_\alpha^{+(m)}, Q_\beta^{+(n)}\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} J^{(m+n)} + H^{(m+n)} \right) - (\gamma^0 C)_{\alpha\beta} \left(U^{(m+n)} + \frac{1}{\ell} T^{(m+n)} \right), \\ \{Q_\alpha^{+(m)}, Q_\beta^{-(n)}\} &= -(\gamma^a C)_{\alpha\beta} \left(\frac{1}{\ell} G_a^{(m+n)} + P_a^{(m+n)} \right), \\ \{Q_\alpha^{-(m)}, Q_\beta^{-(n)}\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} J^{(m+n+1)} + H^{(m+n+1)} \right) + (\gamma^0 C)_{\alpha\beta} \left(U^{(m+n+1)} + \frac{1}{\ell} T^{(m+n+1)} \right). \end{aligned} \quad (3.39)$$

Let us note that the (anti-)commutators are Abelian for $T_A^{(m+n)} = \lambda_{2N+1} T_A$ due to the properties of the semigroup $S_E^{(2N)}$. The $\mathcal{N} = 2$ generalized Newton-Hooke superalgebra (3.38)–(3.39) corresponds to the $\mathcal{N} = 2$ supersymmetric extension of the generalized Newton-Hooke algebra [85] which has been denoted as $\mathfrak{nh}^{(N)}$ in [90]. In the vanishing cosmological constant limit $\ell \rightarrow \infty$, we obtain the $\mathcal{N} = 2$ supersymmetric extension of the generalized Galilean algebras, denoted as $\mathfrak{gal}^{(N)}$ in [90]. The super $\mathfrak{gal}^{(N)}$, obtained after taking the flat limit, can be viewed both as a generalization of the extended Newtonian superalgebra discussed in [95, 103] and as a supersymmetric extension of the extended post-Newtonian symmetries [85, 111]. Let us note that the super $\mathfrak{gal}^{(N)}$ can also be derived as a resonant $S_E^{(2N)}$ -expansion of the $\mathcal{N} = 2$ Poincaré superalgebra (see figure 5).

The $\mathcal{N} = 2$ extended Newton-Hooke superalgebra (3.29)–(3.30) together with the extended Bargmann one are recovered for $N = 1$, namely with $S_E^{(2)}$ as the semigroup involved in the expansion. For $N = 2$, we obtain the exotic Newtonian superalgebra introduced in [103], whose generators obey the $\mathcal{N} = 2$ extended Newton-Hooke (anti-)commutation relations (3.29)–(3.30), together with the following commutation relations:

$$\begin{aligned}
 [\mathbf{G}_a, \mathbf{B}_b] &= -\epsilon_{ab} \mathbf{Z}, & [\mathbf{J}, \mathbf{B}_a] &= \epsilon_{ab} \mathbf{B}_b, & [\mathbf{J}, \mathbf{T}_a] &= \epsilon_{ab} \mathbf{T}_b, \\
 [\mathbf{G}_a, \mathbf{T}_b] &= -\epsilon_{ab} \mathbf{Y}, & [\mathbf{H}, \mathbf{B}_a] &= \epsilon_{ab} \mathbf{T}_b, & [\mathbf{H}, \mathbf{T}_a] &= \frac{1}{\ell^2} \epsilon_{ab} \mathbf{B}_b, \\
 [\mathbf{P}_a, \mathbf{B}_a] &= -\epsilon_{ab} \mathbf{Y}, & [\mathbf{S}, \mathbf{G}_a] &= \epsilon_{ab} \mathbf{B}_b, & [\mathbf{S}, \mathbf{P}_a] &= \epsilon_{ab} \mathbf{T}_b, \\
 [\mathbf{P}_a, \mathbf{T}_a] &= -\frac{1}{\ell^2} \epsilon_{ab} \mathbf{Z}, & [\mathbf{M}, \mathbf{G}_a] &= \epsilon_{ab} \mathbf{T}_b, & [\mathbf{M}, \mathbf{P}_a] &= \frac{1}{\ell^2} \epsilon_{ab} \mathbf{B}_b, \\
 [\mathbf{J}, \mathbf{W}_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^\pm, & [\mathbf{S}, \mathbf{Q}_\alpha^-] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^-, & [\mathbf{S}, \mathbf{R}_\alpha] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^+, \\
 [\mathbf{H}, \mathbf{W}_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^\pm, & [\mathbf{M}, \mathbf{Q}_\alpha^-] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^-, & [\mathbf{M}, \mathbf{R}_\alpha] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^+, \\
 [\mathbf{Z}, \mathbf{Q}_\alpha^+] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^+, & [\mathbf{Y}, \mathbf{Q}_\alpha^+] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^+, & [\mathbf{G}_a, \mathbf{R}_\alpha] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta \mathbf{W}_\beta^-, \\
 [\mathbf{G}_a, \mathbf{W}_\alpha^-] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta \mathbf{W}_\beta^+, & [\mathbf{P}_a, \mathbf{R}_\alpha] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta \mathbf{W}_\beta^-, & [\mathbf{P}_a, \mathbf{W}_\alpha^-] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta \mathbf{W}_\beta^+, \\
 [\mathbf{B}_a, \mathbf{Q}_\alpha^\pm] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta \mathbf{W}_\beta^\mp, & [\mathbf{T}_a, \mathbf{Q}_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta \mathbf{W}_\beta^\mp, & [\mathbf{T}_1, \mathbf{W}_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^\pm, \\
 [\mathbf{T}_2, \mathbf{Q}_\alpha^-] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^-, & [\mathbf{T}_2, \mathbf{R}_\alpha] &= \frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^+, & [\mathbf{T}_3, \mathbf{Q}_\alpha^+] &= \frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{W}_\beta^+, \tag{3.40}
 \end{aligned}$$

and anti-commutators:

$$\begin{aligned}
 \{\mathbf{Q}_\alpha^+, \mathbf{W}_\beta^+\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{Z} + \mathbf{Y} \right) - (\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{T}_3 + \mathbf{U}_3 \right), \\
 \{\mathbf{Q}_\alpha^+, \mathbf{W}_\beta^-\} &= -(\gamma^a C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{B}_a + \mathbf{T}_a \right), \\
 \{\mathbf{Q}_\alpha^-, \mathbf{R}_\beta\} &= -(\gamma^a C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{B}_a + \mathbf{T}_a \right), \\
 \{\mathbf{Q}_\alpha^-, \mathbf{W}_\beta^-\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{Z} + \mathbf{Y} \right) + (\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{T}_3 + \mathbf{U}_3 \right), \\
 \{\mathbf{R}_\alpha, \mathbf{R}_\beta\} &= -(\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{Z} + \mathbf{Y} \right) - (\gamma^0 C)_{\alpha\beta} \left(\frac{1}{\ell} \mathbf{T}_3 + \mathbf{U}_3 \right). \tag{3.41}
 \end{aligned}$$

Here the generators have been identified to the generalized Newton-Hooke ones (3.37) as follows:

$$\begin{aligned}
 \mathbf{J} &= \mathbf{J}^{(0)}, & \mathbf{G}_a &= \mathbf{G}_a^{(0)}, & \mathbf{T}_1 &= \mathbf{T}^{(0)}, & \mathbf{U}_1 &= \mathbf{U}^{(0)}, \\
 \mathbf{S} &= \mathbf{J}^{(1)}, & \mathbf{B}_a &= \mathbf{G}_a^{(1)}, & \mathbf{T}_2 &= \mathbf{T}^{(1)}, & \mathbf{U}_2 &= \mathbf{U}^{(1)}, \\
 \mathbf{Z} &= \mathbf{J}^{(2)}, & \mathbf{P}_a &= \mathbf{P}_a^{(0)}, & \mathbf{T}_3 &= \mathbf{T}^{(2)}, & \mathbf{U}_3 &= \mathbf{U}^{(2)}, \\
 \mathbf{H} &= \mathbf{H}^{(0)}, & \mathbf{T}_a &= \mathbf{P}_a^{(1)}, & \mathbf{Q}_a^+ &= \mathbf{Q}_a^{+(0)}, & \mathbf{R}_a &= \mathbf{Q}_a^{+(1)}, \\
 \mathbf{M} &= \mathbf{H}^{(1)}, & \mathbf{W}_a^+ &= \mathbf{Q}_a^{+(2)}, & \mathbf{Q}_a^- &= \mathbf{Q}_a^{-(0)}, & \mathbf{W}_a^- &= \mathbf{Q}_a^{-(1)}, \\
 \mathbf{Y} &= \mathbf{H}^{(2)}.
 \end{aligned} \tag{3.42}$$

In the vanishing cosmological constant limit $\ell \rightarrow \infty$, we find the extended Newtonian superalgebra introduced in [95]. At the bosonic level, the central charges $\mathbf{Y}$ and $\mathbf{Z}$ are essential to ensure the non-degeneracy of the invariant bilinear trace of the three-dimensional extended Newtonian gravity. In the absence of such central charges, the algebra corresponds to the so-called Newtonian algebra, which has been first encountered as the underlying symmetry of an action principle for Newtonian gravity [113] in four spacetime dimensions. On the other hand, the Newton-Hooke version of the Newtonian algebra has been useful to construct a four-dimensional non-relativistic gravity action based on the MacDowell-Mansouri formalism [89].

3.2.4 Generalized (AdS-)static superalgebras

The $S_E^{(2N)}$ semigroup can also be applied to the Carrollian sector to reproduce generalizations of the (AdS-)static superalgebra. In particular, $\mathcal{N} = 2$ generalized AdS-static superalgebras are obtained by expanding the $\mathcal{N} = 2$ AdS-Carroll superalgebra (3.13) with $S_E^{(2N)}$. Before applying the expansion, we first consider the subspace decomposition of the $\mathcal{N} = 2$ AdS-Carroll superalgebra given in (3.31), which satisfies a $\mathbb{Z}_2$ -graded Lie algebra (2.4). A subset decomposition of the semigroup $S_E^{(2N)} = S_0 \cup S_1$ resonant with (2.4) is given by (3.35). Then, a $\mathcal{N} = 2$ generalized AdS-static superalgebra is derived after performing a resonant $S_E^{(2N)}$ -expansion of the $\mathcal{N} = 2$ AdS-Carroll superalgebra (3.13) and extracting a 0_S -reduced subalgebra. The expanded generators can be expressed in terms of the AdS-Carroll ones through the semigroup elements as

$$\begin{aligned}
 \mathbf{J}^{(m)} &= \lambda_{2m} \mathbf{J}, & \mathbf{P}_a^{(m)} &= \lambda_{2m+1} \mathbf{P}_a, & \mathbf{T}^{(m)} &= \lambda_{2m} \mathbf{T}, & \mathbf{Q}_\alpha^{+(m)} &= \lambda_{2m} \mathbf{Q}_\alpha^+, \\
 \mathbf{H}^{(m)} &= \lambda_{2m} \mathbf{H}, & \mathbf{G}_a^{(m)} &= \lambda_{2m+1} \mathbf{G}_a, & \mathbf{U}^{(m)} &= \lambda_{2m} \mathbf{U}, & \mathbf{Q}_\alpha^{-(m)} &= \lambda_{2m+1} \mathbf{Q}_\alpha^-.
 \end{aligned} \tag{3.43}$$

The explicit (anti-)commutation relations of the $\mathcal{N} = 2$ generalized AdS-static superalgebra are obtained by considering the multiplication laws of $S_E^{(2N)}$ and the original (anti-)commutators of the $\mathcal{N} = 2$ AdS-Carroll superalgebra. Therefore we get:

$$\begin{aligned}
 [\mathbf{J}^{(m)}, \mathbf{G}_a^{(n)}] &= \epsilon_{ab} \mathbf{G}_b^{(m+n)}, & [\mathbf{J}^{(m)}, \mathbf{P}_a^{(n)}] &= \epsilon_{ab} \mathbf{P}_b^{(m+n)}, \\
 [\mathbf{P}_a^{(m)}, \mathbf{P}_b^{(n)}] &= -\frac{1}{\ell^2} \epsilon_{ab} \mathbf{J}^{(m+n+1)}, & [\mathbf{H}^{(m)}, \mathbf{P}_a^{(n)}] &= \frac{1}{\ell^2} \epsilon_{ab} \mathbf{G}_b^{(m+n)}, \\
 [\mathbf{G}_a^{(m)}, \mathbf{P}_a^{(n)}] &= -\epsilon_{ab} \mathbf{H}^{(m+n+1)}, & [\mathbf{J}^{(m)}, \mathbf{Q}_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathbf{Q}_\beta^\pm,
 \end{aligned}$$

$$\begin{aligned}
 [\mathbf{P}_a^{(m)}, \mathbf{Q}_\alpha^{+(n)}] &= -\frac{1}{2\ell} (\gamma_a)_\alpha{}^\beta \mathbf{Q}_\beta^{-(m+n)}, & [\mathbf{P}_a^{(m)}, \mathbf{Q}_\alpha^{-(n)}] &= -\frac{1}{2\ell} (\gamma_a)_\alpha{}^\beta \mathbf{Q}_\beta^{+(m+n+1)}, \\
 [\mathbf{T}^{(m)}, \mathbf{Q}_\alpha^{\pm(n)}] &= \pm \frac{1}{2} (\gamma_0)_\alpha{}^\beta \mathbf{Q}_\beta^{\pm(m+n)}, \\
 \{\mathbf{Q}_\alpha^{+(m)}, \mathbf{Q}_\beta^{+(n)}\} &= -(\gamma^0 C)_{\alpha\beta} (\mathbf{H}^{(m+n)} + \mathbf{U}^{(m+n)}), \\
 \{\mathbf{Q}_\alpha^{+(m)}, \mathbf{Q}_\beta^{-(n)}\} &= -\frac{1}{\ell} (\gamma^a C)_{\alpha\beta} \mathbf{G}_a^{(m+n)}, \\
 \{\mathbf{Q}_\alpha^{-(m)}, \mathbf{Q}_\beta^{-(n)}\} &= -(\gamma^0 C)_{\alpha\beta} (\mathbf{H}^{(m+n+1)} - \mathbf{U}^{(m+n+1)}).
 \end{aligned} \tag{3.44}$$

As in the generalized $\mathfrak{nh}^{(N)}$ superalgebra, the (anti-)commutators vanish for $T_A^{(m+n)} = \lambda_{2N+1} T_A$, due to the 0_S -reduction condition imposed during the expansion. The obtained superalgebra (3.44), which we have denoted as the $\mathcal{N} = 2$ **ad $\mathfrak{s}$ -stat $^{(N)}$** superalgebra, reproduces an $\mathcal{N} = 2$ generalized static superalgebra in the vanishing cosmological constant limit $\ell \rightarrow \infty$. The $\mathcal{N} = 2$ **stat $^{(N)}$** superalgebra can also be recovered after imposing a resonant $S_E^{(2N)}$ -expansion on the $\mathcal{N} = 2$ Carroll superalgebra and applying a 0_S -reduction. Alternatively, the $\mathcal{N} = 2$ **ad $\mathfrak{s}$ -stat $^{(N)}$** superalgebra together with its flat limit can be derived from the $\mathcal{N} = 2$ super $\mathfrak{nh}^{(N)}$ one (3.38)–(3.39) after performing a resonant $S_E^{(2)}$ -expansion and extracting a $\check{S}_E^{(2N)}$ reduced subalgebra (see figure 5). As in the generalized $\mathfrak{nh}^{(N)}$ superalgebra, the (anti-)commutators vanishes for $T_A^{(m+n)} = \lambda_{2N+1} T_A$ due to the 0_S -reduction condition imposed during the expansion. The obtained superalgebra (3.44), which we have denoted as the $\mathcal{N} = 2$ **ad $\mathfrak{s}$ -stat $^{(N)}$** superalgebra, reproduces a $\mathcal{N} = 2$ generalized static superalgebra in the vanishing cosmological constant limit $\ell \rightarrow \infty$. The $\mathcal{N} = 2$ **stat $^{(N)}$** superalgebra can also be recovered after imposing a resonant $S_E^{(2N)}$ -expansion on the $\mathcal{N} = 2$ Carroll superalgebra and applying a 0_S -reduction. Alternatively, the $\mathcal{N} = 2$ **ad $\mathfrak{s}$ -stat $^{(N)}$** superalgebra together with its flat limit can be derived from the $\mathcal{N} = 2$ super $\mathfrak{nh}^{(N)}$ one (3.38)–(3.39) after performing a resonant $S_E^{(2)}$ -expansion and extracting a $\check{S}_E^{(2N)}$ reduced subalgebra (see figure 5).

4 Three-dimensional non-Lorentzian supergravity theories based on extended kinematical superalgebras

In this section, we present three-dimensional NL supergravity actions invariant under NL superalgebras which have been obtained after a suitable sequence of expansions starting from the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra and using the $S_E^{(2)}$ and $\check{S}_E^{(2)}$ semigroups. Our construction is based on the CS formalism, whose action reads

$$I_{\text{CS}} = \frac{k}{4\pi} \int_{\mathcal{M}} \langle AdA + \frac{2}{3} A^3 \rangle, \tag{4.1}$$

where A is the gauge connection one-form, $\langle \cdots \rangle$ denotes the invariant tensor and k corresponds to the CS level of the theory, related to the gravitational constant G through $k = 1/(4G)$. The degeneracy of the invariant bilinear trace prevents to obtain a kinematical term for each gauge field and some fields could not be determined by the field equations. Remarkably, the $\mathcal{N} = 2$ Carrollian superalgebras and the extended kinematical superalgebras discussed in the NR regime admit a non-degenerate invariant metric which, in three spacetime dimensions, implies the vanishing of the curvatures as field equations in the CS formulation.




Figure 5. This cube summarizes the generalized expansion processes starting from the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2,2) \otimes \mathfrak{sp}(2)$ superalgebra.

It is important to emphasize that the non-degeneracy of the original superalgebra does not necessarily imply that the expanded superalgebra also admits a non-degenerate bilinear trace. Nonetheless, the non-degeneracy condition of the relativistic $\mathcal{N} = 2$ superalgebra, given by $\tilde{\alpha}_1 \neq 0$ in (3.2), requires the existence of the relativistic generators $\tilde{P}_A$, $\tilde{J}_A$, $\tilde{T}$, and $\tilde{U}$. Then, considering the subspace decomposition employed in both the UR and the NR expansions (see table 5 and 6), together with the multiplication law of the S_E semigroup (2.8), the expanded superalgebras inherit the non-degeneracy of the relativistic invariant tensor by including at least λ_2 in the chosen semigroup S_E . Thus, the smallest semigroup that satisfies the non-degeneracy criterion for both the NR and the UR cases is $S_E^{(2)} = \{\lambda_0, \lambda_1, \lambda_2, \lambda_3\}$, where λ_3 is the zero element of the semigroup.

Here, we will present the CS supergravity action based on each $\mathcal{N} = 2$ kinematical superalgebra of the cube 4 and their generalizations discussed in the previous section. For this purpose, we will divide our results into three parts. First, we will review the $\mathcal{N} = 2$ AdS-Carroll supergravity and explore novel generalizations. Then, we will study the non-relativistic regime of the $\mathfrak{osp}(2,2) \otimes \mathfrak{sp}(2)$ supergravity along with its vanishing cosmological constant limit. Finally, we will analyze NL supergravity actions based on the $\mathcal{N} = 2$ (AdS-)static superalgebra and its generalization.

4.1 $\mathcal{N} = 2$ AdS-Carroll Chern-Simons supergravity and beyond

The $\mathcal{N} = 2$ AdS-Carroll supergravity defined à la CS in three spacetime dimensions has been first discussed in [99]. Here, we briefly review its construction and present its generalization: the $\mathcal{N} = 2$ $\mathfrak{adS-car}^{(N)}$ supergravity.

Let A be the gauge connection one-form for the $\mathcal{N} = 2$ AdS-Carroll superalgebra (3.13),

$$A = \omega J + \omega^a G_a + \tau H + e^a P_a + t T + u U + \bar{\psi}^+ Q^+ + \bar{\psi}^- Q^-, \quad (4.2)$$

where ω and ω_a are the time and spatial components of the spin-connection. The time and spatial components of the dreibein are given by τ and e^a , respectively. On the other hand, t and u are the gauge field dual to the generators T and U , respectively. The curvature two-form $F = dA + \frac{1}{2} [A, A]$ reads

$$F = F(\omega) J + F^a(\omega^b) G_a + F(\tau) H + F^a(e^b) P_a + F(t) T + F(u) U + \nabla \bar{\psi}^+ Q^+ + \nabla \bar{\psi}^- Q^-, \quad (4.3)$$

whose components are given by

$$\begin{aligned} F(\omega) &= d\omega + \frac{1}{2\ell^2} \epsilon^{ac} e_a e_c = R(\omega) + \frac{1}{2\ell^2} \epsilon^{ac} e_a e_c, \\ F^a(\omega^b) &= d\omega^a + \epsilon^{ac} \omega \omega_c + \frac{1}{\ell^2} \epsilon^{ac} \tau e_c + \frac{1}{\ell} \bar{\psi}^+ \gamma^a \psi^- = R^a(\omega^b) + \frac{1}{\ell^2} \epsilon^{ac} \tau e_c + \frac{1}{\ell} \bar{\psi}^+ \gamma^a \psi^-, \\ F(\tau) &= d\tau + \epsilon^{ac} \omega_a e_c + \frac{1}{2} \bar{\psi}^+ \gamma^0 \psi^+ + \frac{1}{2} \bar{\psi}^- \gamma^0 \psi^- = R(\tau) + \frac{1}{2} \bar{\psi}^+ \gamma^0 \psi^+ + \frac{1}{2} \bar{\psi}^- \gamma^0 \psi^-, \\ F^a(e^b) &= de^a + \epsilon^{ac} \omega e_c = R^a(e^b), \\ F(t) &= dt, \\ F(u) &= du + \frac{1}{2} \bar{\psi}^+ \gamma^0 \psi^+ - \frac{1}{2} \bar{\psi}^- \gamma^0 \psi^-, \\ \nabla \psi^+ &= d\psi^+ + \frac{1}{2} \omega \gamma_0 \psi^+ + \frac{1}{2\ell} e^a \gamma_a \psi^- - \frac{1}{2} t \gamma_0 \psi^+, \\ \nabla \psi^- &= d\psi^- + \frac{1}{2} \omega \gamma_0 \psi^- + \frac{1}{2\ell} e^a \gamma_a \psi^+ + \frac{1}{2} t \gamma_0 \psi^-. \end{aligned} \quad (4.4)$$

Let us note that the curvature two-forms (4.4) reduce to the $\mathcal{N} = 2$ Carroll ones in the vanishing cosmological constant limit $\ell \rightarrow \infty$. The $\mathcal{N} = 2$ AdS-Carroll superalgebra admits a non-degenerate invariant tensor whose non-vanishing components are given by [99]

$$\begin{aligned} \langle JJ \rangle &= -\alpha_0, & \langle P_a P_b \rangle &= \frac{\alpha_0}{\ell^2} \delta_{ab}, \\ \langle JH \rangle &= -\alpha_1, & \langle G_a P_b \rangle &= \alpha_1 \delta_{ab}, \\ \langle TT \rangle &= \alpha_0, & \langle TU \rangle &= \alpha_1, \\ \langle Q_\alpha^+ Q_\beta^+ \rangle &= 2\alpha_1 C_{\alpha\beta}, & \langle Q_\alpha^- Q_\beta^- \rangle &= 2\alpha_1 C_{\alpha\beta}, \end{aligned} \quad (4.5)$$

where α_0 and α_1 are arbitrary constants which can be related to the relativistic ones (3.2) through the $S_E^{(2)}$ elements as

$$\alpha_0 = \lambda_0 \tilde{\alpha}_0, \quad \alpha_1 = \lambda_2 \tilde{\alpha}_1. \quad (4.6)$$

Here, the non-degeneracy of the invariant tensor requires $\alpha_1 \neq 0$.

The CS supergravity action for the $\mathcal{N} = 2$ AdS-Carroll superalgebra (3.13) is obtained considering the non-vanishing components of the invariant tensor (4.5) along the gauge connection one-form (4.2) into the general CS expression (4.1). Thus, we get [99]

$$I_{\text{adS-car}}^{\mathcal{N}=2} = \frac{k}{4\pi} \int \alpha_0 \left[\frac{1}{\ell^2} e_a R^a(e^b) - \omega R(\omega) + t dt \right] + \alpha_1 \left[2e_a R^a(\omega^b) - 2\tau R(\omega) + \frac{1}{\ell^2} \epsilon^{ac} \tau e_a e_c + 2t du - 2\bar{\psi}^+ \nabla \psi^+ - 2\bar{\psi}^- \nabla \psi^- \right]. \quad (4.7)$$

The $\mathcal{N} = 2$ AdS-Carroll CS supergravity action contains two independent terms. The term proportional to α_0 corresponds to the UR version of the exotic Lagrangian [62]. On the other hand, the sector along α_1 corresponds to the Carrollian regime of the $\text{SO}(2)$ extension of the $\mathcal{N} = 2$ AdS supergravity Lagrangian [109]. Let us note that the $\mathcal{N} = 2$ Carroll CS supergravity theory is derived in the vanishing cosmological constant limit $\ell \rightarrow \infty$. Since the supersymmetric terms appear only in the term involving α_1 , we can, without loss of generality, set $\alpha_0 = 0$ as this does not affect the non-degeneracy. In particular, the non-degeneracy ensures that the field equations are given by the vanishing of the curvature two-forms (4.4), which transform covariantly with respect to the supersymmetry transformation rules:

$$\begin{aligned} \delta\omega &= 0, \\ \delta\omega^a &= \frac{1}{\ell} \bar{\varepsilon}^+ \gamma^a \psi^- + \frac{1}{\ell} \bar{\varepsilon}^- \gamma^a \psi^+, \\ \delta\tau &= \bar{\varepsilon}^+ \gamma^0 \psi^+ + \bar{\varepsilon}^- \gamma^0 \psi^-, \\ \delta e^a &= 0, \\ \delta t &= 0, \\ \delta u &= \bar{\varepsilon}^+ \gamma^0 \psi^+ - \bar{\varepsilon}^- \gamma^0 \psi^-, \\ \delta\psi^+ &= d\varepsilon^+ + \frac{1}{2} \omega \gamma_0 \varepsilon^+ + \frac{1}{2\ell} e^a \gamma_a \varepsilon^- - \frac{1}{2} t \gamma_0 \varepsilon^+, \\ \delta\psi^- &= d\varepsilon^- + \frac{1}{2} \omega \gamma_0 \varepsilon^- + \frac{1}{2\ell} e^a \gamma_a \varepsilon^+ + \frac{1}{2} t \gamma_0 \varepsilon^-. \end{aligned} \quad (4.8)$$

Here, $\varepsilon^\pm$ are the gauge parameters related to the fermionic charges $Q^\pm$. The supersymmetry transformation rules for the $\mathcal{N} = 2$ Carroll CS supergravity are recovered in the flat limit $\ell \rightarrow \infty$.

Generalized $\mathcal{N} = 2$ AdS-Carroll supergravity theory and its flat limit

A generalized UR CS supergravity theory can be constructed from the $\mathcal{N} = 2$ generalized AdS-Carroll superalgebra (3.21)–(3.22). To this end, let us consider the gauge connection one-form for the $\mathcal{N} = 2$ $\text{adS-car}^{(N)}$,

$$A = \sum_{m=0}^{N-1} \left(\omega^{(m)} \mathbf{J}^{(m)} + \omega^a{}^{(m)} \mathbf{G}_a^{(m)} + \tau^{(m)} \mathbf{H}^{(m)} + e^a{}^{(m)} \mathbf{P}_a^{(m)} + t^{(m)} \mathbf{T}^{(m)} + u^{(m)} \mathbf{U}^{(m)} + \bar{\psi}^{+(m)} \mathbf{Q}^{+(m)} + \bar{\psi}^{-(m)} \mathbf{Q}^{-(m)} \right). \quad (4.9)$$

The corresponding curvature two-form reads

$$F = \sum_{m=0}^{N-1} \left[F \left(\omega^{(m)} \right) J^{(m)} + F^a \left(\omega^b{}^{(m)} \right) G_a^{(m)} + F \left(\tau^{(m)} \right) H^{(m)} + F^a \left(e^b{}^{(m)} \right) P_a^{(m)} \right. \\ \left. + F \left(t^{(m)} \right) T^{(m)} + F \left(u^{(m)} \right) U^{(m)} + \nabla \bar{\psi}^{+(m)} Q^{+(m)} + \nabla \bar{\psi}^{-(m)} Q^{-(m)} \right], \quad (4.10)$$

where the curvature components are directly derived from the $\mathcal{N} = 2$ **adS-car**^(N) superalgebra and are given by

$$F \left(\omega^{(m)} \right) = d\omega^{(m)} + \frac{1}{2\ell^2} \sum_{n,p=0}^{N-1} \delta_{n+p}^m \epsilon^{ac} e_a^{(n)} e_c^{(p)} + \frac{1}{2\ell} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \bar{\psi}^{\pm(n)} \gamma^0 \psi^{\pm(p)}, \\ + \frac{1}{2} \sum_{n,p=0}^{N-1} \delta_{n+p+2}^m \epsilon^{ac} \omega_a^{(n)} \omega_c^{(p)}, \\ F^a \left(\omega^b{}^{(m)} \right) = d\omega^a{}^{(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p}^m \left(\epsilon^{ac} \omega_a^{(n)} \omega_c^{(p)} + \frac{1}{\ell^2} \epsilon^{ac} \tau^{(n)} e_c^{(p)} + \frac{1}{\ell} \bar{\psi}^{+(n)} \gamma^a \psi^{-(p)} \right), \\ F \left(\tau^{(m)} \right) = d\tau^{(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p}^m \left(\epsilon^{ac} \omega_a^{(n)} e_c^{(p)} + \frac{1}{2} \bar{\psi}^{+(n)} \gamma^0 \psi^{+(p)} + \frac{1}{2} \bar{\psi}^{-(n)} \gamma^0 \psi^{-(p)} \right), \\ F^a \left(e^b{}^{(m)} \right) = de^a{}^{(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p}^m \epsilon^{ac} \omega_a^{(n)} e_c^{(p)} + \sum_{n,p=0}^{N-1} \delta_{n+p+2}^m \epsilon^{ac} \tau^{(n)} \omega_c^{(p)}, \\ F \left(t^{(m)} \right) = dt^{(m)} + \frac{1}{2\ell} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \bar{\psi}^{\pm(n)} \gamma^0 \psi^{\pm(p)}, \\ F \left(u^{(m)} \right) = du^{(m)} + \frac{1}{2} \sum_{n,p=0}^{N-1} \delta_{n+p}^m \bar{\psi}^{\pm(n)} \gamma^0 \psi^{\pm(p)}, \\ \nabla \psi^{\pm(m)} = d\psi^{\pm(m)} + \frac{1}{2} \sum_{n,p=0}^{N-1} \delta_{n+p}^m \left(\omega^{(n)} \gamma_0 \psi^{\pm(p)} - \frac{1}{\ell} e^a{}^{(n)} \gamma_a \psi^{\mp(p)} \mp t^{(n)} \gamma_0 \psi^{\pm(p)} \right) \\ + \frac{1}{2} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \left(\omega^a{}^{(n)} \gamma_a \psi^{\mp(p)} + \frac{1}{\ell} \tau^{(n)} \gamma_0 \psi^{\pm(p)} \right). \quad (4.11)$$

In the vanishing cosmological constant limit $\ell \rightarrow \infty$, we obtain the curvature two-forms of the $\mathcal{N} = 2$ generalized Carroll superalgebra. The $\mathcal{N} = 2$ **adS-car**^(N) superalgebra admits the following components of an invariant tensor:

$$\langle J^{(m)} H^{(n)} \rangle = -\alpha_{m+n+1}, \quad \langle G_a^{(m)} P_b^{(n)} \rangle = \alpha_{m+n+1} \delta_{ab}, \\ \langle T^{(m)} U^{(n)} \rangle = \alpha_{m+n+1}, \quad \langle U^{(m)} U^{(n)} \rangle = -\frac{1}{\ell} \alpha_{m+n+2}, \\ \langle Q_\alpha^{+(m)} Q_\beta^{+(n)} \rangle = 2\alpha_{m+n+1} C_{\alpha\beta}, \quad \langle Q_\alpha^{-(m)} Q_\beta^{-(n)} \rangle = 2\alpha_{m+n+1} C_{\alpha\beta}, \quad (4.12)$$

where $i \leq N$ for α_i . Here, we have intentionally omitted the components related to the exotic sector of the theory. The non-degeneracy of the invariant tensor is guaranteed for $\alpha_N \neq 0$. Let us note that, for $N = 1$, we recover the invariant tensor for the $\mathcal{N} = 2$ AdS-Carroll superalgebra.

Then, by considering the invariant tensor (4.12) and the connection one-form (4.9) in the CS expression (4.1), we get an UR CS supergravity action for the $\mathcal{N} = 2$ generalized AdS-Carroll superalgebra (3.21)–(3.22),

$$I_{\text{adS-car}^{(N)}}^{\mathcal{N}=2} = \frac{k}{4\pi} \int \sum_{p=1}^N \alpha_p \delta_{m+n+1}^p \left[2e_a^{(m)} R^a \left(\omega^{b(n)} \right) - 2\tau^{(m)} R \left(\omega^{(n)} \right) + \frac{\delta_{q+r}^n}{\ell^2} \epsilon^{ac} \tau^{(m)} e_a^{(q)} e_c^{(r)} \right. \\ \left. + 2t^{(m)} du^{(n)} - \frac{\delta_{q+1}^n}{\ell} u^{(m)} du^{(q)} - 2\bar{\psi}^{+(m)} \nabla \psi^{+(n)} - 2\bar{\psi}^{-(m)} \nabla \psi^{-(n)} \right], \quad (4.13)$$

where we have defined

$$R^a \left(\omega^{b(m)} \right) = d\omega^a{}^{(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p}^m \epsilon^{ac} \omega_a^{(n)} \omega_c^{(p)}, \\ R \left(\omega^{(m)} \right) = d\omega^{(m)} + \frac{1}{2} \sum_{n,p=0}^{N-1} \delta_{n+p+2}^m \epsilon^{ac} \omega_a^{(n)} \omega_c^{(p)}. \quad (4.14)$$

Here, we can set $\alpha_i = 0$ for $i \neq p$ without affecting the non-degeneracy condition $\alpha_N \neq 0$. The non-degeneracy of the invariant tensor (4.12) ensures that the field equations are given by the vanishing of the curvature two-forms (4.11). One can notice that the $\mathcal{N} = 2$ **adS-car**^(N) CS supergravity action is invariant under the following supersymmetry transformation rules:

$$\delta\omega^{(m)} = \frac{1}{\ell} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \bar{\varepsilon}^{\pm(n)} \gamma^0 \psi^{\pm(p)}, \\ \delta\omega^a{}^{(m)} = \frac{1}{\ell} \sum_{n,p=0}^{N-1} \delta_{n+p}^m \bar{\varepsilon}^{\pm(n)} \gamma^a \psi^{\mp(p)}, \\ \delta\tau^{(m)} = \sum_{n,p=0}^{N-1} \delta_{n+p}^m \bar{\varepsilon}^{\pm(n)} \gamma^0 \psi^{\pm(p)}, \\ \delta e^a{}^{(m)} = \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \bar{\varepsilon}^{\pm(n)} \gamma^a \psi^{\mp(p)}, \\ \delta t^{(m)} = \pm \frac{1}{\ell} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \bar{\varepsilon}^{\pm(n)} \gamma^0 \psi^{\pm(p)}, \\ \delta u^{(m)} = \pm \frac{1}{\ell} \sum_{n,p=0}^{N-1} \delta_{n+p}^m \bar{\varepsilon}^{\pm(n)} \gamma^0 \psi^{\pm(p)}, \\ \delta\psi^{\pm(m)} = d\varepsilon^{\pm(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p}^m \left(\frac{1}{2} \omega^{(n)} \gamma_0 \varepsilon^{\pm(p)} + \frac{1}{2\ell} e^a{}^{(n)} \gamma_a \varepsilon^{\mp(p)} \mp \frac{1}{2} t^{(n)} \gamma_0 \varepsilon^{\pm(p)} \right) \\ + \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \left(\frac{1}{2} \omega^a{}^{(n)} \gamma_a \varepsilon^{\mp(p)} + \frac{1}{2\ell} \tau^{(n)} \gamma_0 \varepsilon^{\pm(p)} \right), \quad (4.15)$$

with $\varepsilon^{\pm(m)}$ being the gauge parameters related to the fermionic generators $Q^{\pm(m)}$. Let us note that the CS supergravity action (4.13) can be written as the sum of individual Lagrangians based on $\mathcal{N} = 2$ **adS-car**^(p) for $p = 1, 2, \dots, N$,

$$I_{\text{adS-car}^{(N)}}^{\mathcal{N}=2} = \frac{k}{4\pi} \int \sum_{p=1}^N \alpha_p \mathcal{L}_{\text{adS-car}^{(p)}}^{\mathcal{N}=2}. \quad (4.16)$$

In particular, $p = 1$ reproduces the $\mathcal{N} = 2$ AdS-Carroll supergravity Lagrangian proportional to α_1 in (4.7). For $p = 2$, the Lagrangian is invariant under the $\mathcal{N} = 2$ **ads-car**⁽²⁾ superalgebra (3.23)–(3.24),

$$\begin{aligned} \mathcal{L}_{\text{ads-car}^{(2)}}^{\mathcal{N}=2} = & \left[2e_a R^a(k^b) + 2m_a R^a(\omega^b) - 2\tau R(k) - 2hR(\omega) + \frac{2}{\ell^2} \epsilon^{ac} \tau e_a m_c + \frac{1}{\ell^2} \epsilon^{ac} h e_a e_c \right. \\ & \left. + 2t_1 du_2 + 2t_2 du_1 - \frac{1}{\ell} u_1 du_1 - 2\bar{\psi}^\pm \nabla \chi^\pm - 2\bar{\chi}^\pm \nabla \psi^\pm \right], \end{aligned} \quad (4.17)$$

with

$$\begin{aligned} R^a(k^b) &= dk^a + \epsilon^{ac} \omega k_c + \epsilon^{ac} k \omega_c, \\ R(k) &= dk, \\ \nabla \chi^\pm &= d\chi^\pm + \frac{1}{2} \omega \gamma_0 \chi^\pm + \frac{1}{2} \omega^a \gamma_a \psi^\mp + \frac{1}{2} k \gamma_0 \psi^\pm + \frac{1}{2\ell} e^a \gamma_a \chi^{mp} + \frac{1}{2\ell} \tau \gamma_0 \psi^\pm \\ &\quad + \frac{1}{2\ell} m^a \gamma_a \psi^\pm \mp \frac{1}{2} t_1 \gamma_0 \chi^\pm \mp \frac{1}{2} t_2 \gamma_0 \psi^\pm. \end{aligned} \quad (4.18)$$

On the other hand, $R^a(\omega^a)$, $R(\omega)$ and $\nabla \psi^\pm$ are given in (4.4). Here we have defined

$$\begin{aligned} \omega^{(0)} &= \omega, & \omega^{(1)} &= k, & \tau^{(0)} &= \tau, & \tau^{(1)} &= h, \\ \omega_a^{(0)} &= \omega_a, & \omega_a^{(1)} &= k_a, & e_a^{(0)} &= e_a, & e_a^{(1)} &= m_a, \\ t^{(0)} &= t_1, & t^{(1)} &= t_2, & u^{(0)} &= u_1, & u^{(1)} &= u_2, \\ \psi_\alpha^{\pm(0)} &= \psi_\alpha^\pm, & \psi_\alpha^{\pm(1)} &= \chi_\alpha^\pm. \end{aligned} \quad (4.19)$$

In the vanishing cosmological constant limit $\ell \rightarrow \infty$, the theory reduces to the $\mathcal{N} = 2$ **car**⁽²⁾ supergravity, which extends the $\mathcal{N} = 2$ Carroll supergravity appearing along the α_1 term. Post-Carrollian extensions of the $\mathcal{N} = 2$ AdS-Carroll supergravity (4.7) appear at higher order in p .

4.2 $\mathcal{N} = 2$ extended Newton-Hooke Chern-Simons supergravity and its generalization

The three-dimensional $\mathcal{N} = 2$ extended Newton-Hooke supergravity has been first introduced in [97] and subsequently recovered in [102, 103]. Here, we start with a brief review of the extended Newton-Hooke supergravity theory discussed in [102, 103]. Then, we show that the NR expansion, as in the Carrollian case, allows us to derive novel post-Newtonian generalizations of the Newton-Hooke supergravity.

The gauge connection one-form for the $\mathcal{N} = 2$ extended Newton-Hooke reads

$$\begin{aligned} A &= \omega J + \omega^a G_a + sS + \tau H + e^a P_a + mM + t_1 T_1 + t_2 T_2 + u_1 U_1 + u_2 U_2 \\ &\quad + \bar{\psi}^+ Q^+ + \bar{\psi}^- Q^- + \bar{\rho} R, \end{aligned} \quad (4.20)$$

with s and m being the gauge field related to the central charges S and M . The extended Newton-Hooke superalgebra admits a non-degenerate bilinear invariant tensor whose

components are given by

$$\begin{aligned}
 \langle \text{JS} \rangle &= -\sigma_0, & \langle \text{G}_a \text{G}_b \rangle &= \sigma_0 \delta_{ab}, \\
 \langle \text{JM} \rangle &= \langle \text{HS} \rangle = -\beta_1, & \langle \text{G}_a \text{P}_b \rangle &= \beta_1 \delta_{ab}, \\
 \langle \text{HM} \rangle &= -\frac{\sigma_0}{\ell^2}, & \langle \text{P}_a \text{P}_b \rangle &= \frac{\sigma_0}{\ell^2} \delta_{ab}, \\
 \langle \text{T}_1 \text{T}_2 \rangle &= \sigma_0, & \langle \text{T}_1 \text{U}_2 \rangle &= \langle \text{T}_2 \text{U}_1 \rangle = \beta_1, \\
 \langle \text{U}_1 \text{U}_2 \rangle &= -\frac{\beta_1}{\ell}, & \langle \text{Q}_\alpha^- \text{Q}_\beta^- \rangle &= \langle \text{Q}_\alpha^+ \text{R}_\beta \rangle = 2 \left(\beta_1 + \frac{\sigma_0}{\ell} \right) C_{\alpha\beta}.
 \end{aligned} \tag{4.21}$$

Here, σ_0 and β_1 are related to the relativistic $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ constants through the $S_E^{(2)}$ elements:

$$\sigma_0 = \lambda_2 \tilde{\alpha}_0, \quad \beta_1 = \lambda_2 \tilde{\alpha}_1. \tag{4.22}$$

Here, σ_0 is related to an exotic NR sector [102, 103] which can be set to zero since the non-degeneracy is satisfied for $\sigma_0 \neq \pm \ell \beta_1$ and $\beta_1 \neq 0$. Then, the NR CS supergravity action based on the $\mathcal{N} = 2$ extended Newton-Hooke superalgebra (3.29) is given by

$$\begin{aligned}
 I_{\text{extended nh}}^{\mathcal{N}=2} &= \frac{k}{4\pi} \int \beta_1 \left[2e_a R^a(\omega^b) - 2mR(\omega) - 2\tau R(s) + \frac{1}{\ell^2} \epsilon^{ac} \tau e_a e_c + 2t_1 du_2 + 2t_2 du_1 \right. \\
 &\quad \left. - \frac{2}{\ell} u_1 du_2 - 2\bar{\psi}^- \nabla \psi^- - 2\bar{\psi}^+ \nabla \rho - 2\bar{\rho} \nabla \psi^+ \right],
 \end{aligned} \tag{4.23}$$

where

$$\begin{aligned}
 R(\omega) &= d\omega, \\
 R^a(\omega^a) &= d\omega^a + \epsilon^{ac} \omega \omega_c, \\
 R(s) &= ds + \frac{1}{2} \epsilon^{ac} \omega_a \omega_c, \\
 \nabla \psi^+ &= d\psi^+ + \frac{1}{2} \omega \gamma_0 \psi^+ + \frac{1}{2\ell} \tau \gamma_0 \psi^+ - \frac{1}{2} t_1 \gamma_0 \psi^+, \\
 \nabla \psi^- &= d\psi^- + \frac{1}{2} \omega \gamma_0 \psi^- + \frac{1}{2\ell} \tau \gamma_0 \psi^- + \frac{1}{2} \omega^a \gamma_a \psi^+ + \frac{1}{2\ell} e^a \gamma_a \psi^+ + \frac{1}{2} t_1 \gamma_0 \psi^-, \\
 \nabla \rho &= d\rho + \frac{1}{2} \omega \gamma_0 \rho + \frac{1}{2} s \gamma_0 \psi^+ + \frac{1}{2} \omega^a \gamma_a \psi^- + \frac{1}{2\ell} e^a \gamma_a \psi^- + \frac{1}{2\ell} \tau \gamma_0 \rho \\
 &\quad + \frac{1}{2\ell} m \gamma_0 \psi^+ - \frac{1}{2} t_1 \gamma_0 \rho - \frac{1}{2} t_2 \gamma_0 \psi^+.
 \end{aligned} \tag{4.24}$$

In the vanishing cosmological constant limit $\ell \rightarrow \infty$, the CS action reproduces the extended Bargmann theory in the presence of the additional bosonic content t_1, t_2, u_1 , and u_2 [102]. The presence of such gauge fields is not whimsical but is due to the non-degeneracy requirement. Indeed, they are related to the bosonic generators $\{\text{T}_1, \text{T}_2, \text{U}_1, \text{U}_2\}$, which appear as expansions of the $\mathfrak{so}(2)$ automorphism generator T and central charge U of the $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra (3.4). As in the relativistic case [109], the extra generators ensure the non-degeneracy of the bilinear invariant trace, implying the vanishing of the curvature two-forms as equations of motion even in the flat limit. In particular, the bosonic

curvature two-forms read

$$\begin{aligned}
 F(\omega) &= R(\omega) + \frac{1}{2\ell} \bar{\psi}^+ \gamma^0 \psi^+, \\
 F(\omega^a) &= R^a(\omega^b) + \frac{1}{\ell^2} \epsilon^{ac} \tau e_c + \frac{1}{\ell} \bar{\psi}^+ \gamma^a \psi^-, \\
 F(\tau) &= d\tau + \frac{1}{2} \bar{\psi}^+ \gamma^0 \psi^+, \\
 F(e^a) &= de^a + \epsilon^{ac} \omega e_c + \epsilon^{ac} \tau \omega_c + \bar{\psi}^+ \gamma^a \psi^-, \\
 F(s) &= R(s) + \frac{1}{2\ell^2} \epsilon^{ac} e_a e_c + \frac{1}{2\ell} \bar{\psi}^- \gamma^0 \psi^- + \frac{1}{\ell} \bar{\psi}^+ \gamma^0 \rho, \\
 F(m) &= dm + \epsilon^{ac} \omega_a e_c + \frac{1}{2} \bar{\psi}^- \gamma^0 \psi^- + \bar{\psi}^+ \gamma^0 \rho, \\
 F(t_1) &= dt_1 + \frac{1}{2\ell} \bar{\psi}^+ \gamma^0 \psi^+, \\
 F(t_2) &= dt_2 - \frac{1}{2\ell} \bar{\psi}^- \gamma^0 \psi^- + \frac{1}{\ell} \bar{\psi}^+ \gamma^0 \rho, \\
 F(u_1) &= du_1 + \frac{1}{2} \bar{\psi}^+ \gamma^0 \psi^+, \\
 F(u_2) &= du_2 - \frac{1}{2} \bar{\psi}^- \gamma^0 \psi^- + \bar{\psi}^+ \gamma^0 \rho.
 \end{aligned} \tag{4.25}$$

On the other hand, the fermionic curvature two-forms are defined in (4.24).

Generalized $\mathcal{N} = 2$ Newton-Hooke and Galilean CS supergravity theories

A CS supergravity action based on the $\mathcal{N} = 2$ $\mathfrak{nh}^{(N)}$ superalgebra (3.38)–(3.39) can be constructed from the gauge connection one-form,

$$\begin{aligned}
 A &= \sum_{m=0}^N \left(\omega^{(m)} \mathbf{J}^{(m)} + \tau^{(m)} \mathbf{H}^{(m)} + t^{(m)} \mathbf{T}^{(m)} + u^{(m)} \mathbf{U}^{(m)} + \bar{\psi}^{+(m)} \mathbf{Q}^{+(m)} \right) \\
 &+ \sum_{m=0}^{N-1} \left(\omega^{a(m)} \mathbf{G}_a^{(m)} + e^{a(m)} \mathbf{P}_a^{(m)} + \bar{\psi}^{-(m)} \mathbf{Q}^{-(m)} \right),
 \end{aligned} \tag{4.26}$$

and the non-vanishing components of the invariant tensor,

$$\begin{aligned}
 \langle \mathbf{J}^{(m)} \mathbf{H}^{(n)} \rangle &= -\beta_{m+n}, & \langle \mathbf{G}_a^{(m)} \mathbf{P}_b^{(n)} \rangle &= \beta_{m+n+1} \delta_{ab}, \\
 \langle \mathbf{T}^{(m)} \mathbf{U}^{(n)} \rangle &= \beta_{m+n}, & \langle \mathbf{U}^{(m)} \mathbf{U}^{(n)} \rangle &= -\frac{1}{\ell} \beta_{m+n}, \\
 \langle \mathbf{Q}_\alpha^{-(m)} \mathbf{Q}_\beta^{-(n)} \rangle &= 2\beta_{m+n+1} C_{\alpha\beta}, & \langle \mathbf{Q}_\alpha^{+(m)} \mathbf{Q}_\beta^{+(n)} \rangle &= 2\beta_{n+m} C_{\alpha\beta},
 \end{aligned} \tag{4.27}$$

with $1 \leq p \leq N$ for the β_p constants. Here, the β_p constants are related to the relativistic ones (3.2) through the $S_E^{(2N)}$ semigroup elements as

$$\beta_p = \lambda_{2p} \tilde{\alpha}_1. \tag{4.28}$$

Without loss of generality, we can set $\beta_0 = 0$, β_0 being related to the three-dimensional Galilean gravity term [85, 114].

A generalized NR CS supergravity action constructed from the gauge connection one-form (4.26) is then given by

$$I_{\mathfrak{nh}^{(N)}}^{\mathcal{N}=2} = \frac{k}{4\pi} \int \sum_{p=1}^N \beta_p \delta_{m+n}^p \left[-2\tau^{(m)} R(\omega^{(n)}) + \frac{\delta_{q+r+1}^n}{\ell^2} \epsilon^{ac} \tau^{(m)} e_a^{(q)} e_c^{(r)} + 2t^{(m)} du^{(n)} - \frac{1}{\ell} u^{(m)} du^{(n)} - 2\bar{\psi}^{+(m)} \nabla \psi^{+(n)} \right] + \beta_p \delta_{m+n+1}^p \left[2e_a^{(m)} R^a(\omega^{b(n)}) - 2\bar{\psi}^{-(m)} \nabla \psi^{-(n)} \right], \quad (4.29)$$

with

$$\begin{aligned} R(\omega^{(m)}) &= d\omega^{(m)} + \frac{1}{2} \sum_{p,q=0}^{N-1} \delta_{p+q+1}^m \epsilon^{ac} \omega_a^{(p)} \omega_c^{(q)}, \\ R^a(\omega^{a(m)}) &= d\omega^{a(m)} + \sum_{n,p=0}^N \delta_{n+p}^m \epsilon^{ac} \omega^{(n)} \omega_c^{(p)}, \\ \nabla \psi^{+(m)} &= d\psi^{+(m)} + \frac{1}{2} \sum_{n,p=0}^N \delta_{n+p}^m \left(\omega^{(n)} \gamma_0 \psi^{+(p)} + \frac{1}{\ell} \tau^{(n)} \gamma_0 \psi^{+(p)} - t^{(n)} \gamma_0 \psi^{+(p)} \right) \\ &\quad + \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \left(\omega^{a(n)} \gamma_a \psi^{-(p)} + \frac{1}{\ell} e^{a(n)} \gamma_a \psi^{-(p)} \right), \\ \nabla \psi^{-(m)} &= d\psi^{-(m)} + \frac{1}{2} \sum_{n,p=0}^{N-1} \delta_{n+p}^m \left(\omega^{(n)} \gamma_0 \psi^{-(p)} + \frac{1}{\ell} \tau^{(n)} \gamma_0 \psi^{-(p)} + t^{(n)} \gamma_0 \psi^{-(p)} \right) \\ &\quad + \omega^{a(n)} \gamma_a \psi^{+(p)} + \frac{1}{\ell} e^{a(n)} \gamma_a \psi^{+(p)}. \end{aligned} \quad (4.30)$$

The obtained NR CS supergravity action is based on the $\mathcal{N} = 2$ $\mathfrak{nh}^{(N)}$ superalgebra and can be seen as the supersymmetric extension of the generalized Newton-Hooke gravity action introduced in [85]. In the vanishing cosmological constant limit $\ell \rightarrow \infty$ the CS action reproduces a generalized Galilean supergravity. As in the generalized super AdS-Carroll case, the CS supergravity action can be written as the sum of Lagrangians invariant under the $\mathcal{N} = 2$ $\mathfrak{nh}^{(p)}$ superalgebra for $p = 1, 2, \dots, N$. Indeed, the CS action (4.29) can be expressed as follows:

$$I_{\mathfrak{nh}^{(N)}}^{\mathcal{N}=2} = \frac{k}{4\pi} \int \sum_{p=1}^N \beta_p \mathcal{L}_{\mathfrak{nh}^{(p)}}^{\mathcal{N}=2}. \quad (4.31)$$

For each Lagrangian we have set $\beta_i = 0$ for $i \neq p$ since the non-degeneracy condition for $\mathfrak{nh}^{(p)}$ is given by $\beta_p \neq 0$. Then, for an arbitrary p , the equations of motion for the $\mathfrak{nh}^{(p)}$ supergravity are given by the vanishing of the following curvature two-forms (field strengths of the bosonic one-form gauge potentials):

$$\begin{aligned} F(\omega^{(m)}) &= R(\omega^{(m)}) + \frac{1}{2} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \left(\frac{1}{\ell^2} \epsilon^{ac} e_a^{(n)} e_c^{(p)} + \frac{1}{\ell} \bar{\psi}^{-(n)} \gamma^0 \psi^{-(p)} \right) \\ &\quad + \frac{1}{2\ell} \sum_{n,p=0}^N \delta_{n+p}^m \bar{\psi}^{+(n)} \gamma^0 \psi^{+(p)}, \\ F^a(\omega^{b(m)}) &= R^a(\omega^{b(m)}) + \sum_{n,p=0}^{N-1} \delta_{n+p}^m \left(\frac{1}{\ell^2} \epsilon^{ac} \tau^{(n)} e_c^{(p)} + \frac{1}{\ell} \bar{\psi}^{+(n)} \gamma^a \psi^{-(p)} \right), \end{aligned}$$

$$\begin{aligned}
 F\left(\tau^{(m)}\right) &= d\tau^{(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \left(\epsilon^{ac} \omega_a^{(n)} e_c^{(p)} + \frac{1}{2} \bar{\psi}^{-(n)} \gamma^0 \psi^{-(p)} \right) \\
 &\quad + \frac{1}{2} \sum_{n,p=0}^N \delta_{n+p}^m \bar{\psi}^{+(n)} \gamma^0 \psi^{+(p)}, \\
 F^a\left(e^{b(m)}\right) &= de^{a(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p}^m \left(\epsilon^{ac} \omega_c^{(n)} e_c^{(p)} + \epsilon^{ac} \tau^{(n)} \omega_c^{(p)} + \bar{\psi}^{+(n)} \gamma^a \psi^{-(p)} \right), \\
 F\left(t^{(m)}\right) &= dt^{(m)} + \frac{1}{2\ell} \sum_{n,p=0}^N \delta_{n+p}^m \bar{\psi}^{+(n)} \gamma^0 \psi^{+(p)} + \frac{1}{2\ell} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \bar{\psi}^{-(n)} \gamma^0 \psi^{-(p)}, \\
 F\left(u^{(m)}\right) &= du^{(m)} + \frac{1}{2} \sum_{n,p=0}^N \delta_{n+p}^m \bar{\psi}^{+(n)} \gamma^0 \psi^{+(p)} + \frac{1}{2} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \bar{\psi}^{-(n)} \gamma^0 \psi^{-(p)}; \tag{4.32}
 \end{aligned}$$

the fermionic curvature two-forms are defined in (4.30). One can notice that the Lagrangian for $p = 1$ corresponds to the extended Newton-Hooke supergravity Lagrangian obtained in (4.23). For $p \geq 2$, the Lagrangian reproduces post-Newtonian extensions of the extended Newton-Hooke supergravity. For instance, $p = 2$ reproduces the so-called exotic Newtonian supergravity [103]. Indeed, by considering the following redefinition of the gauge fields:

$$\begin{aligned}
 \omega^{(0)} &= \omega, & \omega^{(1)} &= s, & \omega^{(2)} &= z, \\
 \tau^{(0)} &= \tau, & \tau^{(1)} &= m, & \tau^{(2)} &= t, \\
 t^{(0)} &= t_1, & t^{(1)} &= t_2, & t^{(2)} &= t_3, \\
 u^{(0)} &= u_1, & u^{(1)} &= u_2, & u^{(2)} &= u_3, \\
 \psi_\alpha^{+(0)} &= \psi_\alpha^+, & \psi_\alpha^{+(1)} &= \rho_\alpha, & \psi_\alpha^{+(2)} &= \phi_\alpha^+, \\
 \omega_a^{(0)} &= \omega_a, & e_a^{(0)} &= e_a, & \psi_\alpha^{-(0)} &= \psi_\alpha^-, \\
 \omega_a^{(1)} &= b_a, & e_a^{(1)} &= t_a, & \psi_\alpha^{-(1)} &= \phi_\alpha^-, \tag{4.33}
 \end{aligned}$$

we get

$$\begin{aligned}
 \mathcal{L}_{\text{nh}^{(2)}}^{\mathcal{N}=2} &= 2e^a R^a(b^b) + 2t^a R^a(\omega^b) - 2\tau R(z) - 2mR(s) - 2yR(\omega) + \frac{2}{\ell^2} \epsilon^{ac} \tau e_a t_c + \frac{1}{\ell^2} \epsilon^{ac} m e_a e_c \\
 &\quad + 2t_1 du_3 + 2t_2 du_2 + 2t_3 du_1 - \frac{2}{\ell} u_1 du_3 - \frac{1}{\ell} u_2 du_2 - 2\bar{\psi}^\pm \nabla \phi^\pm - 2\bar{\phi}^\pm \nabla \psi^\pm - 2\bar{\rho} \nabla \rho. \tag{4.34}
 \end{aligned}$$

Here, the curvature two-forms read

$$\begin{aligned}
 R^a(b^b) &= db^a + \epsilon^{ac} \omega b_c + \epsilon^{ac} s \omega_c, \\
 R(z) &= dz + \epsilon^{ac} \omega_a b_c, \\
 \nabla \phi^+ &= d\phi^+ + \frac{1}{2} \omega \gamma_0 \phi^+ + \frac{1}{2} s \gamma_0 \rho + \frac{1}{2} \omega^a \gamma_a \phi^- + \frac{1}{2} b^a \gamma_a \psi^- + \frac{1}{2\ell} e^a \gamma_a \phi^- + \frac{1}{2\ell} \tau \gamma_0 \phi^+ \\
 &\quad + \frac{1}{2\ell} m \gamma_0 \rho + \frac{1}{2\ell} t^a \gamma_a \psi^- + \frac{1}{2} z \gamma_0 \psi^+ + \frac{1}{2\ell} y \gamma_0 \psi^+ - \frac{1}{2} t_1 \gamma_0 \phi^+ - \frac{1}{2} t_2 \gamma_0 \rho - \frac{1}{2} t_3 \gamma_0 \psi^+, \\
 \nabla \phi^- &= d\phi^- + \frac{1}{2} \omega \gamma_0 \phi^- + \frac{1}{2} s \gamma_0 \psi^- + \frac{1}{2} \omega^a \gamma_a \rho + \frac{1}{2} b^a \gamma_a \psi^+ + \frac{1}{2\ell} e^a \gamma_a \rho + \frac{1}{2\ell} \tau \gamma_0 \phi^- \\
 &\quad + \frac{1}{2\ell} m \gamma_0 \psi^- + \frac{1}{2\ell} t^a \gamma_a \psi^+ - \frac{1}{2} t_1 \gamma_0 \phi^- - \frac{1}{2} t_2 \gamma_0 \psi^-. \tag{4.35}
 \end{aligned}$$

Moreover, $R(\omega)$, $R^a(\omega^b)$, $R(s)$, $\nabla\psi^\pm$, and $\nabla\rho$ have been defined in (4.24). In the flat limit $\ell \rightarrow \infty$, we recover the extended Newtonian supergravity introduced in [95], which can also be understood as post-Newtonian corrections [115]. At higher order in p , the Lagrangians $\mathcal{L}_{\text{nh}^{(2)}}^{\mathcal{N}=2}$ extends to the supersymmetric case the family of Newton-Hooke gravity theories discussed in [85].

Interestingly, the $\mathcal{N} = 2$ generalized Newton-Hooke CS supergravity action (4.29) can alternatively be obtained from the relativistic one,

$$I_{\text{AdS}}^{\mathcal{N}=2} = \int \tilde{\alpha}_1 \left[2e_A R^A + \frac{1}{3\ell^2} \epsilon_{ABC} e^A e^B e^C + \mathfrak{t} du - \frac{1}{\ell^2} u du - 2\bar{\Psi}^i \nabla \Psi^i \right], \quad (4.36)$$

where

$$\begin{aligned} R^A &= d\omega^A + \frac{1}{2} \epsilon^{ABC} \omega_B \omega_C, \\ T^A &= de^A + \frac{1}{2} \epsilon^{ABC} \omega_B e_C, \\ \nabla \Psi^i &= d\Psi^i + \frac{1}{2} \omega^A \gamma_A \Psi^i + \frac{1}{2\ell} e^A \gamma_A \Psi^i + \mathfrak{t} \epsilon^{ij} \Psi^j. \end{aligned} \quad (4.37)$$

Indeed, the NR supergravity action (4.23) is obtained by expanding the relativistic theory (4.36) after identifying the NR gauge fields in terms of the relativistic ones through the $S_E^{(2)}$ elements as

$$\begin{aligned} \omega^{(m)} &= \lambda_{2m} \omega_0, & \tau^{(m)} &= \lambda_{2m} e_0 & \psi^{+(m)} &= \lambda_{2m} \Psi^+, & t^{(m)} &= \lambda_{2m} \mathfrak{t}, & u^{(m)} &= \lambda_{2m} u, \\ \omega_a^{(m)} &= \lambda_{2m+1} \omega_a, & e_a^{(m)} &= \lambda_{2m+1} e_a & \psi^{-(m)} &= \lambda_{2m+1} \Psi^-, \end{aligned} \quad (4.38)$$

where we have defined

$$\Psi_\alpha^\pm = \frac{1}{\sqrt{2}} \left(\Psi_\alpha^1 \pm (\gamma^0)_{\alpha\beta} \Psi_\beta^2 \right). \quad (4.39)$$

4.3 $\mathcal{N} = 2$ (AdS-)static Chern-Simons supergravity and its generalization

A third class of kinematical superalgebras are the static one which, as we have shown in the previous section, can be either be obtained from the UR superalgebras or from the NR ones. Here, we present the three-dimensional CS supergravity action based on the $\mathcal{N} = 2$ extended AdS-static superalgebra (3.32)–(3.33) and its flat limit.

The gauge connection one-form for the $\mathcal{N} = 2$ AdS-static superalgebra is given by

$$\begin{aligned} A &= \omega J + \omega^a G_a + s S + \tau H + e^a P_a + m M + t_1 T_1 + t_2 T_2 + u_1 U_1 + u_2 U_2 \\ &\quad + \bar{\psi}^+ Q^+ + \bar{\psi}^- Q^- + \bar{\rho} R. \end{aligned} \quad (4.40)$$

The corresponding curvature two-form reads

$$\begin{aligned} F &= \mathcal{F}(\omega) J + \mathcal{F}^a(\omega^b) G_a + \mathcal{F}(\tau) H + \mathcal{F}^a(e^b) P_a + \mathcal{F}(t_1) T_1 + \mathcal{F}(t_2) T_2 \\ &\quad + \mathcal{F}(u_1) U_1 + \mathcal{F}(u_2) U_2 + \nabla \bar{\psi}^+ Q^+ + \nabla \bar{\psi}^- Q^- + \nabla \bar{\rho} R, \end{aligned} \quad (4.41)$$

with

$$\begin{aligned} \mathcal{F}(\omega) &= d\omega = \mathcal{R}(\omega), \\ \mathcal{F}^a(\omega^b) &= d\omega^a + \epsilon^{ac} \omega \omega_c + \frac{1}{\ell^2} \epsilon^{ac} \tau e_c + \frac{1}{\ell} \bar{\psi}^+ \gamma^a \psi^- = \mathcal{R}^a(\omega^b) + \frac{1}{\ell^2} \epsilon^{ac} \tau e_c + \frac{1}{\ell} \bar{\psi}^+ \gamma^a \psi^-, \\ \mathcal{F}(\tau) &= d\tau + \frac{1}{2} \bar{\psi}^+ \gamma^0 \psi^+, \end{aligned}$$

$$\begin{aligned}
 \mathcal{F}(e^a) &= de^a + \epsilon^{ac}\omega e_c, \\
 \mathcal{F}(s) &= ds + \frac{1}{2\ell^2}\epsilon^{ac}e_a e_c = \mathcal{R}(s) + \frac{1}{2\ell^2}\epsilon^{ac}e_a e_c, \\
 \mathcal{F}(m) &= dm + \epsilon^{ac}\omega_a e_c + \frac{1}{2}\bar{\psi}^-\gamma^0\psi^-, \\
 \mathcal{F}(t_1) &= dt_1, \\
 \mathcal{F}(t_2) &= dt_2, \\
 \mathcal{F}(u_1) &= du_1 + \frac{1}{2}\bar{\psi}^+\gamma^0\psi^+, \\
 \mathcal{F}(u_2) &= du_2 + \frac{1}{2}\bar{\psi}^-\gamma^0\psi^- + \bar{\psi}^+\gamma^0\rho, \\
 \nabla\psi^+ &= d\psi^+ + \frac{1}{2}\omega\gamma_0\psi^+ - \frac{1}{2}t_1\gamma_0\psi^+, \\
 \nabla\psi^- &= d\psi^- + \frac{1}{2}\omega\gamma_0\psi^- + \frac{1}{2\ell}e^a\gamma_a\psi^+ + \frac{1}{2}t_1\gamma_0\psi^-, \\
 \nabla\rho &= d\rho + \frac{1}{2}\omega\gamma_0\rho + \frac{1}{2}s\gamma_0\psi^+ + \frac{1}{2\ell}e^a\gamma_a\psi^- - \frac{1}{2}t_1\gamma_0\rho - \frac{1}{2}t_2\gamma_0\psi^+.
 \end{aligned} \tag{4.42}$$

Let us note that the curvature two-forms above are quite different from the extended Newton-Hooke and extended Bargmann ones, although the gauge connection one-form (4.40) is identical to the extended Newton-Hooke one (4.20). The non-vanishing components of the invariant tensor for the $\mathcal{N} = 2$ AdS-static superalgebra are given by

$$\begin{aligned}
 \langle \text{JS} \rangle &= -\nu_0, & \langle \text{P}_a \text{P}_b \rangle &= \frac{\nu_0}{\ell^2} \delta_{ab}, \\
 \langle \text{JM} \rangle &= \langle \text{HS} \rangle = -\mu_1, & \langle \text{G}_a \text{P}_b \rangle &= \mu_1 \delta_{ab}, \\
 \langle \text{T}_1 \text{T}_2 \rangle &= \nu_0, & \langle \text{T}_1 \text{U}_2 \rangle &= \langle \text{T}_2 \text{U}_1 \rangle = \mu_1, \\
 \langle \text{Q}_\alpha^- \text{Q}_\beta^- \rangle &= 2\mu_1 C_{\alpha\beta}, & \langle \text{Q}_\alpha^+ \text{R}_\beta \rangle &= 2\mu_1 C_{\alpha\beta},
 \end{aligned} \tag{4.43}$$

where ν_0 and μ_1 can be written in terms of the $\mathcal{N} = 2$ AdS-Carroll constants (4.5) and the element of the $S_E^{(2)}$ semigroup as

$$\nu_0 = \lambda_2 \alpha_0, \quad \mu_1 = \lambda_2 \alpha_1.$$

Alternatively, ν_0 and μ_1 can also be recovered from the extended Newton-Hooke ones (4.21) with

$$\nu_0 = \lambda_0 \sigma_0, \quad \mu_1 = \lambda_2 \beta_1.$$

The exotic sector proportional to ν_0 can be set to zero without affecting the non-degeneracy criterion, which requires $\mu_1 \neq 0$. The $\mathcal{N} = 2$ AdS-static CS supergravity action constructed from the connection one-form (4.40) and the invariant tensor (4.43) is then given by

$$\begin{aligned}
 I_{\text{ads-stat}}^{\mathcal{N}=2} &= \frac{k}{4\pi} \int \mu_1 \left[2e_a \mathcal{R}^a(\omega^b) - 2m \mathcal{R}(\omega) - 2\tau \mathcal{R}(s) + \frac{1}{\ell^2} \epsilon^{ac} \tau e_a e_c + 2t_1 du_2 + 2t_2 du_1 \right. \\
 &\quad \left. - 2\bar{\psi}^+ \nabla \rho - 2\bar{\rho} \nabla \psi^+ - 2\bar{\psi}^- \nabla \psi^- \right].
 \end{aligned} \tag{4.44}$$

The obtained CS action differs from the $\mathcal{N} = 2$ extended Bargmann one (see (4.23) for $\ell \rightarrow \infty$) in the definition of the curvature two-forms (4.42). The non-degeneracy of the bilinear invariant tensor (4.43) ensures the vanishing of the AdS-static curvature two-forms (4.42) as field equations which are different from the extended Bargmann ones (4.25). Nonetheless, a

reinterpretation of the e^a and ω^a gauge fields allows us to recover the known non-relativistic supergravity action [78, 97, 102]. Indeed, the $\mathcal{N} = 2$ extended Bargmann CS supergravity action appears by interchanging the roles of e^a and $\ell\omega^a$. On the other hand, the vanishing cosmological constant limit $\ell \rightarrow \infty$ reproduces an $\mathcal{N} = 2$ static CS supergravity action. The flat limit is free of degeneracy due to the presence of the bosonic generators $\{T_1, T_2, U_1, U_2\}$, ensuring the vanishing of the proper curvature two-forms as field equations both in the AdS-static and static supergravity. The CS supergravity action (4.44) is invariant under the following supersymmetry transformation rules:

$$\begin{aligned}
 \delta\omega &= 0, \\
 \delta\omega^a &= \frac{1}{\ell}\bar{\varepsilon}^+\gamma^a\psi^- + \frac{1}{\ell}\bar{\varepsilon}^-\gamma^a\psi^+, \\
 \delta\tau &= \bar{\varepsilon}^+\gamma^0\psi^+, \\
 \delta e^a &= 0, \\
 \delta s &= 0, \\
 \delta m &= \bar{\varepsilon}^-\gamma^0\psi^- + \bar{\varepsilon}^+\gamma^0\rho + \bar{\varrho}\gamma^0\psi^+, \\
 \delta t_1 &= 0, \\
 \delta t_2 &= 0, \\
 \delta u_1 &= \bar{\varepsilon}^+\gamma^0\psi^+, \\
 \delta u_2 &= \bar{\varepsilon}^+\gamma^0\rho + \bar{\varrho}^+\gamma^0\psi^+ - \bar{\varepsilon}^-\gamma^0\psi^-, \\
 \delta\psi^+ &= d\varepsilon^+ + \frac{1}{2}\omega\gamma_0\varepsilon^+ + \frac{1}{2\ell}e^a\gamma_a\varepsilon^- - \frac{1}{2}t_1\gamma_0\varepsilon^+, \\
 \delta\psi^- &= d\varepsilon^- + \frac{1}{2}\omega\gamma_0\varepsilon^- + \frac{1}{2\ell}e^a\gamma_a\varepsilon^+ + \frac{1}{2}t_1\gamma_0\varepsilon^-, \\
 \delta\rho &= d\varrho + \frac{1}{2}\omega\gamma_0\varrho + \frac{1}{2}s\gamma_0\varepsilon^+ + \frac{1}{2\ell}e^a\gamma_a\varepsilon^- - \frac{1}{2}t_1\gamma_0\varrho - \frac{1}{2}t_2\gamma_0\varepsilon^+,
 \end{aligned} \tag{4.45}$$

where the gauge parameters $\varepsilon^\pm$ and ϱ are related to the fermionic charges $Q^\pm$ and ρ , respectively. The transformation rules for the $\mathcal{N} = 2$ static supergravity theory are derived in the flat limit $\ell \rightarrow \infty$.

Generalized $\mathcal{N} = 2$ (AdS-)static supergravity

The $\mathcal{N} = 2$ **adS-stat**^(N) superalgebra (3.44) admits a non-degenerate invariant tensor whose non-vanishing components read

$$\begin{aligned}
 \langle J^{(m)} H^{(n)} \rangle &= -\mu_{m+n}, & \langle G_a^{(m)} P_b^{(n)} \rangle &= \mu_{m+n+1}\delta_{ab}, \\
 \langle T^{(m)} U^{(n)} \rangle &= \mu_{m+n}, & \langle Q_\alpha^{-(m)} Q_\beta^{-(n)} \rangle &= 2\mu_{m+n+1}C_{\alpha\beta}, \\
 \langle Q_\alpha^{+(m)} Q_\beta^{+(n)} \rangle &= 2\mu_{n+m}C_{\alpha\beta},
 \end{aligned} \tag{4.46}$$

with $1 \leq p \leq N$ for the μ_p constants.⁵ Here, the μ_p constants can be either be obtained from the $\mathcal{N} = 2$ AdS-Carroll ones (4.5),

$$\mu_p = \lambda_{2p}\alpha_1, \tag{4.47}$$

⁵Here we have intentionally omitted the exotic sector.

or from the $\mathcal{N} = 2 \, \mathfrak{nh}^{(N)}$ constants (4.27),

$$\mu_p = \lambda_2 \beta_p. \quad (4.48)$$

A generalized $\mathcal{N} = 2$ (AdS-)static CS supergravity action can be constructed from the invariant tensor (4.46) and the gauge connection one-form for the $\mathcal{N} = 2 \, \mathfrak{ad}\mathfrak{s}\text{-}\mathfrak{stat}^{(N)}$ superalgebra,

$$\begin{aligned} A = & \sum_{m=0}^N \left(\omega^{(m)} \mathbf{J}^{(m)} + \tau^{(m)} \mathbf{H}^{(m)} + t^{(m)} \mathbf{T}^{(m)} + u^{(m)} \mathbf{U}^{(m)} + \bar{\psi}^{+(m)} \mathbf{Q}^{+(m)} \right) \\ & + \sum_{m=0}^{N-1} \left(\omega^{a(m)} \mathbf{G}_a^{(m)} + e^{a(m)} \mathbf{P}_a^{(m)} + \bar{\psi}^{-(m)} \mathbf{Q}^{-(m)} \right), \end{aligned} \quad (4.49)$$

and the corresponding curvature two-forms,

$$\begin{aligned} \mathcal{F}(\omega^{(m)}) &= d\omega^{(m)} + \frac{1}{2\ell^2} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \epsilon^{ac} e_a^{(n)} e_c^{(p)}, \\ \mathcal{F}^a(\omega^{b(m)}) &= d\omega^{a(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p}^m \left(\epsilon^{ac} \omega^{(n)} \omega_c^{(p)} + \frac{1}{\ell^2} \epsilon^{ac} \tau^{(n)} e_c^{(p)} + \frac{1}{\ell} \bar{\psi}^{+(n)} \gamma^a \psi^{-(p)} \right), \\ \mathcal{F}(\tau^{(m)}) &= d\tau^{(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \left(\epsilon^{ac} \omega_a^{(n)} e_c^{(p)} + \frac{1}{2} \bar{\psi}^{-(n)} \gamma^0 \psi^{-(p)} \right) \\ &\quad + \frac{1}{2} \sum_{n,p=0}^N \delta_{n+p}^m \bar{\psi}^{+(n)} \gamma^0 \psi^{+(p)}, \\ \mathcal{F}^a(e^{b(m)}) &= de^{a(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p}^m \epsilon^{ac} \omega^{(n)} e_c^{(p)}, \\ \mathcal{F}(t^{(m)}) &= dt^{(m)}, \\ \mathcal{F}(u^{(m)}) &= du^{(m)} + \frac{1}{2} \sum_{n,p=0}^N \delta_{n+p}^m \bar{\psi}^{+(n)} \gamma^0 \psi^{+(p)} + \frac{1}{2} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m \bar{\psi}^{-(n)} \gamma^0 \psi^{-(p)}, \\ \nabla \psi^{+(m)} &= d\psi^{+(m)} + \frac{1}{2} \sum_{n,p=0}^N \delta_{n+p}^m \left(\omega^{(n)} \gamma_0 \psi^{+(p)} - t^{(n)} \gamma_0 \psi^{+(p)} \right) \\ &\quad + \frac{1}{\ell} \sum_{n,p=0}^{N-1} \delta_{n+p+1}^m e^{a(n)} \gamma_a \psi^{-(p)}, \\ \nabla \psi^{-(m)} &= d\psi^{-(m)} + \frac{1}{2} \sum_{n,p=0}^{N-1} \delta_{n+p}^m \left(\omega^{(n)} \gamma_0 \psi^{-(p)} + t^{(n)} \gamma_0 \psi^{-(p)} + \frac{1}{\ell} e^{a(n)} \gamma_a \psi^{+(p)} \right). \end{aligned} \quad (4.50)$$

Then, the general expression for the CS supergravity action based on the $\mathcal{N} = 2 \, \mathfrak{ad}\mathfrak{s}\text{-}\mathfrak{stat}^{(N)}$ superalgebra (3.44) is given by

$$\begin{aligned} I_{\mathfrak{ad}\mathfrak{s}\text{-}\mathfrak{stat}^{(N)}}^{\mathcal{N}=2} = & \frac{k}{4\pi} \int \sum_{p=1}^N \mu_p \delta_{m+n+1}^p \left[2e_a^{(m)} \mathcal{R}^a(\omega^{b(n)}) - 2\bar{\psi}^{-(m)} \nabla \psi^{-(n)} \right] \\ & + \mu_p \delta_{m+n}^p \left[-2\tau^{(m)} \mathcal{R}(\omega^{(n)}) + \frac{\delta_{g+r+1}^n}{\ell^2} \epsilon^{ac} \tau^{(m)} e_a^{(q)} e_c^{(r)} + 2t^{(m)} du^{(n)} - 2\bar{\psi}^{+(m)} \nabla \psi^{+(n)} \right], \end{aligned} \quad (4.51)$$

where

$$\begin{aligned}\mathcal{R}\left(\omega^{(n)}\right) &= d\omega^{(n)}, \\ \mathcal{R}^a\left(\omega^{b(n)}\right) &= d\omega^{a(m)} + \sum_{n,p=0}^{N-1} \delta_{n+p}^m \epsilon^{ac} \omega^{(n)} \omega_c^{(p)}.\end{aligned}\tag{4.52}$$

The $\mathcal{N} = 2$ $\mathfrak{adS}\text{-}\mathfrak{stat}^{(N)}$ CS supergravity action (4.51) can be expressed as the sum of individual Lagrangians $\mathcal{L}_{\mathfrak{adS}\text{-}\mathfrak{stat}^{(p)}}^{\mathcal{N}=2}$, each of which is invariant under the $\mathfrak{adS}\text{-}\mathfrak{stat}^{(p)}$ superalgebra,

$$I_{\mathfrak{adS}\text{-}\mathfrak{stat}^{(N)}}^{\mathcal{N}=2} = \frac{k}{4\pi} \int \sum_{p=1}^N \mu_p \mathcal{L}_{\mathfrak{adS}\text{-}\mathfrak{stat}^{(p)}}^{\mathcal{N}=2}.\tag{4.53}$$

For $p = 1$, we recover the $\mathcal{N} = 2$ AdS-static CS Lagrangian (4.44) along the μ_1 constant. Generalized AdS-static supergravity actions are derived for $p \geq 2$. According to figures 4 and 5, the corresponding $\mathcal{N} = 2$ static CS supergravity theory together with its generalizations are obtained either as a vanishing cosmological constant limit $\ell \rightarrow \infty$ of the $\mathfrak{adS}\text{-}\mathfrak{stat}^{(N)}$ theory, or by expanding the $\mathcal{N} = 2$ $\mathfrak{gal}^{(N)}/\mathfrak{car}^{(N)}$ supergravity models. In the expansion procedure, the $\mathcal{N} = 2$ $\mathfrak{stat}^{(N)}$ CS supergravity action can be obtained from the generalized Galilean/Carrollian supergravity action. Indeed, the generalized static gauge fields can be expressed in terms of the $\mathfrak{gal}^{(N)}$ ones and the $S_E^{(2)}$ elements as

$$\begin{aligned}\omega^{(m)} &= \lambda_0 \omega^{(m)}, \quad \tau^{(m)} = \lambda_2 \tau^{(m)} \quad \psi^{+(m)} = \lambda_1 \psi^{+(m)}, \quad t^{(m)} = \lambda_0 t^{(m)}, \quad u^{(m)} = \lambda_2 u^{(m)}, \\ \omega_a^{(m)} &= \lambda_2 \omega_a^{(m)} \quad e_a^{(m)} = \lambda_0 e_a^{(m)} \quad \psi^{-(m)} = \lambda_1 \psi^{-(m)},\end{aligned}\tag{4.54}$$

or in terms of the $\mathfrak{car}^{(N)}$ ones and the $S_E^{(2N)}$ elements,

$$\begin{aligned}\omega^{(m)} &= \lambda_{2m} \omega, \quad \tau^{(m)} = \lambda_{2m} \tau \quad \psi^{+(m)} = \lambda_{2m} \psi^+, \quad t^{(m)} = \lambda_{2m} t, \quad u^{(m)} = \lambda_{2m} u, \\ \omega_a^{(m)} &= \lambda_{2m+1} \omega_a \quad e_a^{(m)} = \lambda_{2m+1} e_a \quad \psi^{-(m)} = \lambda_{2m+1} \psi^-.\end{aligned}\tag{4.55}$$

5 Discussion

In this work, we have presented the supersymmetric extension of kinematical Lie algebras by applying systematic S -expansions starting from a $\mathfrak{so}(2)$ -extension of the $\mathfrak{osp}(2,2) \times \mathfrak{sp}(2)$ superalgebra. We have classified both NR and UR regimes of the relativistic $\mathcal{N} = 2$ AdS and Poincaré superalgebras constrained by the non-degeneracy condition. Such condition ensures that the field equations derived from the corresponding CS supergravity action are given by the vanishing of the curvatures for each supersymmetric (extended) kinematical Lie algebra. Our results can be summarized through the cube 4, in which both the relativistic and the NL superalgebras admit a non-degenerate invariant tensor. To this end, each NL regime requires particular considerations at the level of the semigroups and the subspaces of the original superalgebra. We then extend our results to post-Newtonian and post-Carrollian superalgebras, whose expansion relations have been summarized in cube 5. We have also considered the construction of CS supergravity theories for each (extended) kinematical superalgebra and their generalizations.

The general framework to derive NL regimes of known relativistic supergravity theories presented here could serve as a starting point for approaching various open issues. Of particular interest is the extension of our results to higher spacetime dimensions. In odd spacetime dimensions, the expansion method could be applied to distinct CS (super)gravity models defined in arbitrary higher spacetime dimensions [63]. Nonetheless, a geometric approach à la MacDowell-Mansouri [116] is required to explore the NL counterpart of (super)gravity theories defined in even spacetime dimensions. At the bosonic level, it would be interesting to explore the NL version of the Lovelock gravity [117, 118] which reproduces the CS and Born-Infeld (BI) gravity for particular values of the Lovelock parameters. One could explore whether, as in the relativistic case, the obtained NL CS and BI gravity theories can be recovered from a NL version of the Lovelock gravity model. Our results could be useful to explore the NL regime of the applications of higher-curvature terms in holography [119, 120]. A first step toward holographic developments involves the potential extension to higher dimensions and, above all, the study of the resulting theory in the presence of a non-trivial spacetime boundary, following the approach of [121–125]. This shall be followed by an investigation of the asymptotic symmetries and the theory emerging at the boundary. Further applications in the context of holography and analogue modeling include, for instance, the study of hydrodynamic systems — e.g., the “Carrollian fluids” [126] — within the framework of gauge-gravity duality applied within relativistic and non-relativistic physics.

Another aspect that deserves to be explored is the conformal extension of NL (super)algebras. The S -expansion method employed here could be useful to classify diverse NL (super)algebras and extend to conformal symmetries both the original cube of Bacry and Lévy-Leblond [5] and its supersymmetric extension presented here. One could expect to obtain the Galilean conformal (**gca**) and Carrollian conformal (**cca**) (super)algebras [10, 12, 127, 128] by expanding the conformal one, after considering suitable partitions of the semigroups and decompositions of the relativistic subspaces. In analogy to the results obtained here, bigger semigroups should allow us to derive post-Newtonian and post-Carrollian extensions of the **gca** and **cca** (super)algebras, whose (anti-)commutators should satisfy a Schrödinger-like (super)algebra. Motivated by the $\mathfrak{bms}_3/\mathfrak{gca}_2$ (or $\mathfrak{bms}_3/\mathfrak{cca}_2$) duality [34, 47], it would be interesting to explore the existence of a duality between two-dimensional extended NL conformal (super)algebras and infinite-dimensional asymptotic (super)algebra in higher spacetime dimensions.

It would be also interesting to extend our results to higher-spin gravity and bigger symmetries. The NL regime of gravity consistently coupled to spin-3 gauge fields has already been discussed in [65, 66, 87]. One could go further and study the distinct NL versions of gravity coupled to spin-5/2 gauge fields, also referred to as hypergravity [129–134]. One should expect to find hypersymmetric extensions of the Galilean and Carrollian algebras. Our method should allow us to select the “good” hyperalgebras provided with non-degenerate bilinear invariant traces useful to construct CS hypergravity actions. In this direction, extended kinematical hyperalgebras should appear in the NR counterpart. On the other hand, one could extend the study of kinematical Lie (super)algebras to the Maxwellian case. The Maxwell algebra, first introduced in [135–137] has received a growing interest due to its different applications in the (super)gravity context [138–151]. A Maxwellian generalization of the cube presented in [90] and its supersymmetric extensions introduced here could require a

subtle treatment. In particular, as it was pointed out in [152], the Maxwellian Carroll algebra obtained as a UR limit of the Maxwell algebra suffers from degeneracy, as in the NR limit. We expect to solve the degeneracy issue appearing in the Carrollian regime with a larger semigroup, analogously to the strategy employed here, in the NR expansion.

Acknowledgments

The authors would like to thank Evelyn Rodríguez for valuable comments. This work was funded by the National Agency for Research and Development ANID - FONDECYT grants No. 1211077 and 11220328. This work was supported by FAA 2023 of the Universidad Católica de la Santísima Concepción (P.C.). L.R. would like to thank the DISAT of the Polytechnic of Turin and the INFN for financial support. P.C. and L.R. would like to thank to the Dirección de Investigación and Vice-rectoría de Investigación of the Universidad Católica de la Santísima Concepción, Chile, for their constant support.

Data Availability Statement. This article has no associated data or the data will not be deposited.

Code Availability Statement. This article has no associated code or the code will not be deposited.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E. Cartan, *Sur les variétés à connexion affine et la théorie de la relativité généralisée (première partie)*, *Ann. Ecole Norm. Sup.* **40** (1923) 325.
- [2] E. Cartan, *Sur les variétés à connexion affine et la théorie de la relativité généralisée. (première partie) (Suite)*, *Annales Sci. Ecole Norm. Sup.* **41** (1924) 1 [[INSPIRE](#)].
- [3] J. Lévy-Leblond, *Une nouvelle limite non-relativiste du groupe de poincaré*, *Ann. Inst. H. Poincaré Phys. Theor.* **3** (1965) 1.
- [4] N.D. Sen Gupta, *On an analogue of the Galilei group*, *Nuovo Cim. A* **44** (1966) 512 [[INSPIRE](#)].
- [5] H. Bacry and J. Lévy-Leblond, *Possible kinematics*, *J. Math. Phys.* **9** (1968) 1605 [[INSPIRE](#)].
- [6] D.T. Son, *Toward an AdS/cold atoms correspondence: A geometric realization of the Schrödinger symmetry*, *Phys. Rev. D* **78** (2008) 046003 [[arXiv:0804.3972](#)] [[INSPIRE](#)].
- [7] K. Balasubramanian and J. McGreevy, *Gravity duals for non-relativistic CFTs*, *Phys. Rev. Lett.* **101** (2008) 061601 [[arXiv:0804.4053](#)] [[INSPIRE](#)].
- [8] S. Kachru, X. Liu and M. Mulligan, *Gravity duals of Lifshitz-like fixed points*, *Phys. Rev. D* **78** (2008) 106005 [[arXiv:0808.1725](#)] [[INSPIRE](#)].
- [9] M. Taylor, *Non-relativistic holography*, [arXiv:0812.0530](#) [[INSPIRE](#)].
- [10] A. Bagchi and R. Gopakumar, *Galilean Conformal Algebras and AdS/CFT*, *JHEP* **07** (2009) 037 [[arXiv:0902.1385](#)] [[INSPIRE](#)].

- [11] S.A. Hartnoll, *Lectures on holographic methods for condensed matter physics*, *Class. Quant. Grav.* **26** (2009) 224002 [[arXiv:0903.3246](#)] [[INSPIRE](#)].
- [12] A. Bagchi, R. Gopakumar, I. Mandal and A. Miwa, *GCA in 2d*, *JHEP* **08** (2010) 004 [[arXiv:0912.1090](#)] [[INSPIRE](#)].
- [13] M.H. Christensen, J. Hartong, N.A. Obers and B. Rollier, *Torsional Newton-Cartan Geometry and Lifshitz Holography*, *Phys. Rev. D* **89** (2014) 061901 [[arXiv:1311.4794](#)] [[INSPIRE](#)].
- [14] M.H. Christensen, J. Hartong, N.A. Obers and B. Rollier, *Boundary Stress-Energy Tensor and Newton-Cartan Geometry in Lifshitz Holography*, *JHEP* **01** (2014) 057 [[arXiv:1311.6471](#)] [[INSPIRE](#)].
- [15] J. Hartong, E. Kiritsis and N.A. Obers, *Lifshitz space-times for Schrödinger holography*, *Phys. Lett. B* **746** (2015) 318 [[arXiv:1409.1519](#)] [[INSPIRE](#)].
- [16] J. Hartong, E. Kiritsis and N.A. Obers, *Schrödinger Invariance from Lifshitz Isometries in Holography and Field Theory*, *Phys. Rev. D* **92** (2015) 066003 [[arXiv:1409.1522](#)] [[INSPIRE](#)].
- [17] J. Zaanen, Y.-W. Sun, Y. Liu and K. Schalm, *Holographic Duality in Condensed Matter Physics*, Cambridge Univ. Press (2015) [[DOI:10.1017/CB09781139942492](#)] [[INSPIRE](#)].
- [18] A. Fontanella and J.M. Nieto García, *Constructing nonrelativistic AdS_5/CFT_4 holography*, *Phys. Rev. D* **111** (2025) 026003 [[arXiv:2403.02379](#)] [[INSPIRE](#)].
- [19] A. Fontanella and J.M. Nieto García, *Nonrelativistic Holography from AdS_5/CFT_4* , *Phys. Rev. Lett.* **133** (2024) 151601 [[arXiv:2409.02267](#)] [[INSPIRE](#)].
- [20] A. Fontanella and J.M. Nieto García, *Revisiting the Symmetries of Galilean Electrodynamics*, [arXiv:2411.19217](#) [[INSPIRE](#)].
- [21] P. Hořava, *Quantum Gravity at a Lifshitz Point*, *Phys. Rev. D* **79** (2009) 084008 [[arXiv:0901.3775](#)] [[INSPIRE](#)].
- [22] J. Hartong, E. Kiritsis and N.A. Obers, *Field Theory on Newton-Cartan Backgrounds and Symmetries of the Lifshitz Vacuum*, *JHEP* **08** (2015) 006 [[arXiv:1502.00228](#)] [[INSPIRE](#)].
- [23] J. Hartong and N.A. Obers, *Hořava-Lifshitz gravity from dynamical Newton-Cartan geometry*, *JHEP* **07** (2015) 155 [[arXiv:1504.07461](#)] [[INSPIRE](#)].
- [24] M. Taylor, *Lifshitz holography*, *Class. Quant. Grav.* **33** (2016) 033001 [[arXiv:1512.03554](#)] [[INSPIRE](#)].
- [25] J. Hartong, Y. Lei and N.A. Obers, *Nonrelativistic Chern-Simons theories and three-dimensional Hořava-Lifshitz gravity*, *Phys. Rev. D* **94** (2016) 065027 [[arXiv:1604.08054](#)] [[INSPIRE](#)].
- [26] D.O. Devecioglu, N. Ozdemir, M. Ozkan and U. Zorba, *Scale invariance in Newton-Cartan and Hořava-Lifshitz gravity*, *Class. Quant. Grav.* **35** (2018) 115016 [[arXiv:1801.08726](#)] [[INSPIRE](#)].
- [27] C. Hoyos and D.T. Son, *Hall Viscosity and Electromagnetic Response*, *Phys. Rev. Lett.* **108** (2012) 066805 [[arXiv:1109.2651](#)] [[INSPIRE](#)].
- [28] D.T. Son, *Newton-Cartan Geometry and the Quantum Hall Effect*, [arXiv:1306.0638](#) [[INSPIRE](#)].
- [29] A.G. Abanov and A. Gromov, *Electromagnetic and gravitational responses of two-dimensional noninteracting electrons in a background magnetic field*, *Phys. Rev. B* **90** (2014) 014435 [[arXiv:1401.3703](#)] [[INSPIRE](#)].

- [30] M. Geracie, K. Prabhu and M.M. Roberts, *Curved non-relativistic spacetimes, Newtonian gravitation and massive matter*, *J. Math. Phys.* **56** (2015) 103505 [[arXiv:1503.02682](#)] [[INSPIRE](#)].
- [31] A. Gromov, K. Jensen and A.G. Abanov, *Boundary effective action for quantum Hall states*, *Phys. Rev. Lett.* **116** (2016) 126802 [[arXiv:1506.07171](#)] [[INSPIRE](#)].
- [32] G. Barnich and C. Troessaert, *Aspects of the BMS/CFT correspondence*, *JHEP* **05** (2010) 062 [[arXiv:1001.1541](#)] [[INSPIRE](#)].
- [33] G. Barnich, A. Gomberoff and H.A. Gonzalez, *The flat limit of three dimensional asymptotically anti-de Sitter spacetimes*, *Phys. Rev. D* **86** (2012) 024020 [[arXiv:1204.3288](#)] [[INSPIRE](#)].
- [34] A. Bagchi, *Correspondence between Asymptotically Flat Spacetimes and Nonrelativistic Conformal Field Theories*, *Phys. Rev. Lett.* **105** (2010) 171601 [[arXiv:1006.3354](#)] [[INSPIRE](#)].
- [35] G. Dautcourt, *PostNewtonian extension of the Newton-Cartan theory*, *Class. Quant. Grav.* **14** (1997) A109 [[gr-qc/9610036](#)] [[INSPIRE](#)].
- [36] D. Van den Bleeken, *Torsional Newton-Cartan gravity from the large c expansion of general relativity*, *Class. Quant. Grav.* **34** (2017) 185004 [[arXiv:1703.03459](#)] [[INSPIRE](#)].
- [37] D. Hansen, J. Hartong, N.A. Obers and G. Oling, *Galilean first-order formulation for the nonrelativistic expansion of general relativity*, *Phys. Rev. D* **104** (2021) L061501 [[arXiv:2012.01518](#)] [[INSPIRE](#)].
- [38] G. Gibbons, K. Hashimoto and P. Yi, *Tachyon condensates, Carrollian contraction of Lorentz group, and fundamental strings*, *JHEP* **09** (2002) 061 [[hep-th/0209034](#)] [[INSPIRE](#)].
- [39] A. Bagchi, *Tensionless Strings and Galilean Conformal Algebra*, *JHEP* **05** (2013) 141 [[arXiv:1303.0291](#)] [[INSPIRE](#)].
- [40] A. Bagchi, S. Chakraborty and P. Parekh, *Tensionless Strings from Worldsheet Symmetries*, *JHEP* **01** (2016) 158 [[arXiv:1507.04361](#)] [[INSPIRE](#)].
- [41] A. Bagchi, S. Chakraborty and P. Parekh, *Tensionless Superstrings: View from the Worldsheet*, *JHEP* **10** (2016) 113 [[arXiv:1606.09628](#)] [[INSPIRE](#)].
- [42] A. Bagchi, A. Banerjee, S. Chakraborty and P. Parekh, *Inhomogeneous Tensionless Superstrings*, *JHEP* **02** (2018) 065 [[arXiv:1710.03482](#)] [[INSPIRE](#)].
- [43] A. Bagchi, A. Banerjee, S. Chakraborty and P. Parekh, *Exotic Origins of Tensionless Superstrings*, *Phys. Lett. B* **801** (2020) 135139 [[arXiv:1811.10877](#)] [[INSPIRE](#)].
- [44] A. Pérez, *Asymptotic symmetries in Carrollian theories of gravity*, *JHEP* **12** (2021) 173 [[arXiv:2110.15834](#)] [[INSPIRE](#)].
- [45] A. Pérez, *Asymptotic symmetries in Carrollian theories of gravity with a negative cosmological constant*, *JHEP* **09** (2022) 044 [[arXiv:2202.08768](#)] [[INSPIRE](#)].
- [46] O. Fuentealba, M. Henneaux, P. Salgado-Rebolledo and J. Salzer, *Asymptotic structure of Carrollian limits of Einstein-Yang-Mills theory in four spacetime dimensions*, *Phys. Rev. D* **106** (2022) 104047 [[arXiv:2207.11359](#)] [[INSPIRE](#)].
- [47] C. Duval, G.W. Gibbons and P.A. Horvathy, *Conformal Carroll groups and BMS symmetry*, *Class. Quant. Grav.* **31** (2014) 092001 [[arXiv:1402.5894](#)] [[INSPIRE](#)].
- [48] J. Hartong, *Gauging the Carroll Algebra and Ultra-Relativistic Gravity*, *JHEP* **08** (2015) 069 [[arXiv:1505.05011](#)] [[INSPIRE](#)].

- [49] J. Hartong, *Holographic Reconstruction of 3D Flat Space-Time*, *JHEP* **10** (2016) 104 [[arXiv:1511.01387](#)] [[INSPIRE](#)].
- [50] A. Bagchi, R. Basu, A. Kakkar and A. Mehra, *Flat Holography: Aspects of the dual field theory*, *JHEP* **12** (2016) 147 [[arXiv:1609.06203](#)] [[INSPIRE](#)].
- [51] L. Donnay, A. Fiorucci, Y. Herfray and R. Ruzziconi, *Carrollian Perspective on Celestial Holography*, *Phys. Rev. Lett.* **129** (2022) 071602 [[arXiv:2202.04702](#)] [[INSPIRE](#)].
- [52] A. Saha, *Intrinsic approach to 1 + 1D Carrollian Conformal Field Theory*, *JHEP* **12** (2022) 133 [[arXiv:2207.11684](#)] [[INSPIRE](#)].
- [53] A. Saha, *Carrollian approach to 1 + 3D flat holography*, *JHEP* **06** (2023) 051 [[arXiv:2304.02696](#)] [[INSPIRE](#)].
- [54] A. Saha, *$w_{1+\infty}$ and Carrollian holography*, *JHEP* **05** (2024) 145 [[arXiv:2308.03673](#)] [[INSPIRE](#)].
- [55] L. Donnay and C. Marteau, *Carrollian Physics at the Black Hole Horizon*, *Class. Quant. Grav.* **36** (2019) 165002 [[arXiv:1903.09654](#)] [[INSPIRE](#)].
- [56] L. Ciambelli, R.G. Leigh, C. Marteau and P.M. Petropoulos, *Carroll Structures, Null Geometry and Conformal Isometries*, *Phys. Rev. D* **100** (2019) 046010 [[arXiv:1905.02221](#)] [[INSPIRE](#)].
- [57] D. Grumiller et al., *Spacetime structure near generic horizons and soft hair*, *Phys. Rev. Lett.* **124** (2020) 041601 [[arXiv:1908.09833](#)] [[INSPIRE](#)].
- [58] J. de Boer et al., *Carroll stories*, *JHEP* **09** (2023) 148 [[arXiv:2307.06827](#)] [[INSPIRE](#)].
- [59] F. Ecker et al., *Carroll black holes*, *SciPost Phys.* **15** (2023) 245 [[arXiv:2308.10947](#)] [[INSPIRE](#)].
- [60] M. Banados, C. Teitelboim and J. Zanelli, *The black hole in three-dimensional space-time*, *Phys. Rev. Lett.* **69** (1992) 1849 [[hep-th/9204099](#)] [[INSPIRE](#)].
- [61] A. Achucarro and P.K. Townsend, *A Chern-Simons Action for Three-Dimensional anti-De Sitter Supergravity Theories*, *Phys. Lett. B* **180** (1986) 89 [[INSPIRE](#)].
- [62] E. Witten, *(2+1)-Dimensional Gravity as an Exactly Soluble System*, *Nucl. Phys. B* **311** (1988) 46 [[INSPIRE](#)].
- [63] J. Zanelli, *Lecture notes on Chern-Simons (super-)gravities. Second edition (February 2008)*, in the proceedings of the *7th Mexican Workshop on Particles and Fields*, Merida, Mexico, November 10–17 (1999) [[hep-th/0502193](#)] [[INSPIRE](#)].
- [64] J. Matulich, S. Prohazka and J. Salzer, *Limits of three-dimensional gravity and metric kinematical Lie algebras in any dimension*, *JHEP* **07** (2019) 118 [[arXiv:1903.09165](#)] [[INSPIRE](#)].
- [65] E. Bergshoeff, D. Grumiller, S. Prohazka and J. Rosseel, *Three-dimensional Spin-3 Theories Based on General Kinematical Algebras*, *JHEP* **01** (2017) 114 [[arXiv:1612.02277](#)] [[INSPIRE](#)].
- [66] P. Concha, C. Henríquez-Báez and E. Rodríguez, *Non-relativistic and ultra-relativistic expansions of three-dimensional spin-3 gravity theories*, *JHEP* **10** (2022) 155 [[arXiv:2208.01013](#)] [[INSPIRE](#)].
- [67] J.-R. Derome and J.-G. Dubois, *Hooke's symmetries and nonrelativistic cosmological kinematics. — I*, *Nuovo Cim. B Series 11* **9** (1972) 351.
- [68] J. Gomis, D. Hidalgo and P. Salgado-Rebolledo, *Non-relativistic and Carrollian limits of Jackiw-Teitelboim gravity*, *JHEP* **05** (2021) 162 [[arXiv:2011.15053](#)] [[INSPIRE](#)].
- [69] L. Ravera and U. Zorba, *Carrollian and non-relativistic Jackiw-Teitelboim supergravity*, *Eur. Phys. J. C* **83** (2023) 107 [[arXiv:2204.09643](#)] [[INSPIRE](#)].

- [70] R. Aldrovandi, A.L. Barbosa, L.C.B. Crispino and J.G. Pereira, *Non-Relativistic spacetimes with cosmological constant*, *Class. Quant. Grav.* **16** (1999) 495 [[gr-qc/9801100](#)] [[INSPIRE](#)].
- [71] G.W. Gibbons and C.E. Patricot, *Newton-Hooke space-times, Hpp waves and the cosmological constant*, *Class. Quant. Grav.* **20** (2003) 5225 [[hep-th/0308200](#)] [[INSPIRE](#)].
- [72] J. Bruges, J. Gomis and K. Kamimura, *Newton-Hooke algebras, non-relativistic branes and generalized pp-wave metrics*, *Phys. Rev. D* **73** (2006) 085011 [[hep-th/0603023](#)] [[INSPIRE](#)].
- [73] P.D. Alvarez, J. Gomis, K. Kamimura and M.S. Plyushchay, *(2+1)D Exotic Newton-Hooke Symmetry, Duality and Projective Phase*, *Annals Phys.* **322** (2007) 1556 [[hep-th/0702014](#)] [[INSPIRE](#)].
- [74] G. Papageorgiou and B.J. Schroers, *Galilean quantum gravity with cosmological constant and the extended q-Heisenberg algebra*, *JHEP* **11** (2010) 020 [[arXiv:1008.0279](#)] [[INSPIRE](#)].
- [75] C. Duval and P. Horvathy, *Conformal Galilei groups, Veronese curves, and Newton-Hooke spacetimes*, *J. Phys. A* **44** (2011) 335203 [[arXiv:1104.1502](#)] [[INSPIRE](#)].
- [76] C. Duval, G. Gibbons and P. Horvathy, *Conformal and projective symmetries in Newtonian cosmology*, *J. Geom. Phys.* **112** (2017) 197 [[arXiv:1605.00231](#)] [[INSPIRE](#)].
- [77] G. Papageorgiou and B.J. Schroers, *A Chern-Simons approach to Galilean quantum gravity in 2+1 dimensions*, *JHEP* **11** (2009) 009 [[arXiv:0907.2880](#)] [[INSPIRE](#)].
- [78] E.A. Bergshoeff and J. Rosseel, *Three-Dimensional Extended Bargmann Supergravity*, *Phys. Rev. Lett.* **116** (2016) 251601 [[arXiv:1604.08042](#)] [[INSPIRE](#)].
- [79] M. Hatsuda and M. Sakaguchi, *Wess-Zumino term for the AdS superstring and generalized Inonu-Wigner contraction*, *Prog. Theor. Phys.* **109** (2003) 853 [[hep-th/0106114](#)] [[INSPIRE](#)].
- [80] J.A. de Azcárraga, J.M. Izquierdo, M. Picon and O. Varela, *Generating Lie and gauge free differential (super)algebras by expanding Maurer-Cartan forms and Chern-Simons supergravity*, *Nucl. Phys. B* **662** (2003) 185 [[hep-th/0212347](#)] [[INSPIRE](#)].
- [81] F. Izaurieta, E. Rodríguez and P. Salgado, *Expanding Lie (super)algebras through Abelian semigroups*, *J. Math. Phys.* **47** (2006) 123512 [[hep-th/0606215](#)] [[INSPIRE](#)].
- [82] J.A. de Azcárraga, J.M. Izquierdo, M. Picon and O. Varela, *Expansions of algebras and superalgebras and some applications*, *Int. J. Theor. Phys.* **46** (2007) 2738 [[hep-th/0703017](#)] [[INSPIRE](#)].
- [83] P. Concha and E. Rodríguez, *Non-Relativistic Gravity Theory based on an Enlargement of the Extended Bargmann Algebra*, *JHEP* **07** (2019) 085 [[arXiv:1906.00086](#)] [[INSPIRE](#)].
- [84] D.M. Peñafiel and P. Salgado-Rebolledo, *Non-relativistic symmetries in three space-time dimensions and the Nappi-Witten algebra*, *Phys. Lett. B* **798** (2019) 135005 [[arXiv:1906.02161](#)] [[INSPIRE](#)].
- [85] J. Gomis, A. Kleinschmidt, J. Palmkvist and P. Salgado-Rebolledo, *Newton-Hooke/Carrollian expansions of (A)dS and Chern-Simons gravity*, *JHEP* **02** (2020) 009 [[arXiv:1912.07564](#)] [[INSPIRE](#)].
- [86] E. Bergshoeff, J. Gomis and P. Salgado-Rebolledo, *Non-relativistic limits and three-dimensional coadjoint Poincaré gravity*, *Proc. Roy. Soc. Lond. A* **476** (2020) 20200106 [[arXiv:2001.11790](#)] [[INSPIRE](#)].
- [87] R. Caroca, D.M. Peñafiel and P. Salgado-Rebolledo, *Nonrelativistic spin-3 symmetries in 2+1 dimensions from expanded and extended Nappi-Witten algebras*, *Phys. Rev. D* **107** (2023) 064034 [[arXiv:2208.00602](#)] [[INSPIRE](#)].

- [88] P. Concha, E. Rodríguez, G. Rubio and P. Yañez, *Three-dimensional Newtonian gravity with cosmological constant and torsion*, *Eur. Phys. J. C* **83** (2023) 47 [[arXiv:2204.11763](#)] [[INSPIRE](#)].
- [89] P. Concha, E. Rodríguez and G. Rubio, *Non-relativistic gravity theories in four spacetime dimensions*, *JHEP* **02** (2023) 191 [[arXiv:2210.04101](#)] [[INSPIRE](#)].
- [90] P. Concha, D. Pino, L. Ravera and E. Rodríguez, *Extended kinematical 3D gravity theories*, *JHEP* **01** (2024) 040 [[arXiv:2310.01335](#)] [[INSPIRE](#)].
- [91] E.A. Bergshoeff and J. Rosseel, *Non-Lorentzian Supergravity*, in *Handbook of Quantum Gravity*, C. Bambi et al., Sperling Singapore (2023) [[DOI:10.1007/978-981-19-3079-9_52-1](#)] [[arXiv:2211.02604](#)] [[INSPIRE](#)].
- [92] R. Andringa, E.A. Bergshoeff, J. Rosseel and E. Sezgin, *3D Newton-Cartan supergravity*, *Class. Quant. Grav.* **30** (2013) 205005 [[arXiv:1305.6737](#)] [[INSPIRE](#)].
- [93] E. Bergshoeff, J. Rosseel and T. Zojer, *Newton-Cartan (super)gravity as a non-relativistic limit*, *Class. Quant. Grav.* **32** (2015) 205003 [[arXiv:1505.02095](#)] [[INSPIRE](#)].
- [94] E. Bergshoeff, J. Rosseel and T. Zojer, *Newton-Cartan supergravity with torsion and Schrödinger supergravity*, *JHEP* **11** (2015) 180 [[arXiv:1509.04527](#)] [[INSPIRE](#)].
- [95] N. Ozdemir, M. Ozkan, O. Tunca and U. Zorba, *Three-Dimensional Extended Newtonian (Super)Gravity*, *JHEP* **05** (2019) 130 [[arXiv:1903.09377](#)] [[INSPIRE](#)].
- [96] P. Concha, L. Ravera and E. Rodríguez, *Three-dimensional Maxwellian extended Bargmann supergravity*, *JHEP* **04** (2020) 051 [[arXiv:1912.09477](#)] [[INSPIRE](#)].
- [97] N. Ozdemir, M. Ozkan and U. Zorba, *Three-dimensional extended Lifshitz, Schrödinger and Newton-Hooke supergravity*, *JHEP* **11** (2019) 052 [[arXiv:1909.10745](#)] [[INSPIRE](#)].
- [98] L. Ravera, *AdS Carroll Chern-Simons supergravity in 2 + 1 dimensions and its flat limit*, *Phys. Lett. B* **795** (2019) 331 [[arXiv:1905.00766](#)] [[INSPIRE](#)].
- [99] F. Ali and L. Ravera, *$\mathcal{N}$ -extended Chern-Simons Carrollian supergravities in 2 + 1 spacetime dimensions*, *JHEP* **02** (2020) 128 [[arXiv:1912.04172](#)] [[INSPIRE](#)].
- [100] J.A. de Azcárraga, D. Gútiez and J.M. Izquierdo, *Extended $D = 3$ Bargmann supergravity from a Lie algebra expansion*, *Nucl. Phys. B* **946** (2019) 114706 [[arXiv:1904.12786](#)] [[INSPIRE](#)].
- [101] P. Concha, L. Ravera and E. Rodríguez, *Three-dimensional non-relativistic extended supergravity with cosmological constant*, *Eur. Phys. J. C* **80** (2020) 1105 [[arXiv:2008.08655](#)] [[INSPIRE](#)].
- [102] P. Concha, M. Ipinza, L. Ravera and E. Rodríguez, *Non-relativistic three-dimensional supergravity theories and semigroup expansion method*, *JHEP* **02** (2021) 094 [[arXiv:2010.01216](#)] [[INSPIRE](#)].
- [103] P. Concha, L. Ravera and E. Rodríguez, *Three-dimensional exotic Newtonian supergravity theory with cosmological constant*, *Eur. Phys. J. C* **81** (2021) 646 [[arXiv:2104.12908](#)] [[INSPIRE](#)].
- [104] P. Concha, L. Ravera and E. Rodríguez, *Three-dimensional non-relativistic supergravity and torsion*, *Eur. Phys. J. C* **82** (2022) 220 [[arXiv:2112.05902](#)] [[INSPIRE](#)].
- [105] P. Concha, E. Rodríguez and S. Salgado, *Three-dimensional non-relativistic Hietarinta supergravity*, *Eur. Phys. J. C* **85** (2025) 47 [[arXiv:2409.01298](#)] [[INSPIRE](#)].
- [106] E.A. Bergshoeff et al., *Non-relativistic ten-dimensional minimal supergravity*, *JHEP* **12** (2021) 123 [[arXiv:2107.14636](#)] [[INSPIRE](#)].

- [107] E. Bergshoeff, C. Blair, J. Lahnsteiner and J. Rosseel, *A Consistent Limit of 11D Supergravity*, *PoS CORFU2022* (2023) 162 [INSPIRE].
- [108] E.A. Bergshoeff, C.D.A. Blair, J. Lahnsteiner and J. Rosseel, *The surprising structure of non-relativistic 11-dimensional supergravity*, *JHEP* **12** (2024) 010 [arXiv:2407.21648] [INSPIRE].
- [109] P.S. Howe, J.M. Izquierdo, G. Papadopoulos and P.K. Townsend, *New supergravities with central charges and Killing spinors in (2+1)-dimensions*, *Nucl. Phys. B* **467** (1996) 183 [hep-th/9505032] [INSPIRE].
- [110] E. Bergshoeff, J. Gomis and L. Parra, *The symmetries of the Carroll Superparticle*, *J. Phys. A* **49** (2016) 185402 [arXiv:1503.06083] [INSPIRE].
- [111] J. Gomis, A. Kleinschmidt, J. Palmkvist and P. Salgado-Rebolledo, *Symmetries of post-Galilean expansions*, *Phys. Rev. Lett.* **124** (2020) 081602 [arXiv:1910.13560] [INSPIRE].
- [112] P. Concha, L. Ravera and E. Rodríguez, *Three-dimensional exotic Newtonian gravity with cosmological constant*, *Phys. Lett. B* **804** (2020) 135392 [arXiv:1912.02836] [INSPIRE].
- [113] D. Hansen, J. Hartong and N.A. Obers, *Gravity between Newton and Einstein*, *Int. J. Mod. Phys. D* **28** (2019) 1944010 [arXiv:1904.05706] [INSPIRE].
- [114] E. Bergshoeff et al., *Carroll versus Galilei Gravity*, *JHEP* **03** (2017) 165 [arXiv:1701.06156] [INSPIRE].
- [115] J. Gomis, A. Kleinschmidt and J. Palmkvist, *Galilean free Lie algebras*, *JHEP* **09** (2019) 109 [arXiv:1907.00410] [INSPIRE].
- [116] S.W. MacDowell and F. Mansouri, *Unified Geometric Theory of Gravity and Supergravity*, *Phys. Rev. Lett.* **38** (1977) 739 [Erratum *ibid.* **38** (1977) 1376] [INSPIRE].
- [117] D. Lovelock, *The Einstein tensor and its generalizations*, *J. Math. Phys.* **12** (1971) 498 [INSPIRE].
- [118] D. Lovelock, *The four-dimensionality of space and the einstein tensor*, *J. Math. Phys.* **13** (1972) 874 [INSPIRE].
- [119] M. Brigante et al., *The Viscosity Bound and Causality Violation*, *Phys. Rev. Lett.* **100** (2008) 191601 [arXiv:0802.3318] [INSPIRE].
- [120] X.O. Camanho, J.D. Edelstein and J.M. Sánchez De Santos, *Lovelock theory and the AdS/CFT correspondence*, *Gen. Rel. Grav.* **46** (2014) 1637 [arXiv:1309.6483] [INSPIRE].
- [121] L. Andrianopoli and R. D'Auria, *$N = 1$ and $N = 2$ pure supergravities on a manifold with boundary*, *JHEP* **08** (2014) 012 [arXiv:1405.2010] [INSPIRE].
- [122] P. Concha, L. Ravera and E. Rodríguez, *On the supersymmetry invariance of flat supergravity with boundary*, *JHEP* **01** (2019) 192 [arXiv:1809.07871] [INSPIRE].
- [123] A. Banaudi and L. Ravera, *Generalized AdS-Lorentz deformed supergravity on a manifold with boundary*, *Eur. Phys. J. Plus* **133** (2018) 514 [arXiv:1803.08738] [INSPIRE].
- [124] K. Eder and H. Sahlmann, *Holst-MacDowell-Mansouri action for (extended) supergravity with boundaries and super Chern-Simons theory*, *JHEP* **07** (2021) 071 [arXiv:2104.02011] [INSPIRE].
- [125] L. Andrianopoli and L. Ravera, *On the Geometric Approach to the Boundary Problem in Supergravity*, *Universe* **7** (2021) 463 [arXiv:2111.01462] [INSPIRE].

- [126] L. Ciambelli et al., *Flat holography and Carrollian fluids*, *JHEP* **07** (2018) 165 [[arXiv:1802.06809](#)] [[INSPIRE](#)].
- [127] A. Bagchi and I. Mandal, *Supersymmetric Extension of Galilean Conformal Algebras*, *Phys. Rev. D* **80** (2009) 086011 [[arXiv:0905.0580](#)] [[INSPIRE](#)].
- [128] I. Mandal, *Supersymmetric Extension of GCA in 2d*, *JHEP* **11** (2010) 018 [[arXiv:1003.0209](#)] [[INSPIRE](#)].
- [129] C. Aragone and S. Deser, *Hypersymmetry in $D = 3$ of Coupled Gravity Massless Spin $5/2$ System*, *Class. Quant. Grav.* **1** (1984) L9 [[INSPIRE](#)].
- [130] Y.M. Zinoviev, *Hypergravity in AdS_3* , *Phys. Lett. B* **739** (2014) 106 [[arXiv:1408.2912](#)] [[INSPIRE](#)].
- [131] M. Henneaux, A. Pérez, D. Tempo and R. Troncoso, *Extended $\bar{d}e$ Sitter hypergravity in $2+1$ dimensions and hypersymmetry bounds*, in the proceedings of the *International Workshop on Higher Spin Gauge Theories*, Singapore, Singapore, November 4–6 (2015) [[DOI:10.1142/9789813144101_0009](#)] [[arXiv:1512.08603](#)] [[INSPIRE](#)].
- [132] M. Henneaux, A. Pérez, D. Tempo and R. Troncoso, *Hypersymmetry bounds and three-dimensional higher-spin black holes*, *JHEP* **08** (2015) 021 [[arXiv:1506.01847](#)] [[INSPIRE](#)].
- [133] O. Fuentealba, J. Matulich and R. Troncoso, *Extension of the Poincaré group with half-integer spin generators: hypergravity and beyond*, *JHEP* **09** (2015) 003 [[arXiv:1505.06173](#)] [[INSPIRE](#)].
- [134] O. Fuentealba, J. Matulich and R. Troncoso, *Asymptotically flat structure of hypergravity in three spacetime dimensions*, *JHEP* **10** (2015) 009 [[arXiv:1508.04663](#)] [[INSPIRE](#)].
- [135] R. Schrader, *The maxwell group and the quantum theory of particles in classical homogeneous electromagnetic fields*, *Fortsch. Phys.* **20** (1972) 701 [[INSPIRE](#)].
- [136] H. Bacry, P. Combe and J.L. Richard, *Group-theoretical analysis of elementary particles in an external electromagnetic field. II. the nonrelativistic particle in a constant and uniform field*, *Nuovo Cim. A* **70** (1970) 289 [[INSPIRE](#)].
- [137] J. Gomis and A. Kleinschmidt, *On free Lie algebras and particles in electro-magnetic fields*, *JHEP* **07** (2017) 085 [[arXiv:1705.05854](#)] [[INSPIRE](#)].
- [138] P. Salgado, R.J. Szabo and O. Valdivia, *Topological gravity and transgression holography*, *Phys. Rev. D* **89** (2014) 084077 [[arXiv:1401.3653](#)] [[INSPIRE](#)].
- [139] S. Hoseinzadeh and A. Rezaei-Aghdam, *$(2+1)$ -dimensional gravity from Maxwell and semisimple extension of the Poincaré gauge symmetric models*, *Phys. Rev. D* **90** (2014) 084008 [[arXiv:1402.0320](#)] [[INSPIRE](#)].
- [140] J.A. de Azcárraga, K. Kamimura and J. Lukierski, *Generalized cosmological term from Maxwell symmetries*, *Phys. Rev. D* **83** (2011) 124036 [[arXiv:1012.4402](#)] [[INSPIRE](#)].
- [141] R. Durka, J. Kowalski-Glikman and M. Szczachor, *Gauged AdS -Maxwell algebra and gravity*, *Mod. Phys. Lett. A* **26** (2011) 2689 [[arXiv:1107.4728](#)] [[INSPIRE](#)].
- [142] P.K. Concha and E.K. Rodríguez, *Maxwell Superalgebras and Abelian Semigroup Expansion*, *Nucl. Phys. B* **886** (2014) 1128 [[arXiv:1405.1334](#)] [[INSPIRE](#)].
- [143] P.K. Concha and E.K. Rodríguez, *$N = 1$ Supergravity and Maxwell superalgebras*, *JHEP* **09** (2014) 090 [[arXiv:1407.4635](#)] [[INSPIRE](#)].
- [144] D.M. Peñañiel and L. Ravera, *On the Hidden Maxwell Superalgebra underlying $D = 4$ Supergravity*, *Fortsch. Phys.* **65** (2017) 1700005 [[arXiv:1701.04234](#)] [[INSPIRE](#)].

- [145] L. Ravera, *Hidden role of Maxwell superalgebras in the free differential algebras of $D = 4$ and $D = 11$ supergravity*, *Eur. Phys. J. C* **78** (2018) 211 [[arXiv:1801.08860](#)] [[INSPIRE](#)].
- [146] P. Concha et al., *Asymptotic symmetries of three-dimensional Chern-Simons gravity for the Maxwell algebra*, *JHEP* **10** (2018) 079 [[arXiv:1805.08834](#)] [[INSPIRE](#)].
- [147] P. Concha, D.M. Peñafiel and E. Rodríguez, *On the Maxwell supergravity and flat limit in 2+1 dimensions*, *Phys. Lett. B* **785** (2018) 247 [[arXiv:1807.00194](#)] [[INSPIRE](#)].
- [148] D. Chernyavsky, N.S. Deger and D. Sorokin, *Spontaneously broken 3d Hietarinta/Maxwell Chern-Simons theory and minimal massive gravity*, *Eur. Phys. J. C* **80** (2020) 556 [[arXiv:2002.07592](#)] [[INSPIRE](#)].
- [149] S. Kibaroğlu and O. Cebecioğlu, *Gauge theory of the Maxwell and semi-simple extended (anti) de Sitter algebra*, *Int. J. Mod. Phys. D* **30** (2021) 2150075 [[arXiv:2007.14795](#)] [[INSPIRE](#)].
- [150] R. Caroca et al., *Hypersymmetric extensions of Maxwell-Chern-Simons gravity in 2+1 dimensions*, *Phys. Rev. D* **104** (2021) 064011 [[arXiv:2105.12243](#)] [[INSPIRE](#)].
- [151] J. Matulich and E. Rodríguez, *Enlarged super- $\mathfrak{bms}_3$ algebra and its flat limit*, *Phys. Rev. D* **110** (2024) 064064 [[arXiv:2310.16614](#)] [[INSPIRE](#)].
- [152] P. Concha, D. Peñafiel, L. Ravera and E. Rodríguez, *Three-dimensional Maxwellian Carroll gravity theory and the cosmological constant*, *Phys. Lett. B* **823** (2021) 136735 [[arXiv:2107.05716](#)] [[INSPIRE](#)].