

AN IN SILICO-IN VIVO FRAMEWORK FOR THE ACUTE OCULAR AND CARDIOVASCULAR  
RESPONSE TO 6° HEAD-DOWN TILT

*Original*

AN IN SILICO-IN VIVO FRAMEWORK FOR THE ACUTE OCULAR AND CARDIOVASCULAR RESPONSE TO 6° HEAD-DOWN TILT / Fois, Matteo; Artiles, Ana-Diaz; Zaman, Syeda Yasmin; Ridolfi, Luca; Scarsoglio, Stefania. - ELETTRONICO. - (2023). (Intervento presentato al convegno XII Annual Meeting of the Italian Chapter of the European Society of Biomechanics tenutosi a Turin nel 18-19 Settembre 2023).

*Availability:*

This version is available at: 11583/2982505 since: 2023-09-27T08:03:52Z

*Publisher:*

ESB-ITA

*Published*

DOI:

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

# AN *IN SILICO-*IN VIVO** FRAMEWORK FOR THE ACUTE OCULAR AND CARDIOVASCULAR RESPONSE TO 6° HEAD-DOWN TILT

Matteo Fois (1), Ana Diaz-Artilles (2), Syeda Yasmin Zaman (2),  
Luca Ridolfi (3) and Stefania Scarsoglio (1)

1. Department of Mechanical and Aerospace Engineering, Politecnico di Torino, Turin, Italy; 2. Department of Aerospace Engineering, Texas A&M University, College Station, Texas (USA); 3. Department of Environmental, Land and Infrastructure Engineering, Politecnico di Torino, Turin, Italy

## Introduction

Ensuring the healthiness of astronauts undergoing long-term missions in space is of primary importance for the aerospace community. Spaceflight associated neuro-ocular syndrome (SANS) has been widely acknowledged to cause severe ocular disorders in astronauts returning from long permanence in weightlessness [1].

In this context, head-down tilt (HDT) has gained large popularity to resemble the cardiovascular response to microgravity, as well as to study SANS [2].

We propose a novel multiscale numerical framework to simulate the acute response to 6° HDBR – with *in vivo* validation – to help shed light on SANS onset.

## Methods

We developed a new cardiovascular model integrating our previously validated *in silico* framework [3] with a lumped model of the eye [4] and of the cerebrovascular circulation [5]. The global model presents a 1D description of the arterial tree combined with 0D analogues of the remaining vasculature, accounting for short-term homeostatic control and for the action of gravity during posture changes. Using this model, we simulated a tilt maneuver between 80° head-up tilt (HUT) and 6° HDT.

*In vivo* measures were taken at the Bioastronautics and Human Performance (BHP) Laboratory at Texas A&M University (TX, USA). Six healthy male subjects were positioned upright (80°) on an inversion table, tilted down to -6° HDT for approximately 10 minutes and eventually tilted back upright to 80°. Subjects' arterial pressure, heart rate, cardiac output, stroke volume, and intraocular pressure measures were collected at each position. Seated baseline measures of the same parameters were also acquired.

## Results

The model results in response to acute 6° HDT – and tilt back to 80° – are shown in Figure 1. The corresponding *in vivo* mean and standard deviation ( $\mu \pm \sigma$ ) of subjects' parameters are also depicted in Figure 1.

## Discussion

The model results reproduce the *in vivo* acute ocular and cardiovascular response to 6°HDT. In addition, the model offers novel insights into the hemodynamic mechanisms related to intraocular pressure and




Figure 1: Model results and experimental data for finger mean arterial pressure at brachial level (fMAP, panel a), heart rate (HR, panel b), cardiac output (CO, panel c), stroke volume (SV, panel d) and intraocular pressure (IOP, panel e). ICP (panel f) is the simulated intracranial pressure. S: baseline seated.

intracranial pressure – in terms of average and pulsatile values – both at steady-state and during the transient dynamics.

As also observed upon entering microgravity, the initial fluid shift associated with 6° HDT leads to a sudden increase in SV and CO accompanied by a drop in HR. While arterial pressure does not show wide variations, IOP rises by 4.2 mmHg (model) compared to 80° upright ( $21.0 \pm 3.8$  at 6° HDT vs.  $18.0 \pm 3.2$  at 80° HUT). In addition, the model simulates an ICP increase of 11.3 mmHg in response to 6° HDT [6-7] which is much larger than the observed IOP increase. As a result, our model predicts that the transmural pressure IOP-ICP is markedly reduced during acute HDT (-43%) [6-8]. These results may contribute to the understanding of SANS among astronauts experiencing analog ocular and vascular responses upon long-term space missions.

## References

1. Mader et al, Ophthalmology, 118.10: 2058-2069, 2011.
2. Ong et al, Front Neurol, 12:648958, 2021.
3. Fois et al, Front Physiol, 13:826989, 2022.
4. Petersen et al, J Appl Physiol, 132(1):24-35, 2022.
5. Ursino & Giannessi, Ann Biomed Eng, 38:955-974, 2010
6. Lawley et al, J Physiol, 595: 2115-2127, 2017.
7. Laurie et al, Physiol Rep, 5(11):e13302, 2017
8. Zhang and Hargens, Physiol Rev, 98(1):59-87, 2017

