POLITECNICO DI TORINO Repository ISTITUZIONALE

Real life clinical outcomes of relapsed/refractory diffuse large B cell lymphoma in the rituximab era: The STRIDER study

Original

Real life clinical outcomes of relapsed/refractory diffuse large B cell lymphoma in the rituximab era: The STRIDER study / Dogliotti, Irene; Peri, Veronica; Clerico, Michele; Vassallo, Francesco; Musto, Davide; Mercadante, Silvio; Ragaini, Simone; Botto, Barbara; Levis, Mario; Novo, Mattia; Ghislieri, Marco; Molinaro, Luca; Mortara, Umberto; Consoli, Chiara; Lonardo, Alessio; Bondielli, Giulia; Ferrero, Simone; Freilone, Roberto; Ricardi, Umberto; Bruno, Benedetto; Cavallo, Federica. - In: CANCER MEDICINE. - ISSN 2045-7634. - ELETTRONICO. - 13:14(2024), pp. 1-15. [10.1002/cam4.7448]

Availability:

This version is available at: 11583/2991100 since: 2024-07-22T13:15:42Z

Publisher: Wiley

Published DOI:10.1002/cam4.7448

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

RESEARCH ARTICLE

Real life clinical outcomes of relapsed/refractory diffuse large B cell lymphoma in the rituximab era: The STRIDER study

Irene Dogliotti ¹ Veronica Peri ^{1,2} Michele Clerico ¹ Francesco Vassallo ³
Davide Musto ² Silvio Mercadante ² Simone Ragaini ^{1,2} Barbara Botto ⁴
Mario Levis ⁵ Mattia Novo ⁴ Marco Ghislieri ^{6,7} Luca Molinaro ⁸
Umberto Mortara ⁸ Chiara Consoli ^{1,2} Alessio Lonardo ² Giulia Bondielli ²
Simone Ferrero ^{1,2} Roberto Freilone ⁴ Umberto Ricardi ⁵ Benedetto Bruno ^{1,2} []
Federica Cavallo ^{1,2}

¹Division of Hematology, University Hospital A.O.U. "Città della Salute e della Scienza", Turin, Italy

²Division of Hematology U, Department of Molecular Biotechnologies and Health Sciences, University of Turin, Turin, Italy

³Division of Hematology, Ospedale Santa Croce e Carle, Cuneo, Italy

⁴Division of Hematology, AOU "Città della Salute e della Scienza di Torino", Turin, Italy

⁵Division of Radiotherapy, Department of Oncology., University Hospital A.O.U. "Città della Salute e della Scienza", Turin, Italy

⁶Department of Electronics and Telecommunications, Politecnico di Torino, Turin, Italy

⁷PolitoBIOMed Lab, Politecnico di Torino, Turin, Italy

⁸Division of Pathology, AOU "Città della Salute e della Scienza di Torino", Turin, Italy

Correspondence

Federica Cavallo, Division of Hematology U, Department of Molecular Biotechnologies and Health Sciences, University of Turin, University Hospital A.O.U. "Città della Salute e della Scienza", Via Genova 3, Turin 10100, Italy. Email: f.cavallo@unito.it

Abstract

Background: Relapse and refractory (R/R) rates after first-line R-CHOP in diffuse large B cell lymphomas (DLBCL) are ~40% and ~15% respectively.

Aims: We conducted a retrospective real-world analysis aimed at evaluating clinical outcomes of R/R DLBCL patients.

Material and Methods: Overall, 403 consecutive DLBCL patients treated in two large hematological centers in Torino, Italy were reviewed.

Results: At a median follow up of 50 months, 5-year overall survival from diagnosis (OS-1) was 66.5%, and 2-year progression free survival (PFS-1) was 68%. 134 (34.4%) patients relapsed (n=46, 11.8%) or were refractory (n=88, 22.6%) to R-CHOP. Most employed salvage treatments included platinum salt-based regimens in 38/134 (28.4%), lenalidomide in 14 (10.4%). Median OS and PFS after disease relapse or progression (OS-2 and PFS-2) were 6.7 and 5.1 months respectively. No significant difference in overall response rate, OS-2 or PFS-2 in patients treated with platinum-based regimens versus other regimens was observed. By multivariate

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). Cancer Medicine published by John Wiley & Sons Ltd.

analysis, age between 60 and 80 years, germinal center B cell type cell of origin and extranodal involvement of <2 sites were associated with better OS-2.

Discussion: Our findings confirm very poor outcomes of R/R DLBCL in the rituximab era. Widespread approval by national Medicine Agencies of novel treatments such as CAR-T cells and bispecific antibodies as second-line is eagerly awaited to improve these outcomes.

K E Y W O R D S

chemotherapy, diffuse large B cell lymphoma, real world, refractory disease, relapse, rituximab, transplantation

1 | INTRODUCTION

Clinical outcomes of patients with diffuse large B cell lymphoma (DLBCL) remain suboptimal with relapse and chemo-refractoriness rates up to 40%-50% and 10%-15% respectively.^{1,2} In the relapsed/refractory (R/R) population, disease-specific features such as MYC and BCL2/ BCL6 translocations (i.e., double or triple hit lymphoma) or protein expression (i.e., double/triple expressors), and unfavorable gene expression signatures are associated with very poor outcomes.³⁻⁸ The "classical" firstline treatment backbone, containing an anthracycline and an anti CD20 antibody (commonly "R-CHOP": rituximab, cyclophosphamide, vincristine, doxorubicin, prednisone),⁹⁻¹¹ has recently been challenged by the introduction of novel agents such as first-line polatuzumab vedotin,¹² while salvage treatment in the R/R setting has not yet been standardized, particularly for elderly patients.¹ Even though second-line combinations with platinum salts and cytarabine, or etoposide, and rituximab (i.e., R-DHAP: rituximab, cytarabine, cisplatin and dexamethasone; R-OxDHA: rituximab, oxaliplatin, cytarabine and dexamethasone; R-ICE: rituximab, ifosfamide, etoposide, carboplatin; R-GDP: rituximab, gemcitabine, dexamethasone and cisplatin), followed by autologous stem cell transplantation (ASCT) as consolidation, are commonly employed for medically fit patients,¹³⁻¹⁶ population-based studies report that over half of R/R patients undergo mainly palliation.¹⁷ For subsequent relapses, approved agents in Italy include single agent pixantrone and lenalidomide, with a median OS ranging between 8 and 10 months.^{18,19} Allogeneic stem cell transplantation (allo-SCT) is used only in selected cases with good response to re-induction therapy, with 4-year overall survival (OS) around 20%; its feasibility, however, is limited by patient age, comorbidities, and risk of treatmentrelated toxicities.^{20,21}

Recently, several novel strategies have been developed. However, their approval relies on national Medicine Agencies the policies of which differ significantly from country to country. The U.S. Food and Drug Administration (FDA) approved chimeric antigen receptor (CAR) T cell therapies axicabtagene ciloleucel (axi-cel) and tisagenlecleucel (tisa-cel), in 2017 and 2018, respectively, for adult patients with high-grade B-cell lymphomas or DLBCL R/R to at least 2 therapy lines. In the ZUMA 1 and JULIET studies, overall response rate (ORR) were 82% and 52% with a complete response (CR) rate of 54% and 40% respectively.^{22,23} In the TRANSCEND study,²⁴ similar results were reported with lisocabtagene maraleucel, approved in 2021, and later extended also to refractory patients and to those relapsing within 12 months after first line treatment in June, 2022, according to the results of the TRANSFORM study.²⁵ In long term safety and efficacy analyses median OS at 24 months was not reached for both axi-cel²⁶ and tisa-cel in patients who achieved CR at 3 and 6 months.²⁷ Recently, axi-cel was FDA- approved for DLBCL refractory to first-line chemo-immunotherapy based on the ZUMA-7 study.²⁸ Other recently approved agents include: tafasitamab, an Fc-enhanced, humanized, monoclonal antibody targeting CD19, used in combination with lenalidomide in adult patients not eligible for ASCT²⁹; the antibody-drug conjugate polatuzumab vedotin, a CD79bdirected antibody conjugated with monomethyl auristatin (MMAE), used in combination with bendamustine and rituximab after at least two prior therapies³⁰; loncastuximab tesirine, a CD19-directed antibody and alkylating agent conjugate³¹; and the XPO inhibitor selinexor used as single agent after two to five systemic treatment lines.³²

Here, we present a study designed to determine reallife clinical outcomes of R/R DLBCL in the rituximab era, and to possibly characterize baseline features at diagnosis that may predict poor response to first-line treatment and response to salvage therapies.

2 | MATERIALS AND METHODS

2.1 | Study design

The "STRIDER" ("strategies of treatment in diffuse large B cell lymphoma in the era of rituximab") is a retrospective, observational study designed to evaluate clinical outcomes of R/R DLBCL patients after first-line treatment in the rituximab era in a real-world setting. Between January 2010 and December 2019, patients older than 18 years, consecutively treated at 2 Tertiary Referral Centers (Division of Hematology-University of Torino, Italy, and Division of Hematology, AOU Città della Salute e della Scienza-Torino, Italy), were evaluated for enrollment. The study was proposed and discussed with the patients by the treating hematologist during follow up visits. Patients were enrolled after obtaining informed consent. The study was approved by the local Institutional Review Board (A.O.U. Città della Salute e della Scienza, Torino, Italy) and conducted according to the Declaration of Helsinki. Patient data were obtained from hospital health records and research files. All data were pseudo-anonymized by assignment of a study specific patient code.

2.2 | Inclusion criteria

Major inclusion criteria included initial biopsy-proven diagnosis of either DLBCL or high grade B cell lymphomas (HGBCL). Histological and immune-histochemical diagnosis by tru-cut core-needle biopsies was allowed, while cases diagnosed by fine needle aspiration cytology were excluded. Transformed disease after a previous diagnosis of low grade lymphoma was allowed. R/R disease was documented by biopsy, imaging studies or clinical evaluation; refractoriness to first line treatment was defined as reappearance or progression of DLBCL or HGBCL within 12 months from initial diagnosis or disease-related death (POD-12). For all patients, retrieved data included demographics, whole blood counts, basic metabolic panel; imaging studies (computed tomography, CT, and fluorodeoxyglucose-positron emission tomography, PET, if available); histology studies by lymph node biopsy, bone marrow biopsy and bone marrow aspirate; presence of B symptoms, performance status, prognostic scores [IPI, age-adjusted IPI, Central Nervous S,35ystem International Prognostic Index (CNS-IPI)]; number of therapy lines and regimens employed. Cell of origin (COO) was determined by Hans' algorithm method; bulky disease was defined as any lesion >6 cm by CT.

2.3 | Statistical analysis

Cancer Medicine

Primary endpoint was OS for R/R patients after salvage treatment (OS-2). Secondary endpoints included progression free survival after first line treatment (PFS-1), PFS after salvage treatment defined as PFS-2; OS-1; POD12; determinants for survival outcomes; distribution of second line regimens (descriptive analysis); efficacy outcomes with salvage treatments (response and duration of response). Sample size estimation was not predefined, all consecutive patients meeting the inclusion criteria during the defined study period were eligible for the study though patients with missing data were excluded from the analysis. Response to therapy was determined by the 1999 International Working Group response criteria³³ and by Lugano response criteria for patients diagnosed after 2014.³⁴ Baseline characteristics of R/R patients were compared to those of non R/R (NRR) patients to identify potential prognostic factors. Statistical analyses were carried out using R (v 4.3.1). Survival curves were plotted with Kaplan-Meier method and compared with log-rank test. Medians between groups for continuous variables were compared by the Kruskal-Wallis (for non-normal variables) or the one-way ANOVA test (for normal variables); the chi-squared test or Fisher's exact test for small study samples, were employed for categorical variables. The Cox proportional hazards model was implemented for the univariate and multivariate survival analyses. In particular, an AIC-based backward stepwise algorithm (R function stats::step) was used to perform the variable selection, from 12 initial variables (outcome after firstline, age, hemoglobin, COO, gender, ECOG performance status, stage, extranodal involvement, LDH, ki67, type of first- and second-line treatment) to 6 (age, hemoglobin, COO, extranodal involvement, ki67, type of second line treatment). This method allows to determine the most relevant covariates for the outcomes of interest and to analyze potential confounding factors for each covariate. As required by the algorithm, the dataset was restricted to the R/R patients for whom there were no missing data on the initial 12 variables (n=48). Thus, multivariate analysis for OS-2 was re-run by applying a data imputation algorithm, the "Multiple Imputation by Chained Equations" (MICE), a robust, informative method to analyze datasets with missing data. The procedure "fills in" missing data through an iterative series of predictive models. In each iteration, a specified variable is imputed using the other variables in the dataset. These iterations are run until convergence has been met. POD12 was determined by logistic regression: the model predicts the probability of being refractory/relapsed/dead due to DLBCL within 12 months and the causing factors.

3 of 15

-WILEY

3 | RESULTS

3.1 | Baseline patient characteristics

Four-hundred-three patients met the inclusion criteria, Thirteen were not included in the final analysis because of incomplete data (n=8) or early treatment discontinuation (n=5) (Figure 1). Among patients included in the final analyses, 384 had DLBCL, NOS, while 6 (2%) had HGBCL. At follow up, after first line therapy, 256/390 patients (65.6%) were still in first response, whereas 134(34.4%) had either relapsed (n = 46, 11.8%) or were refractory (n = 88, 22.6%). Patient characteristics at diagnosis are shown in Table 1. Median age was similar in the "response" and in the "relapse" groups (67.7 and 66.3 years respectively), though significantly lower as compared to the "refractory" patient group (75.0 years) (p < 0.001). High baseline IPI score (categorized as 0-1, 2-3, 4-5) (p < 0.001), advanced stage (p < 0.001), B symptoms (p < 0.001), and ≥ 2 extranodal site involvement (p=0.035) were more frequent among patients with R/R disease. At diagnosis, CNS involvement was observed in 7/390 (1.8%) patients; overall, high CNS IPI risk was present in 53 (20.7%) for the NRR

group, while in 12 (26.1%) and 40 (45.5%) for relapsed and refractory patients respectively (p < 0.001). Bulky disease was present in 71 (27.7%), 18 (39.1%) and 42 (47.7%) of NRR, relapsed and refractory patients (p < 0.001) respectively. The expression patterns of conventional immune-histochemical markers (namely CD20, BCL2, BCL6, C-MYC) in the diagnostic lymph node biopsy did not significantly differ between groups.

3.2 | First-line treatments

Most patients (374/390, 95.9%) underwent chemotherapy with a curative intent (Table 1): 315 (80.8%) were treated with standard R-CHOP/COMP (liposomal doxorubicin in case of previous cardiovascular disease), 28 (7.2%) with R-mini-CHOP/COMP, 9 (2.3%) underwent alternating R-CODOX-M (rituximab, cyclophosphamide, vincristine, doxorubicin and methotrexate) and R-IVAC (rituximab, ifosfamide, etoposide, cytarabine), 6 (1.5%) R-DA EPOCH (rituximab, etoposide, vincristine, cyclophosphamide and prednisone), 24 (6.2%) were treated with oral chemotherapy \pm rituximab. CNS prophylaxis with intrathecal methotrexate



FIGURE 1 The STRIDER study: Graphical representation of enrollment, treatment, and follow-up of 403 patients.

			Open Access			
TABLE 1 Patient characteristics and tree	eatments.					
			R/R			
				Refractory		
	All	NRR	Relapsed disease	disease		
	N (%)	N (%)	N (%)			
Total patients	390 (100.0)	256 (65.6)	134 (34.4)			
			46 (11.8)	88 (22.6)		
Baseline characteristics						
Sex						
Male	218 (55.9)	138 (53.9)	33 (71.7)	47 (53.4)		
Female	172 (44.1)	118 (46.1)	13 (28.3)	41 (46.6)		
Median age (years [IQR])	68.6 [58.8-76.7]	67.7 [56.8–74.6]	66.3 [58.7–75.5] 75.0 [63.0)-81.2]		
Stage at diagnosis						
1–2	95 (24.4)	79 (30.9)	8 (17.4)	8 (9.1)		
3–4	291 (74.6)	175 (68.4)	37 (80.4)	79 (89.8)		
Missing	4 (1.0)	2 (0.8)	1 (2.2)	1(1.1)		
B symptoms at diagnosis	131 (33.6)	69 (27.0)	18 (39.1)	44 (50.0)		
Missing	2 (0.5)	0 (0)	1 (2.2)	1 (1.1)		
ECOG_PS						
0–1	318 (81.5)	210 (82.0)	38 (82.6)	70 (79.5)		
2–4	65 (16.7)	44 (17.2)	6 (13.0)	15 (17.0)		
Missing	7 (1.8)	2 (0.8)	2 (4.3)	3 (3.4)		
Extranodal sites involvement (≥2 sites)	173 (44.4)	102 (39.8)	23 (50.0)	48 (54.5)		
Missing	4 (1.0)	2 (0.8)	1 (2.2)	1 (1.1)		
IPI Score						
0–1	81 (20.8)	70 (27.3)	6 (13.0)	5 (5.7)		
2-3	201 (51.5)	132 (51.6)	26 (56.5)	43 (48.9)		
4–5	102 (26.2)	51 (19.9)	12 (26.1)	39 (44.3)		
Missing	6 (1.5)	3 (1.2)	2 (4.3)	1 (1.1)		
Bulky disease						
Yes	131 (33.6)	71 (27.7)	18 (39.1)	42 (47.7)		
Missing	2 (0.5)	0 (0)	1 (2.2)	1(1.1)		
CNS-IPI score high risk (≥4–5-6)	105 (26.9)	53 (20.7)	12 (26.1)	40 (45.5)		
Treatment intent						
Curative	374 (95.9)	251 (98.0)	43 (93.5)	80 (90.9)		
Palliative	13 (3.3)	4 (1.6)	2 (4.3)	7 (8.0)		
Missing	3 (0.8)	1 (0.4)	1 (2.2)	1(1.1)		
First line treatment						
R-CHOP/R-COMP	315 (80.8)	224 (87.5)	38 (82.6)	53 (60.2)		
R-mini CHOP/COMP	28 (7.2)	12 (4.7)	2 (4.3)	14 (15.9)		
R-CODOX-M/R-IVAC	9 (2.3)	4 (1.6)	1 (2.2)	4 (4.5)		
DA-EPOCH	6 (1.5)	3 (1.2)	1 (2.2)	2 (2.3)		
Oral chemotherapy \pm rituximab	24 (6.2)	11 (4.3)	3 (6.5)	10 (11.4)		
Missing	8 (2.1)	2 (0.8)	1 (2.2)	5 (5.7)		

(Continues)

TABLE I (Continued)				
			R/R	
	All	NRR	Relapsed disease	Refractory disease
	N (%)	N (%)	N (%)	
Second line treatment	102 (100.0)			
Platinum based ^a	38 (37.3)	-	16 (39.0)	22 (36.1)
Lenalidomide	14 (13.7)	-	5 (12.2)	9 (14.8)
Clinical trial	2 (2.0)	-	2 (4.9)	0 (0.0)
Other	37 (36.3)	-	15 (36.6)	22 (36.1)
Oral chemotherapy \pm rituximab	11 (10.8)	-	3 (7.3)	8 (13.1)
Second line consolidation				
RT	11 (10.8)	-	6 (14.6)	5 (8.2)
ASCT	6 (5.9)	-	5 (12.2)	1 (1.6)
Missing	49 (48.0)		19 (46.3)	30 (49.2)

Abbreviations: ASCT, autologous stem cell transplant; CNS, central nervous system; ECOG_PS, Eastern Cooperative Oncology Group_performance status; IPI, international prognostic index; NRR, not relapsed and not refractory disease; R/R, relapsed or refractory disease; R-CHOP/COMP, Rituximab, doxorubicin, vincristine, cyclophosphamide and prednisone (with liposomal doxorubicin in case of previous cardiovascular disease); R-CODOX-M, rituximab, cyclophosphamide, vincristine, doxorubicin and methotrexate; R-DA EPOCH, rituximab, etoposide, vincristine, cyclophosphamide and prednisone; R-IVAC, rituximab, ifosfamide, etoposide, cytarabine; RT, radiotherapy.

The significance is p < 0.05 are in bold.

^aPlatinum-based treatments: rituximab + OxDHA or DHAP or GDP or GemOx.

WILEY_Cancer Medicine

(0 (1)

was used in 73 patients (18.7%), while i.v. methotrexate in 20 (5.1%). Following first-line therapy, 101/390 (25.9%) patients received consolidation radiotherapy (30–36Gy) on bulky disease, while 4 (1.0%) received (ASCT).

3.3 Second- and third-line treatments

Overall, 102/134 patients received second-line treatments. Salvage therapies included high dose, platinum-based (rituximab + OxDHA/DHAP/GemOx) regimens in 38/102 (37.3%) and 6 (5.9%) underwent ASCT. Of note, all patients that were deemed eligible for ASCT in terms of age (<70 years for standard dose conditioning with BEAM/ FEAM schemes, <75 for reduced intensity conditioning) and comorbidities received high dose treatments. Other treatments included lenalidomide (n=14, 13.7%), investigational drugs in clinical trials (n=2, 2.0%), and miscellaneous regimens not containing platinum, mainly oral chemotherapy \pm rituximab. Median age was 62 years for patients who received platinum-based regimens versus 75.5 years for those who did not (p < 0.001). Secondline consolidation with radiotherapy was performed in 11 (10.8%) patients (Table 1). Forty-nine patients R/R to second-line therapies received third-line treatments, including lenalidomide (n=13/49, 26.5%), platinum-based regimens (n=2, 4.0%), investigational drugs (n=1, 2.0%)and others (n = 33, 67.3%). For patients who did not receive

any second-line treatment (n=32), main reason was rapidly progressive disease, with a median time from progression to disease-related death of 15 days. Median age of this subset of patients was 79.4 years old, at initial diagnosis 10 patients (31.3%) had ECOG PS 2–4, and 14 patients (43.8%) had IPI score 4–5; 23 patients (71.9%) had elevated LDH.

3.4 Response to treatments

After first-line, ORR was 81.0% (316/390) including 72.6% complete remission (CR) and 8.5% partial remission (PR); 4.6% of patients had incomplete response data. Overall, only 102 (76.1%) of 134 R/R patients underwent secondline therapy (Figure 1). Reasons for not receiving salvage treatment included disease related death in 29/32 (90.6%) patients. After second-line, ORR was 59.8% (61/102), with CR in 48% patients, PR in 11.8%, stable disease in 2.9% and progressive disease in 34.3% patients; in 2.9% patients data were incomplete. Eighty-nine patients (87.3% of 102 who underwent second-line therapy) were R/R to second-line treatment. ORR after second-line platinum based chemotherapy was 60.5%; in particular 18/38 (48.6%) patients achieved CR and 5 (13.5%) PR; ORR with other regimens was similar (59.4% with 50% CR). Outcomes with secondline lenalidomide included 64.3% ORR with 50.0% patients achieving CR, though response duration was short. Third-line therapy was feasible in 49/89 (55.1%), while the

remaining rapidly deceased due to DLBCL progression (n=36/40, 90.0%, Figure 1). Among those who received third-line therapy, only 10/49 (20.4%) were alive at last follow up.

3.5 | Survival outcomes

At a median follow up of 50 months, overall 5-year OS from diagnosis (OS-1) was 66.5% (Figure 2A). However, median OS-1 for NRR patients was not reached with 243/256 patients (94.9%) alive at last follow up, whereas median OS-1 was 40.0 months (IQR 25.2-64.5) for relapsed and 11.6 months (IQR 6.8-20.7) for refractory patients respectively (Figure 3A). Overall median PFS-1 was not reached with a 2-year PFS-1 of 68.8% (CI 63.5%-72.9%, Figure 2B), whereas median PFS-1 was 20.0 months (15.9-30.6) for relapsed and 6.4 months (IQR 4.5-8.5) for refractory patients respectively (Figure 3B). Overall median OS-2 and PFS-2 were 6.7 (IQR 1.8-18.1), and 5.1 (IQR 2.4-15.5) months respectively (Figure 4). OS-2 was 11.6 months (IQR 3.2–29.0) for relapsed and 4 months (IQR 1.8-10.7) for refractory patients (Figure 5A), while median PFS-2 was 10.1 months (IQR 3.8-18.8) and 4 months (IQR 2.2-6.7) respectively (Figure 5B). In R/R patients (n=107) treated with firstline R-CHOP-like (including R-CHOP/COMP and mini-RCHOP/COMP), OS-2 and PFS-2 did not significantly differ (median OS-2: 8, median PFS-2: 5.2 months).

3.6 | Prognostic factors for survival

Among 134 R/R patients, factors significantly associated with OS-2, by univariate analysis, were disease relapse (p=0.027, HR=1.6), age older than 80 years (p=0.002, HR=2.4), non-GCB COO (p=0.004, HR=2.2), male gender

Cancer Medicine

(p=0.011, HR=0.6), ECOG PS 2-4 (p=0.014, HR=1.9),high-dose first-line treatment (p=0.011, HR=2.5), firstline palliative treatment (p=0.011, HR=2.4), and salvage treatment other than platinum-based regimens (p=0.014, HR = 1.8) (Table 2). After adjusting for confounders, significant variables for OS-2 were age between 60 and 80 years (p = 0.009, HR = 0.20), non-GCB COO (p = 0.010, HR=2.89), ECOG PS 2-4 (p=0.016, HR=2.34), extranodal involvement ≥ 2 sites (p = 0.026, HR = 2.34) (Figure 6). Similarly, significant variables for PFS-2 were first-line outcome (p=0.024), extranodal involvement (p=0.031), LDH (p=0.017); after adjusting for confounders in a no-missing dataset (n=55), no variable was statistically significant (p > 0.05, Table 2). In particular, there was no statistically significant difference in PFS-2 or OS-2 between patients who received platinum-based regimens as compared to those who received other therapies (Figure 7).

By univariate analysis, OS-1 was significantly associated with disease relapse (p < 0.0001, HR for relapsed: 26.7, for refractory: 66.2), age (p < 0.0001, HR for patients >80 years old: 5.9), Ann Arbor stage (p = 0.011, HR for stage III-IV: 2.0), LDH (p = 0.001, HR 2.1), type of first-line treatment (p = 0.006, HR for patients receiving intensive regimens: 2.7, Table 1S), while PFS-1 was significantly associated with age (p < 0.0001, HR for patients >80 years old: 4.8), hemoglobin levels (p = 0.031, HR for patients with hemoglobin >10 g/dL: 0.61), stage (p = 0.001, HR for stage III-IV: 2.2), LDH (p = 0.001, HR 2.0), type of first-line treatment (p = 0.045, HR for patients receiving intensive regimens: 2.0, Table S1).

By multivariate analysis, in 48 R/R patients with a complete dataset, among 6 variables (age, hemoglobin, COO, extranodal involvement, ki67, type of second-line treatment), age between 60 and 80 years, GCB-type COO and extranodal involvement of <2 sites were significantly associated with OS-2 (Table 3). By implementing



FIGURE 2 Overall survival (OS1, A) and progression free survival (PFS1, B) from diagnosis (all patients).



FIGURE 3 Overall survival (OS1, A) and progression free survival (PFS1, B) for non-relapsed/refractory patients, relapsed and refractory patients.



FIGURE 4 Overall survival after first relapse/progression (OS2, A) and progression free survival after first relapse/progression (PFS2, B) from diagnosis.



FIGURE 5 Overall survival (OS2, A) and progression free survival (PFS2, B) from second line treatment for relapsed and refractory patients by univariate analysis.

TABLE 2	Univariate analysis for OS2
and PFS2 by	Cox proportional hazards
model.	

DOGLIOTTI ET AL.

	OS2			PFS2		
Characteristics	HR	95% CI	<i>p</i> -Value	HR	95% CI	p-Value
Outcome						
Relapsed						
Refractory	1.6	1.0-2.3	0.027	1.6	1.0 - 2.5	0.025
Age class						
<60 years						
60-80 years	0.8	0.4–1.3	0.349	0.7	0.4–1.2	0.170
>80 years	2.4	1.4-4.2	0.002	1.513	0.8-2.9	0.210
Sex						
Female						
Male	0.6	0.4–0.9	0.011	0.8	0.5-1.2	0.345
Hb						
$\leq 10 \text{g/dL}$						
>10g/dL	0.9	0.6-1.5	0.797	1.8	0.9-3.5	0.095
COO Hans						
GCB						
Non GCB	2.2	1.3-3.9	0.004	1.4	0.8-2.5	0.220
ECOG						
0–1						
2-4	1.8	1.1-3.0	0.014	0.9	0.4–1.7	0.666
Stage						
1–2						
3–4	1.0	0.6–1.9	0.788	1.2	0.6-2.1	0.567
Extranodal disease						
0-1						
≥2	1.2	0.8–1.7	0.442	1.6	1.0-2.4	0.032
Ki-67						
≤70%						
>70%	1.4	0.7–2.2	0.112	1.3	0.8-2.0	0.344
LDH						
0						
1	1.4	0.9–2.3	0.126	1.9	1.1-3.2	0.019
Type of treatment						
Anthracycline						
High doses	2.5	1.2-5.1	0.011	2.2	1.0-4.8	0.054
Palliative	2.4	1.2-4.6	0.011	1.8	0.8-3.9	0.133
Salvage treatment						
Platinum based	1.0	11.00	0.014	1.0	0.7.1.6	0.711
Other	1.8	1.1-2.9	0.014	1.0	0.7 - 1.6	0.744

Cancer Medicine

Abbreviations: COO, cell of origin; ECOG_PS, Eastern Cooperative Oncology Performance Status; GCB, germinal centre B-cell; Hb, hemoglobin; HR, hazard ratio; LDH, lactate dehydrogenase; OS2, overall survival from relapse; PFS2, progression free survival from relapse.

The significance is p < 0.05 are in bold.

the multiple imputation model for missing data of the 12 variables (n=134), results did not significantly differ with age (p=0.014), COO (p=0.016) and number

of extranodal sites (p = 0.034) remaining significantly associated with OS-2. No baseline feature resulted significantly associated with PFS-2 by multivariate analysis

9 of 15

/ILEY



FIGURE 6 Overall survival (OS2) stratified by age (A), cell of origin (COO) (B) and by ECOG PS (C) at initial diagnosis by univariate analysis.



FIGURE 7 Overall survival (OS2, A) and progression free survival (PFS2, B) with platinum-based versus other regimens as second line treatment by univariate analysis.

on 55 patients with complete data (Table 3). Moreover, logistic regression was performed to highlight determinant features for early (<12 months) R/R disease

(POD12), and significant factors were age older than 80 years (p = 0.005, OR 8.7) and stage III-IV (p = 0.011, OR 7.7, Table S2).

			Canc	er Medicin	COpen Access	WILE	Y 11 of 15
analysis oportional		OS2			PFS2		
	Characteristics	HR	95% CI	<i>p</i> -Value	HR	95% CI	<i>p</i> -Value
	Age class						
	<60 years						
	60-80 years	0.2	0.1-0.7	0.009	0.4	0.9–1.0	0.052
	>80 years	0.5	0.1-2.2	0.4	1.5	0.5-4.3	0.464
	Hb						
	$\leq 10 g/dL$						
	>10g/dL	4.9	0.6-40.0	0.135			
	COO Hans						
	GCB						
	Non GCB	2.8	1.3-6.1	0.010	1.6	0.9–2.9	0.134
	Extranodal disease						
	0-1						
	≥2	2.3	1.1, 4.9	0.026			
	Ki67%						
	≤70%						
	>70%	1.7	0.8-3.6	0.167			
	Salvage treatment						
	Platinum based						
	Other	2.0	0.8-5.1	0.113	1.7	0.9–3.3	0.104
	Sex						
	Female						
	Male				1.6	0.8 - 3.1	0.148

Abbreviations: COO, cell of origin; ECOG_PS, Eastern Cooperative Oncology Performance Status; GCB, germinal centre B-cell; Hb, hemoglobin; HR, hazard ratio; LDH, lactate dehydrogenase; OS2, overall survival from relapse; PFS2, progression free survival from relapse.

The significance is p < 0.05 are in bold.

4 | DISCUSSION

Despite the significant improvement with the introduction of rituximab, clinical outcomes of patients with R/R DLBCL remain invariably poor. Recently, novel immunotherapies such as CAR T cells, antibody-drug conjugates (i.e., polatuzumab vedotin, loncastuximab tesirine), and bispecific antibodies (i.e., glofitamab, epcoritamab) have however shown impressive results. Nonetheless, these agents have not yet been largely approved as second-line therapies by many national Medicine Agencies. Thus, effective salvage treatments remain an urgent medical need. The STRIDER study is a large, retrospective, real life study that confirmed the dismal prognosis of R/R patients in the rituximab era. OS-2 was disappointingly short with rates of about 6 and 4 months in relapsed and refractory patients respectively, with no statistically significant differences between the 2 cohorts. The SCHOLAR-1, the largest pooled analysis on R/R DLBCL patients in the rituximab era, reported a

median OS for refractory patients of 6.3 months from the start of salvage treatment.¹

Salvage treatments commonly include platinum-based regimens such as R-ICE, R-DHAP, R-GDP followed by ASCT consolidation.^{13-16,35} Studies with DHAP and GDP reported 4-year OS of 39%,35 and, with R-DHAP and R-ICE, 3-year OS of 49%.¹⁴ Though comparative studies were not designed, ORRs did not differ significantly ranging from 40% to 60%. R-GemOx, investigated in an older population with median age of 69 years, did not include consolidation with ASCT,³⁶ reporting 5-year OS of 14%. In our rituximab-exposed population, ORR with platinum-based regimens were similar to previous reports, though the initial relatively good ORR and CR rates did not translate into a survival advantage, likely due to short response duration. For this reason, many patients undergoing platinumbased regimens did not receive ASCT consolidation, thus explaining the similar results observed with other less intensive second-line salvage treatments. Of note, only 38/102 (37.3%) patients were however treated with

TABLE 3 Multivariate analysis for OS2 and PFS2 by Cox proportiona hazards model. WILEY_Cancer Medicine

platinum-containing regimens, showing that, in a real life setting, the administration of intensive salvage treatments is not feasible in most patients. Moreover, frailty and chemo-refractoriness, in our experience, were also documented by the fact that only 6 (15.8% of the candidates to transplant patients) receiving platinum-based schemes eventually underwent ASCT. In published reports, the "intent-to-salvage transplant" was as low as 30%.^{14,35}

Reliable clinical parameters predicting increased risk of refractoriness prior to R-CHOP are lacking. Given that every treatment cycle may reduce chemosensitivity, it would be important to identify factors that predict survival and response to salvage treatments. Previous studies showed that high-intermediate IPI score at relapse³⁷ and secondary aaIPI³⁸ affected response rate after salvage therapy while factors such as COO by Hans' algorithm at relapse,³⁹ relapse <12 months after initial therapy⁴⁰ and prior rituximab treatment⁴¹ also affected prognosis. We tried to identify parameters, at diagnosis, that could predict inferior outcomes in case of relapse or progression. By multivariate analysis, age category, COO, and number of extranodal sites involved were predictive of OS-2. However, the current role of patient age should be re-assessed in the light of novel treatments with better toxicity profile compared to standard chemotherapy. Interestingly, our patients between 60 and 80 years showed a significant survival advantage over younger patients. This could partly be explained by more aggressive biologic features in younger patients that may be detected with novel molecular signatures. Of note, a prospective study by the Fondazione Italiana Linfomi (FIL) identified age over 80 years as an independent variable correlated with OS. Moreover, a novel prognostic score for elderly patients, the EPI score, by a simplified version of the geriatric assessment (sGA) was proposed. By classifying patients as fit, unfit, and frail, the EPI score risk correlated with OS of 75%, 58%, and 43%, respectively.⁴² As a matter of fact, at our centre, a thorough geriatric assessment is part of the initial clinical work up by which elderly patients are offered treatment based on their fitness. This may be the reason for the low rates of palliative care undergone by our patients at diagnosis (3.3%) and at relapse/refractoriness (20%) compared to other reports.¹⁷ Indeed, patients over 80 are frequently ineligible for chemotherapy due to comorbidities.⁴³ Lenalidomide may be a valid option, especially in non-GCB patients, with reported ORRs ranging from 29% to 37% with up to 20% CR rates.44,45 We report ORR and CR of 64.3% and 50%, respectively, in 14 patients, median age 76 years, treated with second-line lenalidomide, whereas thirdline lenalidomide in 13 patients showed inferior outcomes. Finally, a recent real world experience reported

outcomes and costs associated with CAR-T cell therapy in DLBCL patients older than 65 (of note; however, regulatory approval in Italy actually limit access to CAR-T programs only to patients <75 years old). Overall, median OS was 17.1 months with no difference between age groups (65–69; 70–74; >75 years).⁴⁶ No patient in the STRIDER was treated with CAR-T cells as the study was conducted before CAR-T cell therapies were commercially approved after two treatment lines.

Splitting the COO category in GCB and non-GCB (NOS) patients by immunohistochemical analysis,^{47–49} according to Hans' algorithm, evaluated at baseline in over half of our patients, identified a strong variable associated with both OS-1 and OS-2 by multivariate analysis. Overall, the role of COO, assessed even with more precise tools such as Nanostring platforms, has recently been questioned in favor of more complex gene signature classifications, able to detect DLBCL molecular heterogeneity and to predict clinical outcome. However, these technologies are not routinely available at most centers and are primarily used in the context of clinical trials.^{5–8}

With the emergence of new therapies, and the increasing biological understanding of DLBCL pathogenesis, efforts to re-design first-line treatments beyond R-CHOP are being made. A stringent baseline risk stratification and the availability of a dynamic risk evaluation in the follow up that includes first-line response and disease kinetics (i.e., early vs. late relapse) should allow to promptly identify poor prognosis patients, who may benefit from earlier interventions with novel immunotherapies, including CAR-T cells.

5 | CONCLUSIONS

The STRIDER is a large, retrospective, real life study that confirms the poor prognosis of R/R DLBCL patients in the rituximab era, before the implementation of "next generation" salvage treatments such as CAR-T cells and bispecific antibodies. Efficacy of high dose chemotherapy-based salvage treatments is limited, requiring the urgent and widespread approval of these novel immunotherapies, mainly investigated in clinical trials in Europe.

AUTHOR CONTRIBUTIONS

Irene Dogliotti: Conceptualization (lead); data curation (equal); investigation (lead); writing – original draft (lead); writing – review and editing (equal). **Veronica Peri:** Data curation (lead); writing – original draft (equal); writing – review and editing (lead). **Michele Clerico:** Data curation (equal); writing – review and editing (equal). Francesco Vassallo: Data curation (equal). Davide Musto: Methodology (equal); software (equal). Silvio Mercadante: Methodology (equal); software (lead). Simone Ragaini: Data curation (equal); writing - review and editing (equal). Barbara Botto: Supervision (equal); writing - review and editing (equal). Mario Levis: Data curation (equal); writing review and editing (equal). Mattia Novo: Data curation (equal); writing - review and editing (equal). Marco Ghislieri: Methodology (equal); software (equal). Luca Molinaro: Data curation (equal); investigation (equal). Umberto Mortara: Data curation (supporting). Chiara **Consoli:** Data curation (supporting); writing – review and editing (supporting). Alessio Lonardo: Project administration (equal). Giulia Bondielli: Project administration (equal). Simone Ferrero: Investigation (equal); resources (equal); writing - review and editing

(equal). Roberto Freilone: Supervision (equal); writing - review and editing (equal). Umberto Ricardi: Supervision (equal); writing - review and editing (equal). Benedetto Bruno: Methodology (equal); supervision (lead); writing - original draft (equal). Federica Cavallo: Conceptualization (lead); funding acquisition (lead); methodology (equal); resources (equal); supervision (equal); writing - original draft (equal); writing review and editing (lead).

ACKNOWLEDGMENTS

The STRIDER study was supported by an unconditional grant by Roche Italia. ID was supported by SIE (Società Italiana di Ematologia) and AIL (Associazione Italiana contro le Leucemie, Linfomi e Mieloma).

CONFLICT OF INTEREST STATEMENT

BB has received speaker honoraria from Amgen, Janssen, Novartis, BeiGene, Bristol Myers Squibb, GlaxoSmithKline, Jazz Pharmaceuticals, Astrazeneca, Incyte; has served on advisory boards for Amgen, Jazz Pharmaceuticals. SF has received research funding from Janssen, Morphosys, Gilead, Beigene; has served on advisory boards for EusaPharma, Janssen, Clinigen, Incyte, Italfarmaco, Roche, Abbvie, has served as consultant for EusaPharma, Janssen, Sandoz, Abbvie, has received speaker honoraria from Janssen, EusaPharma, Servier, Gentili, Lilly, Beigene, Gilead. FC has received speaker honoraria from Lilly, Incyte; has served on advisory boards for Roche, Incyte; has served as consultant for Astrazeneca.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Irene Dogliotti D https://orcid.org/0009-0004-0873-1622 Benedetto Bruno Dhttps://orcid. org/0000-0002-7299-6770 Federica Cavallo D https://orcid. org/0000-0003-2047-1099

REFERENCES

- 1. Crump M, Neelapu SS, Farooq U, et al. Outcomes in refractory diffuse large B-cell lymphoma: results from the international SCHOLAR-1 study. Blood. 2017;130:1800-1808. doi:10.1182/ blood-2017-03-769620
- 2. Feugier P, van Hoof A, Sebban C, et al. Long-term results of the R-CHOP study in the treatment of elderly patients with diffuse large B-cell lymphoma: a study by the groupe d'etude des lymphomes de l'adulte. J Clin Oncol. 2005;23:4117-4126. doi:10.1200/JCO.2005.09.131
- 3. Petrich AM, Gandhi M, Jovanovic B, et al. Impact of induction regimen and stem cell transplantation on outcomes in doublehit lymphoma: a multicenter retrospective analysis. Blood. 2014;124:2354-2361. doi:10.1182/blood-2014-05-578963
- 4. Swerdlow SH, Campo E, Pileri SA, et al. The 2016 revision of the World Health Organization classification of lymphoid neoplasms. Blood. 2016;127:2375-2390. doi:10.1182/blood-2016-01-643569
- 5. Chapuy B, Stewart C, Dunford AJ, et al. Molecular subtypes of diffuse large B cell lymphoma are associated with distinct pathogenic mechanisms and outcomes. Nat Med. 2018;24:679-690. doi:10.1038/s41591-018-0016-8
- 6. Lacy SE, Barrans SL, Beer PA, et al. Targeted sequencing in DLBCL, molecular subtypes, and outcomes: a Haematological malignancy research network report. Blood. 2020;135:1759-1771. doi:10.1182/BLOOD.2019003535
- 7. Schmitz R, Wright GW, Huang DW, et al. Genetics and pathogenesis of diffuse large B-cell lymphoma. N Engl J Med. 2018;378:1396-1407. doi:10.1056/nejmoa1801445
- 8. Wright GW, Huang DW, Phelan JD, et al. A probabilistic classification tool for genetic subtypes of diffuse large B cell lymphoma with therapeutic implications. Cancer Cell. 2020;37:551-568. e14. doi:10.1016/j.ccell.2020.03.015
- 9. Coiffier B, Lepage E, Brière J, et al. CHOP chemotherapy plus rituximab compared with CHOP alone in elderly patients with diffuse large-B-cell lymphoma. N Engl J Med. 2002;346:235-242. doi:10.1056/nejmoa011795
- 10. Coiffier B, Thieblemont C, Van Den Neste E, et al. Long-term outcome of patients in the LNH-98.5 trial, the first randomized study comparing rituximab-CHOP to standard CHOP chemotherapy in DLBCL patients: a study by the Groupe d'Etudes des Lymphomes de l'Adulte. Blood. 2010;116(12):2040-2045. doi:10.1182/blood-2010-03-276246
- 11. Pfreundschuh M, Trümper L, Osterborg A, et al. CHOP-like chemotherapy plus rituximab versus CHOP-like chemotherapy alone in young patients with good-prognosis diffuse large-Bcell lymphoma: a randomised controlled trial by the MabThera international trial (MInT) group. Lancet Oncol. 2006;7:379-391. doi:10.1016/S1470-2045(06)70664-7
- Tilly H, Morschhauser F, Sehn LH, et al. Polatuzumab vedotin 12. in previously untreated diffuse large B-cell lymphoma. N Engl J Med. 2022;386:351-363. doi:10.1056/nejmoa2115304

WILEY_Cancer Medicine _

- Tilly H, Gomes da Silva M, Vitolo U, et al. Diffuse large B-cell lymphoma (DLBCL): ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2015;26:v116-v125. doi:10.1093/annonc/mdv304
- 14. Gisselbrecht C, Glass B, Mounier N, et al. Salvage regimens with autologous transplantation for relapsed large B-cell lymphoma in the rituximab era. *J Clin Oncol.* 2010;28:4184-4190. doi:10.1200/JCO.2010.28.1618
- 15. Kewalramani T, Zelenetz AD, Nimer SD, et al. Rituximab and ICE as second-line therapy before autologous stem cell transplantation for relapsed or primary refractory diffuse large B-cell lymphoma. *Blood*. 2004;103(10):3684-3688.
- Crump M, Baetz T, Couban S, et al. Gemcitabine, dexamethasone, and cisplatin in patients with recurrent or refractory aggressive histology B-cell non-Hodgkin lymphoma: a phase II study by the National Cancer Institute of Canada clinical trials group (NCIC-CTG). *Cancer*. 2004;101:1835-1842. doi:10.1002/ cncr.20587
- Arboe B, Olsen MH, Gørløv JS, et al. Treatment intensity and survival in patients with relapsed or refractory diffuse large Bcell lymphoma in Denmark: A Real-Life Populationbased Study. *Clin Epidemiol.* 2019;11:207-216. doi:10.2147/CLEP.S178003
- Pettengell R, Coiffier B, Narayanan G, et al. Pixantrone dimaleate versus other chemotherapeutic agents as a single-agent salvage treatment in patients with relapsed or refractory aggressive non-Hodgkin lymphoma: a phase 3, multicentre, openlabel, randomised trial. *Lancet Oncol.* 2012;13:696-706.
- Czuczman MS, Trněný M, Davies A, et al. A phase 2/3 multicenter, randomized, open-label study to compare the efficacy and safety of lenalidomide versus investigator's choice in patients with relapsed or refractory diffuse large B-cell lymphoma. *Clin Cancer Res.* 2017;23:4127-4137. doi:10.1158/1078-0432. CCR-16-2818
- Klyuchnikov E, Bacher U, Kroll T, et al. Allogeneic hematopoietic cell transplantation for diffuse large B cell lymphoma: who, when and how. *Bone Marrow Transplant*. 2014;49:1-7. doi:10.1038/bmt.2013.72
- Izumi K, Kanda J, Nishikori M, et al. Outcomes of allogeneic stem cell transplantation for DLBCL: a multi-center study from the Kyoto stem cell transplantation group. *Ann Hematol.* 2019;98:2815-2823. doi:10.1007/s00277-019-03835-3
- Neelapu SS, Locke FL, Bartlett NL, et al. Axicabtagene ciloleucel CAR T-cell therapy in refractory large B-cell lymphoma. N Engl J Med. 2017;377:2531-2544. doi:10.1056/nejmoa1707447
- Schuster SJ, Bishop MR, Tam CS, et al. Tisagenlecleucel in adult relapsed or refractory diffuse large B-cell lymphoma. N Engl J Med. 2019;380:45-56. doi:10.1056/nejmoa1804980
- Abramson JS, Palomba ML, Gordon LI, et al. Lisocabtagene maraleucel for patients with relapsed or refractory large Bcell lymphomas (TRANSCEND NHL 001): A Multicentre Seamless Design Study. *Lancet.* 2020;396:839-852. doi:10.1016/ S0140-6736(20)31366-0
- 25. Kamdar M, Solomon SR, Arnason J, et al. Lisocabtagene maraleucel versus standard of care with salvage chemotherapy followed by autologous stem cell transplantation as secondline treatment in patients with relapsed or refractory large B-cell lymphoma (TRANSFORM): results from an interim analysis of an open-label, randomised, phase 3 trial. *Lancet*. 2022;399:2294-2308.

- Locke FL, Ghobadi A, Jacobson CA, et al. Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1–2 trial. *Lancet Oncol.* 2019;20:31-42. doi:10.1016/ S1470-2045(18)30864-7
- Schuster SJ, Tam CS, Borchmann P, et al. Long-term clinical outcomes of tisagenlecleucel in patients with relapsed or refractory aggressive B-cell lymphomas (JULIET): A Multicentre, Open-Label, Single-Arm, Phase 2 Study. *Lancet Oncol.* 2021;22:1403-1415. doi:10.1016/S1470-2045(21)00375-2
- Locke FL, Miklos DB, Jacobson CA, et al. Axicabtagene ciloleucel as second-line therapy for large B-cell lymphoma. *N Engl J Med.* 2022;386:640-654. doi:10.1056/nejmoa2116133
- Salles G, Duell J, González Barca E, et al. Tafasitamab plus lenalidomide in relapsed or refractory diffuse large B-cell lymphoma (L-MIND): A Multicentre, Prospective, Single-Arm, Phase 2 Study. *Lancet Oncol.* 2020;21:978-988. doi:10.1016/ S1470-2045(20)30225-4
- Sehn LH, Herrera AF, Flowers CR, et al. Polatuzumab vedotin in relapsed or refractory diffuse large B-cell lymphoma. *J Clin Oncol.* 2020;38:155-165. doi:10.1200/JCO.19.00172
- Caimi PF, Ai W, Alderuccio JP, et al. Loncastuximab tesirine in relapsed or refractory diffuse large B-cell lymphoma (LOTIS-2): a multicentre, open-label, single-arm, phase 2 trial. *Lancet Oncol.* 2021;22:790-800. doi:10.1016/S1470-2045(21)00139-X
- 32. Kalakonda N, Maerevoet M, Cavallo F, et al. Selinexor in patients with relapsed or refractory diffuse large B-cell lymphoma (SADAL): a single-arm, multinational, multicentre, open-label, phase 2 trial. *Lancet Haematol.* 2020;7:e511-e522. doi:10.1016/S2352-3026(20)30120-4
- Cheson BD, Horning SJ, Coiffier B, et al. Report of an international workshop to standardize response criteria for non-Hodgkin's lymphomas. NCI sponsored international working group. J Clin Oncol. 1999;17:1244.
- Cheson BD, Fisher RI, Barrington SF, et al. Recommendations for initial evaluation, staging, and response assessment of hodgkin and non-hodgkin lymphoma: the lugano classification. J Clin Oncol. 2014;32:3059-3067. doi:10.1200/JCO.2013.54.8800
- 35. Crump M, Kuruvilla J, Couban S, et al. Randomized comparison of gemcitabine, dexamethasone, and cisplatin versus dexamethasone, cytarabine, and cisplatin chemotherapy before autologous stem-cell transplantation for relapsed and refractory aggressive lymphomas: NCIC-CTG LY.12. J Clin Oncol. 2014;32:3490-3496. doi:10.1200/JCO.2013.53.9593
- Mounier N, el Gnaoui T, Tilly H, et al. Rituximab plus gemcitabine and oxaliplatin in patients with refractory/relapsed diffuse large B-cell lymphoma who are not candidates for high-dose therapy. A phase II lymphoma study association trial. *Haematologica*. 2013;98:1726-1731. doi:10.3324/haematol.2013.090597
- Blay JY, Gomez F, Sebban C, et al. The International Prognostic Index correlates to survival in patients with aggressive lymphoma in relapse: analysis of the PARMA trial. *Blood.* 1998;92:3562-3568.
- Hamlin PA, Zelenetz AD, Kewalramani T, et al. Age-adjusted international prognostic index predicts autologous stem cell transplantation outcome for patients with relapsed or primary refractory diffuse large B-cell lymphoma. *Blood*. 2003;102:1989-1996. doi:10.1182/blood-2002-12-3837
- Thieblemont C, Briere J, Mounier N, et al. The germinal center/activated B-cell subclassification has a prognostic impact for response to salvage therapy in relapsed/refractory

14 of 15

diffuse large B-cell lymphoma: A Bio-Coral Study. *J Clin Oncol.* 2011;29:4079-4087. doi:10.1200/JCO.2011.35.4423

- Guglielmi C, Gomez F, Philip T, et al. Time to relapse has prognostic value in patients with aggressive lymphoma enrolled onto the parma trial. *J Clin Oncol.* 1998;16:3264-3269. doi:10.1200/JCO.1998.16.10.3264
- Martín A, Conde E, Arnan M, et al. R-ESHAP as salvage therapy for patients with relapsed or refractory diffuse large B-cell lymphoma: the influence of prior exposure to rituximab on outcome. A GEL/TAMO Study. *Haematologica*. 2008;93:1829-1836. doi:10.3324/haematol.13440
- Merli F, Luminari S, Tucci A, et al. Simplified geriatric assessment in older patients with diffuse large B-cell lymphoma: the prospective elderly project of the Fondazione Italiana Linfomi. *J Clin Oncol.* 2021;39:1214-1222. doi:10.1200/JCO.20.02465
- Wildes TM, Augustin KM, Sempek D, et al. Comorbidities, not age, impact outcomes in autologous stem cell transplant for relapsed non-Hodgkin lymphoma. *Biol Blood Marrow Transplant*. 2008;14:840-846. doi:10.1016/j.bbmt.2008.05.002
- Mondello P, Steiner N, Willenbacher W, et al. Lenalidomide in relapsed or refractory diffuse large B-cell lymphoma: is it a valid treatment option? *Oncologist.* 2016;21:1107-1112. doi:10.1634/ theoncologist.2016-0103
- 45. Broccoli A, Casadei B, Chiappella A, et al. Lenalidomide in pretreated patients with diffuse large B-cell lymphoma: An Italian Observational Multicenter Retrospective Study in daily clinical practice. *Oncologist.* 2019;24:1246-1252. doi:10.1634/ theoncologist.2018-0603

- Chihara D, Liao L, Tkacz J, et al. Real-world experience of CAR T-cell therapy in older patients with relapsed/refractory diffuse large B-cell lymphoma. *Blood*. 2023;142:1047-1055. doi:10.1182/blood.2023020197
- Alizadeh AA, Eisen MB, Davis RE, et al. Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. *Nature*. 2000;403:503-511. doi:10.1038/35000501
- Lenz G, Wright GW, Emre NCT, et al. Molecular subtypes of diffuse large B-cell lymphoma arise by distinct genetic pathways. *Proc Natl Acad Sci USA*. 2008;105:13520-13525. doi:10.1073/pnas.0804295105
- 49. Rosenwald A, Wright G, Chan WC, et al. The use of molecular profiling to predict survival after chemotherapy for diffuse large-B-cell lymphoma. *N Engl J Med.* 2002;346:1937-1947. doi:10.1056/nejmoa012914

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Dogliotti I, Peri V, Clerico M, et al. Real life clinical outcomes of relapsed/refractory diffuse large B cell lymphoma in the rituximab era: The STRIDER study. *Cancer Med.* 2024;13:e7448. doi:10.1002/cam4.7448