

Reconstructing Atmospheric Parameters of Exoplanets Using Deep Learning

*Original*

Reconstructing Atmospheric Parameters of Exoplanets Using Deep Learning / Giobergia, Flavio; Koudounas, Alkis; Baralis, Elena Maria. - (2023). (Intervento presentato al convegno 2023 IEEE 17th International Conference on Application of Information and Communication Technologies (AICT) tenutosi a Baku (AZE) nel 18-20 October 2023) [10.1109/AICT59525.2023.10313185].

*Availability:*

This version is available at: 11583/2982630 since: 2023-09-30T14:02:47Z

*Publisher:*

IEEE

*Published*

DOI:10.1109/AICT59525.2023.10313185

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

IEEE postprint/Author's Accepted Manuscript

©2023 IEEE. Personal use of this material is permitted. Permission from IEEE must be obtained for all other uses, in any current or future media, including reprinting/republishing this material for advertising or promotional purposes, creating new collecting works, for resale or lists, or reuse of any copyrighted component of this work in other works.

(Article begins on next page)

# Reconstructing Atmospheric Parameters of Exoplanets Using Deep Learning

Flavio Giobergia  
Politecnico di Torino  
Turin, Italy  
flavio.giobergia@polito.it

Alkis Koudounas  
Politecnico di Torino  
Turin, Italy  
alkis.koudounas@polito.it

Elena Baralis  
Politecnico di Torino  
Turin, Italy  
elena.baralis@polito.it

**Abstract**—Exploring exoplanets has transformed our understanding of the universe by revealing many planetary systems that defy our current understanding. To study their atmospheres, spectroscopic observations are used to infer essential atmospheric properties that are not directly measurable. Estimating atmospheric parameters that best fit the observed spectrum within a specified atmospheric model is a complex problem that is difficult to model. In this paper, we present a multi-target probabilistic regression approach that combines deep learning and inverse modeling techniques within a multimodal architecture to extract atmospheric parameters from exoplanets. Our methodology overcomes computational limitations and outperforms previous approaches, enabling efficient analysis of exoplanetary atmospheres. This research contributes to advancements in the field of exoplanet research and offers valuable insights for future studies.

**Index Terms**—deep learning, exoplanets, atmospheric retrieval

## I. INTRODUCTION

Exploring exoplanets has revolutionized our knowledge of the universe, unveiling a diverse range of planetary systems that challenge our existing notions [1]. These exoplanets span a wide spectrum, ranging from Earth-like habitable worlds [2] to searing hot-Jupiters [3] and everything in between. Their existence offers profound insights into the mechanisms governing planetary formation and evolution.

To deepen our understanding and contextualize our own solar system within the broader galactic landscape, the European Space Agency (ESA) is collaborating with NASA and JAXA to develop the Ariel Space Mission [4]. This ambitious endeavor aims to extensively investigate the atmospheres of hundreds of exoplanets in close proximity using a meter-class telescope in L2. By analyzing spectra and photometric data in the visible and infrared range, Ariel will extract chemical compositions and study the thermal properties of exoplanetary atmospheres. The mission’s open data policy will allow rapid access to high-quality exoplanet spectra for scientific research.

While the Ariel mission promises groundbreaking discoveries, it faces a formidable hurdle in the complex modeling of exoplanetary atmospheres. These atmospheres exhibit intricate chemistries, cloud formations, and dynamic processes, making their characterization a highly intricate task. The traditional approach of forward modeling, which involves simulating atmospheric spectra based on predefined models, is computationally intensive and unsuitable for efficient fitting or

sampling techniques such as Markov Chain Monte Carlo or Nested Sampling [5].

This paper presents our proposed solution to the Ariel Machine Learning Data Challenge<sup>1</sup>, which involves addressing the multi-target probabilistic regression problem of extracting critical atmospheric parameters of exoplanets on the Ariel Big Challenge (ABC) dataset [6]. We propose an innovative multimodal architecture that leverages deep learning and inverse modeling techniques to map observed atmospheric signatures to the underlying planetary characteristics. In addition, we incorporate a modeling approach where the target output is treated as a Gaussian distribution with a full covariance matrix. This enables us to predict and estimate the parameters  $\mu$  (mean) and  $\Sigma$  (covariance matrix) that define the distribution. By employing a data-driven approach, our method aims to overcome the limitations of conventional forward modeling and expedite the characterization process, enabling rapid analysis of exoplanetary atmospheres.

Our contributions are as follows:

- We introduce a novel deep learning multimodal approach that addresses the multi-regression problem of determining key atmospheric parameters of exoplanets, outperforming previous approaches.
- We overcome the computational limitations of traditional forward modeling techniques by leveraging data-driven methodologies. Our approach allows for faster and more efficient analysis of exoplanetary atmospheres.
- Our work contributes to the broader field of exoplanet research by providing a more efficient and accurate method for characterizing exoplanetary atmospheres.

The insights gained from our approach have practical implications for future space missions, including the Ariel Space Mission. Our methodology, which achieved the 8<sup>th</sup> position in the highly competitive competition with over 250 participants, provides a promising approach to accelerate the analysis of exoplanetary atmospheres. This approach enables quicker and more thorough characterization of numerous exoplanets, contributing to advancements in the field.

The remainder of the paper is organized as follows. We first examine the related works and highlight the difference between the present work and existing approaches (Section II).

<sup>1</sup><https://www.ariel-datachallenge.space/>

We then detail the problem we have to address (Section III) and the dataset utilized for the experiments, and we discuss the methodology employed in our approach (Section IV). In Section V, we present our experimental results and finally highlight the implications and potential applications of our findings (Section VI).

## II. RELATED WORKS

The field of exoplanet studies has evolved from detecting exoplanets to characterizing their atmospheres through retrieval techniques. Retrieval is an inverse modeling approach that compares forward models of a planet’s spectrum to observational data, allowing for the estimation of atmospheric parameters [7], [8]. Atmospheric retrieval is thus crucial in helping astronomers understand and analyze individual observations acquired through spectroscopy during transit, eclipse, and phase curves. This retrieval technique applies to both low and high-resolution data [9]–[13], making it an indispensable tool for astronomers in interpreting the atmospheric properties of exoplanets.

Bayesian methods are commonly used to retrieve noisy exoplanet spectra, providing posterior distributions that constrain model parameters and determine their statistical significance, leveraging sampling techniques such as Markov Chain Monte Carlo (MCMC) or Nested Sampling [5]. However, traditional Bayesian retrieval methods can be computationally expensive and time-consuming [14]. As the volume of exoplanetary data increases, driven by missions like the James Webb Space Telescope (JWST) [15] and Ariel [4], alternative approaches to computing posterior distributions are needed.

Machine learning (ML) and deep learning (DL) techniques have been applied to many areas within exoplanetary science, including data detrending [16], [17], debris removal [18], [19], and planet detection and characterization [20]–[23]. Various ML and DL approaches, such as random forests, generative adversarial networks (GANs), convolutional neural networks (CNNs), and Bayesian neural networks, have been employed to improve retrieval efficiency while significantly reducing computational time. Although these approaches show promise, as computational time is reduced, the accuracy of the posterior estimation decreases due to substantial approximations made during Bayesian inference.

In contrast to previous approaches, our methodology incorporates a multimodal 1D-CNN that combines information from both the spectral data and auxiliary data related to stellar and planetary parameters. This novel approach allows us to achieve favorable results in approximating the posterior distribution while maintaining remarkably low computational time.

## III. PROBLEM STATEMENT

Exoplanets are discovered using various methods, but the most commonly used techniques are radial velocity and transit [6]. Transit is an indirect method to monitor changes in the brightness of the host star. During a transit event, the planet passes in front of the star, causing a reduction in the amount

of light observed from Earth. By studying these transit events at different wavelengths, astronomers can gain insights into the atmospheric properties of the exoplanet.

The exoplanetary atmosphere affects the transit depth by absorbing stellar light wavelength-dependently. This absorption profile is influenced by the composition (molecular species, clouds, hazes) and characteristics (thermal structure) of the atmosphere. Astronomers employ simplified models to analyze the observed signal, known as a *spectrum*, and investigate the underlying atmospheric processes occurring in exoplanet atmospheres.

The study of exoplanetary atmospheres relies on spectroscopic observations to infer fundamental atmospheric properties that cannot be directly measured. This process is known as the inverse problem [24], where one tries to deduce the atmospheric properties based on the observed effects. However, the observed effects are often corrupted due to factors such as noise and limited spectroscopic coverage, leading to a loss of information, which can, in turn, result in model degeneracy.

The main objective of atmospheric retrieval is to estimate the parameters that best explain the observed spectrum under a given atmospheric model. This is typically approached using a forward model, which incorporates assumptions about the atmosphere and an optimizer. In the Bayesian framework, the goal is to determine the posterior distribution, which represents the conditional distribution of the model parameters given the observed data.

In our case, the aim of the Ariel Machine Learning Data Challenge is to predict the conditional joint distribution (or the Bayesian posterior distribution) of 7 atmospheric properties given the observed spectrum, namely planet radius (in Jupiter radii), temperature (in Kelvin), and the log-abundance of five atmospheric gases:  $H_2O$ ,  $CO_2$ ,  $CO$ ,  $CH_4$ , and  $NH_3$ . The probability densities of these variables ( $\theta = \theta_1, \theta_2, \dots, \theta_n$ ) given the observed data ( $D$ ) are collectively referred to as the posterior distribution,  $P(\theta|D)$ .

### A. Dataset

To develop and validate our approach, we utilize the Ariel Big Challenge (ABC) Database [6], encompassing simulated atmospheric spectra, ground-truth models, and Bayesian posterior distributions for a total of 6,766 exoplanets. The database is generated using the powerful Alfnor [25] framework combining the TauREx 3 [26] atmospheric modeling suite and the ArielRad [27] instrument simulator. The dataset comprises two types of data, namely spectral and auxiliary data. Additionally, the target data is available in the form of tracedata.

*Spectral data.* Each entry in the dataset consists of an atmospheric spectrum with 52 data points. Each of them contains information about the intensity measure (transit depth), the corresponding wavelength of light, the spectral resolution (wavelength bin size), and the associated measurement uncertainty. Figure 1 represents the spectrum for one of the exoplanets available in the training set.




Fig. 1. Spectral information available for each exoplanet. The quantity  $(R_p/R_*)^2$  represents the ratio between the planet’s radius, and that of the orbited star since the measured dip can be quantified in terms of a ratio of the areas of the two entities involved.

*Auxiliary data.* Each example further encompasses eight additional stellar and planetary parameters as auxiliary information, including star distance, stellar mass, stellar radius, stellar temperature, planet mass, orbital period, semi-major axis, planet radius, and surface gravity. This information is unique to each planet and is sourced from various exoplanet datasets.

*Tracedata.* The target output is provided in terms of samples from the conditional joint distribution, characterized by the aforementioned 7 atmospheric properties, referred to as tracedata. For each planet, an average of 3,938 samples are available (standard deviation of 654). Figure 2 shows one such distribution as an example.

#### IV. METHODOLOGY

We propose adopting a deep learning solution to solve the Ariel Machine Learning Data Challenge problem. At a high level, we leverage both the spectral data and the auxiliary data that has been made available with the adoption of a multimodal architecture; we additionally model the target output as a full covariance Gaussian distribution, thus making it possible to predict the parameters  $\mu$  and  $\Sigma$  that characterize it. Figure 3 summarizes the proposed approach. The rest of this section analyzes each component in further detail.

##### A. Spectral data

As already argued in Subsection III-A, the spectral data conveys significant information regarding the composition of the atmosphere of an exoplanet. Since the atmospheric spectrum is continuous in nature, we consider adopting a 1-dimensional convolutional (1D-CNN) model to process the spectrum to be able to characterize local behaviors that can be found in the spectrum itself. The adoption of 1D-CNN models has already been adopted in literature with promising results [22]. We build a representation of the spectrogram as




Fig. 2. Samples for the atmospheric parameters for one of the available exoplanets. In blue is the ground truth distribution, in orange the distribution predicted by the proposed approach.

a dense vector by flattening the feature maps obtained as the output of the 1D-CNN.

##### B. Auxiliary data

The auxiliary data available provides information on the planetary system, including the exoplanet under study and the orbited star. Based on the intuition that some of the auxiliary information may have non-linear relationships with the target outputs, we introduce a non-linear transformation of these inputs. In particular, we additionally obtain the inverse of the auxiliary quantities and, from these, we extract the polynomials up to degree 2. Starting from the original 8 features, we obtain a total of 152 transformed auxiliary features. The extracted features are then concatenated to the dense vector that is used to represent the spectral data. In this way, we produce a vector that represents the entire input both in terms of atmospheric spectrum and contextual planetary system information.

##### C. Parameters estimation

The main goal of this work is to produce samples that are drawn from an underlying, unknown data distribution. These samples represent different atmospheric models that make different assumptions (e.g., in terms of atmospheric composition). We frame the problem by modeling the underlying distribution to produce the desired samples as a byproduct.

The target available is provided as samples from the ground truth distribution. Since we aim to predict this distribution, we need to make an assumption regarding the type of probability distribution. We assume the distribution to be a multivariate




Fig. 3. Architecture for the proposed solution. The spectral data is processed through a convolutional model (1D-CNN). The auxiliary data is augmented and merged with the processed spectral data. A final fully-connected model (FC-NN) maps the processed input to the desired probability distribution. The parameters for the target distribution are also produced and used when computing the loss function.

Gaussian one with full covariance: based on the intuition that can be extracted from Figure 2, this is a simplifying assumption. However, it allows for a simple parametrization of the distribution. Additionally, it allows for the modeling of interactions across dimensions thanks to the adoption of a full covariance.

For each planet, we focus on  $n$  target variables  $\theta_1, \theta_2, \dots, \theta_n$  (for this specific challenge – as discussed – we have  $n = 7$ , but the same approach can be adopted for differently framed problems). We can define the mean vector  $\mu \in \mathbb{R}^n$  such that  $\mu_i = E[T_i]$  and the covariance matrix  $\Sigma \in \mathbb{R}^{n \times n}$  such that  $\Sigma_{ij} = \text{cov}(T_i, T_j)$  and  $\Sigma_{ii} = \text{Var}(T_i)$ .

Since the covariance matrix is symmetric, we can build a compressed representation using  $\frac{n(n+1)}{2}$  values. When accounting for  $\mu$ , the overall distribution can be parametrized with a total of  $\frac{n(n+3)}{2}$  values. We refer to the target representation for the  $k^{\text{th}}$  planet as  $y^{(k)} \in \mathbb{R}^{\frac{n(n+3)}{2}}$ . We note that we can freely move from the  $\mu^{(k)}, \Sigma^{(k)}$  representation to the  $y^{(k)}$  one. Similarly, we refer to the predicted parameters as either  $\hat{y}^{(k)}$  or as the corresponding  $\hat{\mu}^{(k)}, \hat{\Sigma}^{(k)}$ .

#### D. Loss function

To compute the proximity of the predicted distribution with the target one, we considered various possible loss functions. We empirically observed that computing the KL divergence leads to a difficult convergence process. In contrast, we obtain better convergence properties by minimizing the L1 loss function between the predicted and ground truth values. In particular, the loss is defined as:

$$\ell(y, \hat{y}) = |y - \hat{y}| \quad (1)$$

This is equivalent, in terms of mean vectors and covariance matrices, as:

$$\ell(\mu_i, \Sigma_i, \hat{\mu}_i, \hat{\Sigma}_i) = \sum_{0 \leq i < n} \sum_{i < j < n} |\Sigma_{ij} - \hat{\Sigma}_{ij}| + \sum_{0 \leq i < n} |\Sigma_{ii} - \hat{\Sigma}_{ii}| + \sum_{0 \leq i < n} |\mu_i - \hat{\mu}_i| \quad (2)$$

#### V. RESULTS

In accordance with the policies of the Ariel Machine Learning Data Challenge, we evaluate the quality of the obtained results using two terms: the posterior score and the spectral score. The two metrics complement each other: the first ensures that the predictions accurately replicate each individual distribution for the variables involved, while the second focuses on preserving the physical laws or, more precisely, the interrelationships (covariance) among different targets.

We employ the 2 Sample Kolmogorov–Smirnov test (K-S test) to evaluate the posterior score scenario. The K-S test is a widely used statistical test determining whether two given samples are derived from the same continuous distribution underlying them. The output is adjusted to range from 0 to 1000, where a score of 1000 indicates the highest similarity.

Regarding the spectral score, we compare the “median” values in spectral space, which consists of the median value for each wavelength bin, along with their uncertainty bounds (interquartile ranges of each wavelength bin), against the corresponding values obtained from Bayesian Nested Sampling. We quantify the differences using an inverse Huber loss. The

spectral score is calculated as a linear combination of the differences between the bounds and the differences between the median spectra. The maximum score is set to 1000, with 0 representing the lowest similarity.

The final score is computed as a weighted sum of the spectral loss and posterior loss:

$$Final = (1 - \beta) \cdot Spectral + \beta \cdot Posterior \quad (3)$$

where  $\beta$  is set to 0.8. The value set for  $\beta$  is in accordance with the one adopted for the Ariel Machine Learning Data Challenge and reflects the greater importance that is generally assigned to the posterior score w.r.t. the spectral one. The final score ranges from a minimum of 0 to a maximum of 1000.

Table I summarizes the results achieved by the baseline model presented in [22] and by the proposed approach on a test set obtained as a 20% hold-out on the available data. For completeness, we include the results obtained by the proposed methodology when changing the loss function to another commonly adopted one, the mean squared error (L2).

We show that the proposed approach consistently outperforms the baseline model in terms of overall (final) score. This is the result of a model that performs consistently well in terms of both posterior and spectral scores. By contrast, the method proposed in [22] is unbalanced toward performing significantly well in terms of spectral score with a consistent drop in performance as far as the posterior score is concerned.

TABLE I  
RESULTS ON THE ARIEL BIG CHALLENGE DATABASE IN TERMS OF POSTERIOR, SPECTRAL, AND FINAL SCORE. BEST RESULTS ARE HIGHLIGHTED IN BOLD.

Method	Posterior Score	Spectral Score	Final Score
Yip et al. [22]	331.58 ± 60.57	<b>880.05 ± 4.75</b>	441.28 ± 48.97
Ours	611.77 ± 14.52	658.57 ± 9.63	<b>621.13 ± 9.95</b>
Ours w/ L2	610.054 ± 11.10	640.35 ± 12.77	616.10 ± 9.31
Ours w/ L1+L2	<b>613.04 ± 15.96</b>	637.07 ± 11.26	617.71 ± 15.33

### A. Qualitative results

Figure 2 shows a ground truth distribution (in blue), along with the predicted one (in orange). We observe how there is a reasonably good fit for most variables, considering the constraints introduced by the Gaussian simplifying assumption. We can also notice how using a full covariance allows for better modeling of variables that show some degree of correlation.

However, we note that the proposed model particularly struggles with predicting the planet’s radius. Indeed, the distribution of samples for that parameter is rather narrow. In contrast, the model makes a high-variance prediction: this produces a non-zero overlap with the target variable, but it is still far from correct or meaningful modeling.

## VI. CONCLUSION AND FUTURE WORKS

In this paper, we presented a possible solution to identifying atmospheric parameters in exoplanets. The proposed

solution leverages both spectral data and auxiliary information available on the planets to produce a meaningful estimate of the parameters of interest. We show that our methodology outperforms the baseline model of reference by producing a balanced prediction regarding both posterior and spectral scores, allowing us to reach 8<sup>th</sup> place in the Ariel Machine Learning Challenge.

By qualitatively inspecting some of the predicted parameters, we observe how the model struggles to reconstruct some of them while performing rather well on others. We aim to improve the performance of the proposed solution by addressing this aspect and by exploring whether consistent patterns exist in the characteristics of the worse reconstructed exoplanets.

## ACKNOWLEDGMENT

This study was carried out within the FAIR - Future Artificial Intelligence Research and received funding from the European Union Next-GenerationEU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.3 – D.D. 1555 11/10/2022, PE00000013), as a part of the MALTO (MACHINE Learning @ poliTO) team, with partial support by SmartData@PoliTO center on Big Data and Data Science. This manuscript reflects only the authors’ views and opinions, neither the European Union nor the European Commission can be considered responsible for them.

## REFERENCES

- [1] N. M. Batalha, “Exploring exoplanet populations with nasa’s kepler mission,” *Proceedings of the National Academy of Sciences*, vol. 111, no. 35, pp. 12 647–12 654, 2014.
- [2] L. Kaltenegger, “How to characterize habitable worlds and signs of life,” *Annual Review of Astronomy and Astrophysics*, vol. 55, pp. 433–485, 2017.
- [3] R. I. Dawson and J. A. Johnson, “Origins of hot jupiters,” *Annual Review of Astronomy and Astrophysics*, vol. 56, pp. 175–221, 2018.
- [4] G. Tinetti, P. Drossart, P. Eccleston, P. Hartogh, A. Heske, J. Leconte, G. Micela, M. Ollivier, G. Pilbratt, L. Puig *et al.*, “The science of ariel (atmospheric remote-sensing infrared exoplanet large-survey),” in *Space Telescopes and Instrumentation 2016: Optical, Infrared, and Millimeter Wave*, vol. 9904. SPIE, 2016, pp. 658–667.
- [5] N. Madhusudhan, *Atmospheric Retrieval of Exoplanets*. Cham: Springer International Publishing, 2018, pp. 2153–2182. [Online]. Available: [https://doi.org/10.1007/978-3-319-55333-7\\_104](https://doi.org/10.1007/978-3-319-55333-7_104)
- [6] Q. Changeat and K. H. Yip, “Esa-ariel data challenge neurips 2022: introduction to exo-atmospheric studies and presentation of the atmospheric big challenge (abc) database,” *RAS Techniques and Instruments*, vol. 2, no. 1, pp. 45–61, 2023.
- [7] L. D. Deming and S. Seager, “Illusion and reality in the atmospheres of exoplanets,” *Journal of Geophysical Research: Planets*, vol. 122, no. 1, pp. 53–75, 2017.
- [8] M. D. Himes, J. Harrington, A. D. Cobb, A. G. Baydin, F. Soboczenski, M. D. O’Beirne, S. Zorzan, D. C. Wright, Z. Scheffer, S. D. Domagal-Goldman *et al.*, “Accurate machine-learning atmospheric retrieval via a neural-network surrogate model for radiative transfer,” *The Planetary Science Journal*, vol. 3, no. 4, p. 91, 2022.
- [9] T. Mikal-Evans, D. K. Sing, J. K. Barstow, T. Kataria, J. Goyal, N. Lewis, J. Taylor, N. J. Mayne, T. Daylan, H. R. Wakeford *et al.*, “Diurnal variations in the stratosphere of the ultrahot giant exoplanet wasp-121b,” *Nature Astronomy*, vol. 6, no. 4, pp. 471–479, 2022.

- [10] M. Mansfield, L. Wiser, K. B. Stevenson, P. Smith, M. R. Line, J. L. Bean, J. J. Fortney, V. Parmentier, E. M.-R. Kempton, J. Arcangeli *et al.*, “Confirmation of water absorption in the thermal emission spectrum of the hot jupiter wasp-77ab with hst/wfc3,” *The Astronomical Journal*, vol. 163, no. 6, p. 261, 2022.
- [11] Q. Changeat, “On spectroscopic phase-curve retrievals: H<sub>2</sub> dissociation and thermal inversion in the atmosphere of the ultrahot jupiter wasp-103 b,” *The Astronomical Journal*, vol. 163, no. 3, p. 106, 2022.
- [12] A. Boucher, A. Darveau-Bernier, S. Pelletier, D. Lafreniere, E. Artigau, N. J. Cook, R. Allart, M. Radica, R. Doyon, B. Benneke *et al.*, “Characterizing exoplanetary atmospheres at high resolution with spirou: Detection of water on hd 189733 b,” *The Astronomical Journal*, vol. 162, no. 6, p. 233, 2021.
- [13] P. Mollière, T. Stolker, S. Lacour, G. Otten, J. Shangquan, B. Charnay, T. Molyarova, M. Nowak, T. Henning, G.-D. Marleau *et al.*, “Retrieving scattering clouds and disequilibrium chemistry in the atmosphere of hr 8799e,” *Astronomy & Astrophysics*, vol. 640, p. A131, 2020.
- [14] C. Zhang, J. Bütepage, H. Kjellström, and S. Mandt, “Advances in variational inference,” *IEEE transactions on pattern analysis and machine intelligence*, vol. 41, no. 8, pp. 2008–2026, 2018.
- [15] J. P. Gardner, J. C. Mather, M. Clampin, R. Doyon, M. A. Greenhouse, H. B. Hammel, J. B. Hutchings, P. Jakobsen, S. J. Lilly, K. S. Long *et al.*, “The james webb space telescope,” *Space Science Reviews*, vol. 123, pp. 485–606, 2006.
- [16] M. Morvan, A. Tsiaras, N. Nikolaou, and I. P. Waldmann, “Pylightcurve-torch: a transit modeling package for deep learning applications in pytorch,” *Publications of the Astronomical Society of the Pacific*, vol. 133, no. 1021, p. 034505, 2021.
- [17] J. E. Krick, J. Fraine, J. Ingalls, and S. Deger, “Random forests applied to high-precision photometry analysis with spitzer irac,” *The Astronomical Journal*, vol. 160, no. 3, p. 99, 2020.
- [18] S. Lawler and B. Gladman, “Debris disks in kepler exoplanet systems,” *The Astrophysical Journal*, vol. 752, no. 1, p. 53, 2012.
- [19] A. Koudounas, F. Giobergia, and E. Baralis, “Time-of-flight cameras in space: Pose estimation with deep learning methodologies,” in *2022 IEEE 16th International Conference on Application of Information and Communication Technologies (AICT)*. IEEE, 2022, pp. 1–6.
- [20] F. A. Martinez, M. Min, I. Kamp, and P. I. Palmer, “Convolutional neural networks as an alternative to bayesian retrievals,” *arXiv preprint arXiv:2203.01236*, 2022.
- [21] K. H. Yip, Q. Changeat, B. Edwards, M. Morvan, K. L. Chubb, A. Tsiaras, I. P. Waldmann, and G. Tinetti, “On the compatibility of ground-based and space-based data: Wasp-96 b, an example,” *The Astronomical Journal*, vol. 161, no. 1, p. 4, 2020.
- [22] K. H. Yip, Q. Changeat, N. Nikolaou, M. Morvan, B. Edwards, I. P. Waldmann, and G. Tinetti, “Peeking inside the black box: Interpreting deep-learning models for exoplanet atmospheric retrievals,” *The Astronomical Journal*, vol. 162, no. 5, p. 195, 2021.
- [23] K. H. Yip, Q. Changeat, A. Al-Refaie, and I. Waldmann, “To sample or not to sample: Retrieving exoplanetary spectra with variational inference and normalising flows,” *arXiv preprint arXiv:2205.07037*, 2022.
- [24] R. Potthast, “A survey on sampling and probe methods for inverse problems,” *Inverse Problems*, vol. 22, no. 2, p. R1, 2006.
- [25] L. V. Mugnai, A. Al-Refaie, A. Bocchieri, Q. Changeat, E. Pascale, and G. Tinetti, “Alfnoor: Assessing the information content of ariel’s low-resolution spectra with planetary population studies,” *The Astronomical Journal*, vol. 162, no. 6, p. 288, 2021.
- [26] A. F. Al-Refaie, Q. Changeat, I. P. Waldmann, and G. Tinetti, “Taurex 3: A fast, dynamic, and extendable framework for retrievals,” *The Astrophysical Journal*, vol. 917, no. 1, p. 37, 2021.
- [27] L. V. Mugnai, E. Pascale, B. Edwards, A. Papageorgiou, and S. Sarkar, “Arielrad: the ariel radiometric model,” *Experimental Astronomy*, vol. 50, pp. 303–328, 2020.