

Normalized positive solutions for Schrödinger equations with potentials in unbounded domains

*Original*

Normalized positive solutions for Schrödinger equations with potentials in unbounded domains / Lancelotti, S.; Molle, R..  
- In: PROCEEDINGS OF THE ROYAL SOCIETY OF EDINBURGH. SECTION A. MATHEMATICS. - ISSN 0308-2105. -  
(2023), pp. 1-34. [10.1017/prm.2023.78]

*Availability:*

This version is available at: 11583/2984196 since: 2023-12-04T16:06:02Z

*Publisher:*

Cambridge University Press

*Published*

DOI:10.1017/prm.2023.78

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

Cambridge University Press postprint/Author's Accepted Manuscript

This article has been published in a revised form in PROCEEDINGS OF THE ROYAL SOCIETY OF EDINBURGH. SECTION A. MATHEMATICS <http://doi.org/10.1017/prm.2023.78>. This version is free to view and download for private research and study only. Not for re-distribution or re-use. © copyright holder.

(Article begins on next page)

# Normalized positive solutions for Schrödinger equations with potentials in unbounded domains

Sergio Lancelotti

*Dipartimento di Scienze Matematiche, Politecnico di Torino*

*Corso Duca degli Abruzzi n. 24, 10129 Torino, Italy*

`sergio.lancelotti@polito.it`

Riccardo Molle

*Dipartimento di Matematica, Università di Roma "Tor Vergata"*

*Via della Ricerca Scientifica n. 1, 00133 Roma, Italy*

`molle@mat.uniroma2.it`

July 3, 2023

## Abstract

The paper deals with the existence of positive solutions with prescribed  $L^2$  norm for the Schrödinger equation

$$-\Delta u + \lambda u + V(x)u = |u|^{p-2}u, \quad u \in H_0^1(\Omega), \quad \int_{\Omega} u^2 dx = \rho^2, \quad \lambda \in \mathbb{R},$$

where  $\Omega = \mathbb{R}^N$  or  $\mathbb{R}^N \setminus \Omega$  is a compact set,  $\rho > 0$ ,  $V \geq 0$  (also  $V \equiv 0$  is allowed),  $p \in (2, 2 + \frac{4}{N})$ . The existence of a positive solution  $\bar{u}$  is proved when  $V$  verifies a suitable decay assumption  $(D_\rho)$ , or if  $\|V\|_{L^q}$  is small, for some  $q \geq \frac{N}{2}$  ( $q > 1$  if  $N = 2$ ). No smallness assumption on  $V$  is required if the decay assumption  $(D_\rho)$  is fulfilled. There are no assumptions on the size of  $\mathbb{R}^N \setminus \Omega$ . The solution  $\bar{u}$  is a bound state and no ground state solution exists, up to the autonomous case  $V \equiv 0$  and  $\Omega = \mathbb{R}^N$ .

*A.M.S. Subject classification:* 35J10; 35J60; 35J20.

*Keywords:* Nonlinear Schrödinger equations; Normalized solutions; Exterior domains; Positive solutions.

## 1 Introduction and main results

In this paper we study a class of problems of the form

$$\begin{cases} -\Delta u + \lambda u + V(x)u = |u|^{p-2}u & \text{in } \Omega, \\ \lambda \in \mathbb{R}, \quad u \in H_0^1(\Omega), \quad \int_{\Omega} u^2 dx = \rho^2, \end{cases} \quad (P)$$

where  $\Omega = \mathbb{R}^N$  or  $\Omega$  is an exterior domain, that is  $\mathbb{R}^N \setminus \Omega$  is a compact set,  $N \geq 1$ ,  $V$  is a given potential and  $2 < p < 2 + 4/N$ , namely the nonlinearity is superlinear and *mass subcritical*. Here  $\lambda$

will arise as a Lagrange multiplier related to the *mass constraint*  $\|u\|_{L^2} = \rho > 0$ . We will focus on potentials that verify

$$\begin{aligned} V &\in L^q(\mathbb{R}^N), \quad q \in [\max(N/2, 1), \infty], \quad \text{with } q \neq 1 \text{ if } N = 2 \\ V(x) &\geq 0 \text{ a.e. in } \mathbb{R}^N. \end{aligned} \tag{1.1}$$

Problems of the form **(P)** arise from the nonlinear Schrödinger and Klein-Gordon equations

$$i\Phi_t + \Delta\Phi + f(\Phi) = 0 \tag{S}$$

$$\Phi_{tt} - \Delta\Phi + m\Phi = f(\Phi), \tag{KG}$$

where  $\Phi : \mathbb{R}^N \times (0, +\infty) \rightarrow \mathbb{C}$ . If  $f(re^{i\theta}) = f(r)e^{i\theta}$ ,  $r, \theta \in \mathbb{R}$ , then one can look for standing wave solutions of **(S)** and **(KG)**, namely solutions of the form

$$\Phi(x, t) = e^{i\lambda t}u(x), \quad x \in \mathbb{R}^N, \quad t > 0, \tag{1.2}$$

where  $u$  is a real function. In the model case  $f(\Phi) = |\Phi|^{p-2}\Phi$  we consider,  $u$  has to solve the equation in **(P)**. We refer the reader to [8–10] for more detailed physical motivations.

If the frequency  $\lambda$  in the ansatz (1.2) is fixed, and we consider the pure power model case with  $p \in (2, 2^*)$ , where  $2^* = \frac{2N}{(N-2)^+}$ , then looking for solutions of **(S)** and **(KG)** corresponds to looking for critical points of the action functional

$$E_\lambda(u) = \frac{1}{2} \int_{\mathbb{R}^N} (|\nabla u|^2 + [\lambda + V(x)]u^2) dx - \frac{1}{p} \int_{\mathbb{R}^N} |u|^p dx, \quad u \in H^1(\mathbb{R}^N).$$

A very large number of works are devoted to this unconstrained problem, we only refer the reader to [2, 11, 39] for a survey on almost classical results, and to the recent papers [16, 33] and references therein for new contributions.

Another point of view is to fix a priori the  $L^2$  norm of the solution. This point of view is particularly relevant because this quantity (the *mass* or the *charge* of the particle) is preserved along the time evolution. In this case, the solutions  $u$  correspond to the critical points of the energy functional

$$E(u) = \frac{1}{2} \int_{\mathbb{R}^N} [|\nabla u|^2 + V(x)u^2] dx - \frac{1}{p} \int_{\mathbb{R}^N} |u|^p dx \quad u \in H^1(\Omega),$$

constrained on

$$S_\rho := \{u \in H^1(\mathbb{R}^N) : \|u\|_2 = \rho\},$$

and the frequency  $\lambda$  arises as a Lagrange multiplier. Even if this fixed mass problem is classic (see for example [10]), only in the last decade it has been studied extensively and, in particular, very little has been done in the non-autonomous case (see [1, 5, 6, 24–26, 34, 41] and references therein).

We refer the reader to [17] for a recent analysis on the connections between the fixed mass and the fixed frequency points of view (see also [35] and references therein).

If  $2 + \frac{4}{N} < p < 2^*$ , the so called mass-supercritical regime, then  $E$  is not bounded from below on  $S_\rho$ , as follows by evaluating the functional over the fibers on  $S_\rho$  of the type  $u_t := t^{N/2}u_1(t)$ , for fixed  $u_1 \in S_\rho$  and  $t > 0$ . So the problem cannot be addressed by minimization. Jeanjean in the pioneering paper [25] analyzed the autonomous problem, for more general mass-supercritical nonlinearities, and he recognized a mountain pass structure, related to the above-introduced fibers. The method developed in [25] does not work in the non-autonomous case, even if the potential is radially symmetric. The non-autonomous case in the mass-supercritical regime has been studied in [6, 34], for not necessarily symmetrical potentials. In [6] the authors found a mountain-pass solution if  $V \geq 0$  is suitably small,

while [34] concerns the case  $V \leq 0$  and the existence of two solutions is proved when  $V$  is suitably small and the operator  $-\Delta + V$  is not positive-definite.

In the mass-subcritical regime  $2 < p < 2 + \frac{4}{N}$ , the functional  $E$  is bounded from below on  $S_\rho$  by the Gagliardo-Nirenberg inequality (see (3.4), (3.5)). Ikoma e Miyamoto in [24] considered more general non-autonomous problems of the type

$$\begin{cases} -\Delta u + \lambda u + V(x)u = f(x, u) & \text{(eq)} \\ u \in H_0^1(\Omega), u \in S_\rho, \lambda \in \mathbb{R}, \end{cases} \quad (1.3)$$

where  $\Omega = \mathbb{R}^N$ , and the nonlinearity  $f(x, u) = f(u)$  has sub-linear growth in 0 and mass-subcritical growth at infinity, that is

$$\lim_{s \rightarrow 0} \frac{f(s)}{s} = \lim_{|s| \rightarrow \infty} \frac{f(s)}{|s|^{1+4/N}} = 0.$$

If the shape of the potential is  $V \leq 0$  and  $V(x) \rightarrow 0$  as  $|x| \rightarrow \infty$ , and both  $f$  and  $V$  satisfy various technical assumptions, then the authors prove, by concentration-compactness arguments, that there exists  $\rho_0 > 0$  such that the energy functional related to (1.3) has a global minimum in  $S_\rho$  for  $\rho > \rho_0$ , and the global minimum is not attained if  $0 < \rho < \rho_0$ . Moreover, some sufficient conditions on  $f$  and  $V$  are provided to get  $\rho_0 = 0$ .

In the more recent paper [1], Alves and Ji considered in problem (P) some classes of potentials  $V$  where a global minimum for  $E$  on  $S_\rho$  exists, for suitable  $\rho$ . Namely, in [1] the authors work on potentials  $V$ , not required vanishing at infinity, that verify one of these assumptions:  $(V_1)$   $\inf_{\mathbb{R}^N} V < \liminf_{|x| \rightarrow \infty} V(x)$ ,  $(V_2)$  there exists  $\mu_0 > 0$  such that  $\text{meas}\{V > \mu_0\} < \infty$  and  $\text{int}(V^{-1}(0)) \neq \emptyset$ ,  $(V_3)$   $V$  is 1-periodic in each variable,  $(V_4)$   $V$  is asymptotically 1-periodic, that is there exists a 1-periodic function  $V_p : \mathbb{R}^N \rightarrow \mathbb{R}$  such that

$$V(x) \leq V_p(x), \quad \forall x \in \mathbb{R}^N, \quad \text{and} \quad \lim_{|x| \rightarrow \infty} |V_p(x) - V(x)| = 0.$$

After submitting this paper, the interesting works [37, 40] have been brought to our attention. In those papers, non homogeneous nonlinearities as in (1.3) are studied.

Yang, Qi and Zou in [40] improve the result of [24]. They consider problem (1.3) with  $\Omega = \mathbb{R}^N$ ,  $V(x) \leq \lim_{|x| \rightarrow \infty} V(x) =: V_\infty \in (-\infty, +\infty]$  and  $f \in \mathcal{C}(\Omega \times \mathbb{R}, \mathbb{R})$  satisfies Berestycki-Lions type conditions with mass sub-critical growth. Moreover, some technical assumptions used in [24] are removed. In the coercive case  $V_\infty = +\infty$ , the existence of a ground state solution is proved for every  $\rho > 0$ , while for  $V_\infty < +\infty$  the authors show that still  $\rho_0 \geq 0$  exists such that there is a ground state solution for  $\rho > \rho_0$  and it does not exist for  $0 < \rho < \rho_0$ .

Some results in [37] concern symmetric exterior domains in some mass-supercritical regimes. Therein, Song studies (1.3) when  $\Omega$  is a bounded starshaped domain or the complementary of a ball in  $\mathbb{R}^N$  with  $N \geq 2$ ,  $V \equiv 0$  and  $f \in \mathcal{C}^1(\Omega \times \mathbb{R}, \mathbb{R})$  is a superlinear and mass-supercritical nonlinearity. When  $\Omega$  is the complementary of a ball and  $f(\cdot, u)$  is radially symmetric, the author proves that there exists a sequence  $((\lambda_k, u_k))_k$  in  $H_0^1(\Omega) \times \mathbb{R}$  of solutions of (1.3)(eq) such that  $\lambda_k \rightarrow 0$  and  $|u_k|_2 \rightarrow \infty$ , while there exists no sequence  $((\lambda_k, u_k))_k$  in  $H_0^1(\Omega) \times \mathbb{R}$  of solutions of (1.3)(eq) such that  $\lambda_k \rightarrow 0$  and  $|u_k|_2 \rightarrow 0$ . Moreover, if (1.3)(eq) has at most one positive radial solution in  $H_0^1(\Omega)$  for every  $\lambda > 0$ , then (1.3) has a positive radial solution for large  $\rho$ .

Here, we are concerned with the mass-subcritical case when  $V \geq 0$  so no minimum solution exists, up to the autonomous case  $V \equiv 0$  and  $\Omega = \mathbb{R}^N$  (see Proposition 1.4). Moreover, we focus on domains that can be not only  $\mathbb{R}^N$  but also general exterior domains, answering the question raised in [34] for mass-supercritical problems, whether the existence of bound state solutions can be treated in exterior domains as in the whole space. To the best of our knowledge, these issues are only partially addressed in previous papers. About exterior domains in the mass-subcritical case, it is worth mentioning [41],

where the authors consider the autonomous problem  $V \equiv 0$  in dimension  $N \geq 3$  and they found the existence of a bound state solution if the size of “hole”  $\mathbb{R}^N \setminus \Omega$  is small.

The main results are the following:

**Theorem 1.1.** *Let  $N \geq 2$ ,  $\Omega = \mathbb{R}^N$  or  $\mathbb{R}^N \setminus \Omega$  compact,  $\rho > 0$ . If  $V$  satisfies (1.1) and*

$$\int_{\mathbb{R}^N} V(x) |x|^{N-1} e^{d_\rho |x|} dx < \infty, \quad (D_\rho)$$

where

$$d_\rho = \left[ 2^{(1 - \frac{p-2}{4-N(p-2)})} \sqrt{\lambda_1} \right] \rho^{\frac{p-2}{4-N(p-2)}}, \quad (1.4)$$

(see (2.5) for the constant  $\lambda_1$ ), then there exists a solution  $(\lambda, \bar{u})$  of (P) such that  $\lambda > 0$  and  $\bar{u} \geq 0$ .

**Theorem 1.2.** *Let  $N \geq 2$ ,  $\Omega = \mathbb{R}^N$  or  $\mathbb{R}^N \setminus \Omega$  compact,  $\rho > 0$ . There exists  $L = L(q, \Omega, \rho) > 0$  such that if  $V$  satisfies (1.1), with*

$$V(x) \longrightarrow 0 \quad \text{as } |x| \rightarrow \infty \quad \text{if } V \in L^\infty(\Omega), \quad (1.5)$$

and  $\|V\|_q < L$ , then problem (P) has a solution  $(\lambda, \bar{u})$ , verifying  $\lambda > 0$  and  $\bar{u} \geq 0$ .

The case  $N = 1$  has its own specificity and will be dealt with in Section 6.

A priori, a nonnegative solution  $\bar{u}$  of (P) belongs to  $H^1(\Omega)$ , so we cannot say that  $\bar{u} > 0, \forall x \in \Omega$ . Anyway, under mild assumptions,  $\bar{u}$  turns out to be sufficiently regular to apply the Harnack inequality and therefore to be pointwise positive. In the following Proposition, we collect some regularity properties. Its proof is almost standard, so we only outline it in the Appendix.

**Proposition 1.3.** *Let  $\bar{u}$  be a solution of (P). If  $N \geq 2$ , then*

- a) *if  $V \in L^q_{\text{loc}}(\Omega)$  for some  $q > \frac{N}{2}$ , then  $\bar{u} \in \mathcal{C}^{0,\alpha}_{\text{loc}}(\Omega)$ , and if  $\bar{u}$  is nonnegative then  $\bar{u}(x) > 0 \forall x \in \Omega$ ;*
- b) *if  $V \in L^q_{\text{loc}}(\Omega)$  for some  $q > N$ , then  $\bar{u} \in \mathcal{C}^{1,\alpha}_{\text{loc}}(\Omega)$ ;*
- c) *if  $V \in \mathcal{C}^{0,\alpha}_{\text{loc}}(\Omega)$ , then  $\bar{u} \in \mathcal{C}^2(\Omega)$  and it is a classical solution.*

If  $N = 1$ , then

- d) *if  $V \in L^1_{\text{loc}}(\Omega)$ , then  $\bar{u}$  is continuously differentiable and if  $\bar{u}$  is nonnegative then  $\bar{u}(x) > 0 \forall x \in \Omega$ ;*
- e) *if  $V \in L^q_{\text{loc}}(\Omega)$  for some  $q > 1$ , then  $\bar{u} \in \mathcal{C}^{1,\alpha}_{\text{loc}}(\Omega)$ .*

By Proposition 1.3, the solutions given by Theorem 1.1 and 1.2 are positive if assumption (1.1) holds with  $q > \frac{N}{2}$ .

The solution we find is a bound state solution. Indeed, in Section 2 we verify that no ground state solution can exist:

**Proposition 1.4.** *Assume that  $V$  satisfies the assumptions of Theorem 1.1 or of Theorem 1.2. If  $V \not\equiv 0$  or  $\Omega \neq \mathbb{R}^N$ , then problem (P) has no ground state solution.*

Actually, if  $V$  satisfies both the assumptions of Theorem 1.1 and of Theorem 1.2, then the solutions provided by the Theorems are found exploiting the same topological configuration so it is reasonable to expect that they are the same solution. Moreover, the topological characterization suggests that they have Morse index  $N$ .

In (1.1), the assumption that  $V$  vanishes at infinity is necessary to get the compactness condition and cannot be dropped, by the following nonexistence result (see [19], [34, Proposition 1.10], [12, Theorem 1.1]).

**Proposition 1.5.** *Let  $p \in (2, 2^*)$ ,  $V \in L^\infty(\mathbb{R}^N)$  and assume that there exists  $\frac{\partial V}{\partial \nu} \in L^\ell(\mathbb{R}^N)$  for some  $\nu \in \mathbb{R}^N \setminus \{0\}$  and  $\ell \in [\max(1, \frac{N}{2}), +\infty]$ ,  $\ell \neq 1$  if  $N = 2$ . If  $\frac{\partial V}{\partial \nu} \geq 0$  and  $\frac{\partial V}{\partial \nu} \not\equiv 0$ , then problem*

$$-\Delta u + \lambda u - V(x)u = |u|^{p-2}u \quad u \in S_\rho, \quad \lambda \in \mathbb{R} \quad (1.6)$$

has no solutions in  $C^1(\mathbb{R}^N) \cap W^{2,2}(\mathbb{R}^N)$ .

See Remark 6.2 and Proposition 6.3 for nonexistence results in the 1-dimensional case.

Some remarks are in order, concerning the decay assumption  $(D_\rho)$  in Theorem 1.1. It does not require any smallness assumption on  $V$  and moreover does not depend on the domain, hence if it is verified then problem  $(P)$  has a solution for every exterior domain  $\Omega$ .

On the other hand,  $(D_\rho)$  depends on  $\rho$  by (1.4), where  $d_\rho \rightarrow \infty$  as  $\rho \rightarrow \infty$  because  $p < 2 + \frac{4}{N}$ . As a consequence, a potential that verifies  $(D_\rho)$  for every  $\rho > 0$  has to decay faster than any exponential.

*Remark 1.6.* By the proof of Theorem 1.2 we verify the existence of a constant  $\bar{L} = \bar{L}(N) > 0$  such that

$$L(N/2, \mathbb{R}^N, \rho) = \bar{L} \quad \forall \rho > 0. \quad (1.7)$$

Instead, for  $q > \frac{N}{2}$ ,

$$\lim_{\rho \rightarrow 0} L(q, \mathbb{R}^N, \rho) = 0, \quad \lim_{\rho \rightarrow \infty} L(q, \mathbb{R}^N, \rho) = \infty. \quad (1.8)$$

As a consequence, if  $\|V\|_{N/2}$  is suitably small, then problem  $(P)$  in  $\mathbb{R}^N$  has a solution for every  $\rho > 0$ , while if  $V \in L^q(\mathbb{R}^N)$ , for  $q > N/2$ , then there exists  $\bar{\rho} = \bar{\rho}(\|V\|_q) \geq 0$  such that problem  $(P)$  has a solution for every  $\rho > \bar{\rho}$  (see (5.14)).

In Theorems 1.1. and 1.2 of [34] a mountain pass solution is found in the mass supercritical case with negative potential, under smallness assumptions on  $V$  similar to the ones considered here. In that case, if  $q = N/2$  there is a uniform bound as in (1.7) while for  $q > N/2$  the limits in (1.8) are reversed.

If  $\rho$  is fixed, we will observe that

$$\lim_{r(\Omega) \rightarrow \infty} L(q, \Omega, \rho) = 0,$$

where

$$r(\Omega) = \sup\{r \in \mathbb{R}^+ : B_r(y) \subset \mathbb{R}^N \setminus \Omega \text{ for some } y \in \mathbb{R}^N\} \quad (1.9)$$

(see Remark 5.1). Hence, there is no potential  $V \not\equiv 0$  such that Theorem 1.2 applies for every exterior domain  $\Omega$ , for  $\rho > 0$  fixed.

If we consider  $V \equiv 0$ , then both Theorems 1.1 and 1.2 apply and provide the following result.

**Corollary 1.7.** *Let  $N \geq 2$ ,  $\mathbb{R}^N \setminus \Omega$  compact and  $\rho > 0$ , then there exist  $\lambda > 0$  and  $\bar{u} \in C^2(\Omega)$ ,  $\bar{u}(x) > 0$ ,  $\forall x \in \Omega$ , such that*

$$\begin{cases} -\Delta \bar{u} + \lambda \bar{u} = \bar{u}^{p-1} & \text{in } \Omega, \\ \lambda \in \mathbb{R}, \quad \bar{u} \in H_0^1(\Omega), \quad \int_{\Omega} \bar{u}^2 dx = \rho^2. \end{cases}$$

Corollary 1.7 extends the result of [41] to the dimension 2 and to every exterior domains. The proof in [41] cannot be extended to this more general framework because it does not work in dimension 2 and when the size of  $\mathbb{R}^N \setminus \Omega$  is large.

We prove Theorems 1.1 and 1.2 by variational methods, looking for bound state solutions. The analysis of the compactness presents a lot of difficulties related to the unboundedness of the domain, that is not assumed to be symmetric. In order to recover a local compactness condition, we first see that the Lagrange multiplier  $\lambda$  related to a Palais-Smale sequence in a negative range is positive, and then we employ a splitting Lemma from [7] for PS sequences of  $E_\lambda$ . With this decomposition in hands,

we perform in Proposition 3.1 a sharp fine estimate of the first energy interval  $I$  above the infimum where the compactness condition holds (see Remark 3.4).

The topological argument relies on min-max techniques that make use of a barycentric map and the Brouwer degree. Since we have no smallness assumption on  $\mathbb{R}^N \setminus \Omega$ , and in Theorem 1.1 we have no bound on any Lebesgue norm of the potential, a major difficulty is to work in the compactness interval  $I$ , in the min-max procedure. To overcome this problem, we will proceed by analyzing the energy interaction of positive solutions  $w_1$  and  $w_2$  of some suitable “problems at infinity”, such that  $\|w_1\|_{L^2}^2 + \|w_2\|_{L^2}^2 = \rho^2$ . This idea is inspired by the unconstrained case (see [14, 28]), where the problem at infinity is univocally determined by the choice of  $\lambda$ . Here, the need to choose different functions  $w_1, w_2$  prevents the use of the arguments developed in [14, 28]) and requires the introduction of different and more refined estimates.

To verify that the solutions we find do not change sign, we prove in Proposition 2.2 that the energy of every solution that changes sign is not in the energy interval we are working in. In particular, Proposition 2.2 and Corollary 2.3 give information also on changing sign solutions of the autonomous problem in  $\mathbb{R}^N$ . We point out that to get nonnegative solutions here we could exploit the symmetry of the functional and work near the cone of the positive functions, by using [21, Theorem 4.5] and proceeding as in [6]. The advantage of this other approach is a simplification of the proof of the compactness condition, because the lack of compactness in such a case comes only from the positive solution of the problem at infinity. On the other hand, the approach we employ here allows us both to bound from below the energy of the changing sign solutions and to gain a more general analysis of the Palais-Smale sequences.

The paper is organized as follows: in Section 2 we introduce some preliminary results, in Section 3 we prove the local compactness condition and Section 4 is devoted to the sharp energy estimates that are necessary in Section 5 to prove Theorems 1.1 and 1.2, Section 6 concerns the case  $N = 1$  and in the Appendix we give a sketch of the proof of the regularity proposition, with some references for detailed proofs.

## 2 Notations, variational framework and preliminary results

Throughout the paper we make use of the following notation:

- $2_c := 2 + \frac{4}{N}$ ,  $s = \frac{2}{N} \frac{p-2}{2_c-p}$ .
- $L^q(\mathcal{O})$ ,  $1 \leq q \leq \infty$ ,  $\mathcal{O} \subseteq \mathbb{R}^N$  a measurable set, denotes the Lebesgue space, the norm in  $L^q(\mathcal{O})$  is denoted by  $|\cdot|_q$  if there is no ambiguity.
- For  $u \in H_0^1(\Omega)$  we denote by  $u$  also the function in  $H^1(\mathbb{R}^N)$  obtained setting  $u \equiv 0$  in  $\mathbb{R}^N \setminus \Omega$ .
- For any  $R > 0$  and for any  $z \in \mathbb{R}^N$ ,  $B_R(z)$  denotes the closed ball of radius  $R$  centered at  $z$ , and for any measurable set  $\mathcal{O} \subset \mathbb{R}^N$ ,  $|\mathcal{O}|$  denotes its Lebesgue measure.
- $H^1(\mathbb{R}^N)$  is the usual Sobolev space endowed with the standard norm

$$\|u\| := \left[ \int_{\mathbb{R}^N} (|\nabla u|^2 + u^2) dx \right]^{\frac{1}{2}}.$$

- $c, c', C, C', C_i, \dots$  denote various positive constants that can also vary from one line to another.
- $o(f)$  and  $O(f)$  denote the usual Landau notations:  $\frac{o(f)}{f} \rightarrow 0$  as  $f \rightarrow 0$  and  $|O(f)| \leq C|f|$  for some positive constant  $C$ .

We will find solutions  $\bar{u}$  of problem  $(P)$  as critical points of the functional  $E$  constrained on  $S_\rho$ . If  $\lambda$  is the Lagrange multiplier related to  $\bar{u}$ , then  $\bar{u}$  is also a free critical point of the related free functional  $E_\lambda$

Let us assume  $\lambda > 0$  and recall some well-known properties of the limit problem, for  $V \equiv 0$  and  $\Omega = \mathbb{R}^N$ ,

$$\begin{cases} -\Delta u + \lambda_\infty u = |u|^{p-2}u & \text{in } \mathbb{R}^N, \\ \lambda_\infty \in \mathbb{R}, \quad u \in S_\rho. \end{cases} \quad (P_\infty)$$

$(P_\infty)$  has a unique positive solution  $w \in H^1(\mathbb{R}^N)$ , up to translations, which is radial, radially decreasing, and belongs to  $C^2(\mathbb{R}^N)$ . The function  $w$  verifies the minimality property

$$m := E_\infty(w) = \min_{u \in S_\rho} E_\infty(u), \quad (2.1)$$

where

$$E_\infty(u) = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 dx - \frac{1}{p} \int_{\mathbb{R}^N} |u|^p dx.$$

Correspondingly, the solutions of  $(P_\infty)$  are also free critical points of the limit functional

$$E_{\lambda,\infty}(u) = E_\infty(u) + \frac{\lambda}{2} \int_{\mathbb{R}^N} u^2 dx \quad u \in H^1(\mathbb{R}^N).$$

Moreover,

$$m < 0 \quad \text{and} \quad \lambda_\infty > 0 \quad (2.2)$$

and there exists  $c_1 > 0$  such that

$$w(|x|) e^{\sqrt{\lambda_\infty}|x|} |x|^{\frac{N-1}{2}} \rightarrow c_1 \quad \text{as } |x| \rightarrow \infty, \quad (2.3)$$

$$w'(|x|) e^{\sqrt{\lambda_\infty}|x|} |x|^{\frac{N-1}{2}} \rightarrow -c_1 \sqrt{\lambda_\infty} \quad \text{as } |x| \rightarrow \infty. \quad (2.4)$$

Inequality  $m < 0$  in (2.2) follows choosing  $u_1 \in S_\rho$  and testing (2.1) on  $u_t := t^{N/2} u_1(t) \in S_\rho$ ,  $t > 0$ , taking into account  $p < 2_c$ . The positivity of  $\lambda_\infty$  comes from Pohozaev and Nehari identities (see also Proposition 2.1). For the properties of  $w$  we refer the reader to [4, 8, 22, 27, 38].

For any  $k > 0$ , let us denote by  $w_{k\rho^2}$  the positive solution of

$$\begin{cases} -\Delta u + \lambda_{k\rho^2} u = |u|^{p-2}u & \text{in } \mathbb{R}^N, \\ \lambda_{k\rho^2} \in \mathbb{R}, \quad u \in S_{\sqrt{k}\rho}, \end{cases} \quad (2.5)$$

where  $w_{k\rho^2}$  verifies

$$m_{k\rho^2} := E_\infty(w_{k\rho^2}) = \inf_{u \in S_{\sqrt{k}\rho}} E_\infty(u). \quad (2.6)$$

It turns out that

$$w_{k\rho^2}(x) = k^{\frac{s}{p-2}} w(k^{\frac{s}{2}} x), \quad m_{k\rho^2} = k^{1+s} m < 0, \quad \lambda_{k\rho^2} = k^s \lambda_\infty > 0 \quad (2.7)$$

where

$$s := \frac{2}{N} \frac{p-2}{2_c-p}$$

(see notations). From  $p < 2_c$  it follows  $s > 0$ . By (2.7), in particular we have

$$\lambda_{\rho^2} = \lambda_\infty = \rho^s \lambda_1. \quad (2.8)$$

For  $k = 0$  we set  $w_0 = 0$  and for  $k = 1$  we simply write  $w_{\rho^2} = w$ .

Moreover, for any  $k > 0$ , by (2.3), (2.4) and (2.7) we have

$$w_{k\rho^2}(|x|) e^{\sqrt{k^s \lambda_\infty}|x|} |x|^{\frac{N-1}{2}} \rightarrow c_k \quad \text{as } |x| \rightarrow +\infty, \quad (2.9)$$

$$w'_{k\rho^2}(|x|) e^{\sqrt{k^s \lambda_\infty} |x|} |x|^{\frac{N-1}{2}} \rightarrow -c_k \sqrt{k^s \lambda_\infty} \quad \text{as } |x| \rightarrow +\infty \quad (2.10)$$

where

$$c_k = c_1 k^{s(\frac{1}{p-2} - \frac{N-1}{4})}.$$

From  $p < 2_c$  there follows  $\frac{1}{p-2} - \frac{N-1}{4} > 0$ , so that

$$c_k \rightarrow 0 \quad \text{as } k \rightarrow 0, \quad c_k \rightarrow c_1 \quad \text{as } k \rightarrow 1.$$

*Proof of Proposition 1.4.* Let us prove that

$$\inf_{S_\rho} E = m \quad (2.11)$$

and the infimum is not achieved.

Obviously we have that  $\inf_{S_\rho} E \geq m$ . To show that the equality holds, let us consider the sequence  $(u_n)_n$  defined by  $u_n = w(x - y_n)$ , where  $(y_n)_n$  is a sequence in  $\mathbb{R}^N$  such that  $|y_n| \rightarrow +\infty$ . By (1.1) and (2.3), and taking into account  $(D_\rho)$  or (1.5), we have that

$$\lim_{n \rightarrow +\infty} \int_{\mathbb{R}^N} V(x) u_n^2(x) dx = 0,$$

which implies  $\lim_{n \rightarrow +\infty} E(u_n) = m$ .

Now, assume by contradiction that the infimum  $\inf_{S_\rho} E$  is achieved by a function  $\bar{u}$ . Then, by

$$m \leq E_\infty(\bar{u}) \leq E(\bar{u}) = m, \quad \bar{u} \in S_\rho, \quad (2.12)$$

and by the uniqueness of the minimizers of (2.1), we should have  $\bar{u} = \pm w(x - \bar{y})$  for a suitable  $\bar{y} \in \mathbb{R}^N$ . Since  $w(x - \bar{y}) > 0$  for all  $x \in \mathbb{R}^N$ , we can deduce that  $\Omega$  must be the entire space  $\mathbb{R}^N$  and by (2.12) we get

$$\int_{\mathbb{R}^N} V(x) \bar{u}^2(x) dx = 0,$$

that implies  $V \equiv 0$  because  $V \geq 0$ . So we are in contradiction with  $\Omega \neq \mathbb{R}^N$  or  $V \not\equiv 0$ . □

Next proposition states the positivity of the Lagrange multipliers in (P).

**Proposition 2.1.** *Assume that  $u \in S_\rho$  and  $\lambda \in \mathbb{R}$  solve (P). If  $E(u) < 0$ , then  $\lambda > 0$ .*

*Proof.* We have

$$\int_{\Omega} \nabla u \cdot \nabla v dx + \lambda \int_{\Omega} uv dx + \int_{\Omega} V(x) uv dx - \int_{\Omega} |u|^{p-2} uv dx = 0 \quad \forall v \in H_0^1(\Omega).$$

Then,

$$\lambda \rho^2 = |u|_p^p - |\nabla u|_2^2 - \int_{\Omega} V(x) u^2 dx = -pE(u) + \frac{p-2}{2} |\nabla u|_2^2 + \frac{p-2}{2} \int_{\Omega} V(x) u^2 dx > 0. \quad (2.13)$$

□

Before concluding this section, let us find out some features of the changing sign solutions.

**Proposition 2.2.** *Let  $u \in S_\rho$  be a solution of (P) ((P<sub>∞</sub>)). If  $u$  changes sign, then  $E(u) > 2^{-s}m$  ( $E_\infty(u) > 2^{-s}m$ ).*

*Proof.* Since  $m < 0$ , we can assume  $E(u) < 0$ . According to Proposition 2.1, let  $\lambda > 0$  be the Lagrange multiplier corresponding to the solution  $u$  and let  $w_\infty \in H^1(\mathbb{R}^N)$  be the positive solution of  $-\Delta v + \lambda v = v^{p-1}$ . By (2.7),

$$E_\infty(w_\infty) = \min_{S_{|w_\infty|_2}} E_\infty = \left( \frac{|w_\infty|_2^2}{\rho^2} \right)^{1+s} m. \quad (2.14)$$

Let us write  $u = u^+ - u^-$ , where  $u^\pm = \max\{\pm u, 0\}$ ,  $\rho^2 = |u^+|_2^2 + |u^-|_2^2$ . Since  $u^+$  and  $u^-$  are on the Nehari manifold  $\mathcal{N}$  corresponding to  $E_\lambda$ , then  $(E_\lambda)|_{\mathcal{N}}(u^\pm) > E_{\lambda,\infty}(w_\infty)$  (see [7] and [31, Proposition 2.1]). Namely, by (2.14),

$$E(u^\pm) > \left( \frac{|w_\infty|_2^2}{\rho^2} \right)^{1+s} m + \frac{\lambda}{2} (|w_\infty|_2^2 - |u^\pm|_2^2).$$

Hence, the claim read as

$$E(u) = E(u^+) + E(u^-) > 2 \left( \frac{|w_\infty|_2^2}{\rho^2} \right)^{1+s} m + \frac{\lambda}{2} (2|w_\infty|_2^2 - \rho^2) \geq 2^{-s} m,$$

and, equivalently, as

$$\left[ \left( \frac{2|w_\infty|_2^2}{\rho^2} \right)^{1+s} - 1 \right] 2^{-s} m + \frac{\lambda}{2} \rho^2 \left[ \left( \frac{2|w_\infty|_2^2}{\rho^2} \right) - 1 \right] \geq 0. \quad (2.15)$$

Since  $(\lambda, w_\infty)$  solves (2.5) for  $k = \frac{|w_\infty|_2^2}{\rho^2}$ , the following system is fulfilled

$$\begin{cases} \frac{1}{2} |\nabla w_\infty|_2^2 - \frac{1}{p} |w_\infty|_p^p = m_{|w_\infty|_2^2} & \text{Energy of } w_\infty \\ |\nabla w_\infty|_2^2 + \lambda |w_\infty|_2^2 - |w_\infty|_p^p = 0 & \text{Nehari} \\ \frac{N-2}{2} |\nabla w_\infty|_2^2 + \frac{N}{2} \lambda |w_\infty|_2^2 - \frac{N}{p} |w_\infty|_p^p = 0 & \text{Pohozaev.} \end{cases} \quad (2.16)$$

Solving (2.16) and taking into account (2.7) we get

$$\frac{\lambda}{2} \rho^2 = - \frac{2p - N(p-2)}{4 - N(p-2)} \left( \frac{|w_\infty|_2^2}{\rho^2} \right)^s m. \quad (2.17)$$

Putting (2.17) in (2.15), we have to verify

$$\left[ \left( \frac{2|w_\infty|_2^2}{\rho^2} \right)^{1+s} - 1 \right] - \frac{2p - N(p-2)}{4 - N(p-2)} \left( \frac{2|w_\infty|_2^2}{\rho^2} \right)^s \left[ \left( \frac{2|w_\infty|_2^2}{\rho^2} \right) - 1 \right] \leq 0.$$

Then the assertion is proved because  $\frac{2p-N(p-2)}{4-N(p-2)} = 1 + s$  and

$$[t^{1+s} - 1] - (1+s)t^s[t-1] \leq 0 \quad \forall t \geq 0.$$

□

**Corollary 2.3.** *If  $u$  is a changing sign solution of*

$$-\Delta v + \lambda v = |v|^{p-2}v \quad v \in H^1(\mathbb{R}^N), \quad (2.18)$$

*then  $|u|_2^2 > 2|w_\infty|_2^2$ , where  $w_\infty$  denotes the positive solution of (2.18).*

Clearly, here  $\lambda > 0$  by Pohozaev identity.

*Proof.* Let us call  $E(u) = M$ . From system (2.16) applied to  $u$  and  $w_\infty$  it follows

$$|u|_2^2 = 2 \frac{1+s}{\lambda} (-M), \quad |w_\infty|_2^2 = 2 \frac{1+s}{\lambda} \left( -m_{|w_\infty|_2^2} \right). \quad (2.19)$$

By Proposition 2.2,  $M > 2^{-s} m_{|u|_2^2}$ , then, by (2.7) and (2.19),

$$|u|_2^2 < 2 \frac{1+s}{\lambda} 2^{-s} \left( -m_{|u|_2^2} \right) = 2 \frac{1+s}{\lambda} 2^{-s} \left( \frac{|u|_2^2}{|w_\infty|_2^2} \right)^{1+s} \left( -m_{|w_\infty|_2^2} \right) = 2^{-s} |w_\infty|_2^2 \left( \frac{|u|_2^2}{|w_\infty|_2^2} \right)^{1+s},$$

that completes the proof.  $\square$

### 3 The compactness condition

**Proposition 3.1.** *Let  $(u_n)_n$  be a Palais-Smale sequence at the level  $c$  for  $E$  constrained on  $S_\rho$ . If  $c \in (m, 2^{-s}m)$  then there exists a critical point  $u_0 \in S_\rho$  such that  $u_n \rightarrow u_0$ , as  $n \rightarrow \infty$ .*

To prove Proposition 3.1, we will use the well known splitting lemma of Benci and Cerami for the unconstrained problem ([7, Lemma 3.1]).

**Lemma 3.2.** *Let  $\lambda > 0$  and let  $(u_n)_n$  in  $H^1(\mathbb{R}^N)$  be a Palais-Smale sequence for  $E_\lambda$ . Then there exist a critical point  $u_0$  of  $E_\lambda$ , an integer  $h \geq 0$ ,  $h$  non-trivial solutions  $w^1, \dots, w^h \in H^1(\mathbb{R}^N)$  to the limit equation*

$$-\Delta v + \lambda v = |v|^{p-2}v$$

and  $h$  sequences  $(y_n^j)_n \subset \mathbb{R}^N$ ,  $1 \leq j \leq h$ , such that  $|y_n^j| \rightarrow \infty$  as  $n \rightarrow \infty$ , and

$$u_n = u_0 + \sum_{j=1}^h w^j(\cdot - y_n^j) + o(1) \quad \text{strongly in } H^1(\mathbb{R}^N), \quad (3.1)$$

up to a subsequence. Moreover, we have

$$E_\lambda(u_n) \rightarrow E_\lambda(u_0) + \sum_{j=1}^h E_{\lambda, \infty}(w^j) \quad \text{as } n \rightarrow \infty, \quad (3.2)$$

and

$$|u_n|_2^2 = |u_0|_2^2 + \sum_{j=1}^h |w^j|_2^2 + o(1). \quad (3.3)$$

*Proof of Proposition 3.1.* We claim that  $(u_n)_n$  is bounded in  $H_0^1(\Omega)$ . Indeed,  $u_n \in S_\rho$ ,  $\forall n \in \mathbb{N}$ , and by the Gagliardo-Nirenberg inequality

$$|u|_p^p \leq C_{GN} |u|_2^{p - \frac{N(p-2)}{2}} |\nabla u|_2^{\frac{N(p-2)}{2}}. \quad (3.4)$$

(see [29, Theorem 12.83]) we have

$$c + o(1) = E(u_n) \geq \frac{1}{2} |\nabla u_n|_2^2 - C |\nabla u_n|_2^{\frac{N(p-2)}{2}}, \quad (3.5)$$

with  $\frac{N(p-2)}{2} < 2$  because  $p < 2_c$ .

Since  $(u_n)_n$  is a constrained PS-sequence, there exists a sequence  $(\lambda_n)_n$  in  $\mathbb{R}$  such that

$$\int_{\Omega} \nabla u_n \cdot \nabla v \, dx + \lambda_n \int_{\Omega} u_n v \, dx + \int_{\Omega} V(x) u_n v \, dx - \int_{\Omega} |u_n|^{p-2} u_n v \, dx = o(1) \|v\| \quad \forall v \in H_0^1(\Omega). \quad (3.6)$$

Setting  $v = u_n$  in (3.6), and taking into account that  $(u_n)_n$  is bounded in  $H^1$ , we can argue as in (2.13) obtaining

$$\lambda_n \rho^2 = -pE(u_n) + \frac{p-2}{2} |\nabla u_n|_2^2 + \frac{p-2}{2} \int_{\Omega} V(x) u_n^2 \, dx + o(1) \geq -pc + o(1) > 0, \quad (3.7)$$

for large  $n$ . Since  $(u_n)_n$  is bounded in  $H_0^1(\Omega)$ , the first relation in (3.7) implies that the sequence  $(\lambda_n)_n$  is bounded. Moreover, from (3.7) we infer also that  $\lambda_n \geq c > 0$ , for a suitable constant  $c$ . Then we can assume that  $\lambda_n \rightarrow \lambda > 0$ . Hence, by (3.6) we are in position to apply Lemma 3.2 and we can decompose  $u_n$  according to (3.1).

(I) If  $h = 0$ , we are done.

So, we assume by contradiction  $h \geq 1$  and we are going to show that then

$$E(u_0) + \sum_{j=1}^h E_{\infty}(w^j) \geq 2^{-s} m, \quad (3.8)$$

up to the case  $u_0 = 0$ ,  $h = 1$  and  $w^1 > 0$ , that arises for

$$\lim_{n \rightarrow \infty} E(u_n) = m. \quad (3.9)$$

Once (3.8) is proved, the contradiction comes out, because (3.2) and (3.3) provide  $c = E(u_0) + \sum_{j=1}^h E_{\infty}(w^j)$ , and  $c \in (m, 2^{-s} m)$  by assumption.

(II) If  $u_0 \equiv 0$  and  $h = 1$  occurs, then  $|w^1|_2 = \rho$ , by (3.3). Hence, if  $w^1 > 0$  then  $c = m$  by (3.2) and (2.1), so (3.9) is proved. We observe that this is the only case in which (3.9) holds, up to the autonomous case  $\Omega = \mathbb{R}^N$  and  $V \equiv 0$ , when also the case  $u_0 > 0$  and  $h = 1$  verifies (3.9). On the other hand, if  $w^1$  is a changing sign solution, then  $c > 2^{-s} m$  by Proposition 2.2.

(III) If  $u_0 \equiv 0$  and  $h \geq 2$ , we proceed by induction. For  $h = 2$ , we get (3.8) arguing exactly as in the proof of Proposition 2.2, with  $w^1$  in place of  $u^+$  and  $w^2$  in place of  $u^-$ . Observe that the case  $w^1 = w^2 > 0$  is the only case when the equality holds in (3.8).

Suppose now  $h \geq 3$  and (3.8) holds for  $h - 1$ , namely for every  $\rho_1 > 0$

$$\sum_{j=1}^{h-1} E_{\infty}(w^j) \geq 2^{-s} m_{\rho_1^2}, \quad (3.10)$$

whenever  $w^j \not\equiv 0$ , for all  $j \in \{1, \dots, h-1\}$ , and  $\sum_{j=1}^{h-1} |w^j|_2^2 = \rho_1^2$ . Then, let us prove (3.8) for  $h$ . We can assume that  $|w^h|_2^2 \leq \rho^2/3$ . Taking into account (3.10) and (2.7), we have

$$\begin{aligned} \sum_{j=1}^h E_{\infty}(w^j) &\geq 2^{-s} m_{(\rho^2 - |w^h|_2^2)} + E_{\infty}(w^h) \\ &\geq 2^{-s} m_{(\rho^2 - |w^h|_2^2)} + m_{|w^h|_2^2} \\ &= 2^{-s} \left( \frac{\rho^2 - |w^h|_2^2}{\rho^2} \right)^{1+s} m + \left( \frac{|w^h|_2^2}{\rho^2} \right)^{1+s} m. \end{aligned}$$

Hence, it is sufficient to verify that

$$2^{-s} \left( \frac{\rho^2 - |w^h|_2^2}{\rho^2} \right)^{1+s} m + \left( \frac{|w^h|_2^2}{\rho^2} \right)^{1+s} m \geq 2^{-s} m. \quad (3.11)$$

Inequality (3.11) is equivalent to

$$2^s t^{1+s} + (1-t)^{1+s} \leq 1, \quad (3.12)$$

where  $t := \frac{|w^h|_2^2}{\rho^2} \in (0, 1/3]$ . Since inequality (3.12) holds for every  $t \in [0, 1/3]$ , estimate (3.8) is proved for  $u_0 \equiv 0$ .

(IV) If  $u_0 \not\equiv 0$ , we can proceed as in the previous steps: first by considering the case  $h = 1$  and arguing as in the proof of Proposition 2.2, and then finishing the proof by induction.  $\square$

**Corollary 3.3.** *If  $(u_n)_n$  in  $S_\rho$  satisfies  $\lim_{n \rightarrow \infty} E(u_n) = m$ , then there exists a sequence  $(y_n)_n$  in  $\mathbb{R}^N$  such that*

$$u_n(x) = w(x - y_n) + o(1) \quad \text{in } H^1. \quad (3.13)$$

If  $\Omega \neq \mathbb{R}^N$  or  $V \not\equiv 0$ , then  $|y_n| \rightarrow \infty$ , as  $n \rightarrow \infty$ .

*Proof.* By the Ekeland variational principle, there exists a PS sequence  $(v_n)_n$  for  $E$  constrained on  $S_\rho$  such that  $\lim_{n \rightarrow \infty} E(v_n) = m$  and  $u_n = v_n + o(1)$  in  $H^1$  (see ([18, Proposition 5.1]), then we can assume that  $(u_n)_n$  is PS sequence. Then (3.13) is a direct consequence of (3.9) and part (II) in the proof of Proposition 3.1.

If  $y_n \rightarrow \bar{y} \in \mathbb{R}^N$ , up to a subsequence, then  $u_n \rightarrow w(\cdot - \bar{y})$  in  $H^1$  and a.e., so that  $\Omega = \mathbb{R}^N$  because  $w(x) > 0, \forall x \in \mathbb{R}^N$ . Moreover

$$\begin{aligned} m &= \lim_{n \rightarrow \infty} E(u_n) = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla w(x - \bar{y})|^2 dx + \frac{1}{2} \int_{\mathbb{R}^N} V(x) w(x - \bar{y})^2 dx - \frac{1}{p} \int_{\mathbb{R}^N} w^p(x - \bar{y}) dx \\ &= m + \frac{1}{2} \int_{\mathbb{R}^N} V(x) w(x - \bar{y})^2 dx \end{aligned}$$

implies  $V \equiv 0$ , again because  $w(x) > 0, \forall x \in \mathbb{R}^N$ .

So the proof is completed.  $\square$

*Remark 3.4.* By (III) in the proof of Proposition 3.1 we see that, for every sequence  $(y_n)_n$  in  $\mathbb{R}^N$  such that  $\lim_{n \rightarrow \infty} |y_n| = \infty$ ,

$$E(w_{\rho^2/2}(\cdot - y_n) + w_{\rho^2/2}(\cdot + y_n)) = 2^{-s} m.$$

Then the sequence  $(w_{\rho^2/2}(\cdot - y_n) + w_{\rho^2/2}(\cdot + y_n))_n$  turns out to be a not relatively compact PS-sequence at the level  $2^{-s} m$ , showing that the compactness interval  $(m, 2^{-s} m)$  cannot be extended.

## 4 Energy estimates

### 4.1 An upper bound

If  $\mathbb{R}^N \setminus \Omega$  is a non empty compact set contained in  $B_{R-1}(0)$ , let us introduce the cut-off function  $\vartheta$ , verifying

$$\vartheta \in C^\infty(\mathbb{R}^N, [0, 1]), \quad \begin{cases} \vartheta(x) = 1 & \text{if } |x| \geq R \\ \vartheta(x) = 0 & \text{if } x \in \mathbb{R}^N \setminus \Omega. \end{cases} \quad (4.1)$$

If  $\Omega = \mathbb{R}^N$ , we agree that  $\vartheta \equiv 1$  on  $\mathbb{R}^N$ . Let us set  $\Sigma = \partial B_2(e_1)$ , where  $e_1$  is the first element of the canonical basis of  $\mathbb{R}^N$ , and for any  $r > 0$  define the map  $\psi_r : [0, 1] \times \Sigma \rightarrow H_0^1(\Omega)$  by

$$\psi_r[t, z](x) = \rho \frac{\vartheta(x) [w_{t\rho^2}(x - rz) + w_{(1-t)\rho^2}(x - re_1)]}{\left| \vartheta(\cdot) [w_{t\rho^2}(\cdot - rz) + w_{(1-t)\rho^2}(\cdot - re_1)] \right|_2}. \quad (4.2)$$

**Proposition 4.1.** *Suppose that  $V$  verifies (1.1) and  $(D_\rho)$ , then*

(a) *there exist  $\bar{r} > 0$  such that for any  $r > \bar{r}$*

$$\mathcal{A}_r = \max \{E(\psi_r[t, z]) : t \in [0, 1], z \in \Sigma\} < 2^{-s} m; \quad (4.3)$$

(b) *for every  $\varepsilon > 0$  there exists  $r_\varepsilon > 0$  such that for any  $r > r_\varepsilon$*

$$\mathcal{L}_r = \max \{E(\psi_r[1, z]) : z \in \Sigma\} \leq m + \varepsilon. \quad (4.4)$$

*Remark 4.2.* In Proposition 4.1,  $V \equiv 0$  is allowed.

Before proving Proposition 4.1, let us recall two technical lemmas. For the proof of Lemma 4.3 we refer to [14] while the proof of Lemma 4.4 is in [3] (see also Lemma 2.9 in [13]).

**Lemma 4.3.** *For all  $a, b \geq 0$ , for all  $p \geq 2$ , the following relation holds true*

$$(a + b)^p \geq a^p + b^p + (p - 1)(a^{p-1}b + ab^{p-1}).$$

**Lemma 4.4.** *If  $g \in L^\infty(\mathbb{R}^N)$  and  $h \in L^1(\mathbb{R}^N)$  are such that, for some  $\alpha \geq 0$ ,  $b \geq 0$ ,  $\gamma \in \mathbb{R}$*

$$\lim_{|x| \rightarrow \infty} g(x) e^{\alpha|x|} |x|^b = \gamma \quad (4.5)$$

and

$$\int_{\mathbb{R}^N} |h(x)| e^{\alpha|x|} |x|^b dx < \infty, \quad (4.6)$$

then, for every  $z \in \mathbb{R}^N \setminus \{0\}$ ,

$$\lim_{r \rightarrow \infty} \left( \int_{\mathbb{R}^N} g(x + rz) h(x) dx \right) e^{\alpha|rz|} |rz|^b = \gamma \int_{\mathbb{R}^N} h(x) e^{-\alpha \frac{r \cdot z}{|z|}} dx.$$

**Lemma 4.5.** *Let  $z \in \Sigma$ . For every  $t \in [0, 1/2]$  and  $r > 0$ , let us set*

$$\delta_t(r) = \left( r^{\frac{N-1}{2}} e^{2\sqrt{t^s \lambda_\infty} r} \right)^{-1}, \quad (4.7)$$

$$\tau_t(r) = \frac{2}{\rho^2} \int_{\mathbb{R}^N} w_{t\rho^2}(x - rz) w_{(1-t)\rho^2}(x - re_1) dx, \quad (4.8)$$

$$\sigma_t(r) = \int_{\mathbb{R}^N} \left[ w_{t\rho^2}^{p-1}(x - rz) w_{(1-t)\rho^2}(x - re_1) + w_{t\rho^2}(x - rz) w_{(1-t)\rho^2}^{p-1}(x - re_1) \right] dx. \quad (4.9)$$

Then the following facts hold:

$$1) \text{ if } t \in [0, \frac{1}{2}), \text{ then } \frac{\tau_t(r)}{\delta_t(r)} \rightarrow c_{1,t} := c_t \int_{\mathbb{R}^N} w_{(1-t)\rho^2}(y) e^{-\sqrt{t^s \lambda_\infty} \frac{y \cdot (e_1 - z)}{2}} dy, \quad \text{as } r \rightarrow \infty,$$

$$2) \text{ if } t \in [0, \frac{1}{2}], \text{ then } \frac{\sigma_t(r)}{\delta_t(r)} \rightarrow c_{2,t} := 2c_t \int_{\mathbb{R}^N} w_{(1-t)\rho^2}^{p-1}(y) e^{-\sqrt{t^s \lambda_\infty} \frac{y \cdot (e_1 - z)}{2}} dy, \quad \text{as } r \rightarrow \infty$$

where  $c_t = c_1 t^s \left( \frac{1}{p-2} - \frac{N-1}{4} \right)$  (see (2.3)). Moreover,

3)  $c_{1,t} \cdot (\frac{1}{2} - t) \leq C$  for every  $t \in [0, \frac{1}{2}]$ .

*Remark 4.6.* The definition of  $\tau_t(r), \sigma_t(r)$  is independent of  $z$ , by symmetry. Moreover,  $c_{1,t} \rightarrow \infty$  as  $t \rightarrow \frac{1}{2}$  and, clearly,  $\tau_0(r) = \sigma_0(r) \equiv 0$ .

*Proof.* Assertions 1), 2) easily follow using (2.9) and by Lemma 4.4. Let us prove assertion 3). Without loss of generality we may assume  $z = -e_1$ , so that  $\frac{z-e_1}{2} = -e_1$ , and to simplify the notations we consider  $\lambda_\infty = 1$ . Moreover, since  $t \mapsto c_{1,t}$  is a continuous function on  $[0, 1/2]$ , it is sufficient to prove

$$\lim_{t \rightarrow \frac{1}{2}} \left( \frac{1}{2} - t \right) \int_{\mathbb{R}^N} w_{(1-t)\rho^2}(y) e^{t^{s/2} y_1} dy < \infty,$$

where  $y_1$  is the first component of  $y \in \mathbb{R}^N$ . Moreover, by (2.9) we are left to prove that

$$\begin{aligned} & \left( \frac{1}{2} - t \right) \int_{\mathbb{R}^N \setminus B_1(0)} |y|^{-\frac{N-1}{2}} e^{-(1-t)^{s/2}|y|} e^{t^{s/2} y_1} dy = \\ & = \left( \frac{1}{2} - t \right) \int_{\mathbb{R}^N \setminus B_1(0)} |y|^{-\frac{N-1}{2}} e^{-[(1-t)^{s/2} - t^{s/2}]|y|} e^{-t^{s/2}(|y|-y_1)} dy \leq C < \infty \quad \text{as } t \rightarrow (1/2)^-. \end{aligned} \quad (4.10)$$

If we set  $a = \frac{1}{2} - t$ , then  $a \rightarrow 0^+$  as  $t \rightarrow (1/2)^-$ , and since  $[(1-t)^{s/2} - t^{s/2}] \geq ca$  as  $a \rightarrow 0$ , with  $c > 0$ , (4.10) can be estimated by

$$a \int_{\mathbb{R}^N \setminus B_1(0)} |y|^{-\frac{N-1}{2}} e^{-ca|y|} e^{-(\frac{1}{2}-a)^{s/2}(|y|-y_1)} dy \leq a \int_{\mathbb{R}^N \setminus B_1(0)} |y|^{-\frac{N-1}{2}} e^{-ca|y|} e^{-(\frac{1}{4})^{s/2}(|y|-y_1)} dy.$$

Making use of spherical coordinates in the subspace  $e_1^\perp = \{v \in \mathbb{R}^N : v \cdot e_1 = 0\}$ , denoting by  $b = (\frac{1}{4})^{s/2}$  and  $\mathbb{R}_+^2 = \mathbb{R} \times [0, +\infty)$ , we have to analyze

$$\begin{aligned} & a \int_{\mathbb{R}_+^2 \setminus B_1(0)} \frac{r^{N-2}}{(r^2 + y_1^2)^{\frac{N-1}{4}} e^{ca(r^2+y_1^2)^{1/2}} e^{b[(r^2+y_1^2)^{1/2}-y_1]}} dr dy_1 = \\ & = a \int_0^\pi \left( \int_1^{+\infty} \frac{\rho^{(N-1)/2}}{e^{[ca+b(1-\cos\vartheta)]\rho}} d\rho \right) (\sin\vartheta)^{N-2} d\vartheta. \end{aligned}$$

Setting  $k = ca + b(1 - \cos\vartheta)$ , we can estimate

$$\int_1^{+\infty} \frac{\rho^{(N-1)/2}}{e^{[ca+b(1-\cos\vartheta)]\rho}} d\rho = \frac{1}{k^{(N-1)/2}} \int_1^{+\infty} \frac{(k\rho)^{(N-1)/2}}{e^{k\rho}} d\rho \leq \frac{1}{k^{(N+1)/2}} \int_0^{+\infty} \frac{\mu^{(N-1)/2}}{e^\mu} d\mu.$$

Then,

$$a \int_0^\pi \left( \int_1^{+\infty} \frac{\rho^{(N-1)/2}}{e^{[ca+b(1-\cos\vartheta)]\rho}} d\rho \right) (\sin\vartheta)^{N-2} d\vartheta \leq Ca \int_0^\pi \frac{(\sin\vartheta)^{N-2}}{[ca + b(1 - \cos\vartheta)]^{(N+1)/2}} d\vartheta.$$

Since  $1 - \cos\vartheta \geq \frac{1}{4} \sin^2\vartheta$  for small  $\vartheta$ , if we set  $u = \sin\vartheta$  and  $\beta = b/(4c)$ , then it is sufficient to evaluate

$$\begin{aligned} & a \int_0^{\frac{1}{2}} \frac{u^{N-2}}{[a + \beta u^2]^{(N+1)/2}} \cdot \frac{1}{(1-u^2)^{1/2}} du \leq \widehat{C} a^{-1/2} \int_0^{\frac{1}{2}} \frac{(\sqrt{\beta/a} u)^{N-2}}{\left[1 + \left(\sqrt{\beta/a} u\right)^2\right]^{(N+1)/2}} du \\ & = \widetilde{C} \int_0^{\frac{1}{2}} \frac{t^{N-2}}{(1+t^2)^{(N+1)/2}} dt \rightarrow C \quad \text{as } a \rightarrow 0, \end{aligned}$$

and the assertion follows. □

**Proof of Proposition 4.1** In this proof we shall consider  $R > 1$  fixed such that  $\mathbb{R}^N \setminus \Omega \subset B_{R-1}(0)$ . In order to simplify the notations, we often omit  $t, z$  and write  $\psi_r = \psi_r[t, z]$ ,  $\delta(r) = \delta_t(r)$ ,  $\sigma(r) = \sigma_t(r)$ ,  $\tau(r) = \tau_t(r)$  (see (4.2), (4.7), (4.8), (4.9)).

We have that

$$E(\psi_r) = \frac{1}{2} \int_{\mathbb{R}^N} [|\nabla \psi_r|^2 + V(x)\psi_r^2] dx - \frac{1}{p} \int_{\mathbb{R}^N} |\psi_r|^p dx.$$

So, to get the statement of the Lemma, we need to estimate this two integrals.

Let us consider  $0 \leq t \leq 1/2$ . In an entirely analogous way we may treat also the case  $1/2 < t \leq 1$ .

Let us set:

$$w_1 = w_{t\rho^2}, \quad w_2 = w_{(1-t)\rho^2}, \quad \lambda_1 = \lambda_{t\rho^2} = t^s \lambda_\infty, \quad \lambda_2 = \lambda_{(1-t)\rho^2} = (1-t)^s \lambda_\infty,$$

and for any  $i = 1, 2$

$$A_i = |\nabla w_i|_2^2, \quad B_i = |w_i|_p^p.$$

Recall that

$$-\Delta w_i + \lambda_i w_i = w_i^{p-1},$$

namely

$$\int_{\mathbb{R}^N} \nabla w_i(x) \cdot \nabla v(x) dx + \lambda_i \int_{\mathbb{R}^N} w_i(x) v(x) dx = \int_{\mathbb{R}^N} w_i^{p-1}(x) v(x) dx, \quad \forall v \in H^1(\mathbb{R}^N), \quad (4.11)$$

and so

$$A_i - B_i = -\lambda_i |w_i|_2^2. \quad (4.12)$$

Evidently for  $t = \frac{1}{2}$  we have  $w_1 = w_2$ ,  $\lambda_1 = \lambda_2$ ,  $A_1 = A_2$ ,  $B_1 = B_2$ . Moreover, we recall that

$$\frac{1}{2} A_1 - \frac{1}{p} B_1 = m_{t\rho^2} = t^{1+s} m, \quad \frac{1}{2} A_2 - \frac{1}{p} B_2 = m_{(1-t)\rho^2} = (1-t)^{1+s} m. \quad (4.13)$$

With these notations,

$$\psi_r(x) = \rho \frac{\vartheta(x) [w_1(x - rz) + w_2(x - re_1)]}{|\vartheta(\cdot) [w_1(\cdot - rz) + w_2(\cdot - re_1)]|_2}.$$

Estimate of  $|\vartheta(\cdot) [w_1(\cdot - rz) + w_2(\cdot - re_1)]|_2$ .

From above

$$\begin{aligned}
|\vartheta(\cdot) [w_1(\cdot - rz) + w_2(\cdot - re_1)]|_2^2 &\leq |w_1(\cdot - rz) + w_2(\cdot - re_1)|_2^2 \\
&= |w_1|_2^2 + |w_2|_2^2 + 2 \int_{\mathbb{R}^N} w_1(x - rz) w_2(x - re_1) dx \\
&= \rho^2 + 2 \int_{\mathbb{R}^N} w_1(x - rz) w_2(x - re_1) dx \\
&= \rho^2(1 + \tau(r)).
\end{aligned} \tag{4.14}$$

From below,

$$\begin{aligned}
|\vartheta(\cdot) [w_1(\cdot - rz) + w_2(\cdot - re_1)]|_2^2 &= \int_{\mathbb{R}^N} |\vartheta(x) [w_1(x - rz) + w_2(x - re_1)]|^2 dx \\
&\geq \int_{\mathbb{R}^N} \left| w_1(x - rz) + w_2(x - re_1) \right|^2 dx \\
&\quad - \int_{B_R(0)} \left| w_1(x - rz) + w_2(x - re_1) \right|^2 dx.
\end{aligned} \tag{4.15}$$

By the asymptotic behavior of  $w_1$  and  $w_2$  (see (2.9)), for any  $q \geq 2$  we have

$$\int_{B_R(0)} \left| w_1(x - rz) + w_2(x - re_1) \right|^q dx \leq 2^{q-1} \int_{B_R(0)} [|w_1(x - rz)|^q + |w_2(x - re_1)|^q] dx = o(\delta(r)). \tag{4.16}$$

Therefore, by (4.14), (4.15) and (4.16) we get

$$|\vartheta(\cdot) [w_1(\cdot - rz) + w_2(\cdot - re_1)]|_2^2 \geq \rho^2(1 + \tau(r)) + o(\delta(r)). \tag{4.17}$$

Estimate of  $\int_{\mathbb{R}^N} [|\nabla \psi_r|^2 + V(x)\psi_r^2] dx$ .

Now, let us estimate

$$\begin{aligned}
&\int_{\mathbb{R}^N} \left| \nabla \left( \vartheta(x) [w_1(x - rz) + w_2(x - re_1)] \right) \right|^2 dx \\
&\leq \int_{\mathbb{R}^N} \left( |\nabla [w_1(x - rz) + w_2(x - re_1)]|^2 \right. \\
&\quad \left. + 2 \int_{\mathbb{R}^N} \left[ \vartheta(x) \nabla \vartheta(x) \right] \cdot \left( [w_1(x - rz) + w_2(x - re_1)] \nabla [w_1(x - rz) + w_2(x - re_1)] \right) dx \right. \\
&\quad \left. + \int_{\mathbb{R}^N} |\nabla \vartheta(x)|^2 [w_1(x - rz) + w_2(x - re_1)]^2 dx \right) dx.
\end{aligned} \tag{4.18}$$

By direct computation, (4.11) and (4.12), we obtain

$$\begin{aligned}
& \int_{\mathbb{R}^N} |\nabla [w_1(x - rz) + w_2(x - re_1)]|^2 dx \\
&= |\nabla w_1|_2^2 + |\nabla w_2|_2^2 + 2 \int_{\mathbb{R}^N} \nabla w_1(x - rz) \cdot \nabla w_2(x - re_1) dx \\
&= |\nabla w_1|_2^2 + |\nabla w_2|_2^2 - (\lambda_1 + \lambda_2) \int_{\mathbb{R}^N} w_1(x - rz) w_2(x - re_1) dx \\
&\quad + \int_{\mathbb{R}^N} \left[ w_1^{p-1}(x - rz) w_2(x - re_1) + w_1(x - rz) w_2^{p-1}(x - re_1) \right] dx \\
&= A_1 + A_2 - (\lambda_1 + \lambda_2) \frac{\rho^2}{2} \tau(r) + \sigma(r).
\end{aligned} \tag{4.19}$$

Since  $\nabla \vartheta \equiv 0$  on  $\mathbb{R}^N \setminus B_R(0)$ , by using (2.9) (2.10) we get

$$\begin{aligned}
\int_{\mathbb{R}^N} |\nabla \vartheta(x)|^2 [w_1(x - rz) + w_2(x - re_1)]^2 dx &\leq c \int_{B_R(0)} [w_1(x - rz) + w_2(x - re_1)]^2 dx \\
&= o(\delta(r))
\end{aligned} \tag{4.20}$$

and

$$\begin{aligned}
& \left| \int_{\mathbb{R}^N} \left[ \vartheta(x) \nabla \vartheta(x) \right] \cdot \left( [w_1(x - rz) + w_2(x - re_1)] \nabla [w_1(x - rz) + w_2(x - re_1)] \right) dx \right| \\
&\leq c \int_{B_R(0)} \left| [w_1(x - rz) + w_2(x - re_1)] \nabla [w_1(x - rz) + w_2(x - re_1)] \right| dx \\
&= o(\delta(r)).
\end{aligned} \tag{4.21}$$

According to the contribution of the potential, by ( $D_\rho$ ), (2.9) and by Lemma 4.4 we have

$$\begin{aligned}
& \int_{\mathbb{R}^N} V(x) |\vartheta(x) [w_1(x - rz) + w_2(x - re_1)]|^2 dx \\
&\leq \int_{\mathbb{R}^N} V(x) [w_1(x - rz) + w_2(x - re_1)]^2 dx = o(\delta(r)).
\end{aligned} \tag{4.22}$$

By (4.17)–(4.22) we deduce

$$\int_{\mathbb{R}^N} [|\nabla \psi_r|^2 + V(x) \psi_r^2] dx \leq \frac{1}{1 + \tau(r) + o(\delta(r))} \left[ A_1 + A_2 - (\lambda_1 + \lambda_2) \frac{\rho^2}{2} \tau(r) + \sigma(r) + o(\delta(r)) \right]. \tag{4.23}$$

Estimate of  $\int_{\mathbb{R}^N} |\psi_r|^p dx$ .

Since  $0 \leq \vartheta(x) \leq 1$  in  $\mathbb{R}^N$  and  $\vartheta \equiv 1$  in  $\mathbb{R}^N \setminus B_R(0)$ , by (4.16) and by Lemma 4.3, we get

$$|\vartheta(\cdot) [w_1(\cdot - rz) + w_2(\cdot - re_1)]|_p^p = \int_{\mathbb{R}^N} \left| \vartheta(x) [w_1(x - rz) + w_2(x - re_1)] \right|^p dx$$

$$\begin{aligned}
&\geq \int_{\mathbb{R}^N} \left| w_1(x - rz) + w_2(x - re_1) \right|^p dx - \int_{B_R(0)} \left| w_1(x - rz) + w_2(x - re_1) \right|^p dx \\
&\geq |w_1|_p^p + |w_2|_p^p \\
&\quad + (p-1) \int_{\mathbb{R}^N} \left[ w_1^{p-1}(x - rz) w_2(x - re_1) + w_1(x - rz) w_2^{p-1}(x - re_1) \right] dx \\
&\quad - \int_{B_R(0)} \left| w_1(x - rz) + w_2(x - re_1) \right|^p dx \\
&= B_1 + B_2 + (p-1)\sigma(r) - \int_{B_R(0)} \left| w_1(x - rz) + w_2(x - re_1) \right|^p dx \\
&\geq B_1 + B_2 + (p-1)\sigma(r) + o(\delta(r)).
\end{aligned} \tag{4.24}$$

Taking into account (4.14) and (4.24) we have that

$$|\psi_r|_p^p \geq \left( \frac{1}{1 + \tau(r)} \right)^{p/2} [B_1 + B_2 + (p-1)\sigma(r) + o(\delta(r))]. \tag{4.25}$$

Estimate of  $E(\psi_r)$ .

Therefore

$$\begin{aligned}
E(\psi_r) &= \frac{1}{2} \int_{\mathbb{R}^N} [|\nabla \psi_r|^2 + V(x)\psi_r^2] dx - \frac{1}{p} |\psi_r|_p^p \leq \\
&\leq \frac{1}{2} \left( \frac{1}{1 + \tau(r) + o(\delta(r))} \right) \left[ A_1 + A_2 - (\lambda_1 + \lambda_2) \frac{\rho^2}{2} \tau(r) + \sigma(r) + o(\delta(r)) \right] \\
&\quad - \frac{1}{p} \left( \frac{1}{1 + \tau(r)} \right)^{p/2} [B_1 + B_2 + (p-1)\sigma(r) + o(\delta(r))].
\end{aligned}$$

Observe that for every  $\sigma, \tau > 0$  and  $\delta \rightarrow 0$  we have

$$\begin{aligned}
&\frac{1}{2} \left( \frac{1}{1 + \tau + o(\delta)} \right) \left[ A_1 + A_2 - (\lambda_1 + \lambda_2) \frac{\rho^2}{2} \tau + \sigma + o(\delta) \right] \\
&\quad - \frac{1}{p} \left( \frac{1}{1 + \tau} \right)^{\frac{p}{2}} [B_1 + B_2 + (p-1)\sigma + o(\delta)] = \varphi(\tau, \sigma) + o(\delta),
\end{aligned}$$

where

$$\varphi(\tau, \sigma) = \frac{1}{2} \left( \frac{1}{1 + \tau} \right) \left[ A_1 + A_2 - (\lambda_1 + \lambda_2) \frac{\rho^2}{2} \tau + \sigma \right] - \frac{1}{p} \left( \frac{1}{1 + \tau} \right)^{\frac{p}{2}} [B_1 + B_2 + (p-1)\sigma],$$

and we have performed a Taylor expansion with respect to  $o(\delta)$ . We will write  $\varphi_t$  in place of  $\varphi$  when we want to emphasize the role of  $t$ .

Now, we are going to consider the Taylor expansion of  $\varphi$ . Observe that these expansions are consistent by (1) and (2) in Lemma 4.5.

By (4.13), and taking into account (4.12), we have that

$$\varphi(0,0) = \left(\frac{1}{2}A_1 - \frac{1}{p}B_1\right) + \left(\frac{1}{2}A_2 - \frac{1}{p}B_2\right) = [t^{1+s} + (1-t)^{1+s}] m, \quad (4.26)$$

$$\begin{aligned} \frac{\partial\varphi}{\partial\tau}(0,0) &= -\frac{1}{2} \left[ A_1 + A_2 + (\lambda_1 + \lambda_2) \frac{\rho^2}{2} - B_1 - B_2 \right] \\ &= -\frac{1}{2} \left[ A_1 - B_1 + \lambda_1 \frac{\rho^2}{2} \right] - \frac{1}{2} \left[ A_2 - B_2 + \lambda_2 \frac{\rho^2}{2} \right] \\ &= -\frac{1}{2} \left[ -\lambda_1 t \rho^2 + \lambda_1 \frac{\rho^2}{2} \right] - \frac{1}{2} \left[ -\lambda_2 (1-t) \rho^2 + \lambda_2 \frac{\rho^2}{2} \right] \\ &= -\frac{1}{2} \lambda_1 \rho^2 \left( \frac{1}{2} - t \right) + \frac{1}{2} \lambda_2 \rho^2 \left( \frac{1}{2} - t \right) \\ &= \frac{1}{2} \left( \frac{1}{2} - t \right) (\lambda_2 - \lambda_1) \rho^2. \end{aligned} \quad (4.27)$$

$$\frac{\partial\varphi}{\partial\sigma}(0,0) = -\left(\frac{1}{2} - \frac{1}{p}\right). \quad (4.28)$$

Hence, we obtain

$$\varphi(\tau, \sigma) = \varphi(0,0) + \frac{1}{2} \left( \frac{1}{2} - t \right) (\lambda_2 - \lambda_1) \rho^2 \tau - \left( \frac{1}{2} - \frac{1}{p} \right) \sigma + o\left(\sqrt{\tau^2 + \sigma^2}\right), \quad \text{as } (\tau, \sigma) \rightarrow (0,0).$$

Now, we want to analyze the asymptotic behaviour of  $\varphi(\tau(r), \sigma(r))$  and, as a consequence, of  $E(\psi_r)$ , as  $r \rightarrow \infty$ .

By Lemma 4.5, for every  $t \in [0, \frac{1}{2})$  we have

$$\varphi(\tau_t(r), \sigma_t(r)) = \varphi(0,0) + \left[ \frac{1}{2} \left( \frac{1}{2} - t \right) (\lambda_2 - \lambda_1) \rho^2 c_{1,t} - \left( \frac{1}{2} - \frac{1}{p} \right) c_{2,t} \right] \delta_t(r) + o(\delta_t(r)),$$

as  $r \rightarrow \infty$ . Taking into account Lemma 4.5 and  $\lambda_1 \rightarrow \lambda_2$  as  $t \rightarrow \frac{1}{2}$ , we get

$$\left[ \frac{1}{2} \left( \frac{1}{2} - t \right) (\lambda_2 - \lambda_1) \rho^2 c_{1,t} - \left( \frac{1}{2} - \frac{1}{p} \right) c_{2,t} \right] \rightarrow -\left( \frac{1}{2} - \frac{1}{p} \right) C_2 < 0, \quad \text{as } t \rightarrow \left( \frac{1}{2} \right)^-.$$

Then, for suitable constants  $C > 0$  and  $\mu \in (0, \frac{1}{2})$

$$E(\psi_r[t, z]) \leq [t^{1+s} + (1-t)^{1+s}] m - C \delta_t(r) + o(\delta_t(r)) \quad \forall t \in [\mu, 1/2)$$

(see (4.26). Since  $\delta_t(r) \rightarrow 0$  as  $r \rightarrow \infty$ , uniformly in  $t$  (see (4.7)), we can conclude that there exists  $r_1 > 0$  such that

$$E(\psi_r[t, z]) < \max_{t \in [0,1]} [t^{1+s} + (1-t)^{1+s}] m = 2^{-s} m \quad \forall t \in [\mu, 1/2), \quad \forall r > r_1. \quad (4.29)$$

If  $t = \frac{1}{2}$ , for suitable  $\alpha, \beta, \gamma \in \mathbb{R}$  we infer

$$\varphi(\tau, \sigma) = \varphi(0,0) - \left( \frac{1}{2} - \frac{1}{p} \right) \sigma + \alpha \tau^2 + \beta \sigma^2 + \gamma \sigma \tau + o(\tau^2 + \sigma^2) \quad \text{as } (\tau, \sigma) \rightarrow (0,0). \quad (4.30)$$

Now, consider that by Lemma 4.5

$$\sigma_{1/2}(r) = (c_{2, \frac{1}{2}} + o(1))\delta_{1/2}(r) \quad (4.31)$$

and that, fixed  $\eta \in \left(0, \frac{\sqrt{2^{-s}\lambda_\infty}}{2}\right]$ , by Lemma 4.4

$$\tau_{1/2}(r) = o\left(e^{-2(\sqrt{2^{-s}\lambda_\infty} - \eta)r} r^{-\frac{N-1}{2}}\right), \quad \text{as } r \rightarrow +\infty,$$

so that

$$(\tau_{1/2}(r))^2 = o(\delta_{1/2}(r)). \quad (4.32)$$

By (4.30), (4.31), (4.32), we can conclude that there exists  $r_2 > 0$  such that

$$E(\psi_r[1/2, z]) = \varphi(0, 0) - C\left(\frac{1}{2} - \frac{1}{p}\right)\delta_{1/2}(r) + o(\delta_{1/2}(r)) < 2^{-s}m \quad \forall r > r_2. \quad (4.33)$$

If  $t \in [0, \mu]$ , then, taking into account that  $\delta_t(r) \leq \delta_0(r)$  for every  $t \in [0, 1/2]$ , we get

$$\begin{aligned} E(\psi_r[t, z]) &\leq \varphi_t(0, 0) + O(\delta_0(r)) \\ &\leq [t^{1+s} + (1-t)^{1+s}]m + O(\delta_0(r)) \\ &\leq [\mu^{1+s} + (1-\mu)^{1+s}]m + O(\delta_0(r)) \quad \forall t \in [0, \mu]. \end{aligned}$$

Since  $[\mu^{1+s} + (1-\mu)^{1+s}]m < 2^{-s}m$ , there exists  $r_3 > 0$  such that

$$E(\psi_r[t, z]) < 2^{-s}m \quad \forall t \in [0, \mu], \quad \forall r > r_3. \quad (4.34)$$

So, assertion (a) follows from (4.29), (4.33), (4.34), for every  $r > \bar{r} := \max\{r_1, r_2, r_3\}$ .

The estimates developed above also show that

$$E(\psi_r[0, z]) \leq \varphi_0(0, 0) + o(\delta_0(r)) = m + o(\delta_0(r))$$

so that  $E(\psi_r[0, z]) \rightarrow m$ , as  $r \rightarrow \infty$ . The same arguments work to evaluate

$$E(\psi_r[1, z]) \leq m + o(\delta_0(r)),$$

uniformly in  $z \in \Sigma$ . So, also (b) is proved.  $\square$

## 4.2 Other estimates

In this subsection we consider the nonautonomous case  $\Omega \neq \mathbb{R}^N$  or  $V \neq 0$ .

The following definition of barycenter of a function  $u \in H^1(\mathbb{R}^N) \setminus \{0\}$ , has been introduced in [15]. We set

$$\mu(u)(x) = \frac{1}{|B_1(0)|} \int_{B_1(x)} |u(y)| dy \quad (4.35)$$

and we remark that  $\mu(u)$  is bounded and continuous, so we can introduce the function

$$\hat{u}(x) = \left[ \mu(u)(x) - \frac{1}{2} \max \mu(u) \right]^+, \quad (4.36)$$

that is continuous and has compact support. Thus, we can set  $\beta : H^1(\mathbb{R}^N) \setminus \{0\} \rightarrow \mathbb{R}^N$  as

$$\beta(u) = \frac{1}{|\hat{u}|_1} \int_{\mathbb{R}^N} \hat{u}(x) x \, dx.$$

The map  $\beta$  has the following properties:

$$\beta \text{ is continuous in } H^1(\mathbb{R}^N) \setminus \{0\}; \quad (4.37)$$

$$\text{if } u \text{ is a radial function, then } \beta(u) = 0; \quad (4.38)$$

$$\beta(tu) = \beta(u) \quad \forall t \in \mathbb{R} \setminus \{0\}, \quad \forall u \in H^1(\mathbb{R}^N) \setminus \{0\}; \quad (4.39)$$

$$\beta(u(x-z)) = \beta(u) + z \quad \forall z \in \mathbb{R}^N \quad \forall u \in H^1(\mathbb{R}^N) \setminus \{0\}. \quad (4.40)$$

Let us set

$$C_0 = \inf\{E(u) : u \in H_0^1(\Omega), |u|_2 = \rho, \beta(u) = 0\}.$$

**Lemma 4.7.** *We have that  $C_0 > m$ .*

*Proof.* Of course we have that  $C_0 \geq m$ . Assume by contradiction that  $C_0 = m$ . Then by Corollary 3.3 there exists a sequence  $(y_n)_n$  in  $\mathbb{R}^N$  such that  $|y_n| \rightarrow \infty$ , as  $n \rightarrow \infty$ , and

$$u_n(x) = w(x - y_n) + \phi_n(x), \quad \phi_n \rightarrow 0 \text{ strongly in } H^1(\mathbb{R}^N).$$

By (4.37), (4.38) and (4.40) we have

$$0 = \beta(u_n) = \beta(w(\cdot - y_n) + \phi_n) = \beta(w + \phi(\cdot + y_n)) + y_n = y_n + o(1),$$

contrary to  $|y_n| \rightarrow \infty$ , as  $n \rightarrow \infty$ .

So the proof is completed □

**Lemma 4.8.** *Let  $\mathcal{A}_r$  and  $\mathcal{L}_r$  be as in Proposition 4.1. Then  $\hat{r} > 0$  exists such that*

$$\mathcal{L}_r < C_0 \leq \mathcal{A}_r, \quad \text{for all } r \geq \hat{r}. \quad (4.41)$$

*Proof.* Inequality  $\mathcal{L}_r < C_0$ , for large  $r$ , follows from (4.4) and Lemma 4.7.

To get the second inequality in (4.41), we claim that  $\beta(\vartheta(\cdot)w(\cdot - rz)) \cdot z > 0$  for all  $z \in \Sigma$ , for large  $r$ . Indeed, by (4.37), (4.38) and (4.40) we have

$$\left| \beta(\vartheta(\cdot)w(\cdot - rz)) - rz \right| = \left| \beta(\vartheta(\cdot + rz)w) \right| \xrightarrow{r \rightarrow \infty} 0,$$

because  $\vartheta(\cdot + rz)w \rightarrow w$  in  $H^1$  as  $r \rightarrow \infty$ , by (2.3) and (2.4). Hence

$$\beta(\vartheta(\cdot)w(\cdot - rz)) = rz + o(1) \quad \text{as } r \rightarrow \infty,$$

as asserted. So, for  $r$  large, the deformation  $\mathcal{G} : [0, 1] \times \Sigma \rightarrow \mathbb{R}^N \setminus \{0\}$  given by

$$\mathcal{G}(s, z) = s\beta(\psi_r[1, z]) + (1-s)z \quad (4.42)$$

is well defined. Then we claim that there exists  $(t_r, z_r) \in [0, 1] \times \Sigma$  such that

$$\beta(\psi_r[t_r, z_r]) = 0. \quad (4.43)$$

Indeed, by the continuity of the maps  $\beta$  and  $\psi_r$ , by  $\mathcal{G}(s, z) \neq 0$ ,  $\forall (s, z) \in [0, 1] \times \Sigma$ , and by the invariance of the topological degree by homotopy, we have

$$0 \neq d(\text{Id}, \Sigma \times [0, 1], 0) = d(\beta \circ \psi_r, \Sigma \times [0, 1], 0).$$

From (4.43) there follows  $C_0 \leq E(\psi_r[t_r, z_r]) \leq \mathcal{A}_r$ , that completes the proof. □

## 5 Proof of the main results

The existence of a positive solution for the autonomous case  $\Omega = \mathbb{R}^N$  and  $V \equiv 0$  is well known, so we prove our results when  $\Omega = \mathbb{R}^N$  and  $V \equiv 0$  does not occur.

**Proof of Theorem 1.1** Let us recall the values

$$\mathcal{A}_r = \max \{E(\psi_r[t, z]) : t \in [0, 1], z \in \Sigma\}, \quad (5.1)$$

$$\mathcal{L}_r = \max \{E(\psi_r[1, y]) : z \in \Sigma\}, \quad (5.2)$$

$$C_0 = \inf \{E(u) : u \in H_0^1(\Omega), |u|_2 = \rho, \beta(u) = 0\}. \quad (5.3)$$

By Propositions 4.1 and 1.4 (see (2.11)), and by Lemma 4.8, we have that for all  $r > \max\{\bar{r}, \widehat{r}\}$

$$m < \mathcal{L}_r < C_0 \leq \mathcal{A}_r < 2^{-s}m. \quad (5.4)$$

We *claim* that the functional  $E$  has a (PS)-sequence in  $[C_0, \mathcal{A}_r]$ . This done, the existence of a critical point  $\bar{u}$  of  $E$  on  $S_\rho$  with  $E(\bar{u}) \leq \mathcal{A}_r$  follows from Proposition 3.1.

Assume, by contradiction, that no (PS)-sequence exists in  $[C_0, \mathcal{A}_r]$ . Then, usual deformation arguments imply the existence of  $\eta > 0$  such that the sublevel  $E^{C_0-\eta} := \{u \in H_0^1(\Omega) : |u|_2^2 = \rho^2, E(u) \leq C_0 - \eta\}$  is a deformation retract of the sublevel  $E^{\mathcal{A}_r} := \{u \in H_0^1(\Omega) : |u|_2^2 = \rho^2, E(u) \leq \mathcal{A}_r\}$ , namely there exists a continuous function  $\varphi : E^{\mathcal{A}_r} \rightarrow E^{C_0-\eta}$  such that

$$\varphi(u) = u \quad \text{for any } u \in E^{C_0-\eta}. \quad (5.5)$$

Furthermore, by (5.4) we can also assume  $\eta$  so small that

$$C_0 - \eta > \mathcal{L}_r. \quad (5.6)$$

Let us define the map  $\mathcal{H} : [0, 1] \times \Sigma \rightarrow \mathbb{R}^N$  by

$$\mathcal{H}(s, y) = \beta(\varphi(\psi_r[t, z])).$$

By (5.6), (5.5), and by using the map  $\mathcal{G}$  introduced in (4.42), we deduce that  $\mathcal{H}$  maps  $\{1\} \times \Sigma$  in a set homotopically equivalent to  $\Sigma$  in  $\mathbb{R}^N \setminus \{0\}$ . Since  $\mathcal{H}$  is a continuous map, and arguing exactly as for (4.43), we get the existence of a point  $(\tilde{t}, \tilde{z}) \in [0, 1] \times \Sigma$  such that

$$0 = \mathcal{H}(\tilde{t}, \tilde{z}) = \beta(\varphi(\psi_r[\tilde{t}, \tilde{z}])).$$

Then by the definition of  $C_0$  we see  $E(\varphi(\psi_r[\tilde{t}, \tilde{z}])) \geq C_0$ , contrary to  $\varphi(\psi_r[t, z]) \in E^{C_0-\eta}$  for every  $(t, z) \in [0, 1] \times \Sigma$ , so the claim must be true.

Finally, since  $E(\bar{u}) \in (m, 2^{-s}m)$  then  $\bar{u}$  has constant sign by Proposition 2.2. Observe that since  $\bar{u}$  solves (P) if and only if  $-\bar{u}$  solves (P), then we have a nonnegative solution.  $\square$

**Proof of Theorem 1.2** This proof proceed in two different ways, according to  $\Omega \neq \mathbb{R}^N$  or  $\Omega = \mathbb{R}^N$ . In both cases, we identify a topological configuration analogous to the one employed in the proof of Theorem 1.1 to get the solution. We only outline the procedure, because the argument is the same already developed in details.

Case  $\Omega \neq \mathbb{R}^N$ .

Let us apply Proposition 4.1, Lemma 4.7 and Lemma 4.8 with  $V \equiv 0$ . Then for a fixed  $r > 0$ , large enough, we get

$$m < \mathcal{L}_{r,0} < C_{0,0} \leq \mathcal{A}_{r,0} < 2^{-s}m, \quad (5.7)$$

where  $\mathcal{L}_{r,0}$ ,  $C_{0,0}$ ,  $\mathcal{A}_{r,0}$  are defined as in (5.1), (5.2), (5.3), with the functional  $E$  replaced by

$$E_0(u) := \frac{1}{2} \int_{\Omega} |\nabla u|^2 dx - \frac{1}{p} \int_{\Omega} |u|^p dx, \quad u \in H_0^1(\Omega).$$

The configuration in (5.7) depends on  $\Omega$  and  $\rho$ .

Now, observe that  $\mathcal{L}_{r,0} \leq \mathcal{L}_r$ ,  $C_{0,0} \leq C_0$ ,  $\mathcal{A}_{r,0} \leq \mathcal{A}_r$ , by (1.1). Moreover, by the Hölder inequality and (2.9), it follows that

$$\lim_{|V|_q \rightarrow 0} \int_{\mathbb{R}^N} V(x) (\psi_r[t, z])^2 dx \leq \lim_{|V|_q \rightarrow 0} |V|_q \max_{t \in [0,1], z \in \Sigma} |(\psi_r[t, z])^2|_{q'} = 0,$$

uniformly in  $t \in [0, 1]$  and  $z \in \Sigma$ , that implies

$$\lim_{|V|_q \rightarrow 0} \mathcal{L}_r = \mathcal{L}_{r,0}, \quad \lim_{|V|_q \rightarrow 0} \mathcal{A}_r = \mathcal{A}_{r,0}.$$

Hence, taking also into account Lemma 4.8, we see that there exists  $L = L(\Omega, \rho)$  such that if  $|V|_q < L$  then the configuration (5.4) is restored. In particular,

$$L < \frac{1}{\max_{t \in [0,1], z \in \Sigma} |(\psi_r[t, z])^2|_{q'}} (2^{-s}m - \mathcal{A}_{r,0}). \quad (5.8)$$

As a consequence, if  $|V|_q < L$  we get a critical value for  $E$  constrained on  $S_\rho$ , in the energy range  $(m, 2^{-s}m)$ .

Again, since the solution  $\bar{u}$  we have found verifies  $E(\bar{u}) \in (m, 2^{-s}m)$ , it is a constant sign solution by Proposition 2.2.

Case  $\Omega = \mathbb{R}^N$ .

Let  $r > 0$  and let us introduce the values

$$\tilde{\mathcal{A}}_r = \max \{E(w(\cdot - y)) : y \in B_r(0)\}, \quad (5.9)$$

$$\tilde{\mathcal{L}}_r = \max \{E(w(\cdot - y)) : y \in \partial B_r(0)\}. \quad (5.10)$$

Then it turns out that for every fixed  $V$  there exists  $r_V > 0$  such that for every  $r > r_V$

$$m < \tilde{\mathcal{L}}_r < C_0 \leq \tilde{\mathcal{A}}_r. \quad (5.11)$$

By Hölder inequality, for every  $y \in \mathbb{R}^N$

$$E(w(\cdot - y)) = m + \frac{1}{2} \int_{\mathbb{R}^N} V(x) [w(\cdot - y)]^2 dx \leq m + \frac{1}{2} |V|_q |w^2|_{q'}.$$

Hence,  $\tilde{\mathcal{A}}_r < 2^{-s}m$  provided

$$\frac{1}{2} |V|_q |w^2|_{q'} < \left(1 - \frac{1}{2^s}\right) (-m). \quad (5.12)$$

After some computation, by (2.7) with 1 in place of  $\rho$  and  $\rho^2$  in place of  $k$ , we see that (5.12) is equivalent to

$$|V|_q \rho^{-\frac{2s}{q}(q - \frac{N}{2})} < c, \quad (5.13)$$

for a suitable constant  $c > 0$  depending on  $N, p$  and  $q$ . Hence, setting for example

$$L = \frac{1}{2} c \rho^{\frac{2s}{q}(q - \frac{N}{2})}, \quad (5.14)$$

if  $|V|_q < L$  then the inequalities (5.11) can be completed as

$$m < \tilde{\mathcal{L}}_r < C_0 \leq \tilde{\mathcal{A}}_r < 2^{-s}m, \quad \forall r > r_V. \quad (5.15)$$

Moreover, by (5.13) we obtain (1.7) and (1.8). Finally, by (5.15) we can argue as in the proof of Theorem 1.1 to get the solution we are looking for.  $\square$

*Remark 5.1.* Let  $\rho > 0$  be fixed. Arguing as in [32, Theorem 3.1], it is possible to verify that

$$\lim_{r(\Omega) \rightarrow \infty} C_{0,0} = 2^{-s}m$$

(see(1.9)). Then by  $C_{0,0} \leq \mathcal{A}_{r,0} < 2^{-s}m$  and (5.8) we obtain

$$\lim_{r(\Omega) \rightarrow \infty} L = 0.$$

## 6 The case $N=1$

In this section we consider the 1-dimensional case

$$\begin{cases} -u'' + \lambda u + V(x)u = |u|^{p-2}u & \text{in } I, \\ \lambda \in \mathbb{R}, \quad u \in H_0^1(I), \quad \int_I u^2 dx = \rho^2, \end{cases} \quad (P_1)$$

where we can consider  $I = \mathbb{R}$  or  $I = (0, \infty)$ ,  $V \in L^q(I)$  for some  $q \in [1, \infty]$ ,  $V \geq 0$  and  $2 < p < 6$ .

First, let us consider the entire case. For  $N = 1$ , in the asymptotic behaviour of the limit function  $w$  (see (2.3) and (2.4)) we cannot take advantage of the polinomial contribution provided by  $|x|^{\frac{N-1}{2}}$ . As a consequence, the key estimate (4.3) in Proposition 4.1 does not hold and Theorem 1.1 fails.

On the other hand, Theorem 1.2 does not need Proposition 4.1 and it can be again stated:

**Theorem 6.1.** *Let  $I = \mathbb{R}$ ,  $\rho > 0$ ,  $V \in L^q(\mathbb{R})$ , for  $q \in [1, +\infty]$ , and  $V \geq 0$  in  $\mathbb{R}$ , then there exists a constant  $L = L(\rho) > 0$  such that if  $|V|_q < L$  then problem  $(P_1)$  has a positive solution.*

*Sketch of the proof.* This theorem can be proved exactly as Theorem 1.2-case  $\Omega = \mathbb{R}^N$ : we introduce  $\tilde{\mathcal{A}}_r$ ,  $C_0$  and  $\tilde{\mathcal{L}}_r$  (see (5.9), (5.3) and (5.10)) and then observe that for  $|V|_q$  small  $m < \tilde{\mathcal{L}}_r < C_0 \leq \tilde{\mathcal{A}}_r < 2^{-s}m$ . So, by Proposition 3.1, we can argue by deformation as in the proof of Theorem 1.1, getting the existence of a nonnegative solution  $\bar{u}$ . By Proposition 1.3, the solution  $\bar{u}$  is positive.  $\square$

According to the exterior case, that is  $I = (0, \infty)$ , first let us state some nonexistence results.

*Remark 6.2.* If  $I = (0, +\infty)$  and  $V(x) \equiv 0$ , then the autonomous problem  $(P_1)$  has no solution.

Indeed, if  $u$  is a solution of  $(P_1)$ , then it is a regular free solution of an equation of the form  $-u'' = f(u)$ ,  $u(0) = 0$ , where  $f(u) = -\lambda u + |u|^{p-2}u$ . Then  $u \equiv 0$  by [20, Remark I.3], contrary to  $u \in S_\rho$ .

**Proposition 6.3.** *Let  $I = \mathbb{R}$  or  $I = (0, \infty)$ . If  $V \in L^\infty(I)$  is a monotone locally Lipschitz function such that  $V' \neq 0$  on a set of positive measure, then problem  $(P_1)$  has no solution.*

Proposition 6.3 is a simplified version of Proposition 1.5 in the 1-dimensional case, adapted also to half lines.

*Proof.* We are assuming  $V$  non increasing and  $I = (0, \infty)$ .

Suppose by contradiction that there exists a solution  $u$  of  $(P_1)$ . Since the solutions of  $(P_1)$  are in  $H^2(I)$ , we can consider a sequence  $(u_n)_n$  in  $C_0^\infty((1/n, \infty))$  such that  $u_n \rightarrow u$  in  $H^2(I)$ . For every  $n \in \mathbb{N}$ , the map  $t \mapsto u_n(x+t)$ ,  $t \in (-1/n, \infty)$ , turns out to be a smooth curve in  $H^1(I)$  and we can consider the regular map  $t \mapsto f_n(t) := E(u_n(x+t))$ . Since  $u$  is a solution of  $(P_1)$ , we have

$$f'_n(0) = E'(u_n)[u'_n] = O(1). \quad (6.1)$$

From the monotonicity of  $V$  and Fatou's Lemma we infer

$$\begin{aligned} f'_n(0) &= \lim_{t \rightarrow 0} \frac{1}{t} [E(u_n(\cdot+t)) - E(u_n)] \\ &= \lim_{t \rightarrow 0^+} \frac{1}{2t} \int_{\frac{1}{n}}^{\infty} V(x) [u_n^2(x+t) - u_n^2(x)] dx \\ &= \lim_{t \rightarrow 0^+} \frac{1}{2} \int_{\frac{1}{n}}^{\infty} \frac{V(x-t) - V(x)}{t} u_n^2(x) dx \\ &\geq \frac{1}{2} \int_0^{\infty} (-V'(x)) u_n^2(x) dx \\ &\geq 0. \end{aligned} \quad (6.2)$$

Letting  $n \rightarrow \infty$  in (6.1) and (6.2), we obtain

$$\int_0^{\infty} V'(x) u^2(x) dx = 0,$$

that is a contradiction. Indeed, by assumption  $V' \leq 0$  a.e. in  $I$  and  $V' < 0$  on a set of positive measure, while  $|\{x \in I : u(x) = 0\}| = 0$  because  $u(x) = 0$  implies  $u'(x) \neq 0$  otherwise  $u \equiv 0$  by the Cauchy-Lipschitz theorem.

If  $I = \mathbb{R}$  we can proceed in a similar way. □

*Remark 6.4.* In the proof of Theorem 6.3, we cannot consider directly the curve in  $H_0^1(I)$  defined by  $\gamma(t) := u(\cdot+t)$ ,  $t \geq 0$ . Indeed,  $u'(0) \neq 0$  prevents  $u' \in H_0^1(I)$ , so  $\gamma$  would be not a regular curve in  $H_0^1(I)$ .

Finally, let us state an existence result on half lines, that inherits the topological structure of the entire case.

**Theorem 6.5.** *Let  $I = (0, \infty)$ ,  $\rho > 0$ ,  $V \in L^q(\mathbb{R})$ , for some  $q \in [1, +\infty]$  and  $V \geq 0$ . If  $|V|_q < L$ , with  $L$  as in Theorem 6.1, then there exists  $\bar{R} = \bar{R}(V, \rho) > 0$  such that problem  $(P_1)$  with  $V(x-R)$  has a positive solution for every  $R > \bar{R}$ .*

*Proof.* In order to get a solution, we are going to solve  $(P_1)$  with  $V$  fixed, on  $H_0^1((-R, \infty))$ .

Let us consider  $(P_1)$  on  $H^1(\mathbb{R})$ , with  $V$  fixed, and let us define  $\tilde{\mathcal{A}}_r$ ,  $C_0$  and  $\tilde{\mathcal{L}}_r$  as in the proof of Theorem 6.1. Moreover, let us fix a cut-off function  $\tilde{\vartheta} \in C^\infty(\mathbb{R}, [0, 1])$  such that  $\tilde{\vartheta}(x) = 0 \forall x \in (-\infty, 0)$ ,  $\tilde{\vartheta}(x) = 1 \forall x \in (1, \infty)$  and introduce

$$\begin{aligned} \tilde{\mathcal{A}}_{r,R} &= \max \left\{ E \left( \tilde{\vartheta}(\cdot+R) w(\cdot-y) \right) : y \in [-r, r] \right\}, \\ C_{0,R} &= \inf \{ E(u) : u \in H^1(-R, \infty), |u|_2 = \rho, \beta(u) = 0 \}, \\ \tilde{\mathcal{L}}_{r,R} &= \max \left\{ E \left( \tilde{\vartheta}(\cdot+R) w(\cdot-y) \right) : y \in \{-r, r\} \right\}. \end{aligned}$$

Then,

$$\lim_{R \rightarrow \infty} \tilde{\mathcal{A}}_{r,R} = \tilde{\mathcal{A}}_r, \quad \lim_{R \rightarrow \infty} C_{0,R} = C_0, \quad \lim_{R \rightarrow \infty} \tilde{\mathcal{L}}_{r,R} = \tilde{\mathcal{L}}_r.$$

Hence, for large  $R$ , we get

$$m < \tilde{\mathcal{L}}_{r,R} < C_{0,R} \leq \tilde{\mathcal{A}}_{r,R} < 2^{-s}m$$

and we can argue as in the proof of Theorem 1.1, by Proposition 3.1.  $\square$

## Appendix

The following proof could be shortened by taking into account, for example, Lemma B.3 in [39]. We develop some more details on the bootstrap procedure, for the sake of completeness.

For classical regularity results, we refer the reader for example to [30, §10], and in particular Theorem 10.2 therein, or to [23, §8].

*Proof of Proposition 1.3.* (a) Once proved  $\bar{u} \in C_{\text{loc}}^{0,\alpha}(\Omega)$  we are done. Indeed, when  $q > \frac{N}{2}$  we can apply the Harnack inequality and conclude that the nonnegative solutions are actually strictly positive in  $\Omega$  (see [36, Theorem 7.2.1]).

To verify that  $\bar{u} \in C_{\text{loc}}^{0,\alpha}(\Omega)$ , we begin by observing that the function  $\bar{u}$  verifies

$$-\Delta \bar{u} = \psi(x)\bar{u} \quad \text{for } \psi(x) = -\lambda - V(x) + |\bar{u}|^{p-2}, \quad x \in \Omega. \quad (\text{A.1})$$

Since  $V \in L_{\text{loc}}^q(\Omega)$  with  $q > N/2$ , we have that  $\psi \in L_{\text{loc}}^{q_1}(\Omega)$  with

$$q_1 = \min \left\{ \frac{2N}{(N-2)(p-2)}, q \right\}.$$

Notice also that  $q_1 > \frac{N}{2}$  because  $p < 2^*$ . By Hölder inequality,  $\psi\bar{u} \in L_{\text{loc}}^{r_1}(\Omega)$  with  $r_1$  defined by

$$\frac{1}{r_1} = \frac{1}{q_1} + \frac{N-2}{2N}.$$

By regularity results, we have that:

- 1) if  $r_1 > \frac{N}{2}$ , then  $\bar{u} \in C_{\text{loc}}^{0,\alpha}(\Omega)$  with  $\alpha < 2 - \frac{N}{r_1}$ ;
- 2) if  $r_1 = \frac{N}{2}$ , then  $\bar{u} \in L_{\text{loc}}^s(\Omega) \forall s > 1$ ;
- 3) if  $r_1 < \frac{N}{2}$ , then  $\bar{u} \in L_{\text{loc}}^{r_1 N / (N - 2r_1)}(\Omega)$ .

In case (1) the assertion is proved.

In case (2) it is readily seen that  $\psi\bar{u} \in L_{\text{loc}}^{s_1}(\Omega)$  for every  $1 \leq s_1 < q$ . Hence we can conclude as in the previous case, choosing  $N/2 < s_1 < q$ .

If (3) holds, then  $\psi \in L_{\text{loc}}^{q_2}(\Omega)$ , with

$$q_2 = \min \left\{ \frac{r_1 N}{(N - 2r_1)(p-2)}, q \right\}.$$

Since  $q_1 > \frac{N}{2}$ , we get  $\frac{r_1 N}{N - 2r_1} > \frac{2N}{N-2}$  and hence  $q_2 \geq q_1$ .

Now, repeating the same argument of the previous step, by Hölder inequality  $\psi\bar{u} \in L_{\text{loc}}^{r_2}(\mathbb{R}^N)$  with  $r_2$  defined by

$$\frac{1}{r_2} = \frac{1}{q_2} + \frac{N - 2r_1}{r_1 N},$$

and again we have

- 4) if  $r_2 \geq \frac{N}{2}$ , then  $\bar{u} \in \mathcal{C}_{\text{loc}}^{0,\alpha}(\Omega)$ ;  
5) if  $r_2 < \frac{N}{2}$ , then  $\bar{u} \in L_{\text{loc}}^{r_2 N / (N - 2r_2)}(\Omega)$ .

Observe that  $r_2 > r_1$  because  $q_2 \geq q_1$  and  $\frac{r_1 N}{N - 2r_1} > \frac{2N}{N - 2}$ .

Iterating this bootstrap argument, we claim that, after  $k$  steps,  $r_k \geq \frac{N}{2}$ , so that we are done. If, by contradiction, the claim is false, then for every integer  $k$  we define

$$q_k = \min \left\{ \frac{r_{k-1} N}{(N - 2r_{k-1})(p - 2)}, q \right\}, \quad \frac{1}{r_k} = \frac{1}{q_k} + \frac{N - 2r_{k-1}}{r_{k-1} N}. \quad (\text{A.2})$$

Inductively, it turns out that  $q_k \geq q_{k-1}$  and  $r_k > r_{k-1}$ , for any  $k \in \mathbb{N}$ , with  $q_k \leq q$  and  $r_k < \frac{N}{2}$ . Let us set

$$\lim_{k \rightarrow +\infty} r_k = R \quad \lim_{k \rightarrow +\infty} q_k = Q.$$

Getting  $k \rightarrow \infty$  in (A.2) we obtain

$$\frac{1}{R} = \frac{1}{Q} + \frac{N - 2R}{RN}$$

that implies  $Q = \frac{N}{2}$ , contrary to  $Q \geq q_1 > \frac{N}{2}$ . So, a contradiction arises and (a) is proved.

(b) The function  $\bar{u}$  verifies

$$-\Delta \bar{u} = \phi \quad \text{for } \phi(x) = -\lambda \bar{u} - V(x)\bar{u} + |\bar{u}|^{p-2}\bar{u}, \quad x \in \Omega.$$

If  $V \in L_{\text{loc}}^q(\Omega)$ , with  $q > N$ , then  $\bar{u} \in L_{\text{loc}}^\infty(\Omega)$ , by (a), and so  $\phi \in L_{\text{loc}}^q(\Omega)$ , that allows us to deduce  $\bar{u} \in \mathcal{C}^{1,\alpha}(\Omega)$ .

(c) This point follows by classical regularity results.

(d) By classical regularity results,  $\bar{u}$  is continuously differentiable. If  $\bar{u}$  is nonnegative, then it turns out to be positive, by the Harnack inequality (see [36, Theorem 7.2.1]).

(e) By a direct verification on  $(P_1)$ , we see that there exists  $\bar{u}''$  in  $L^q(\Omega)$ , in a weak sense. Hence, if  $q < 2$  then  $\bar{u}' \in W_{\text{loc}}^{1,q}(\Omega)$  and if  $q \geq 2$  then  $\bar{u}' \in H_{\text{loc}}^1(\Omega)$ . In any case, we can conclude that  $\bar{u}' \in \mathcal{C}_{\text{loc}}^{0,\alpha}(\Omega)$ , so the proof is complete.  $\square$

**Acknowledgments:** The authors thank the anonymous referee for bringing to their attention some recent papers and for useful suggestions.

**Funding:** The authors have been supported by the INdAM-GNAMPA group; R.M. acknowledges also the MIUR Excellence Department Project awarded to the Department of Mathematics, University of Rome ‘‘Tor Vergata’’, CUP E83C23000330006.

**Compliance with Ethical Standards:** We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

**Data availability:** Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

## References

- [1] C. O. Alves and C. Ji. Normalized solutions for the Schrödinger equations with  $L^2$ -subcritical growth and different types of potentials. *J. Geom. Anal.*, 32(5):Paper No. 165, 25, 2022.
- [2] A. Ambrosetti and A. Malchiodi. *Perturbation methods and semilinear elliptic problems on  $\mathbf{R}^n$* , volume 240 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2006.
- [3] A. Bahri and Y. Y. Li. On a min-max procedure for the existence of a positive solution for certain scalar field equations in  $\mathbf{R}^N$ . *Rev. Mat. Iberoamericana*, 6(1-2):1–15, 1990.
- [4] A. Bahri and P.-L. Lions. On the existence of a positive solution of semilinear elliptic equations in unbounded domains. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 14(3):365–413, 1997.
- [5] T. Bartsch and S. de Valeriola. Normalized solutions of nonlinear Schrödinger equations. *Arch. Math. (Basel)*, 100(1):75–83, 2013.
- [6] T. Bartsch, R. Molle, M. Rizzi, and G. Verzini. Normalized solutions of mass supercritical Schrödinger equations with potential. *Comm. Partial Differential Equations*, 46(9):1729–1756, 2021.
- [7] V. Benci and G. Cerami. Positive solutions of some nonlinear elliptic problems in exterior domains. *Arch. Rational Mech. Anal.*, 99(4):283–300, 1987.
- [8] H. Berestycki and P.-L. Lions. Nonlinear scalar field equations. I. Existence of a ground state. *Arch. Rational Mech. Anal.*, 82(4):313–345, 1983.
- [9] T. Cazenave. *Semilinear Schrödinger equations*, volume 10 of *Courant Lecture Notes in Mathematics*. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003.
- [10] T. Cazenave and P.-L. Lions. Orbital stability of standing waves for some nonlinear Schrödinger equations. *Comm. Math. Phys.*, 85(4):549–561, 1982.
- [11] G. Cerami. Some nonlinear elliptic problems in unbounded domains. *Milan J. Math.*, 74:47–77, 2006.
- [12] G. Cerami and R. Molle. On some Schrödinger equations with non regular potential at infinity. *Discrete Contin. Dyn. Syst.*, 28(2):827–844, 2010.
- [13] G. Cerami and R. Molle. Positive bound state solutions for some Schrödinger-Poisson systems. *Nonlinearity*, 29(10):3103–3119, 2016.
- [14] G. Cerami and D. Passaseo. Existence and multiplicity results for semilinear elliptic Dirichlet problems in exterior domains. *Nonlinear Anal.*, 24(11):1533–1547, 1995.
- [15] G. Cerami and D. Passaseo. The effect of concentrating potentials in some singularly perturbed problems. *Calc. Var. Partial Differential Equations*, 17(3):257–281, 2003.
- [16] G. Devillanova and S. Solimini. The role of planar symmetry and of symmetry constraints in the proof of existence of solutions to some scalar field equations. *Nonlinear Anal.*, 201:112060, 13, 2020.
- [17] S. Dovetta, E. Serra, and P. Tilli. Action versus energy ground states in nonlinear Schrödinger equations. *Math. Ann.*, 385(3-4):1545–1576, 2023.
- [18] I. Ekeland. On the variational principle. *J. Math. Anal. Appl.*, 47:324–353, 1974.
- [19] M. J. Esteban and P.-L. Lions. Existence and nonexistence results for semilinear elliptic problems in unbounded domains. *Proc. Roy. Soc. Edinburgh Sect. A*, 93(1-2):1–14, 1982/83.
- [20] M. J. Esteban and P.-L. Lions. Existence and nonexistence results for semilinear elliptic problems in unbounded domains. *Proc. Roy. Soc. Edinburgh Sect. A*, 93(1-2):1–14, 1982/83.

- [21] N. Ghoussoub. *Duality and perturbation methods in critical point theory*, volume 107 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1993. With appendices by David Robinson.
- [22] B. Gidas, W. M. Ni, and L. Nirenberg. Symmetry of positive solutions of nonlinear elliptic equations in  $\mathbf{R}^n$ . In *Mathematical analysis and applications, Part A*, volume 7 of *Adv. in Math. Suppl. Stud.*, pages 369–402. Academic Press, New York-London, 1981.
- [23] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [24] N. Ikoma and Y. Miyamoto. Stable standing waves of nonlinear Schrödinger equations with potentials and general nonlinearities. *Calc. Var. Partial Differential Equations*, 59(2):Paper No. 48, 20 pp., 2020.
- [25] L. Jeanjean. Existence of solutions with prescribed norm for semilinear elliptic equations. *Nonlinear Anal.*, 28(10):1633–1659, 1997.
- [26] L. Jeanjean and S.-S. Lu. Nonradial normalized solutions for nonlinear scalar field equations. *Nonlinearity*, 32(12):4942–4966, 2019.
- [27] M. K. Kwong. Uniqueness of positive solutions of  $\Delta u - u + u^p = 0$  in  $\mathbf{R}^n$ . *Arch. Rational Mech. Anal.*, 105(3):243–266, 1989.
- [28] S. Lancelotti and R. Molle. Positive solutions for autonomous and non-autonomous nonlinear critical elliptic problems in unbounded domains. *NoDEA Nonlinear Differential Equations Appl.*, 27(1):Paper No. 8, 28, 2020.
- [29] G. Leoni. *A first course in Sobolev spaces*, volume 181 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2017.
- [30] E. H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001.
- [31] R. Molle, M. Musso, and D. Passaseo. Positive solutions for a class of nonlinear elliptic problems in  $\mathbf{R}^N$ . *Proc. Roy. Soc. Edinburgh Sect. A*, 130(1):141–166, 2000.
- [32] R. Molle and D. Passaseo. On the behaviour of the solutions for a class of nonlinear elliptic problems in exterior domains. *Discrete Contin. Dynam. Systems*, 4(3):445–454, 1998.
- [33] R. Molle and D. Passaseo. Infinitely many positive solutions of nonlinear Schrödinger equations. *Calc. Var. Partial Differential Equations*, 60(2):Paper No. 79, 35, 2021.
- [34] R. Molle, G. Riey, and G. Verzini. Normalized solutions to mass supercritical Schrödinger equations with negative potential. *J. Differential Equations*, 333:302–331, 2022.
- [35] B. Noris, H. Tavares, and G. Verzini. Existence and orbital stability of the ground states with prescribed mass for the  $L^2$ -critical and supercritical NLS on bounded domains. *Anal. PDE*, 7(8):1807–1838, 2014.
- [36] P. Pucci and J. Serrin. *The maximum principle*, volume 73 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Verlag, Basel, 2007.
- [37] L. Song. Existence and orbital stability/instability of standing waves with prescribed mass for the  $l^2$ -supercritical nls in bounded domains and exterior domains. *Calculus of Variations and Partial Differential Equations*, 62(6):Paper No. 176, 27, 2023.
- [38] W. A. Strauss. Existence of solitary waves in higher dimensions. *Comm. Math. Phys.*, 55(2):149–162, 1977.
- [39] M. Struwe. *Variational methods*, volume 34. Springer-Verlag, Berlin, fourth edition, 2008. Applications to nonlinear partial differential equations and Hamiltonian systems.
- [40] Z. Yang, S. Qi, and W. Zou. Normalized solutions of nonlinear Schrödinger equations with potentials and non-autonomous nonlinearities. *J. Geom. Anal.*, 32(5):Paper No. 159, 27, 2022.

- [41] Z. Zhang and Z. Zhang. Normalized solutions of mass subcritical Schrödinger equations in exterior domains. *NoDEA Nonlinear Differential Equations Appl.*, 29(3):Paper No. 32, 25, 2022.