

First order expansion in the semiclassical limit of the Levy-Lieb functional

Original

First order expansion in the semiclassical limit of the Levy-Lieb functional / Colombo, M., Di Marino, S., Stra, F.. - In: ARCHIVE FOR RATIONAL MECHANICS AND ANALYSIS. - ISSN 1432-0673. - 249:(2025), pp. 1-57. [10.1007/s00205-025-02143-7]

Availability:

This version is available at: 11583/3012801 since: 2026-07-07T15:32:26Z

Publisher:

Springer

Published

DOI:10.1007/s00205-025-02143-7

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



First Order Expansion in the Semiclassical Limit of the Levy–Lieb Functional

MARIA COLOMBO , SIMONE DI MARINO & FEDERICO STRA

Communicated by G. FRIESECKE

Abstract

We prove the conjectured first order expansion of the Levy–Lieb functional in the semiclassical limit, arising from Density Functional Theory (DFT). In particular, we prove a general asymptotic first order lower bound in terms of the zero point oscillation functional and the corresponding asymptotic upper bound in the case of two electrons in one dimension. This is accomplished by interpreting the problem as the singular perturbation of an Optimal Transport problem via a Dirichlet penalization.

Mathematics Subject Classification: 81Q05 82C70 49Q20

Contents

1. Introduction	
2. Γ -limit of the unconstrained problem	
3. Rectangular truncation of Gaussians	
4. Construction of the recovery sequence	
4.1. Structure of optimal plans, maps and potentials in one dimension	
4.2. Choice of the parameters	
4.3. Construction of $\tilde{\gamma}_\varepsilon$	
5. Deconvolution of plans	
6. Potential cost of the remaining mass and kinetic energy of its marginals	
6.1. Marginals of kernels of product type along a linear map	
6.2. Wasserstein estimate of Proposition 6.2	
6.3. Kinetic energy of the remaining mass	
7. Proof of Theorems 1.1 and 1.3	
References	

1. Introduction

A revolutionary approach to finding the ground state energy of a many-body electron system was developed in the 19s by Hohenberg and Kohn [24]: The Density Functional Theory (DFT). Their idea, subsequently formalized mathematically by Lieb and Levy [30], can be seen as breaking up the minimization over all wave functions into a first minimization over those wavefunctions having a given one-particle density and then minimizing the resulting function over the one-particle density. In particular we define that

$$F^\varepsilon(\rho) = \inf \left\{ \int_{\mathbb{R}^{dN}} \left(\frac{\varepsilon}{2} |\nabla \psi|^2 + V_{\text{ee}} |\psi|^2 \right) dx_1 \dots dx_N \right\} \psi \in H^1(\mathbb{R}^{dN}), \psi \mapsto \rho, \quad (1.1)$$

where $\psi \mapsto \rho$ means that the one electron density of ψ is ρ , that is, for all $i = 1, \dots, N$, we require $\rho(x_i) = \int \int |\psi|^2(x_1, \dots, x_n) dx_1 \dots \widehat{dx_i} \dots dx_N$, and, moreover,

$$V_{\text{ext}}(x_1, \dots, x_n) = \sum_{i < j} \frac{1}{|x_i - x_j|}.$$

Then, for any external potential V_{ext} the corresponding energy of the ground state will be equal to

$$E(V_{\text{ext}}) = \inf \left\{ F^{\hbar^2}(\rho) + \int_{\mathbb{R}^d} V_{\text{ext}} d\rho \right\} \rho \in \mathcal{M}_+(\mathbb{R}^d), \rho(\mathbb{R}^d) = N.$$

In particular, $F^{\hbar^2}(\rho)$ is a *universal functional* in the sense that depends only on the number of electrons, whereas the dependence on the external field appears only in the term $\int V_{\text{ext}} d\rho$ in the outer minimization. It becomes then fundamental for applications to approximate $F^{\hbar^2}(\rho)$ and compute this value efficiently. In recent years a new approach gained importance, relying on the analysis of the Strictly Correlated Electrons (SCE) case. The interest in this approach resides in two main reasons: it is mathematically rigorous since it is a limiting procedure starting from the exact functional, and it is highly non-local in nature, thus it can be thought as complementary in some sense to the more classical Local Density Approximation (LDA) approach [27] (see also [28] for other universal approximating functionals). The idea is to have a parameter $\beta > 0$ which tunes the strength of the interaction between the electrons and the functional

$$F_\beta^{\hbar^2}(\rho) = \inf \left\{ \int_{\mathbb{R}^{dN}} \left(\frac{\hbar^2}{2} |\nabla \psi|^2 + \beta V_{\text{ee}} |\psi|^2 \right) dx_1 \dots dx_N \right\} \psi \in H^1(\mathbb{R}^{dN}), \psi \mapsto \rho.$$

The SCE limit considers the case $\beta \rightarrow \infty$, the asymptotic expansion of $F_\beta^{\hbar^2}(\rho)$ in β , to then deduce information about $\beta < \infty$. This limiting procedure has more equivalent formulations. Using the homogeneity of V_{ee} , $F_\beta^{\hbar^2}$ is computed from $F^{\hbar^2}$ by scaling, namely $F_\beta^{\hbar^2}(\rho) = \beta^2 F^{\hbar^2}(\rho_\beta)$ where $\rho_\beta(x) = \beta^{-d} \rho(x/\beta)$. Another equivalent approach considers a varying kinetic energy coefficient: since

$F_{\beta}^{\hbar^2}(\rho) = \beta F^{\hbar^2/\beta}(\rho)$, the asymptotic expansion of $F_{\beta}^{\hbar^2}$ as $\beta \rightarrow +\infty$ can be studied by means of the asymptotic expansion of $F^{\varepsilon}(\rho)$ as $\varepsilon \rightarrow 0$, which is mathematically convenient since the zeroth-order term of the expansion does not need to be renormalized. This is formally the same as changing the value of $\hbar$ in the original definition of the Levy–Lieb functional, but the rigorous justification of this non-physical procedure (since $\hbar$ is a physical constant) relies on the previous explanation.

Seidl was the first to conjecture the SCE limit case for $\varepsilon \rightarrow 0$ in [39] and further investigated in [23]. This new functional found its use, for example in [6, 32, 33, 43] it states that

$$\lim_{\varepsilon \rightarrow 0} F^{\varepsilon}(\rho) = F_{OT}(\rho) := \inf \left\{ \int_{\mathbb{R}^{3N}} V_{ee} d\gamma : \gamma \in \Pi_N(\rho) \right\},$$

where $\Pi_N(\rho)$ is the set of probabilities in $\mathbb{R}^{dN}$ which induce ρ , namely such that its push-forward through any projection $e_i : (x_1, \dots, x_N) \mapsto x_i$ is ρ . We denote with $\Pi_0(\rho)$ the set of $\gamma \in \Pi_N(\rho)$ which are minimizers for $F_{OT}(\rho)$. Moreover, in the physics literature, ansatz of minimizers in the 1D case and radial case were conjectured in [39, 40]: in the 1D case the conjecture was proven to hold in [9], while in the radial case various counterexamples were found in [3, 11, 38]. The functional F_{OT} has been studied in the last years also with regard to the limit of infinitely many particles: the zeroth order expansion for $N \rightarrow \infty$ was investigated in [15], proving the mean field limit, while the first order was proven independently in [26] and [16, 17], with ideas coming from the seminal papers [36, 37], and in connection with the Lieb–Oxford inequality [29, 31]. We refer the reader also to the recent review [20].

Already in [39], the next order in the asymptotic expansion of F^{ε} for $\varepsilon \rightarrow 0$ was conjectured in some cases, while the most general formulation appears in [22]:

$$F^{\varepsilon}(\rho) = F_{OT}(\rho) + \sqrt{\varepsilon} F_{ZPO}(\rho) + O(\varepsilon^{3/4}). \quad (1.2)$$

In both [22, 39] there is an explicit conjecture for the *zero point oscillation* functional $F_{ZPO}(\rho)$ in the case $d = 1$, involving the eigenvalues of the Hessian of an effective potential described in the following. As in [8], we introduce the potential u which solves the dual problem for F_{OT} ; its existence and regularity have been studied in [7, 10, 18, 21]. This is known to be unique if the support of ρ is connected, u is Lipschitz, $F_{OT}(\rho) = N \int u d\rho$,

$$V(x_1, \dots, x_N) = V_{ee}(x_1, \dots, x_N) - u(x_1) - \dots - u(x_N) \geq 0 \quad \forall x_1, \dots, x_N \in \mathbb{R}^d, \quad (1.3)$$

and the optimal plans are supported in the set $\{V = 0\}$. In particular, D^2V is well defined γ -almost everywhere for every optimal plan γ and so we can define that

$$F_{ZPO}(\rho) = \inf \left\{ \frac{1}{2} \int_{\mathbb{R}^{dN}} \text{tr} \left(\sqrt{D^2V} \right) d\gamma \right\} \gamma \in \Pi_0(\rho).$$

In the physics literature V is called the effective potential; u is the Lagrange multiplier associated with the constraint on the marginals, thought as a confining potential

which keeps the optimal plan concentrated on minimal energy configurations with respect to the effective potential; $\frac{1}{2} \operatorname{tr} \left(\sqrt{D^2 V} \right)$ is the lowest eigenvalue of the harmonic oscillator associated with the quadratic form given by $D^2 V$.

From the mathematical viewpoint, the 0-th order approximation was obtained in [13] for the bosonic case and for the fermionic case only when $N = 2$. Then it was settled in [2] for $N = 3$ with the help of spins; in [25] with a different approach involving mixed states (that is, considering the convex relaxation $\bar{F}_\varepsilon(\rho)$ of $F_\varepsilon(\rho)$); finally, [14] provided a proof using only pure states for every N .

Regarding the 1st order, the best known bound in [25] proves that $\bar{F}_\varepsilon(\rho) \leq F_{OT}(\rho) + \sqrt{\varepsilon} C_\rho$ for every admissible ρ , namely an upper bound on the conjectured order of the next term. Our main contribution can be summarised as follows:

Theorem 1.1. *Let $N \geq 2$ and $d \geq 1$ be fixed integers. Let $\rho \in \mathcal{P}(\mathbb{R}^d)$ be such that $F^1(\rho) < \infty$, u the optimal Kantorovich potential associated to the problem $F_{OT}(\rho)$ and suppose that V as in (1.3) is such that $V \in C_{loc}^2(\{V = 0\})$. Then*

- we always have

$$\liminf_{\varepsilon \rightarrow 0} \frac{F^\varepsilon(\rho) - F_{OT}(\rho)}{\sqrt{\varepsilon}} \geq F_{ZPO}(\rho); \quad (1.4)$$

- if in addition $N = 2$, $d = 1$ and $\rho \in C^1(\mathbb{R})$ is a positive probability density, we also have

$$\limsup_{\varepsilon \rightarrow 0} \frac{F^\varepsilon(\rho) - F_{OT}(\rho)}{\sqrt{\varepsilon}} \leq F_{ZPO}(\rho). \quad (1.5)$$

A few comments on the previous statement are listed below. Firstly, the regularity condition on V is natural when computing linear expansions and is automatically guaranteed under assumptions on ρ , namely the finiteness of its kinetic energy and a condition on its decay, see the observation following Theorem 1.3. Secondly, the restriction to $N = 2$ in the upper bound (1.5) is just used to simplify technical details and not make the notation heavier, the general case $N \geq 2$ will be treated in a forthcoming note. In turn, a different strategy would be needed in dimension $d \geq 2$. In fact we rely heavily on the one dimensional setting for the existence, regularity and structure of optimal maps; even the existence of optimal maps is an open problem if $d \geq 2$ and $N \geq 3$. Thirdly, in $d \geq 2$, one may wish to study the problem with the addition of the constraint of the antisymmetry on ψ (instead in $d = 1$ it is sufficient to consider carefully the signs): however we believe that, as in [25], this issue is just of technical nature and can be circumvented considering mixed states.

We now describe the approach to Theorem 1.1 and provide in Theorems 1.2 and 1.3 below two strengthened statements, which easily imply respectively the lower bound (1.4) and the upper bound (1.5). Our approach is based on the variational formulation for the quantity $\frac{F^\varepsilon(\rho) - F_{OT}(\rho)}{\sqrt{\varepsilon}}$. Using that $F_{OT}(\rho) = N \int u \, d\rho = \int_{\mathbb{R}^{dN}} (u(x_1) + \dots + u(x_N)) |\psi|^2(x_1, \dots, x_N) \, dx$ for every $\psi \mapsto \rho$, we obtain that

$$\frac{F^\varepsilon(\rho) - F_{OT}(\rho)}{\sqrt{\varepsilon}} = \min \{E_\varepsilon(\psi)\} \quad \psi \mapsto \rho, \quad (1.6)$$

$$E_\varepsilon(\psi) = \int_{\mathbb{R}^{dN}} \varepsilon^{1/2} \frac{|\nabla \psi(x)|^2}{2} + \varepsilon^{-1/2} V(x) |\psi(x)|^2 dx. \quad (1.7)$$

The core of Theorem 1.1 relies in a precise convergence result on E_ε . Firstly we study the convergence when we *drop the marginals constraint*, which simplifies our analysis but is not the one of physical interest, and we show, roughly speaking, that

$$E_\varepsilon(\psi) \xrightarrow{\Gamma} \begin{cases} \frac{1}{2} \int_{\mathbb{R}^{dN}} \text{tr} \left(\sqrt{D^2 V} \right) d\gamma & \text{if } \gamma(\{V > 0\}) = 0 \\ +\infty & \text{otherwise.} \end{cases} \quad (1.8)$$

The notion of convergence in (1.8) is the so called Γ -convergence, the right variational tool to take the limit in sequences of minimizers, as it happens in Theorem 1.1. For brevity, we don't introduce this abstract notion here but we just present its outcomes in Theorem 1.2. For $\mu_n, \mu \in \mathcal{M}(\mathbb{R}^d)$, we denote with $\mu_n \rightharpoonup \mu$ the weak-* convergence in duality with $C_0(\mathbb{R}^d)$ functions (continuous functions tending to zero at infinity).

Theorem 1.2. (Γ -convergence with free marginals) *Let $V \in C^0(\mathbb{R}^d)$ with $V \geq 0$ and V locally C^2 in a neighborhood of $\{V = 0\}$. Then we have*

(a) *for every sequence $\psi_\varepsilon \in H^1(\mathbb{R}^d)$ with $\int |\psi_\varepsilon|^2 = 1$ and $\liminf E_\varepsilon(\psi_\varepsilon) < \infty$, there exists $\gamma \in \mathcal{M}(\mathbb{R}^d)$ with $\gamma(\{V > 0\}) = 0$ and $\gamma(\mathbb{R}^d) \leq 1$ such that $|\psi_\varepsilon|^2 \rightharpoonup \gamma$ up to subsequences and*

$$\liminf_{\varepsilon \rightarrow 0} E_\varepsilon(\psi_\varepsilon) \geq \frac{1}{2} \int_{\mathbb{R}^{dN}} \text{tr} \left(\sqrt{D^2 V} \right) d\gamma;$$

(b) *for every $\gamma \in \mathcal{P}(\mathbb{R}^d)$ with $\gamma(\{V > 0\}) = 0$, there exists a recovery sequence $\psi_\varepsilon \in H^1(\mathbb{R}^d)$ with $\int |\psi_\varepsilon|^2 = 1$ such that $|\psi_\varepsilon|^2 \rightharpoonup \gamma$ and*

$$\limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\psi_\varepsilon) \leq \frac{1}{2} \int_{\mathbb{R}^{dN}} \text{tr} \left(\sqrt{D^2 V} \right) d\gamma.$$

In order to deal with the actual first order expansion of $F^\varepsilon(\rho)$ we have to enforce the marginal constraint. For the lower bound we can use Theorem 1.2 (a), since the weak-* convergence preserves marginal constraints. In order to prove the upper bound we need to construct a recovery sequence $\tilde{\psi}_\varepsilon$ with an almost optimal energy, but with the additional constraint on the marginals. This is in general nontrivial, since the Lavrentiev gap phenomenon could appear as happens for instance in an example of [14], where the constrained limit energy is smaller than the unconstrained one. This result is established in Theorem 1.3.

Theorem 1.3. *Let $\rho \in C^1(\mathbb{R})$ be a strictly positive probability density with finite kinetic energy*

$$\text{KE}(\rho) = \frac{1}{2} \int_{\mathbb{R}} \left| \partial_x \sqrt{\rho(x)} \right|^2 dx = \int_{\mathbb{R}} \frac{|\partial_x \rho(x)|^2}{8\rho(x)} dx, \quad (1.9)$$

let γ be the optimal transport plan for the problem with Coulomb cost and marginals ρ , let u be its Kantorovich potential, and assume that $u'' \in L^\infty(\mathbb{R})$. Let $V(x, y) = |x - y|^{-1} - u(x) - u(y)$.

Then there exists a recovery sequence $\psi_\varepsilon \in H^1(\mathbb{R}^2)$ such that $\int |\psi_\varepsilon|^2 = 1$, $|\psi_\varepsilon|^2 \rightharpoonup \gamma$, $|\psi_\varepsilon|^2$ has the same marginals as γ and

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\psi_\varepsilon) &= \limsup_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^2} \left(\varepsilon^{1/2} \frac{|\nabla \psi_\varepsilon|^2}{2} + \varepsilon^{-1/2} V |\psi_\varepsilon|^2 \right) dx \\ &\leq \frac{1}{2} \int_{\mathbb{R}^2} \text{tr}(\sqrt{D^2 V}) d\gamma. \end{aligned}$$

Theorem 1.3 is stated with a rather implicit assumption, namely that $u'' \in L^\infty(\mathbb{R})$. This is essentially equivalent to the condition (1.10) on the growth of the tails of ρ . For instance, we show in Lemma 4.1 that Theorem 1.3 can be applied to any strictly positive C^1 probability density ρ such that $\text{KE}(\rho) < \infty$ and

$$\liminf_{|x| \rightarrow \infty} |x|^3 \rho(x) > 0. \quad (1.10)$$

It would be interesting to prove the ZPO conjectured asymptotics for every ρ and in any dimension, in particular removing the heavy tails assumption (1.10) since one expects physical densities to have exponential tails.

Singular perturbations of optimal transport. Since, without loss of generality, we can assume that $\psi > 0$, we rewrite the functional $F^\varepsilon(\rho)$ in terms of $\gamma = |\psi|^2$ as

$$F^\varepsilon(\rho) = \inf \left\{ \int_{\mathbb{R}^{dN}} V_{ee}(x_1, \dots, x_n) d\gamma + \varepsilon \int_{\mathbb{R}^{dN}} \frac{|\nabla \gamma|^2}{\gamma} dx : \gamma \in \Pi_N(\rho) \right\}.$$

We think F^ε as a penalized multi-marginal optimal transport problem, where the penalization on the plan γ is the Fisher information with respect to the Lebesgue measure $I(\gamma) = \int \frac{|\nabla \gamma|^2}{\gamma} dx$. The ZPO conjecture is about the asymptotic expansion of this penalized OT problem. Other results in a similar spirit regard only the case $N = 2$ with the entropic penalization $\text{Ent}(\gamma) = \int \gamma \ln(\gamma) dx$ and either the Monge cost ([19], only in 1D), or the quadratic distance ([12] in a general Riemannian manifold) or a smooth convex cost ([35], in $\mathbb{R}^d$). However, in [12] and in [35] the peculiar structure of the problem and an additivity property of the entropy are heavily used. The approach closest to ours is [19], but the construction of the recovery sequence is heavily simplified by taking advantage of working in the region $\{T = \text{id}\}$. To our knowledge, the present result is the first next-order asymptotics with marginal constraint proven with a rather general strategy.

Theorem 1.2 works in a rather general setting and for any nonnegative potential which is locally C^2 around its zero set (we recall that here ρ does not play a role). Its proof is also short and flexible, sharing analogies with other previous results in the literature. For instance, similar studies look at the spectrum of the operator $-\Delta + \varepsilon^{-1} V$ as $\varepsilon \rightarrow 0$ (see for example [41, 42] and subsequent literature), but our point of view is slightly different. Instead of focusing on the behavior of the eigenvalues we want to know what is the minimal asymptotic energy given the

asymptotic behavior of the wavefunction (that is, their weak limit γ). This is in view of applying the result to a generic $\gamma \in \Pi_N(\rho)$ which might not come from a sequence of eigenvectors.

Theorem 1.2 builds on the idea that, to have the correct minimal energy, the wavefunction must localize around $\{V = 0\}$ and behave locally like a gaussian with variance depending on D^2V . Moreover this minimal energy can be achieved by a recovery sequence built superposing these gaussians, which will be elongated along the zero set of V . In the proof of Theorem 1.2 we construct a wavefunction for $\varepsilon > 0$ which has the proven almost correct energy and therefore can be of interest also for numerical purposes.

Ansatz on wavefunctions with asymptotically minimal energy. The ansatz of a recovery sequence obtained as a superposition of Gaussians is already present in [22], but then, instead of superposing elongated Gaussians, a local nonlinear change of variables is considered, as well as Gaussians in the variable perpendicular to the manifold $\{V = 0\}$. In our opinion, there are no guarantees on the smallness of the kinetic energy in the direction tangent to the manifold. In [22] the constraint on the marginals is also discussed via the introduction of a Lagrange multiplier. However, the latter has no reason to exist for $\varepsilon > 0$ and it is not clear the claim that it intervenes in the asymptotic expansion only at higher order terms.

Outline of the paper. The main idea in Theorem 1.2, once we show that every finite energy sequence needs to concentrate on the set $\{V = 0\}$, is to analyze the energy locally around the zero set of V : being at a minimum, the first non-zero term in the Taylor expansion of V is the quadratic form corresponding to the second derivative. But then the energy is analogous to that of an anisotropic harmonic oscillator, of which we know the ground state (a Gaussian with proper covariance) and ground energy (see Lemma 2.1 and Lemma 2.2). On one hand, an approximation of V with its Taylor expansion will provide the lower bound, while in order to get an upper bound, we use Gaussians elongated along the zero set of V , and then we superpose them to obtain

$$f(x) = \int f_{\varepsilon, x'}(x - x') d\gamma(x'), \quad (1.11)$$

where $f_{\varepsilon, x'}(x)$ is a suitable truncation of the Gaussian $C_{\varepsilon, x'} \exp(-\varepsilon^{-1/2} x' \sqrt{\nabla^2 V(x')x})$. The convexity of E_ε when written in terms of $\gamma = |\psi|^2$ lets us conclude the upper bound with no marginal constraint. Along our recovery sequence the contribution of the two competing functionals, namely the rescaled kinetic and potential energies, is asymptotically equal. This equipartition of energy is expected since, in the case of convergence to a single δ_0 and V quadratic, the kinetic energy rescales as $\text{KE}(\gamma_\lambda) = \lambda^2 \text{KE}(\gamma)$ and the potential energy as $\int V d\gamma_\lambda = \lambda^{-2} \int V d\gamma$, where $\gamma(x) = \lambda \gamma(\lambda x)$.

To tackle Theorem 1.3, namely to modify our ansatz (1.11) to keep the marginals constraint, we split the total mass in two components: a main portion of the mass and a remaining mass. The main part consists of most of the mass and must have the correct asymptotic energy. In Sect. 4 we prove that a suitable superposition of truncated Gaussians (introduced in Sect. 3), similar to the one adopted for Theorem 1.2, produces a sequence with asymptotically optimal energy. The super-

position is carefully chosen so as to leave a remaining mass with good properties for the subsequent construction.

The remaining mass is used to correct the marginals (see Sect. 5) with asymptotically vanishing cost (see Sect. 6). In the literature general methods for re-instating the marginals have been proposed, but they usually miss the correct order of magnitude for the kinetic energy or the potential energy: the only construction that controls in a fine way both of them has been proposed by [1], and then used then by [25] to retrieve in fact an upper bound with the correct order of magnitude. However the problem of this approximation is that it does not catch the correct local behavior because it is a superposition of gaussians with a fixed variance throughout the space. The latter issue motivates our choice of applying the deconvolution procedure in an unconventional way, namely only to the remaining mass, to connect the remaining marginals. Here we use the construction by Bindini in [1] since we are not interested in the sharp local energy but only the correct order of magnitude: in this way we have better control on the kinetic energy, which is bounded by the kinetic energy of the remaining marginals. This construction is presented in Sect. 5, where we show that this deconvolution procedure does not degrade too much the energy estimates. In particular, we estimate the potential energy of the deconvolved plan in terms of the potential energy of the original plan connecting the remaining marginals up to the correct order in ε . This section unveils the most delicate issue of the paper, needed to meet the assumption of the deconvolution. In Sect. 6 we prove that there exists $\tilde{\gamma}_H$ such that $\int V d\tilde{\gamma} = o(\sqrt{\varepsilon})$.

This last estimate of the potential energy means that the remaining marginals $\rho - \rho_\varepsilon^1$ and $\rho - \rho_\varepsilon^2$ can be connected by a plan which is concentrated near $\{V = 0\} = \text{graph } T$. To prove it, we show that $T_\#(\rho - \rho_\varepsilon^1)$ is close to $\rho - \rho_\varepsilon^2$ in the Wasserstein sense. The marginals are obtained by convolution with very anisotropic kernels with size $\varepsilon^{1/4}$ by $\beta^{1/2}$, which is much larger. A naive argument about balance of mass can show that the Wasserstein distance is at most $\beta^{1/2}$, but this is not sufficient. To show the desired estimate, we therefore need to employ a delicate analysis based on the Benamou-Brenier estimate of the transport cost, the linearization of the map T and a technical Proposition 6.3 to take advantage of the special structure of the projections of our truncated Gaussian kernels, introduced in Sect. 3.

2. Γ -limit of the unconstrained problem

In this section we prove Theorem 1.2. The strategy is to prove that for a finite energy sequence, the measure $|\psi_\varepsilon|^2 dx$ will concentrate around $\{V = 0\}$, and here the potential term can be well approximated by its second order Taylor expansion. More precisely, at each minimum point V will have some degenerate directions, and some other nondegenerate directions along which it is bounded from below by a suitable quadratic form.

We can thus look at the linearized problem: using the explicit solution (a gaussian with known variance) when V is a positive definite quadratic form, we estimate locally from below the energy around minimal points (Lemma 2.1), proving in particular Theorem 1.2(a) for V a nondegenerate quadratic potential. In Lemma 2.2

for any semidefinite quadratic form we build a sequence that attains the minimum energy in the limit, while concentrating the mass at a point. This proves Theorem 1.2(b) for $\gamma = \delta_0$ and the construction is inspired by the equality cases in Lemma 2.1, but needs to deal with the possible degeneracy of V at 0 and with a localization, since we have no information on the potential outside the origin.

Lemma 2.1. *Let A be a positive definite symmetric matrix such that $\lambda Id \leq A \leq \Lambda Id$ for some $\Lambda > \lambda > 0$. Whenever $B_{2r}(0) \subset \Omega \subseteq \mathbb{R}^n$, there exist a constant C depending only on λ and r , such that if $\sqrt{\varepsilon} \leq \lambda^2 r^2 / n \Lambda$, we have*

$$\int_{\Omega} \left(\varepsilon |\nabla \psi|^2 + |Ax|^2 |\psi|^2 \right) dx \geq \frac{\text{tr}(A) \sqrt{\varepsilon}}{1 + C \sqrt{\varepsilon}} \int_{\Omega} |\psi|^2 dx \quad \forall \psi \in H^1(\Omega). \quad (2.1)$$

Moreover, if $\Omega = \mathbb{R}^n$ we can take $C = 0$ in the previous formula and equality holds if and only if ψ is a multiple of $e^{-x^t Ax / 2\sqrt{\varepsilon}}$.

Proof. First let us prove that we can take $C = 0$ if $\Omega = \mathbb{R}^n$. In fact, for every $\psi \in C_c^\infty$ we can define $g(x) = \psi(x) e^{x^t Ax / 2}$. Then we have

$$\begin{aligned} |\nabla \psi|^2 &= \left| \nabla g e^{-x^t Ax / 2} - g A x e^{-x^t Ax / 2} \right|^2 \\ &= \left(|\nabla g|^2 - \nabla(g^2) \cdot Ax + g^2 |Ax|^2 \right) e^{-x^t Ax}. \end{aligned}$$

Integrating this identity $\mathbb{R}^n$ and then using the integration by parts formula we get

$$\begin{aligned} \int_{\mathbb{R}^n} |\nabla \psi|^2 dx &= \int_{\mathbb{R}^n} |\nabla g|^2 e^{-x^t Ax} dx + \int_{\mathbb{R}^n} |Ax|^2 g^2 e^{-x^t Ax} dx \\ &\quad + \int_{\mathbb{R}^n} g^2 \nabla \cdot (Ax e^{-x^t Ax}) dx \\ &\geq \int_{\mathbb{R}^n} |Ax|^2 g^2 e^{-x^t Ax} dx + \text{tr}(A) \int_{\mathbb{R}^n} g^2 e^{-x^t Ax} dx \\ &\quad - 2 \int_{\mathbb{R}^n} |Ax|^2 g^2 e^{-x^t Ax} dx \\ &= - \int_{\mathbb{R}^n} |Ax|^2 |\psi|^2 dx + \text{tr}(A), \end{aligned}$$

where we used that $|\nabla g|^2$ is nonnegative and that $g^2 e^{-x^t Ax} = |\psi|^2$. Using this inequality with $A/\sqrt{\varepsilon}$ and using the density of C_c^∞ in $H_0^1(\mathbb{R}^n)$ we have that

$$\int_{\mathbb{R}^n} \left(\varepsilon |\nabla \psi|^2 + |Ax|^2 |\psi|^2 \right) dx \geq \text{tr}(A) \sqrt{\varepsilon} \int_{\mathbb{R}^n} |\psi|^2 dx \quad \forall \psi \in H^1(\mathbb{R}^n); \quad (2.2)$$

moreover we can have equality in (2.2) using $g \equiv C_\varepsilon$. In particular the previous computation shows that $\psi_\varepsilon = C_\varepsilon e^{-x^t Ax / 2\sqrt{\varepsilon}}$ is the minimizer for $E_{\mathbb{R}^n}^\varepsilon$ (here C_ε is the normalization constant such that $\int |\psi_\varepsilon|^2 = 1$).¹

¹ This is a standard fact about the ground state energy of an n -dimensional harmonic oscillator: in fact the ground state energy of the 1-dimensional harmonic oscillator Hamiltonian

In the general case let us consider the cut off function $f_r(x) = \min\{1, \left(\frac{|x|}{r} - 2\right)_+\}$. We have that $|\nabla f_r| \leq \frac{1}{r} \chi_{B_{2r} \setminus B_r}$ and $f_r = 1$ on B_r while $f_r = 0$ on B_{2r}^c . In particular, for every function $\psi \in H^1(\Omega)$, we have $f_r \psi \in H_0^1(\mathbb{R}^n)$. However, we can then apply (2.2) in order to obtain

$$\int_{\Omega} \left(\varepsilon |\nabla(\psi f_r)|^2 + |Ax|^2 |\psi|^2 f_r^2 \right) dx \geq \sqrt{\varepsilon} \operatorname{tr}(A) \int_{\Omega} |\psi|^2 f_r^2. \quad (2.3)$$

Now, by Young's inequality, we have that, for every $\delta < 1$,

$$\begin{aligned} |\nabla(\psi f_r)|^2 &= |\nabla \psi|^2 f_r^2 + |\nabla f_r|^2 |\psi|^2 + 2 \nabla f_r \cdot \nabla \psi f_r \psi \\ &\leq (1 + \delta) |\nabla \psi|^2 f_r^2 + (1 + \delta^{-1}) |\nabla f_r|^2 |\psi|^2. \end{aligned} \quad (2.4)$$

Notice also that, since $|Ax|^2 \geq \lambda |x|^2 \geq \lambda^2 r^2$ if $|x| \geq r$, we have

$$(1 + \delta^{-1}) |\nabla f_r|^2 = \frac{\delta + 1}{\delta r^2} \chi_{B_{2r} \setminus B_r} \leq \frac{2}{\delta r^2} \cdot \frac{|Ax|^2}{\lambda^2 r^2}; \quad (2.5)$$

moreover we also have that

$$\operatorname{tr}(A)(1 - f_r^2) \leq n \Lambda \chi_{B_{2r} \setminus B_r} (1 - f_r^2) \leq \frac{n \Lambda |Ax|^2}{\lambda^2 r^2} (1 - f_r^2). \quad (2.6)$$

Now, starting from Equation (2.3) and then adding the term $\sqrt{\varepsilon} \operatorname{tr}(A) \int_{\Omega} |\psi|^2 (1 - f_r^2)$ on both sides, we can use (2.6), (2.4) and (2.5) to obtain

$$\begin{aligned} &\int_{\Omega} \varepsilon (1 + \delta) |\nabla \psi|^2 f_r^2 + \left(f_r^2 + \frac{n \Lambda \sqrt{\varepsilon}}{\lambda^2 r^2} (1 - f_r^2) + \frac{2\varepsilon}{\delta r^4 \lambda^2} \right) |Ax|^2 |\psi|^2 dx \\ &\geq \sqrt{\varepsilon} \operatorname{tr}(A) \int_{\Omega} |\psi|^2 dx; \end{aligned}$$

In particular, as long as $r^2 \lambda^2 \geq n \Lambda \sqrt{\varepsilon}$ we have $f_r^2 + \frac{n \Lambda \sqrt{\varepsilon}}{\lambda^2 r^2} (1 - f_r^2) \leq 1$ and then we optimize in δ , choosing $\delta = \frac{\sqrt{2\varepsilon}}{r^2 \lambda}$, getting

$$\left(1 + \frac{\sqrt{2\varepsilon}}{r^2 \lambda} \right) \cdot \int_{\Omega} \left(\varepsilon |\nabla \psi|^2 + |Ax|^2 |\psi|^2 \right) dx \geq \sqrt{\varepsilon} \operatorname{tr}(A) \int_{\Omega} |\psi|^2 dx,$$

which is precisely (2.1), with $C = \frac{\sqrt{2}}{r^2 \lambda}$. $\square$

Lemma 2.2. *Let us consider a continuous potential $V \geq 0$ such that $V(0) = 0$ and V is twice differentiable in 0. Then there exist a sequence ψ_{ε} such that $|\psi_{\varepsilon}|^2 \rightharpoonup \delta_0$ and*

$$\limsup_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^d} \left(\varepsilon^{1/2} \frac{|\nabla \psi_{\varepsilon}|^2}{2} + \varepsilon^{-1/2} V(x) |\psi_{\varepsilon}|^2 \right) dx \leq \frac{1}{2} \operatorname{tr} \left(\sqrt{D^2 V(0)} \right). \quad (2.7)$$

$h_i = -\varepsilon \partial_{x_i}^2 + (a_i x_i)^2$ is $\sqrt{\varepsilon} a_i$ and the corresponding wave function is $\psi_i = e^{-2a_i x_i^2 / (2\sqrt{\varepsilon})}$. After an orthonormal change of variables, any quadratic potential is equivalent to the n -dimensional Hamiltonian $\sum_i h_i$, which has ground state energy $\sqrt{\varepsilon} \sum_i a_i$ and ground state $\otimes_i \psi_i$.

Proof. First, for a positive definite symmetric matrix A , let us consider consider the function

$$f_\varepsilon = \left(e^{-\frac{x^t A x}{2\sqrt{\varepsilon}}} - e^{-N} \right)_+.$$

Then, letting $\Omega_\varepsilon = \{f_\varepsilon \neq 0\} = \{x^t A x < 2N\sqrt{\varepsilon}\}$, we claim that

$$\frac{\int_{\mathbb{R}^d} \varepsilon |\nabla f_\varepsilon|^2 + |Ax|^2 |f_\varepsilon|^2 dx}{\int_{\mathbb{R}^d} |f_\varepsilon|^2 dx} = \sqrt{\varepsilon} \operatorname{tr}(A) \cdot h(N) \quad (2.8)$$

for some universal function $h(N)$ such that $h(N) \rightarrow 1$ as $N \rightarrow \infty$. We postpone to the end the proof of the claim.

Now, since V has an absolute minimum at $x = 0$ we have that $D^2V(0)$ is a positive semidefinite symmetric matrix.

Since V is twice differentiable at 0, for $\eta > 0$ there exists an increasing continuous function $\delta(\eta) > 0$ with $\delta(\eta) \rightarrow 0$ for $\eta \rightarrow 0$ such that

$$2V(x) \leq x^t D^2V(0)x + \eta|x|^2 \leq |(\Lambda + \sqrt{\eta}\operatorname{Id})x|^2 \quad \forall x \in B_{\delta_\eta}(0).$$

Since the function $\frac{\sqrt{\eta}}{-\log \eta}$ is continuous and strictly increasing, for every ε sufficiently small there exists a unique $\eta = \eta(\varepsilon)$ such that $\sqrt{\varepsilon} = \frac{\sqrt{\eta}\delta(\eta)^2}{-2\log \eta}$. Now, let $A = \sqrt{D^2V(0)} + \sqrt{\eta}\operatorname{Id}$, $N = -\log \eta$ and $\psi_\varepsilon = f_\varepsilon / \sqrt{\int |f_\varepsilon|^2}$. With this choice of parameters, we get $\Omega_\varepsilon \subseteq B_\delta(0)$ (using that $A \geq \sqrt{\eta}\operatorname{Id}$). Since ψ_ε is also normalized, we get that $|\psi_\varepsilon|^2 \rightharpoonup \delta_0$. Moreover $2V(x) \leq |Ax|^2$ in Ω_ε and hence, we can say that

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(\varepsilon^{1/2} \frac{|\nabla \psi_\varepsilon|^2}{2} + \varepsilon^{-1/2} V(x) |\psi_\varepsilon|^2 \right) dx \\ & \leq \frac{1}{2\sqrt{\varepsilon}} \int_{\mathbb{R}^d} \left(\varepsilon |\nabla \psi_\varepsilon|^2 + |Ax|^2 |\psi_\varepsilon|^2 \right) dx \\ & = \frac{1}{2} \operatorname{tr}(A) \cdot h(-\log \eta) \\ & = \frac{1}{2} \left(\operatorname{tr}(\sqrt{D^2V(0)}) + d \cdot \sqrt{\eta} \right) \cdot h(-\log \eta). \end{aligned}$$

Letting $\varepsilon \rightarrow 0$, hence $\eta(\varepsilon) \rightarrow 0$, we deduce (2.7).

We now prove the claim (2.8). First of all,

$$\begin{aligned} |\nabla f_\varepsilon|^2 &= \frac{1}{\varepsilon} |Ax|^2 e^{-\frac{x^t A x}{\sqrt{\varepsilon}}} \chi_{\Omega_\varepsilon}, \\ |f_\varepsilon|^2 &= \chi_{\Omega_\varepsilon} \left(e^{-\frac{x^t A x}{\sqrt{\varepsilon}}} - 2e^{-\frac{x^t A x}{2\sqrt{\varepsilon}}} e^{-N} + e^{-2N} \right). \end{aligned}$$

We make the change of variable $z = \frac{\sqrt{A}}{\sqrt[4]{\varepsilon}} x$ which maps Ω_ε in $B_{\sqrt{2N}}(0) = B$. We obtain

$$\begin{aligned}
& \frac{\det \sqrt{A}}{\varepsilon^{1/2}} \int_{\Omega_\varepsilon} \varepsilon |\nabla f_\varepsilon|^2 + |Ax|^2 |f_\varepsilon|^2 dx \\
&= \int_B \sqrt{\varepsilon} z^t A z \cdot (2e^{-|z|^2} - 2e^{-\frac{|z|^2}{2}-N} + e^{-2N}) dz \\
&= \sqrt{\varepsilon} \cdot \frac{\operatorname{tr}(A)}{d} \int_B |z|^2 (2e^{-|z|^2} - 2e^{-\frac{|z|^2}{2}-N} + e^{-2N}) dz,
\end{aligned}$$

where in the second step we used that in $z^t A z = \sum_{i,j} A_{ij} z_j z_i$ every term with $i \neq j$ integrates to 0 (when multiplied by the function in brackets) because the function is antisymmetric and every term z_i^2 gives the same integral as $|z|^2/d$ by symmetry. Performing a similar calculation for $\int |f_\varepsilon|^2$, we find that

$$\frac{\int_{\Omega_\varepsilon} \varepsilon |\nabla f_\varepsilon|^2 + \frac{1}{\varepsilon} |Ax|^2 |f_\varepsilon|^2 dx}{\int_{\Omega_\varepsilon} |f_\varepsilon|^2 dx} = \frac{\operatorname{tr}(A)}{d} \frac{\int_B z^2 (2e^{-z^2} - 2e^{-\frac{z^2}{2}-N} + e^{-2N}) dz}{\int_B (e^{-z^2} - 2e^{-\frac{z^2}{2}-N} + e^{-2N}) dz}.$$

Since $\int_{\mathbb{R}^d} 2z^2 e^{-z^2} dz = d \int_{\mathbb{R}^d} e^{-z^2}$ it is easy to see that $h(N) := \frac{\int_B z^2 (2e^{-z^2} - 2e^{-\frac{z^2}{2}-N} + e^{-2N}) dz}{d \int_B (e^{-z^2} - 2e^{-\frac{z^2}{2}-N} + e^{-2N}) dz}$ is such that $h(N) \rightarrow 1$ as $N \rightarrow \infty$. $\square$

Proof of Theorem 1.2. Starting from point (a) first of all we have that $|\psi_\varepsilon|^2 \rightharpoonup \gamma$ for some $\gamma \in \mathcal{M}_+(\mathbb{R}^n)$ with $\gamma(\mathbb{R}^d) \leq 1$. Then, assuming $\liminf_{\varepsilon \rightarrow 0} E_\varepsilon(\psi_\varepsilon) = C < \infty$, we have that for every $\delta > 0$, as $\{V > \delta\}$ is an open set

$$\begin{aligned}
\gamma(\{V > \delta\}) &\leq \liminf_{\varepsilon \rightarrow 0} \int_{\{V > \delta\}} |\psi_\varepsilon|^2 \\
&\leq \liminf_{\varepsilon \rightarrow 0} \frac{\sqrt{\varepsilon}}{\delta} \cdot \frac{1}{\sqrt{\varepsilon}} \int_{\mathbb{R}^d} V(x) |\psi_\varepsilon|^2 \\
&\leq \lim_{\varepsilon \rightarrow 0} \frac{\varepsilon}{\delta} C = 0.
\end{aligned}$$

By the arbitrariness of δ we conclude that $\gamma(\{V > 0\}) = 0$.

Now we define the energy density $\mu_\varepsilon = \left(\varepsilon^{1/2} \frac{|\nabla \psi_\varepsilon|^2}{2} + \varepsilon^{-1/2} V(x) |\psi_\varepsilon|^2 \right) \mathcal{L}^d$; by hypothesis we know that the sequence μ_ε is uniformly bounded and in particular, up to subsequences, there exists a weak limit μ . By the semicontinuity of the mass, it will be sufficient to prove that the Radon-Nikodym derivative of μ with respect to γ is greater or equal than $\frac{1}{2} \operatorname{tr}(\sqrt{D^2 V})$. In particular, using a result by [34] (see Theorem 5.3 and Theorem 5.7 in [4] for a clearer explanation) we have that

$$\frac{d\mu}{d\gamma}(v_0) = \lim_{r \rightarrow 0} \frac{\mu(C_r)}{\gamma(C_r)},$$

where C_r are cylinders centered in v_0 , that is a set of the form $B_r^X(x_0) \times B_r^Y(y_0)$ for an orthogonal splitting $\mathbb{R}^d = X \times Y$ and $v_0 = (x_0, y_0)$. It is crucial to notice that any such cylinder is convex and $B_r(v_0) \subseteq C_r \subseteq B_{\sqrt{2}r}(v_0)$: in particular $C_r \in \mathcal{S}_{\sqrt{2}}(v_0)$ in the notation of [4].

Thus we are done if for every $v_0 \in \{V = 0\}$ and every $0 < \delta < 1$, for $r > 0$ small enough there is an open cylinder C_r such that

$$\mu(\overline{C_r}) \geq \liminf_{\varepsilon \rightarrow 0} \mu_\varepsilon(C_r) = \liminf_{\varepsilon \rightarrow 0} \left\{ \int_{C_r} \varepsilon^{1/2} \frac{|\nabla \psi_\varepsilon|^2}{2} + \varepsilon^{-1/2} V(x) |\psi_\varepsilon|^2 dx \right\} \quad (2.9)$$

$$\geq \frac{\sqrt{1-\delta}}{2} \operatorname{tr} \left(\sqrt{D^2 V(v_0)} \right) \gamma(C_r). \quad (2.10)$$

In order to do this let us suppose that $D^2 V \neq 0$, otherwise the statement is trivial. In that case, denote $A = \sqrt{D^2 V(v_0)}$ and make an orthonormal change of coordinates such that $D^2 V(v_0)$ becomes diagonal with the eigenvalues in decreasing order. Then, letting $n = \operatorname{rk}(D^2 V(v_0))$ we consider X the span of the first n eigenvectors and Y the remaining ones. In particular we have $c \operatorname{Id} \leq D^2 V(v_0)|_X \leq C \operatorname{Id}$ for some $C > c > 0$ (depending on v_0) and $D^2 V|_Y = 0$. Moreover, we have that $\nabla_X V(x_0, y_0) = 0$ and $\nabla_X \nabla_X V(x_0, y_0) = D^2 V(v_0)|_X$ is invertible, therefore by the implicit function theorem there exists $r > 0$ and a C^1 function $m : B_r^y(y_0) \rightarrow B_r^x(x_0)$ such that $\nabla_X V(m(y), y) = 0$ for every $y \in B_r^y(y_0)$. The gradient of the implicit function is

$$\nabla_Y m(y) = -(\nabla_{XX} V(m(y), y))^{-1} \nabla_{XY} V(m(y), y),$$

which vanishes for $y = y_0$ since $\nabla_{XY} V(v_0) = 0$. Therefore, up to possibly reducing r , we have that $|m(y) - x_0| \leq \delta |y - y_0|$ for every $y \in B_r^y(y_0)$. By the continuity of $D^2 V$, up to reducing r a second time, we have also $D^2 V(x, y)|_X \geq (1 - \delta) D^2 V(x_0, y_0)|_X$ for every $(x, y) \in B_r^x(x_0) \times B_r^y(y_0)$. Therefore, for every $(x, y) \in B_r^x(x_0) \times B_r^y(y_0)$, we consider the second order Taylor expansion of V at $(y, m(y))$ (recall that $V \geq 0$ and that $\nabla_X V$ vanishes at this point) to find that

$$V(x, y) \geq \frac{1-\delta}{2} (x - m(y))^T D^2 V(v_0)|_X (x - m(y)) = \frac{1-\delta}{2} |A(x - m(y))|^2,$$

where we denoted that $Ax = A(x, 0)$; since A is non degenerate only in the x directions in particular, we have $|Ax| \geq c|x|$ for some $c > 0$ and $\operatorname{tr}_x(A) = \operatorname{tr}(A)$.

Since $B_{r(1-\delta)}^x(m(y)) \subseteq B_r^x(x_0)$, we can use Lemma 2.1 in order to say that there exists a constant C such that, for every $y \in B_r^y(y_0)$,

$$\begin{aligned} & \int_{B_r^x(x_0)} \varepsilon \frac{|\nabla_x \psi_\varepsilon(x, y)|^2}{2} + V(x, y) |\psi_\varepsilon(x, y)|^2 dx \\ & \geq \frac{1}{2} \int_{B_r^x(x_0)} \varepsilon |\nabla_x \psi_\varepsilon|^2 + (1-\delta) |A(x - m(y))|^2 |\psi_\varepsilon|^2 dx \\ & \geq \sqrt{\varepsilon} \frac{\sqrt{1-\delta}}{1 + C\sqrt{\varepsilon}} \cdot \frac{\operatorname{tr}(A)}{2} \int_{B_r^x(x_0)} |\psi_\varepsilon(x, y)|^2 dx. \end{aligned}$$

Integrating with respect to y and then passing to the limit as $\varepsilon \rightarrow 0$ we obtain (2.9).

In order to prove part (b) we use Lemma 2.2 in case $\gamma = \delta_{v_0}$ and in the rest of the cases we argue by convexity: let us consider $\psi_\varepsilon^v(x) := \psi_\varepsilon(x - v)$ a recovery sequence for δ_v and let us define $f_\varepsilon^v = |\psi_\varepsilon^v|^2$. We notice that

$$\frac{1}{\sqrt{\varepsilon}} \int_{\mathbb{R}^d} \left(\varepsilon \frac{|\nabla \psi_\varepsilon^v|^2}{2} + V(x) |\psi_\varepsilon^v|^2 \right) dx = \frac{1}{\sqrt{\varepsilon}} \int_{\mathbb{R}^d} \left(\varepsilon \frac{|\nabla f_\varepsilon^v|^2}{8 f_\varepsilon^v} + V(x) f_\varepsilon^v \right) dx$$

denoting by $H_\varepsilon(f_\varepsilon^v)$ the functional in the right-hand side, we observe that H_ε is convex. Let us consider moreover the set $V_\varepsilon \subseteq \{V = 0\}$ defined as

$$V_\varepsilon = \left\{ v \in \{V = 0\} : H_\varepsilon(f_\varepsilon^v) \leq \frac{1}{2} \operatorname{tr}(\sqrt{D^2 V}(v)) + 1 \right\}.$$

By construction we have that $\limsup_\varepsilon H_\varepsilon(f_\varepsilon^v) \leq \frac{1}{2} \operatorname{tr}(\sqrt{D^2 V}(v))$ for every v ; in particular this implies that $\chi_{V_\varepsilon} \rightarrow \chi_{\{V=0\}}$ pointwise.

Now given a measure $\gamma \in \mathcal{P}(\mathbb{R}^d)$ with $\gamma(\mathbb{R}^d)$ supported in $\{V = 0\}$, by dominated convergence we get $\gamma(V_\varepsilon) \rightarrow 1$. In particular for ε sufficiently small $\gamma(V_\varepsilon) \geq \frac{1}{2}$ and for such ε we define the functions

$$f_\varepsilon^\gamma(x) = \frac{1}{\gamma(V_\varepsilon)} \int_{V_\varepsilon} f_\varepsilon^v(x) d\gamma.$$

Consider now $\psi_\varepsilon(x) = \sqrt{f_\varepsilon^\gamma(x)}$. By construction we have $\int |\psi_\varepsilon|^2 = 1$; moreover, by testing with a continuous function we see that $|\psi_\varepsilon|^2 \rightharpoonup \gamma$ and thanks to the convexity of H_ε we get that

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\psi_\varepsilon) &= \limsup_{\varepsilon \rightarrow 0} H_\varepsilon(f_\varepsilon^\gamma) \leq \limsup_{\varepsilon \rightarrow 0} \frac{1}{\gamma(V_\varepsilon)} \int_{V_\varepsilon} H_\varepsilon(f_\varepsilon^v) d\gamma \\ &= \limsup_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^d} \frac{\chi_{V_\varepsilon}}{\gamma(V_\varepsilon)} E_\varepsilon(\psi_\varepsilon^v) d\gamma. \end{aligned}$$

Now let us consider the functions $g_\varepsilon(v) = \frac{\chi_{V_\varepsilon}(v)}{\gamma(V_\varepsilon)} E_\varepsilon(\psi_\varepsilon^v)$ and $g(v) = \operatorname{tr}(\sqrt{D^2 V}(v))$; if g is not γ -integrable there is nothing to prove; so we can assume $g \in L^1(\gamma)$. By the definition of V_ε , for ε sufficiently small we have that

$$\frac{\chi_{V_\varepsilon}(v)}{\gamma(V_\varepsilon)} E_\varepsilon(\psi_\varepsilon^v) \leq \operatorname{tr}(\sqrt{D^2 V}(v)) + 1,$$

and we notice that the right-hand side is γ -integrable, otherwise there is nothing to prove. By Fatou's lemma, using also that $\gamma(V_\varepsilon) \rightarrow 1$, $\chi_{V_\varepsilon} \rightarrow 1$, we have

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\psi_\varepsilon) &\leq \limsup_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^d} \frac{\chi_{V_\varepsilon}}{\gamma(V_\varepsilon)} E_\varepsilon(\psi_\varepsilon^v) d\gamma \\ &\leq \int_{\mathbb{R}^d} \limsup_{\varepsilon \rightarrow 0} \frac{\chi_{V_\varepsilon}}{\gamma(V_\varepsilon)} E_\varepsilon(\psi_\varepsilon^v) d\gamma \\ &\leq \frac{1}{2} \int_{\mathbb{R}^d} \operatorname{tr}(\sqrt{D^2 V}) d\gamma. \end{aligned}$$

□

3. Rectangular truncation of Gaussians

As described in the introduction, Gaussian densities are asymptotically optimal for the energy of a single delta, therefore, to approach the general case, it is natural to construct a recovery sequence which is a superposition of Gaussian kernels. However, since we do not have global estimates on the potential V , only short-range interactions can be allowed, hence we need the kernels to have compact support. To achieve this, we truncate the densities with a slowly growing parameter N . The construction we present in Sect. 4.3 of the recovery sequence for Theorem 1.3 is then given by a convolution of x -dependent suitably rescaled and truncated Gaussian-like kernels, whose major axis is parallel at each point to graph T .

The requirements discussed above leads us to introduce some kernels which resemble Gaussian densities but have compact support, built in a suitable way that guarantees good properties for both the kinetic and potential energy.

Let M be a positive definite symmetric 2×2 matrix with eigenvalues $a \geq b > 0$ and let w and z be the orthonormal coordinates in the direction of the corresponding eigenvectors. For $N \in \mathbb{R}$ we define the unnormalized Gaussian and unnormalized truncated Gaussian

$$\begin{aligned}\tilde{\Gamma}_{M,\infty}(\mathbf{x}) &= e^{-\mathbf{x}^T M \mathbf{x}} = e^{-aw^2} e^{-bz^2}, \\ \tilde{\Gamma}_{M,N}(\mathbf{x}) &= \left(e^{-aw^2/2} - e^{-N/2} \right)_+^2 \left(e^{-bz^2/2} - e^{-N/2} \right)_+^2.\end{aligned}$$

In the definition of these truncated kernels, we adopt a rectangular truncation, which leads to kernels of product form; we call it rectangular because the support of the truncated Gaussian is a tilted rectangle, aligned with the eigenvectors of the matrix. This structure is crucial to do the computations in Sect. 6.1. Moreover, instead of the obvious truncation, we have to truncate the square root of the Gaussian, and then put a square outside, to guarantee the finiteness of the kinetic energy.

In the proof of the main theorem we will apply the results of this section to the matrix

$$M = M_{\varepsilon,\beta}(x) = \frac{A(x)}{\varepsilon^{1/2}} + \frac{I}{\beta},$$

where $A(x) = \sqrt{\nabla^2 V(x, T(x))}$, hence $a = q(x)/\varepsilon^{1/2} + 1/\beta$ and $b = 1/\beta$, with $q(x)$ being the non-zero eigenvalue of $A(x)$. The parameter $\beta \gg \varepsilon^{1/2}$ is required to ensure that the resulting Gaussian has compact support, since $\nabla^2 V(x, T(x))$ is a singular matrix, hence the natural choice of $M = A(x)/\varepsilon^{1/2}$ wouldn't work properly. We will discuss more about β when we present all the parameters in Sect. 4.2.

Define now the one dimensional integrals

$$\begin{aligned}G_{\alpha,\infty} &= \int_{\mathbb{R}} e^{-\alpha t^2} dt = \frac{\sqrt{\pi}}{\sqrt{\alpha}}, \\ G_{\alpha,N} &= \int_{\mathbb{R}} \left(e^{-\alpha t^2/2} - e^{-N/2} \right)_+^2 dt = \int_{-\sqrt{N/\alpha}}^{\sqrt{N/\alpha}} \left(e^{-\alpha t^2/2} - e^{-N/2} \right)_+^2 dt,\end{aligned}$$

and the integrals of the two dimensional densities

$$G_{M,\infty} = \int_{\mathbb{R}^2} \tilde{\Gamma}_{M,\infty}(\mathbf{x}) d\mathbf{x} = G_{a,\infty} G_{b,\infty} = \frac{\pi}{\sqrt{\det M}},$$

$$G_{M,N} = \int_{\mathbb{R}^2} \tilde{\Gamma}_{M,N}(\mathbf{x}) d\mathbf{x} = G_{a,N} G_{b,N}.$$

We can now introduce the normalized Gaussian and truncated Gaussian, which are probability densities, given by

$$\Gamma_{M,N}(\mathbf{x}) = \frac{\tilde{\Gamma}_{M,N}(\mathbf{x})}{G_{M,N}}, \quad N \in \mathbb{R} \cup \{\infty\}. \quad (3.1)$$

Finally, let $\eta_{M,\infty}$ and $\eta_{M,N}$ be the first marginal of $\Gamma_{M,\infty}$ and $\Gamma_{M,N}$, respectively.

We present here two lemmas, but we postpone their proof to Sect. 7.

Lemma 3.1. *With the definition above, there exists a constant $C > 0$ such that, for every $N \geq 3$, we have*

$$G_{M,N} < G_{M,\infty} < G_{M,N} + C e^{-N/2} G_{M,\infty}, \quad (3.2)$$

$$\|\Gamma_{M,N} - \Gamma_{M,\infty}\|_\infty \leq C \sqrt{\det M} e^{-N/2}, \quad (3.3)$$

$$\|\Gamma_{M,N} - \Gamma_{M,\infty}\|_1 \leq C N e^{-N/2}, \quad (3.4)$$

$$\|\eta_{M,N} - \eta_{M,\infty}\|_\infty \leq C \sqrt{a} \sqrt{N} e^{-N/2}. \quad (3.5)$$

Let us now compute the kinetic and potential energies of a Gaussian. In analogy with (1.9) we define the kinetic energy of a plan $\Gamma \in \mathcal{P}(\mathbb{R}^2)$ as

$$\text{KE}(\Gamma) = \frac{1}{2} \int_{\mathbb{R}^2} |\nabla \sqrt{\Gamma(\mathbf{x})}|^2 d\mathbf{x} = \int_{\mathbb{R}^2} \frac{|\nabla \Gamma(\mathbf{x})|^2}{8\Gamma(\mathbf{x})} d\mathbf{x}. \quad (3.6)$$

A direct computation shows that, if $Be_w = fe_w$ and $Be_z = ge_z$, then

$$\begin{aligned} \int_{\mathbb{R}^2} |B\mathbf{x}|^2 \Gamma_{M,\infty}(\mathbf{x}) d\mathbf{x} &= \int_{\mathbb{R}^2} (f^2 w^2 + g^2 z^2) \frac{\sqrt{ab}}{\pi} e^{-aw^2 - bz^2} dw dz \\ &= \frac{1}{2} \left(\frac{f^2}{a} + \frac{g^2}{b} \right). \end{aligned}$$

Therefore, using $|\nabla \sqrt{\Gamma_{M,\infty}}|^2 = \frac{1}{2} |\nabla [\log \Gamma_{M,\infty}]|^2 \Gamma_{M,\infty} = |M\mathbf{x}|^2 \Gamma_{M,\infty}(\mathbf{x})$ and the previous identity with $B = M$ we get

$$\text{KE}(\Gamma_{M,\infty}) = \frac{1}{2} \int_{\mathbb{R}^2} |M\mathbf{x}|^2 \Gamma_{M,\infty}(\mathbf{x}) d\mathbf{x} = \frac{\text{tr } M}{4}. \quad (3.7)$$

The following lemma compares the potential energy associated to the quadratic potential induced by the matrix M and the kinetic energy of the Gaussian $\Gamma_{M,\infty}$ and the truncated Gaussian $\Gamma_{M,N}$.

Lemma 3.2. *There is a universal constant $C > 0$ such that*

$$\int_{\mathbb{R}^2} |M\mathbf{x}|^2 |\Gamma_{M,N}(\mathbf{x}) - \Gamma_{M,\infty}(\mathbf{x})| d\mathbf{x} \leq C \text{tr}(M) N e^{-N/2}, \quad (3.8)$$

$$|\text{KE}(\Gamma_{M,N}) - \text{KE}(\Gamma_{M,\infty})| \leq C \text{KE}(\Gamma_{M,\infty}) e^{-N/2} = C \text{tr}(M) e^{-N/2}. \quad (3.9)$$

4. Construction of the recovery sequence

4.1. Structure of optimal plans, maps and potentials in one dimension

In this section we comment on the assumption on the boundedness of the second derivative of the Kantorovich potential u in Theorem 1.3 and we provide a class of ρ for which this assumption is satisfied and the main Theorem 1.3 is applicable: we will use the structural results for the 1D Coulomb multimarginal optimal transport problem contained in [9], which allows us to transfer the regularity of ρ to information on V and the optimal maps.

Let $\rho \in C^1_{\text{loc}}(\mathbb{R})$ be a strictly positive probability density. Without loss of generality, by translating we may assume that the origin 0 is the median of the probability ρ , i.e. $\rho((-\infty, 0]) = \rho([0, \infty)) = 1/2$. By the result in [9], the unique optimal plan is induced by a map T from ρ to itself which can be written explicitly in terms of the repartition function of ρ , which is C^2 with strictly positive first derivative. As a consequence, we have that $T \in C^2(\mathbb{R} \setminus \{0\})$, T is increasing in $\mathbb{R}_-$ and $\mathbb{R}_+$, $T(\mathbb{R}_-) = \mathbb{R}_+$, $T(\mathbb{R}_+) = \mathbb{R}_-$ and

$$\lim_{x \rightarrow -\infty} T(x) = 0, \quad \lim_{x \rightarrow 0^-} T(x) = \infty, \quad \lim_{x \rightarrow 0^+} T(x) = -\infty, \quad \lim_{x \rightarrow +\infty} T(x) = 0.$$

The Kantorovich potential $u : \mathbb{R} \rightarrow \mathbb{R}$ satisfies by definition $u(x) + u(y) \leq |x - y|^{-1}$ with equality if and only if $y = T(x)$; hence, using the first order condition of the maximality of $u(x) + u(y) - 1/|x - y|$ at points $(x, y) = (x, T(x))$, we get

$$u'(x) = -\frac{\text{sign}(x - T(x))}{(T(x) - x)^2} = -\frac{\text{sign}(x)}{(T(x) - x)^2} \quad \forall x \in \mathbb{R} \setminus \{0\}, \quad (4.1)$$

therefore u is determined up to a constant. Let V be as in (1.3), which in $d = 1$ and $N = 2$ reads as

$$V(x, y) = |x - y|^{-1} - u(x) - u(y) \quad \text{for } x, y \in \mathbb{R}. \quad (4.2)$$

Assuming that $x < 0$ and $y > 0$ (the other case is analogous), we can compute the gradient of the potential

$$\nabla V(x, y) = \left(\frac{1}{(y-x)^2} - u'(x), -\frac{1}{(y-x)^2} - u'(y) \right);$$

notice that $\nabla V(x, T(x)) = 0$. Differentiating again, we get the Hessian

$$\nabla^2 V(x, y) = \frac{2}{(y-x)^3} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} - \begin{pmatrix} u''(x) & 0 \\ 0 & u''(y) \end{pmatrix}. \quad (4.3)$$

In the proof of Theorem 1.3 we are interested in its behavior in a neighborhood of the graph of T , i.e. when $y = T(x)$.

The aim of the next lemma is twofold: first, we provide an assumption on the tails of ρ which is sufficient to obtain boundedness of the second derivatives of u . Second, we show how the assumption on the boundedness of u'' will be used in

the rest of the paper: through the computation of the Hessian of V , it will provide a sufficient condition to control the growth of V with a (uniform) parabola around the graph of T . We expect Theorem 1.3 to hold even without the uniform growth condition (4.4) below, but at the price of several technical complications that we don't address here.

Lemma 4.1. *Let $\rho \in C_{\text{loc}}^1(\mathbb{R})$ be a strictly positive probability density, let u be its Kantorovich potential and let V as in (4.2).*

(i) *If $u'' \in L^\infty(\mathbb{R})$, then there exists $\varepsilon_0, C > 0$ such that*

$$V(x, y) \leq Cd((x, y), \text{graph}(T))^2 \quad \forall (x, y) \text{ s.t. } d((x, y), \text{graph}(T)) < \varepsilon_0; \quad (4.4)$$

(ii) *if ρ satisfies also (1.10), then $u'' \in L^\infty(\mathbb{R})$ and $\nabla^2 V$ is locally Lipschitz in a neighborhood of $\text{graph}(T)$.*

Proof. Since $\rho \in L^1(\mathbb{R})$, there exists $\delta > 0$ such that every interval I of length $< \delta$ has $\int_I \rho(x) dx < 1/2$. Since $\int_x^{T(x)} \rho(x') dx' = 1/2$ for every $x < 0$ and $\int_{T(x)}^x \rho(x') dx' = 1/2$ for every $x > 0$, we obtain that $|T(x) - x| > \delta$, namely that $\text{graph}(T)$ has positive distance from the diagonal. The hessian of V is given by (4.3) and,

in a neighborhood of $\text{graph}(T)$, the first term is bounded because we are far from the diagonal $y = x$, whereas the second term is bounded everywhere since we have proved that $u'' \in L^\infty(\mathbb{R})$. This implies that V has controlled quadratic growth in a neighborhood of $\text{graph}(T)$, namely that $V(x, y) \leq Cd((x, y), \text{graph}(T))^2$ in $\mathbb{R}^2 \setminus \{|x - y| < \varepsilon\}$ for any $\varepsilon > 0$. The size ε of the neighborhood only depends on ρ (and in particular on the positive distance between the graph of the optimal map T and the diagonal $\{x = y\}$). This establishes (i).

Let us now prove (ii). From the formula (4.1) and the fact that $T \in C_{\text{loc}}^1(\mathbb{R} \setminus \{0\})$ we deduce that $u \in C_{\text{loc}}^2(\mathbb{R} \setminus \{0\})$. We need to check that $u''(x)$ stays bounded as $x \rightarrow 0^\pm$ and $x \rightarrow \pm\infty$. For $x < 0$, we have that

$$u''(x) = \frac{d}{dx} \left[\frac{1}{(T(x) - x)^2} \right] = 2 \frac{1}{(T(x) - x)^3} - 2 \frac{T'(x)}{(T(x) - x)^3}.$$

In both cases when $x \rightarrow -\infty$ or $x \rightarrow 0^-$, the first fraction goes to 0, therefore $u''(x)$ has the same asymptotic behavior as $T'(x)/(T(x) - x)^3$, let it be having a limit, being bounded or diverging. By the change of variable formula, we have moreover that

$$\frac{T'(x)}{(T(x) - x)^3} = \frac{\rho(x)}{(T(x) - x)^3 \rho(T(x))},$$

so our goal is to show that this fraction stays bounded as $x \rightarrow -\infty$ or $x \rightarrow 0^-$. The quantity is clearly non-negative, so we need to prove that the lim sup is finite. It is immediately apparent that

$$\begin{aligned}\limsup_{x \rightarrow 0^-} \frac{T'(x)}{(T(x) - x)^3} &= \limsup_{x \rightarrow 0^-} \frac{\rho(x)}{T(x)^3 \rho(T(x))} = \limsup_{y \rightarrow \infty} \frac{\rho(0)}{y^3 \rho(y)} \\ &= \frac{\rho(0)}{\liminf_{y \rightarrow \infty} y^3 \rho(y)} < \infty,\end{aligned}$$

because of (1.10). Let us now turn to studying the limit for $x \rightarrow -\infty$. Since $T(x) \rightarrow 0^+$ for $x \rightarrow -\infty$, we have

$$\limsup_{x \rightarrow -\infty} \frac{T'(x)}{(T(x) - x)^3} = \limsup_{x \rightarrow -\infty} \frac{\rho(x)}{-x^3 \rho(T(x))} = \limsup_{x \rightarrow -\infty} \frac{\rho(x)}{-x^3 \rho(0)} < \infty,$$

because $\rho \in L^\infty(\mathbb{R})$.

Exactly the same can be said for $x \rightarrow 0^+$ and $x \rightarrow \infty$, and therefore we have that $u \in C^2(\mathbb{R})$ and $u'' \in L^\infty(\mathbb{R})$.

Notice moreover that by (4.1) and the fact that $T \in C^2(\mathbb{R} \setminus \{0\})$ it follows that $u \in C^3(\mathbb{R} \setminus \{0\})$. Hence, denoting $D \subset \mathbb{R}^2$ the diagonal, by (4.3) we have that $\nabla^2 V$ is locally Lipschitz in $(\mathbb{R} \setminus \{0\})^2 \setminus D$, which is a neighborhood of $\text{graph}(T)$. $\square$

Observe that (1.10) and $\text{KE}(\rho) < \infty$ are simultaneously satisfied if

$$\liminf_{|x| \rightarrow \infty} |x|^{\alpha+1} \rho'(x) > 0 \quad \text{and} \quad \limsup_{|x| \rightarrow \infty} |x|^{\beta+1} \rho'(x) < \infty$$

for $\alpha, \beta \in (1, 3)$ with $1 < \beta \leq \alpha < 2\beta + 1$.

4.2. Choice of the parameters

The construction of $\tilde{\gamma}_\varepsilon$ depends on the choice of many parameters. Here we introduce them and specify which inequalities between them are necessary for the construction. First of all, we fix a parameter $H > 1$. All the constructions in the following sections depend on H , which is considered as a fixed parameter, but to ease the notation we don't always make this dependence explicit. This parameter H will play a role in the proof of Theorem 1.3 where we then use a diagonal argument to extract a recovery sequence.

Our construction relies on several structural assumptions. First of all, we need to split the mass in two components: the main part, where most of the energy comes from, and a remaining mass, used to fix the marginals of the recovery sequence. In order to do so, we need to lower the mass ρ by some constant, and for this reason we have to restrict ourselves to the bulk of the mass and avoid the tails at $\pm\infty$. Moreover, in order to be able to estimate the potential energy, we require a linearization of the map T (for the application of Proposition 6.3), whose error depends on T'' . Again for the application of Proposition 6.3, we want the slope of the linearized map to be far from 0 and ∞ . Since T has asymptotes at $\pm\infty$ and 0, we want then to work in a domain that stays away from those regions. Finally, we need a control on the eigenvalue of $\nabla^2 V$ evaluated on $\text{graph } T$, as this will determine the width of the Gaussians used in the construction.

All these reasons motivate the introduction of a set where we have good estimates of the objects intervening in the construction. The invariant domain that we will use is the following. Given $H > 0$ such that $\rho([H, \infty)) < 1/4$, define $0 < r_H < H$ such that $\rho([0, r_H]) = \rho([H, \infty))$ and consider the set

$$\Omega_H = [T(r_H), T(H)] \cup [r_H, H] \quad (4.5)$$

and the enlarged set

$$\Omega'_H = [T(r_{H+1}), T(H+1)] \cup [r_{H+1}, H+1] \supset \Omega_H. \quad (4.6)$$

We have that $T(\Omega_H) = T^{-1}(\Omega_H) = \Omega_H$ and $\Omega_H \nearrow \mathbb{R} \setminus \{0\}$, hence $\int_{\Omega_H^c} \rho(x) dx \rightarrow 0$. Moreover $\rho(\Omega_H \cap (-\infty, 0]) = \rho(\Omega_H \cap ([0, \infty))$ and similarly for Ω'_H .

Let $q(x) = \lambda_1(\nabla^2 V(x, T(x)))$ the positive eigenvalue. Notice that $q(x) > 0$ because, as already mentioned, from (4.3) follows that $\nabla^2 V$ cannot be the zero matrix. Moreover, q is a continuous function because the roots of a polynomial are continuous with respect to the coefficients.

As mentioned above, we want a control on some crucial quantities that influence the estimates of our construction. We therefore introduce the constant

$$\begin{aligned} L = L(H) = \max \Big\{ & \|\rho\|_{L^\infty(\Omega'_H)}, \|1/\rho\|_{L^\infty(\Omega'_H)}, \|1/T'\|_{L^\infty(\Omega'_H)}, \|T\|_{C^2(\Omega'_H)}, \\ & \|u\|_{C^2(\Omega'_H)}, \|q\|_{L^\infty(\Omega'_H)}, \|1/q\|_{L^\infty(\Omega'_H)}, \\ & \text{Lip}(\rho|_{\Omega'_H}), \text{Lip}(u''|_{\Omega'_H}) \Big\}. \end{aligned} \quad (4.7)$$

We now introduce the parameters on which our construction depends. They all depend on $\varepsilon > 0$, the variable that indexes the sequence of energies and the recovery sequence. First of all, we fix a parameter $N = N(\varepsilon) \rightarrow \infty$ faster than $\log \varepsilon$, which will be used to truncate the Gaussian-like kernels so that they have compact support such that

$$N(\varepsilon) = |\log \varepsilon|^{5/4}; \quad (4.8)$$

then a parameter $\beta = \beta(\varepsilon) \rightarrow 0$ such that

$$\varepsilon^{1/2} N \ll \beta \ll \varepsilon^{2/5}, \quad (4.9)$$

used to desingularize the matrix $\sqrt{\nabla^2 V(x, T(x))}$; then a parameter $\delta = \delta(\varepsilon) \rightarrow 0$ such that

$$(\beta N)^{1/2} \ll \delta \ll \varepsilon^{1/8} N^{-3/5}, \quad (4.10)$$

which controls the resolution of the linearization of the map T ; and finally a parameter $\tau = \tau(\varepsilon) \rightarrow 0$ such that

$$\tau \gg \frac{\delta^2}{\varepsilon^{1/4}}, \quad \tau \gg \frac{\varepsilon^{1/2} N^2}{\beta}, \quad \tau \gg \frac{\beta \delta N}{\varepsilon^{1/2}} \gg \frac{\beta \delta^2}{\varepsilon^{1/2}}, \quad \tau \gg \beta^{1/2} N^{1/2} \gg \varepsilon^{1/4} \gg \beta, \quad (4.11)$$

which controls the amount of mass that we subtract from ρ in order to subdivide it in main mass and remaining mass.

In the following sections, we will write many of the estimates relying only on these inequalities, so that their use will be easier in the future. Finally, we provide a choice of the parameters that fulfills the previous inequalities:

$$\begin{aligned} N(\varepsilon) &= |\log \varepsilon|^{5/4}, \quad \beta(\varepsilon) = \varepsilon^{1/2} |\log(\varepsilon)|^3, \quad \delta(\varepsilon) = \varepsilon^{1/8} |\log \varepsilon|^{-1}, \\ \tau(\varepsilon) &= |\log \varepsilon|^{-1/3}. \end{aligned} \quad (4.12)$$

4.3. Construction of $\bar{\gamma}_\varepsilon$

Let T be the optimal map between ρ and itself with respect to the Coulomb cost, which induces the plan γ . For every $(x, T(x)) \in \mathbb{R}^2$ on the graph of T , let us consider the truncated Gaussian $\Gamma_{M_{\varepsilon, \beta(x)}, N}$ given by (3.1) with covariance matrix $M_{\varepsilon, \beta}(x) = \sqrt{D^2 V(x, T(x))}/\varepsilon^{1/2} + \text{Id}/\beta$ and truncated through N .

We subdivide Ω_H into intervals $I_i^1 = [a_i, b_i]$ of length $\delta/2 < b_i - a_i < \delta$. Let T_δ be the piecewise linear interpolation of T on the intervals $I_i^1 = [a_i, b_i]$, namely a map that is affine in each I_i^1 and which coincides with T at the boundary of each interval. We define also $T_\delta = T$ outside of Ω_H . With this definition, writing $T(a_i)$ and $T(b_i)$ in terms of the Taylor expansion of T at x , we observe that

$$\begin{aligned} |T_\delta(x) - T(x)| &\leq \left| \frac{x - a_i}{b_i - a_i} T(a_i) + \frac{b_i - x}{b_i - a_i} T(b_i) - T(x) \right| \\ &\leq \|T''\|_{L^\infty(\Omega_H)} \delta^2 \leq L \delta^2 \end{aligned} \quad (4.13)$$

for every $x \in I_i^1$ and for every i . Let $I_i^2 = T(I_i^1) = T_\delta(I_i^1)$. Fix also points $x_i \in I_i^1$ such that $T'(x_i) = T'_\delta(x_i)$ that will be used to freeze the coefficients of the truncated Gaussians. The existence of such a point is ensured by Lagrange Theorem.

For convenience, in the following we use the notation $\Gamma_{M, N}^{\mathbf{x}}(\mathbf{x}') = \Gamma_{M, N}(\mathbf{x}' - \mathbf{x})$ to denote the Gaussian with covariance matrix M , truncated at level N , centered at $\mathbf{x}$.

Ideally, we would like to build a recovery sequence which is a superposition of Gaussians on $\text{supp } \gamma = \text{graph } T$ as

$$\gamma_\varepsilon(\mathbf{x}') = \int_{\mathbb{R}^2} \Gamma_{M_{\varepsilon, \beta(x)}, N}^{\mathbf{x}}(\mathbf{x}') d\gamma(\mathbf{x}) = \int_{\mathbb{R}} \Gamma_{M_{\varepsilon, \beta(x)}, N}^{(x, T(x))}(\mathbf{x}') d\rho(x).$$

Unfortunately, this plan has the wrong marginals, and there isn't a simple way to fix them without changing too much the energy. For instance one can notice that the construction in [1] pays an excessive price in the deconvolution step (this is analogous to applying Proposition 5.1 to the whole transport plan). We follow therefore a different route.

The actual construction of the recovery sequence is more involved. We split the total mass ρ in two components: a main portion of the mass and a remaining mass. The main part is dealt with in this section by superposing some truncated Gaussian kernels, whereas the remaining mass is used to bring back the marginals to what they need to be (see Sect. 5) without disrupting the energy estimates (see Sect. 6).

The main part of the recovery sequence is a suitable approximation of γ_ε , which is given by the plan

$$\bar{\gamma}_\varepsilon(\mathbf{x}') = \sum_{i=1} \bar{\gamma}_{\varepsilon,i}(\mathbf{x}'), \quad (4.14)$$

where each piece is built by superposing along the graph of T_δ Gaussians with frozen covariance matrix $M_{\varepsilon,\beta}(x_i)$ with $x_i \in I_i^1$, weighing them with the density $\rho - \tau$ (which is positive in Ω'_H for τ , and hence ε , sufficiently small):

$$\begin{aligned} \bar{\gamma}_{\varepsilon,i}(\mathbf{x}') &= \int_{I_i^1} \Gamma_{M_{\varepsilon,\beta}(x_i), N}^{(x, T_\delta(x))}(\mathbf{x}')(\rho(x) - \tau) dx \\ &= \int_{I_i^1} \Gamma_{M_{\varepsilon,\beta}(x_i), N}(\mathbf{x}' - (x, T_\delta(x)))(\rho(x) - \tau) dx \\ &= (\text{Id}, T_\delta)_\#((\rho - \tau)\mathbf{1}_{I_i^1}) * \Gamma_{M_{\varepsilon,\beta}(x_i), N}(\mathbf{x}'). \end{aligned} \quad (4.15)$$

We denote that two marginals by

$$\rho_\varepsilon^1 = \text{proj}_\#^1 \bar{\gamma}_\varepsilon, \quad \rho_\varepsilon^2 = \text{proj}_\#^2 \bar{\gamma}_\varepsilon. \quad (4.16)$$

The next proposition collects the properties of $\bar{\gamma}_\varepsilon$. First, it shows that its marginals are quantitatively below ρ , a property which crucially relies on the presence of τ in the definition of $\bar{\gamma}_\varepsilon$ and on the regularity of ρ and V . Second, it proves that this is still a recovery sequence (without prescribed marginals) as the one build in the proof of Theorem 1.2 was.

Proposition 4.2. *Under the same assumptions of Theorem 1.3, let $\bar{\gamma}_\varepsilon$ be as defined in (4.14), with $\bar{\gamma}_\varepsilon$ constructed with parameter H . Then,*

$$\lim_{\varepsilon \rightarrow 0} E_\varepsilon(\sqrt{\bar{\gamma}_\varepsilon}) \leq \frac{1}{2} \int_{\Omega_H} \text{tr} \left(\sqrt{\nabla^2 V(x, T(x))} \right) \rho(x) dx. \quad (4.17)$$

Moreover, there exists $0 < c_H < 1$ such that, for ε sufficiently small,

$$\rho_\varepsilon^i(x) \leq \rho(x) - c_H \tau \quad \text{for } i = 1, 2 \text{ and } x \in \Omega'_H, \quad (4.18)$$

and

$$(\rho - \rho_\varepsilon^i)(\mathbb{R}) = 1 - \bar{\gamma}_\varepsilon(\mathbb{R}^2) = \rho(\Omega_H^c) + \tau |\Omega_H|. \quad (4.19)$$

Proof. We use here the notation introduced in Sects. 3 and 4.

Step 1: proof of (4.18). Let us deal first with the case of ρ_ε^1 . First of all, notice that $\text{supp } \rho_\varepsilon^1 \subset \Omega'_H$ and $\rho > \tau$ on Ω'_H .

We have that

$$\rho_\varepsilon^1(x) = \sum_i \left([(\rho - \tau)\mathbf{1}_{I_i^1}] * \eta_i^1 \right)(x), \quad (4.20)$$

where $\eta_i^1 = \text{proj}_\#^1 \Gamma_{M_{\varepsilon,\beta}(x_i), N}$. For every x there can be at most two non zero terms contributing to the sum, because the convolutions kernels η_i^1 have support with diameter $\text{diam}(\text{supp } \eta_i^1) \leq C\beta^{1/2}N^{1/2}$, whereas two non-consecutive intervals are further away than $\text{d}(I_{i-1}^1, I_{i+1}^1) \geq C\delta \gg \beta^{1/2}N^{1/2}$, thanks to (4.10).

If $d(x, (I_i^1)^c) \geq C\beta^{1/2}N^{1/2}$, there is actually only the i -th term in the sum. Since the functions $(\rho - \tau)\mathbf{1}_{I_i^1}$ and $(\rho - \tau)$ coincide in $B(x, C\beta^{1/2}N^{1/2})$, when we convolve them with η_i^1 they give the same value at x , and therefore we can compute that

$$\begin{aligned}\rho_\varepsilon^1(x) &= \left([(\rho - \tau)\mathbf{1}_{I_i^1}] * \eta_i^1 \right)(x) = \left((\rho - \tau) * \eta_i^1 \right)(x) \\ &\leq \rho(x) - \tau + C \operatorname{Lip}(\rho|_{\Omega'_H}) \beta^{1/2} N^{1/2}.\end{aligned}$$

Otherwise, let x be a point where the two non zero terms in the sum (4.20) are $i - 1$ and i . Then, since the functions $\mathbf{1}_{I_{i-1}^1 \cup I_i^1}$ and 1 coincide in $B(x, C\beta^{1/2}N^{1/2})$, we have

$$\begin{aligned}\rho_\varepsilon^1(x) &= \left([(\rho - \tau)\mathbf{1}_{I_{i-1}^1}] * \eta_{i-1}^1 \right)(x) + \left([(\rho - \tau)\mathbf{1}_{I_i^1}] * \eta_i^1 \right)(x) \\ &= \left([(\rho - \tau)\mathbf{1}_{I_{i-1}^1 \cup I_i^1}] * \eta_{i-1}^1 \right)(x) + \left([(\rho - \tau)\mathbf{1}_{I_i^1}] * (\eta_i^1 - \eta_{i-1}^1) \right)(x) \\ &= \left((\rho - \tau) * \eta_{i-1}^1 \right)(x) + \left([(\rho - \tau)\mathbf{1}_{I_i^1}] * (\eta_i^1 - \eta_{i-1}^1) \right)(x) \\ &\leq \rho(x) - \tau + C \operatorname{Lip}(\rho|_{\Omega'_H}) \beta^{1/2} N^{1/2} + \|\rho\|_\infty \|\eta_i^1 - \eta_{i-1}^1\|_1 \\ &\leq \rho(x) - \tau + CL\beta^{1/2}N^{1/2} + C_H\varepsilon^{-1/4}\delta^2 + C_H\varepsilon^{-1/2}\beta\delta\end{aligned}$$

because $\|\eta_i^1 - \eta_{i-1}^1\|_1 \leq \|\Gamma_{M_{\varepsilon,\beta}(x_i),N} - \Gamma_{M_{\varepsilon,\beta}(x_{i-1}),N}\|_1 \leq C_H\varepsilon^{-1/4}\delta^2 + C_H\varepsilon^{-1/2}\beta\delta$ by Lemma 4.3 (with $\mathbf{x}_1 = \mathbf{x}_2 = 0$). Therefore (4.18) follows (with any $c_H < 1$) because, from (4.8)–(4.11), we have $\tau \gg \varepsilon^{-1/2}\beta\delta N \gg \varepsilon^{-1/2}\beta\delta$, $\tau \gg \beta^{1/2}N^{1/2}$ and $\tau \gg \varepsilon^{-1/4}\delta^2$.

Let us now deal with the case of ρ_ε^2 . From the definitions (4.14)–(4.16), we have

$$\begin{aligned}\rho_\varepsilon^2(y) &= \sum_i \operatorname{proj}_\#^2 \tilde{\gamma}_{\varepsilon,i}(y) = \sum_i \operatorname{proj}_\#^2 \left((\operatorname{Id}, T_\delta)_\# ((\rho - \tau)\mathbf{1}_{I_i^1}) * \Gamma_{M_{\varepsilon,\beta}(x_i),N} \right)(y) \\ &= \sum_i \left(\operatorname{proj}_\#^2 (\operatorname{Id}, T_\delta)_\# ((\rho - \tau)\mathbf{1}_{I_i^1}) \right) * \left(\operatorname{proj}_\#^2 \Gamma_{M_{\varepsilon,\beta}(x_i),N} \right)(y) \\ &= \sum_i \left([T_\delta \# ((\rho - \tau)\mathbf{1}_{I_i^1})] * \eta_i^2 \right)(y) = \sum_i \left([T_\delta \# (\rho - \tau)\mathbf{1}_{I_i^2}] * \eta_i^2 \right)(y),\end{aligned}$$

where $\eta_i^2 = \operatorname{proj}_\#^2 \Gamma_{M_{\varepsilon,\beta},N}$. For $y \in \Omega_H$ we have

$$\begin{aligned}|T_\delta \# \rho(y) - \rho(y)| &= |T_\delta \# \rho(y) - T_\# \rho(y)| = \left| \frac{\rho(T_\delta^{-1}(y))}{T'_\delta(T_\delta^{-1}(y))} - \frac{\rho(T^{-1}(y))}{T'(T^{-1}(y))} \right| \\ &\leq \frac{|\rho(T_\delta^{-1}(y)) - \rho(T^{-1}(y))|}{T'(T^{-1}(y))} \\ &\quad + \rho(T_\delta^{-1}(y)) \frac{|T'_\delta(T_\delta^{-1}(y)) - T'(T^{-1}(y))|}{T'_\delta(T_\delta^{-1}(y))T'(T^{-1}(y))} \\ &\leq \operatorname{Lip}(\rho|_{\Omega'_H})C\delta L + L \operatorname{Lip}(T'|_{\Omega'_H})C\delta L^2 \leq C_H\delta,\end{aligned}$$

where we used (4.7) to bound the norms of ρ and T' and the Lipschitz constants. In particular, if $y \in I_i^2$, then $T_{\delta\#}(\rho - \tau)(y) = T_{\delta\#}\rho(y) - \tau/T'(x_i) \leq \rho(y) + C_H\delta - \tau/L$.

We have, therefore, that

$$\rho_\varepsilon^2(y) \leq \sum_i \left([(\rho - \tau/L + C_H\delta)\mathbf{1}_{I_i^2}] * \eta_i^2 \right)(y).$$

We observe the similarity with (4.20), where $\rho - \tau$, I_i^1 and η_i^1 have been replaced by $\rho - \tau/L + C_H\delta$, I_i^2 and η_i^2 respectively. We have $\text{diam}(I_i^2) \geq C\delta/L \gg \beta^{1/2}N^{1/2}$ and $\text{diam}(\text{supp } \eta_i^2) \leq C\beta^{1/2}N^{1/2}$, so with the same argument as before we obtain

$$\rho_\varepsilon^2(y) \leq \rho(y) - \tau/L + C_H(\delta + \beta^{1/2}N^{1/2} + \varepsilon^{-1/4}\delta^2 + \varepsilon^{-1/2}\beta\delta),$$

from which the thesis follows with any $c_H < 1/L(H)$ because τ is asymptotically larger than all the error terms.

Step 2: proof of (4.19).

Since $\Gamma_{M_{\varepsilon,\beta}(x_i),N}^{(x,T_\delta(x))}$ is a probability measure, we have

$$\begin{aligned} \bar{\gamma}_\varepsilon(\mathbb{R}^2) &= \sum_i \int_{\mathbb{R}} \int_{I_i^1} \Gamma_{M_{\varepsilon,\beta}(x_i),N}^{(x,T_\delta(x))}(\mathbf{x}')(\rho(x) - \tau) d\mathbf{x}' \\ &= \sum_i \int_{I_i^1} (\rho(x) - \tau) dx = \int_{\Omega_H} (\rho(x) - \tau) dx = \rho(\Omega_H) - \tau|\Omega_H|, \end{aligned}$$

from which

$$1 - \bar{\gamma}_\varepsilon(\mathbb{R}^2) = 1 - [\rho(\Omega_H) - \tau|\Omega_H|] = \rho(\Omega_H^c) + \tau|\Omega_H|.$$

Step 3.

For every i and $x \in I_i^1$, we claim the estimate

$$\begin{aligned} E_\varepsilon \left(\sqrt{\Gamma_{M_{\varepsilon,\beta}(x_i),N}^{(x,T_\delta(x))}} \right) - E_\varepsilon \left(\sqrt{\Gamma_{M_{\varepsilon,\beta}(x),N}^{(x,T(x))}} \right) \\ \leq C_H(\varepsilon^{-3/4}\delta^2 + \varepsilon^{-1}\beta\delta)(\delta^4 + \varepsilon^{1/2}N + \beta N\delta^2) + C_H\varepsilon^{1/2}e^{-N/2} + C_H\delta. \end{aligned} \quad (4.21)$$

Notice that the right hand side of (4.21) goes to 0. In fact, thanks to (4.10) and the fact that $N \rightarrow \infty$ we have

$$\beta N\delta^2 \ll \delta^4 \ll \varepsilon^{1/2} \ll \varepsilon^{1/2}N,$$

therefore, thanks to (4.8)–(4.10), the right hand side of (4.21) is less than

$$\begin{aligned} C_H(\varepsilon^{-3/4}\delta^2 + \varepsilon^{-1}\beta\delta)\varepsilon^{1/2}N + C_H\varepsilon^{1/2}e^{-N/2} + C_H\delta \\ \leq C_H(\varepsilon^{-1/4}\delta^2 + \varepsilon^{-1/2}\beta\delta)N + C_H\varepsilon^{1/2}e^{-N/2} + C_H\delta \\ \ll N|\log \varepsilon|^{-2} + \varepsilon^{1/40}N^{2/5} + e^{-N/2} + \delta \ll 1. \end{aligned}$$

We observe that by (4.13) the points $\mathbf{x}_1 = (x, T_\delta(x))$, $\mathbf{x}_2 = (x, T(x)) \in \mathbb{R}^2$ and the symmetric matrices $A_1 = \sqrt{\nabla^2 V(x_i, T(x_i))}$, $A_2 = \sqrt{\nabla^2 V(x_2)}$ which generate $M_1 = M_{\varepsilon, \beta}(x_i)$ and $M_2 = M_{\varepsilon, \beta}(x)$ satisfy

$$|\mathbf{x}_1 - \mathbf{x}_2| \leq L\delta^2 \quad \text{and} \quad |A_1 - A_2| \leq C_H\delta. \quad (4.22)$$

In fact, $(x_i, T(x_i))$ and $(x, T(x))$ belong to a region of $\text{graph}(T)$ where $\nabla^2 V$ is Lipschitz. As a consequence,

$$|M_1 - M_2| = \left| \left(\frac{A_1}{\varepsilon^{1/2}} - \frac{I}{\beta} \right) - \left(\frac{A_2}{\varepsilon^{1/2}} - \frac{I}{\beta} \right) \right| = \varepsilon^{-1/2} |A_1 - A_2| \leq C_H \varepsilon^{-1/2} \delta. \quad (4.23)$$

As regards the potential energy, denoting by E the union of the supports of the probability measures appearing in (4.21)

$$E = \text{supp } \Gamma_{M_{\varepsilon, \beta}(x_i), N}^{(x, T_\delta(x))} \cup \text{supp } \Gamma_{M_{\varepsilon, \beta}(x), N}^{(x, T(x))}$$

we have

$$\begin{aligned} & \int_{\mathbb{R}^2} V(\mathbf{x}') \Gamma_{M_{\varepsilon, \beta}(x_i), N}^{(x, T_\delta(x))}(\mathbf{x}') d\mathbf{x}' - \int_{\mathbb{R}^2} V(\mathbf{x}') \Gamma_{M_{\varepsilon, \beta}(x), N}^{(x, T(x))}(\mathbf{x}') d\mathbf{x}' \\ & \leq \left(\sup_E V \right) \int_{\mathbb{R}^2} \left| \Gamma_{M_{\varepsilon, \beta}(x_i), N}^{(x, T_\delta(x))}(\mathbf{x}') - \Gamma_{M_{\varepsilon, \beta}(x), N}^{(x, T(x))}(\mathbf{x}') \right| d\mathbf{x}' \end{aligned} \quad (4.24)$$

Observe that, by Lemma 4.3, we have

$$\int_{\mathbb{R}^2} \left| \Gamma_{M_{\varepsilon, \beta}(x_i), N}^{(x, T_\delta(x))}(\mathbf{x}') - \Gamma_{M_{\varepsilon, \beta}(x), N}^{(x, T(x))}(\mathbf{x}') \right| d\mathbf{x}' \leq C_H \varepsilon^{-1/4} \delta^2 + C_H \varepsilon^{-1/2} \beta \delta. \quad (4.25)$$

We claim that E is contained in a strip of size $C(\delta^2 + \varepsilon^{1/4} N^{1/2} + \beta^{1/2} N^{1/2} \delta \text{Lip}(T'))$ around the graph of the affine map which equals T_δ in I_i^1

$$E \subseteq S = \{\mathbf{x} \in \mathbb{R}^2 : d(\mathbf{x}, \text{graph}(T_\delta)) \leq C(\delta^2 + \varepsilon^{1/4} N^{1/2} + \beta^{1/2} N^{1/2} \delta \text{Lip}(T'))\}. \quad (4.26)$$

Indeed, by construction $\text{supp } \Gamma_{M_{\varepsilon, \beta}(x_i), N}^{(x, T_\delta(x))}$ is a rectangle centered on the graph of T_δ and with the longer side aligned with the graph of T_δ . Moreover, its width is $\varepsilon^{1/4} N^{1/2}$; hence the claim (4.26) is proved for the first rectangle in the definition of E .

On the other side, we observe that the rectangle $\text{supp } \Gamma_{M_{\varepsilon, \beta}(x), N}^{(x, T(x))}$ is centered in the point $(x, T(x))$, whose distance from the graph of T_δ is estimated by $\|T''\|_{L^\infty} \delta^2$ by (4.13); moreover, this rectangle is tilted with respect to the previous one proportionally to $|T'(x) - T'_\delta(x)| \leq \delta \text{Lip}(T')$, which in turn gives a contribution of $\beta^{1/2} N^{1/2} \delta \text{Lip}(T')$. This proves (4.26).

Since for every x the point $(x, T(x))$ and the point are at distance at most $(x, T_\delta(x)) \|T''\|_{L^\infty} \delta^2$ by (4.13), from (4.26) we deduce that S is contained in the strip of width $C(1 + \|T''\|_{L^\infty})(\delta^2 + \varepsilon^{1/4} N^{1/2} + \beta^{1/2} N^{1/2} \delta)$ around the graph of T . Hence, recalling that both V and ∇V vanish along the graph of T , and more

precisely recalling that V has uniform quadratic growth around the graph of T has stated in we obtain that

$$\sup_E V \leq C(1 + \|T''\|_{L^\infty})^2(\delta^2 + \varepsilon^{1/4}N^{1/2} + \beta^{1/2}N^{1/2}\delta)^2. \quad (4.27)$$

Hence we deduce by (4.24), (4.25), and (4.27) that

$$\begin{aligned} \int V(\mathbf{x}') \Gamma_{M_{\varepsilon,\beta}(x_i),N}^{(x,T_\delta(x))}(\mathbf{x}') d\mathbf{x}' - \int V(\mathbf{x}') \Gamma_{M_{\varepsilon,\beta}(x),N}^{(x,T(x))}(\mathbf{x}') d\mathbf{x}' \\ \leq C_H(\varepsilon^{-1/4}\delta^2 + \varepsilon^{-1/2}\beta\delta)(\delta^2 + \varepsilon^{1/4}N^{1/2} + \beta^{1/2}N^{1/2}\delta)^2 \\ \leq C_H(\varepsilon^{-1/4}\delta^2 + \varepsilon^{-1/2}\beta\delta)(\delta^4 + \varepsilon^{1/2}N + \beta N\delta^2). \end{aligned} \quad (4.28)$$

In view of (4.28), we deduce the corresponding statement to (4.21) for the potential energy.

As regards the kinetic energy, defined by (3.6), we observe that it equals

$$\begin{aligned} \text{KE}(\Gamma_{M_1,N}^{x_1}) - \text{KE}(\Gamma_{M_2,N}^{x_2}) &= \text{KE}(\Gamma_{M_1,\infty}^{x_1}) - \text{KE}(\Gamma_{M_2,\infty}^{x_2}) \\ &\quad + [\text{KE}(\Gamma_{M_1,N}^{x_1}) - \text{KE}(\Gamma_{M_1,\infty}^{x_1})] \\ &\quad - [\text{KE}(\Gamma_{M_2,N}^{x_2}) - \text{KE}(\Gamma_{M_2,\infty}^{x_2})]. \end{aligned}$$

The first difference is

$$\text{KE}(\Gamma_{M_1,\infty}^{x_1}) - \text{KE}(\Gamma_{M_2,\infty}^{x_2}) = \frac{\text{tr } M_1}{4} - \frac{\text{tr } M_2}{4} = \frac{q_1 - q_2}{4\varepsilon^{1/2}} \leq \frac{\delta}{4\varepsilon^{1/2}}$$

because, by the variational characterization of the largest eigenvalue, we have

$$q_1 = \max_{|v| \leq 1} v^T A_1 v \leq \max_{|v| \leq 1} v^T A_2 v + \max_{|v| \leq 1} v^T (A_1 - A_2) v \leq q_2 + C_H \delta.$$

Using (3.9) for the two terms in square brackets, we get

$$\text{KE}(\Gamma_{M_1,N}^{x_1}) - \text{KE}(\Gamma_{M_2,N}^{x_2}) \leq C(\text{tr } M_1 + \text{tr } M_2)e^{-N/2} + \frac{C_H \delta}{\varepsilon^{1/2}} \leq C_H e^{-N/2} + \frac{C_H \delta}{\varepsilon^{1/2}}.$$

Step 4. We conclude the proof. By convexity of the kinetic energy, since the potential energy is linear and thanks to (4.21), with the remark following it, we have that

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\sqrt{\tilde{\gamma}_\varepsilon}) &\leq \limsup_{\varepsilon \rightarrow 0} \sum_{i=1} \int_{I_i^1} E_\varepsilon \left(\sqrt{\Gamma_{M_{\varepsilon,\beta}(x_i),N}^{(x,T_\delta(x))}} \right) \rho(x) dx \\ &\leq \limsup_{\varepsilon \rightarrow 0} \int_{\Omega_H} E_\varepsilon \left(\sqrt{\Gamma_{M_{\varepsilon,\beta}(x),N}^{(x,T(x))}} \right) \rho(x) dx. \end{aligned}$$

We now estimate the integrand in the right-hand side with $\text{tr} \left(\sqrt{\nabla^2 V(x, T(x))} \right) \rho(x)$ up to small errors. To this end, we bound the kinetic energy by passing to the non-truncated Gaussians with the estimate (3.9) and by the exact computation (3.7). We

obtain

$$\begin{aligned} \varepsilon^{1/2} K E \left(\sqrt{\Gamma_{M_{\varepsilon,\beta}(x),N}^{(x,T(x))}} \right) &= \varepsilon^{1/2} \text{KE} \left(\sqrt{\Gamma_{M_{\varepsilon,\beta}(x),N}} \right) \\ &\leq \varepsilon^{1/2} \text{KE} \left(\sqrt{\Gamma_{M_{\varepsilon,\beta}(x),\infty}} \right) + C \varepsilon^{1/2} \text{tr}(M_{\varepsilon,\beta}(x)) e^{-N/2} \\ &= \varepsilon^{1/2} \frac{\text{tr} M_{\varepsilon,\beta}(x)}{4} + C e^{-N/2} = \frac{1}{4} \text{tr} \left(\sqrt{\nabla^2 V(x, T(x))} \right) + \frac{\varepsilon^{1/2}}{2\beta} + C e^{-N/2}. \end{aligned}$$

In order to estimate the potential energy, we need to compare the V with its second order Taylor expansion. Since $\nabla^2 V$ is locally Lipschitz in a neighborhood of graph T , there is $r > 0$ such that for every $x \in \text{supp } \gamma \cap \Omega_H^2$ and $y \in B_r(0)$ we have

$$\left| V(x+y) - \frac{1}{2} \left| \sqrt{\nabla^2 V(x)} y \right|^2 \right| \leq \|\nabla^3 V\|_{L^\infty} \cdot |y|^3 \leq C_H |y|^3,$$

where the L^∞ norm is taken in a neighborhood of size $(\beta N)^{1/2}$ of $\text{supp } \gamma \cap \Omega_H^2$. Then, thanks to (3.8)² and (3.7), and since $|y| \leq (\beta N)^{1/2}$ for $y \in \text{supp } \Gamma_{M_{\varepsilon,\beta}(x),N}$, we have

$$\begin{aligned} \varepsilon^{-1/2} \int V(y) \Gamma_{M_{\varepsilon,\beta}(x),N}^{(x,T(x))}(y) dy &\leq \frac{\varepsilon^{-1/2}}{2} \int_{\mathbb{R}^2} \left| \sqrt{\nabla^2 V(x, T(x))} y \right|^2 \Gamma_{M_{\varepsilon,\beta}(x),N}(y) dy \\ &\quad + C_H \varepsilon^{-1/2} \int_{\mathbb{R}^2} |y|^3 \Gamma_{M_{\varepsilon,\beta}(x),N}(y) dy \\ &\leq \frac{\varepsilon^{-1/2}}{2} \int_{\mathbb{R}^2} \left| \sqrt{\nabla^2 V(x, T(x))} y \right|^2 \Gamma_{M_{\varepsilon,\beta}(x),\infty}(y) dy \\ &\quad + C \varepsilon^{-1/2} N e^{-N/2} + C_H \varepsilon^{-1/2} \beta^{3/2} N^{3/2} \\ &\leq \frac{1}{4} \text{tr} \left(\sqrt{\nabla^2 V(x, T(x))} \right) + C \varepsilon^{-1/2} N e^{-N/2} + C_H \varepsilon^{-1/2} \beta^{3/2} N^{3/2}. \end{aligned}$$

Overall, from the estimates in Step 4 we obtain that

$$\begin{aligned} E_\varepsilon(\sqrt{\gamma_\varepsilon}) &\leq \frac{1}{2} \int_{\Omega_H} \text{tr} \left(\sqrt{\nabla^2 V(x, T(x))} \right) \rho(x) dx \\ &\quad + C_H \left(\frac{\varepsilon^{1/2}}{\beta} + C e^{-N/2} + C \varepsilon^{-1/2} N e^{-N/2} + C_H \varepsilon^{-1/2} \beta^{3/2} N^{3/2} \right), \end{aligned}$$

which concludes the proof of (4.17) since by (4.8) and (4.9) we have

$$\begin{aligned} \frac{\varepsilon^{1/2}}{\beta} + C e^{-N/2} + C \varepsilon^{-1/2} N e^{-N/2} + C_H \varepsilon^{-1/2} \beta^{3/2} N^{3/2} \\ \ll \frac{1}{N} + C e^{-N/2} + \varepsilon^{1/10} N^{3/2} \ll 1. \end{aligned}$$

² When applying (3.8) to the integral of $\left| \sqrt{\nabla^2 V(x)} y \right|^2$ times the difference of the Gaussians, we use the fact that the matrix $\sqrt{\nabla^2 V(x)}$ is bounded from above by $\varepsilon^{1/2} M_{\varepsilon,\beta}(x)$.

As a side note, it could be easily seen by means of the kind of computations performed in this proof that (4.17) is actually an equality, but since we will not need this in the sequel we do not pursue this matter here. $\square$

Lemma 4.3. *Let $\mathbf{x}_1, \mathbf{x}_2 \in \mathbb{R}^2$ with $|\mathbf{x}_1 - \mathbf{x}_2| \leq C_0 \delta^2$ and let A_1, A_2 be two degenerate positive-semidefinite symmetric 2×2 matrices with $|A_i| \leq L$ and $|A_1 - A_2| \leq C_0 \delta$; finally let $M_i = M_{\varepsilon, \beta}(A_i) = \frac{A_i}{\varepsilon^{1/2}} + \frac{I}{\beta}$. Then there exists a positive constant C_H only depending on H and C_0 such that*

$$\|\Gamma_{M_1, N}^{\mathbf{x}_1} - \Gamma_{M_2, N}^{\mathbf{x}_2}\|_1 \leq C_H \varepsilon^{-1/4} \delta^2 + C_H \varepsilon^{-1/2} \beta \delta.$$

Observe that, under the assumptions (4.8)–(4.10), the right hand side goes to 0 when $\varepsilon \rightarrow 0$.

Proof. Thanks to (3.4) we have

$$\|\Gamma_{M_1, N}^{\mathbf{x}_1} - \Gamma_{M_2, N}^{\mathbf{x}_2}\|_1 \leq \|\Gamma_{M_1, \infty}^{\mathbf{x}_1} - \Gamma_{M_2, \infty}^{\mathbf{x}_2}\|_1 + C_N e^{-N/2}.$$

Notice that, by (4.8)–(4.10), $N e^{-N/2} \ll N^2 \varepsilon^{1/4} = \varepsilon^{1/2} N^2 \varepsilon^{-1/4} \ll \beta N \varepsilon^{-1/4} \ll \delta^2 \varepsilon^{-1/4}$, so we can forget about this error term because it is dominated by the right hand side of the thesis.

We split the estimate into first changing only the center point, and then changing the matrix. By the triangle inequality we have

$$\|\Gamma_{M_1, \infty}^{\mathbf{x}_1} - \Gamma_{M_2, \infty}^{\mathbf{x}_2}\|_1 \leq \|\Gamma_{M_1, \infty}^{\mathbf{x}_1} - \Gamma_{M_1, \infty}^{\mathbf{x}_2}\|_1 + \|\Gamma_{M_1, \infty}^{\mathbf{x}_2} - \Gamma_{M_2, \infty}^{\mathbf{x}_2}\|_1. \quad (4.29)$$

The first norm can be estimated in the following way. If a and b are the eigenvalues of M_1 , we have $\nabla \Gamma_{M_1, \infty}(w, z) = 2(aw, bz)^T \Gamma_{M_1, \infty}(w, z)$, therefore

$$\begin{aligned} \|\nabla \Gamma_{M_1, \infty}\|_1 &\leq G_{M_1, \infty}^{-1} \int_{\mathbb{R}^2} (2a|w| + 2b|z|) e^{-aw^2 - bz^2} dw dz \\ &= \frac{\sqrt{ab}}{\pi} \left(\frac{2\sqrt{\pi}}{\sqrt{b}} + \frac{2\sqrt{\pi}}{\sqrt{a}} \right) = \frac{2}{\sqrt{\pi}} (\sqrt{a} + \sqrt{b}) \\ &\leq C q(x)^{1/2} \varepsilon^{-1/4} \leq C_H \varepsilon^{-1/4}. \end{aligned}$$

From this we get

$$\begin{aligned} \|\Gamma_{M_1, \infty}^{\mathbf{x}_1} - \Gamma_{M_1, \infty}^{\mathbf{x}_2}\|_1 &= \int_{\mathbb{R}^2} |\Gamma_{M_1, \infty}(\mathbf{y} + \mathbf{x}_2 - \mathbf{x}_1) - \Gamma_{M_1, \infty}(\mathbf{y})| d\mathbf{y} \\ &= \int_{\mathbb{R}^2} \left| \int_0^1 \nabla \Gamma_{M_1, \infty}(\mathbf{y} + s(\mathbf{x}_2 - \mathbf{x}_1)) \cdot (\mathbf{x}_2 - \mathbf{x}_1) ds \right| d\mathbf{y} \\ &\leq \int_{\mathbb{R}^2} \int_0^1 |\nabla \Gamma_{M_1, \infty}(\mathbf{y} + s(\mathbf{x}_2 - \mathbf{x}_1))| \cdot |\mathbf{x}_2 - \mathbf{x}_1| ds d\mathbf{y} \\ &\leq C \delta^2 \|\nabla \Gamma_{M_1, \infty}\|_1 \leq C_H \delta^2 \varepsilon^{-1/4}. \end{aligned}$$

Let's now turn to the second norm in (4.29). For $t \in [0, 1]$, define the matrix $B_t = M_1 + t(M_2 - M_1)$. Notice that

$$B_t = \frac{1}{\varepsilon^{1/2}} \left(A_1 + \frac{\varepsilon^{1/2}}{\beta} I + t(A_2 - A_1) \right)$$

and $|A_2 - A_1| < \delta \ll \varepsilon^{1/2}/\beta$, because $\beta\delta \ll \varepsilon^{2/5+1/8}N^{-3/5} \ll \varepsilon^{21/40} \ll \varepsilon^{1/2}$ by (4.9)–(4.10). Therefore we have that for ε small $1/2 \det M_1 \leq \det B_t \leq 2 \det M_1$ and $1/2 \operatorname{tr} M_1 \leq \operatorname{tr} B_t \leq 2 \operatorname{tr} M_1$. Moreover,

$$\begin{aligned} \left| \frac{d}{dt} \log \det B_t \right| &= \frac{1}{\det B_t} \left| \operatorname{tr} \left(\operatorname{adj}(B_t) \frac{dB_t}{dt} \right) \right| = \frac{|\operatorname{tr}(\operatorname{adj}(B_t)(M_2 - M_1))|}{\det B_t} \\ &\leq C_H \frac{|B_t| \cdot |M_2 - M_1|}{\det B_t} \leq C_H \frac{\varepsilon^{-1/2} \cdot \delta \varepsilon^{-1/2}}{\varepsilon^{-1/2} \beta^{-1}} = C_H \frac{\beta \delta}{\varepsilon^{1/2}} \end{aligned}$$

and

$$\begin{aligned} \frac{d}{dt} \Gamma_{B_t, \infty}(\mathbf{x}) &= \frac{d}{dt} \left(\frac{\sqrt{\det B_t}}{\pi} e^{-\mathbf{x}^T B_t \mathbf{x}} \right) \\ &= \frac{\frac{d}{dt} \det B_t}{2\pi \sqrt{\det B_t}} e^{-\mathbf{x}^T B_t \mathbf{x}} - \mathbf{x}^T \frac{dB_t}{dt} \mathbf{x} \frac{\sqrt{\det B_t}}{\pi} e^{-\mathbf{x}^T B_t \mathbf{x}} \\ &= \frac{1}{2} \left(\frac{d}{dt} \log \det B_t \right) \Gamma_{B_t, \infty} - \mathbf{x}^T (M_2 - M_1) \mathbf{x} \Gamma_{B_t, \infty}. \end{aligned}$$

Therefore,

$$\begin{aligned} \|\Gamma_{M_1, \infty}^{x_2} - \Gamma_{M_2, \infty}^{x_2}\|_1 &= \|\Gamma_{M_1, \infty} - \Gamma_{M_2, \infty}\|_1 \leq \int_0^1 \int_{\mathbb{R}^2} \left| \frac{d}{dt} \Gamma_{B_t, \infty}(\mathbf{x}) \right| d\mathbf{x} dt \\ &\leq C_H \int_0^1 \left| \frac{d}{dt} \log \det B_t \right| \int_{\mathbb{R}^2} \Gamma_{B_t, \infty} d\mathbf{x} dt \\ &\quad + \int_0^1 |M_2 - M_1| \frac{\sqrt{\det B_t}}{\pi} \int_{\mathbb{R}^2} |\mathbf{x}|^2 e^{-\mathbf{x}^T B_t \mathbf{x}} d\mathbf{x} dt \\ &= C_H \frac{\beta \delta}{\varepsilon^{1/2}} + \frac{\delta}{\varepsilon^{1/2}} \int_0^1 \frac{\operatorname{tr} B_t}{2 \det B_t} dt \\ &\leq C_H \frac{\beta \delta}{\varepsilon^{1/2}} + C_H \frac{\delta}{\varepsilon^{1/2}} \beta = C_H \frac{\beta \delta}{\varepsilon^{1/2}}. \end{aligned}$$

□

5. Deconvolution of plans

Let $\sigma^1, \sigma^2 \in \mathcal{M}_+(\mathbb{R})$ with the same mass and $\Pi_0 \in \Pi(\sigma^1, \sigma^2)$ be a transport plan. Fix a radial convolution kernel $\Theta \in C^\infty(\mathbb{R}^2; [0, 1])$ with $\operatorname{supp} \Theta \subset B(0, 1)$. Define the rescaling $\Theta_\varepsilon(x) = \varepsilon^{-1/2} \Theta(\varepsilon^{-1/4} x)$ and the marginals $\theta(x) = (\operatorname{proj}_\#^1 \Theta)(x)$, $\theta_\varepsilon(x) = (\operatorname{proj}_\#^1 \Theta_\varepsilon)(x) = \varepsilon^{-1/4} \theta(\varepsilon^{-1/4} x)$. Define the convolved plan

$$\Pi_\varepsilon = \Pi_0 * \Theta_\varepsilon.$$

Let $\sigma_\varepsilon^i = \operatorname{proj}_\#^i \Pi_\varepsilon = \sigma^i * \theta_\varepsilon$, for $i = 1, 2$, be the two marginals of the convolved plan.

We now define the deconvolved plan as introduced in [1, Theorem 6.3]:

$$\begin{aligned}\tilde{\Pi}_\varepsilon(x, y) &= \int_{\mathbb{R}^2} \frac{\sigma^1(x)\theta_\varepsilon(x' - x)}{\sigma_\varepsilon^1(x')} \frac{\sigma^2(y)\theta_\varepsilon(y' - y)}{\sigma_\varepsilon^2(y')} \Pi_\varepsilon(x', y') dx' dy' \\ &= \int_{\mathbb{R}^2} P(x, y, x', y') dx' dy'.\end{aligned}$$

We verify that $\tilde{\Pi}_\varepsilon dx dy \in \Pi(\sigma^1, \sigma^2)$. Indeed, for any integrable function $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ and for every $x \in \mathbb{R}$ we perform this useful computation, integrating first the variable y and then y'

$$\begin{aligned}& \int_{\mathbb{R}^3} F(x, x') P(x, y, x', y') dx' dy' dy \\ &= \int_{\mathbb{R}^2} F(x, x') \frac{\sigma^1(x)\theta_\varepsilon(x' - x)}{\sigma_\varepsilon^1(x')} \frac{\cancel{\sigma_\varepsilon^2(y')}}{\cancel{\sigma_\varepsilon^2(y')}} \Pi_\varepsilon(x', y') dx' dy' \quad (5.1) \\ &= \int_{\mathbb{R}} F(x, x') \frac{\sigma^1(x)\theta_\varepsilon(x' - x)}{\cancel{\sigma_\varepsilon^1(x')}} \cancel{\sigma_\varepsilon^1(x')} dx'.\end{aligned}$$

By taking the function $F \equiv 1$ this reads

$$\int_{\mathbb{R}} \tilde{\Pi}_\varepsilon(x, y) dy = \sigma^1(x).$$

Analogously, one can show that the second marginal of $\tilde{\Pi}_\varepsilon$ is σ^2 . Moreover, $\tilde{\Pi}_\varepsilon$ is supported in a neighborhood of size $\varepsilon^{1/4}$ of the support of Π_ε because, in the definition of $\tilde{\Pi}_\varepsilon$, (x', y') is in the support of Π_ε and $x' - x$ is less than $\varepsilon^{1/4}$. Therefore,

$$\tilde{\Pi}_\varepsilon \text{ is supported in a neighborhood of size } 2\varepsilon^{1/4} \text{ of the support of } \Pi_0. \quad (5.2)$$

We claim in the next proposition that the operation of deconvolving a transport plan Π_0 worsens its potential energy in a controlled way, namely proportionally to the total mass of Π_0 , while on the other hand the kinetic energy of the deconvolution is controlled only in terms of the kinetic energy of the marginals of Π_0 . The loss in potential energy proportional to the mass of Π_0 is the reason why in the proof of Theorem 1.3 we use this deconvolution procedure only on the remaining mass, rather than on the full plan $\tilde{\gamma}_\varepsilon$. Let us also point out that, in the proof of Theorem 1.3, Π_0 will in turn be an ε -dependent plan, built in Sect. 6 below. For this reason we keep explicit all dependences on Π_0 in the next proposition.

Proposition 5.1. *Let V be as in Theorem 1.3, $H > 0$ and Ω_H, Ω'_H be the sets introduced in (4.5), (4.6). Let $\sigma^1, \sigma^2 \in \mathcal{M}_+(\mathbb{R})$ with the same mass, let $\Pi_0 \in \Pi(\sigma^1, \sigma^2)$, and let $\tilde{\Pi}_\varepsilon$ be the plan introduced at the beginning of this section. Then there exists a universal constant $C > 0$.*

$$\text{KE}(\tilde{\Pi}_\varepsilon) \leq \text{KE}(\sigma^1) + \text{KE}(\sigma^2) + \frac{C}{\sqrt{\varepsilon}} \|\Pi_0\|_1 \quad \text{for every } \varepsilon > 0. \quad (5.3)$$

Moreover, there exists a constant $c_H > 0$ depending only on H such that if

$$d((x, y), \text{graph}(T)) \leq c_H \mathbf{1}_{\Omega'_H}(x) \quad \text{for every } (x, y) \in \text{supp } \Pi_0, \quad (5.4)$$

then there exists a universal constant $C > 0$ and a constant $C_H > 0$ depending only on H such that for ε sufficiently small

$$\frac{1}{\sqrt{\varepsilon}} \int_{\mathbb{R}^2} V d\tilde{\Pi}_\varepsilon \leq \frac{C_H}{\sqrt{\varepsilon}} \int_{\mathbb{R}^2} V d\Pi_0 + C \|\Pi_0\|_1. \quad (5.5)$$

We notice that the dependence of the constants in (5.5) is of fundamental importance: in particular the first term will go to 0 (at H fixed) by the particular choice of Π_0 , while the second term will be estimated with (4.19) by the mass of ρ outside Ω_H (times the constant C independent of H); hence this will be small sending $H \rightarrow \infty$ uniformly in ε .

Remark 5.2. Another estimate of the potential energy of the deconvolved plan in terms of the original one was proposed by Lewin [25].

In our notations, tracing in his proof the dependence on $\|\Pi_0\|_1$, we would get an estimate of the form

$$\left| \int_{\mathbb{R}^2} \Phi d\tilde{\Pi}_\varepsilon - \int_{\mathbb{R}^2} \Phi d\Pi_0 \right| \leq \sqrt{\varepsilon} \left(\|\nabla \Phi\|_\infty \int |\nabla \sigma_1| + \|D^2 \Phi\|_\infty \|\Pi_0\|_1 \right),$$

where the norms of $\nabla \Phi$ and $D^2 \Phi$ are calculated in the neighbourhood of the graph of T where $\tilde{\Pi}_\varepsilon$ is supported. Since $\tilde{\Pi}_\varepsilon$ and Π_0 have the same marginals, we notice that

$$\int_{\mathbb{R}^2} V d\tilde{\Pi}_\varepsilon - \int_{\mathbb{R}^2} V d\Pi_0 = \int_{\mathbb{R}^2} V_{ee} d\tilde{\Pi}_\varepsilon - \int_{\mathbb{R}^2} V_{ee} d\Pi_0,$$

Applying the estimate above for $\Phi = V_{ee}$ and using that $|x - y| \geq 1/(2\|\rho\|_\infty)$ on the graph of T , we would get that for ε small enough

$$\left| \int_{\mathbb{R}^2} V d\tilde{\Pi}_\varepsilon - \int_{\mathbb{R}^2} V d\Pi_0 \right| \leq C\sqrt{\varepsilon} \|\rho\|_\infty^3 \sqrt{\|\Pi_0\|_1} \cdot \left(\int |\nabla \sigma_1| + \sqrt{\|\Pi_0\|_1} \right).$$

However, this estimate is not good enough for our application since the BV norm of σ_1 does not go to 0 as $\varepsilon \rightarrow 0$ (recall that σ_1 coincides with ρ in a neighborhood of 0).

Proof. We have that

$$\text{KE}(\tilde{\Pi}_\varepsilon) = \frac{1}{2} \int_{\mathbb{R}^2} \left| \nabla \sqrt{\tilde{\Pi}_\varepsilon(x)} \right|^2 dx = \int_{\mathbb{R}^2} \frac{|\partial_x \tilde{\Pi}_\varepsilon|^2 + |\partial_y \tilde{\Pi}_\varepsilon|^2}{8\tilde{\Pi}_\varepsilon} dx.$$

Let us deal with the first term; the second one is treated analogously. By Hölder inequality and since $\int_{\mathbb{R}^2} P dx' dy' = \tilde{\Pi}_\varepsilon(x, y)$, we have

$$\begin{aligned} |\partial_x \tilde{\Pi}_\varepsilon(x, y)|^2 &= \left| \int_{\mathbb{R}^2} \frac{\partial}{\partial x} \left(\frac{\sigma^1(x) \theta_\varepsilon(x' - x)}{\sigma_\varepsilon^1(x')} \right) \frac{\sigma^2(y) \theta_\varepsilon(y' - y)}{\sigma_\varepsilon^2(y')} \Pi_\varepsilon(x', y') dx' dy' \right|^2 \\ &= \left| \int_{\mathbb{R}^2} \left(\frac{\partial_x \sigma^1(x)}{\sigma^1(x)} + \frac{\partial_x \theta_\varepsilon(x' - x)}{\theta_\varepsilon(x' - x)} \right) P(x, y, x', y') dx' dy' \right|^2 \\ &\leq \tilde{\Pi}_\varepsilon(x, y) \int_{\mathbb{R}^2} \left[\left(\frac{\partial_x \sigma^1(x)}{\sigma^1(x)} \right)^2 + \left(\frac{\partial_x \theta_\varepsilon(x' - x)}{\theta_\varepsilon(x' - x)} \right)^2 \right] \\ &\quad P(x, y, x', y') dx' dy'. \end{aligned}$$

Then

$$\begin{aligned} &\int_{\mathbb{R}^2} \frac{|\partial_x \tilde{\Pi}_\varepsilon(x, y)|^2}{8 \tilde{\Pi}_\varepsilon(x, y)} dx dy \\ &= \int_{\mathbb{R}^4} \frac{1}{8} \left[\left(\frac{\partial_x \sigma^1(x)}{\sigma^1(x)} \right)^2 + \left(\frac{\partial_x \theta_\varepsilon(x' - x)}{\theta_\varepsilon(x' - x)} \right)^2 \right] P(x, y, x', y') dx dy dx' dy'. \end{aligned}$$

Applying (5.1) with $F(x, x')$ given by the expression in square brackets above and we find that

$$\begin{aligned} \int_{\mathbb{R}^2} \frac{|\partial_x \tilde{\Pi}_\varepsilon(x, y)|^2}{8 \tilde{\Pi}_\varepsilon(x, y)} dx dy &= \int_{\mathbb{R}^4} \frac{1}{8} \left[\left(\frac{\partial_x \sigma^1(x)}{\sigma^1(x)} \right)^2 + \left(\frac{\partial_x \theta_\varepsilon(x' - x)}{\theta_\varepsilon(x' - x)} \right)^2 \right] \\ &\quad \times \sigma^1(x) \theta_\varepsilon(x' - x) dx' dx \\ &= \int_{\mathbb{R}} \frac{|\partial_x \sigma^1(x)|^2}{8 \sigma^1(x)} dx + \|\sigma^1\|_1 \int_{\mathbb{R}} \frac{|\partial_x \theta_\varepsilon(z)|^2}{8 \theta_\varepsilon(z)} dz \\ &= \text{KE}(\sigma_1) + \text{KE}(\theta_\varepsilon) \|\Pi_0\|_1. \end{aligned}$$

The kinetic energy of the marginal of the rescaled convolution kernel scales as

$$\begin{aligned} \text{KE}(\theta_\varepsilon) &= \frac{1}{2} \int_{\mathbb{R}} \left| \partial_x \sqrt{\theta_\varepsilon(x)} \right|^2 dx = \int_{\mathbb{R}} \frac{|\theta'_\varepsilon(x)|^2}{8 \theta_\varepsilon(x)} dx \\ &= \int_{\mathbb{R}} \varepsilon^{-3/4} \frac{|\theta'(y)|^2}{8 \theta(y)} \varepsilon^{1/4} dy = \varepsilon^{-1/2} \text{KE}(\theta), \end{aligned}$$

and the kinetic energy of θ is a universal constant since θ is fixed. In conclusion, putting everything together, we obtain (5.3).

Let $c_{0,H}$ be any constant strictly less than the distance of graph T from the diagonal $\{x = y\}$. For ε sufficiently small, $\text{supp } \Pi_\varepsilon$ will be supported at positive distance from the diagonal thanks to (5.2), as soon as the constant c_0 appearing there is less or equal than $c_{0,H}$. By using the Taylor expansion of V centered at (x', y') and the fact that thanks to Lemma 4.1 $\nabla^2 V$ is bounded by a universal constant C

independent of H in the region of interest because we are far from the diagonal, we have that

$$\begin{aligned}
 & \int_{\mathbb{R}^2} V d\Pi_\varepsilon \\
 & \leq \int_{\mathbb{R}^4} V(x', y') \Theta_\varepsilon(x - x', y - y') d\Pi_0(x', y') dx dy \\
 & \quad + \int_{\mathbb{R}^4} \nabla V(x', y') \cdot (x - x', y - y') \Theta_\varepsilon(x - x', y - y') d\Pi_0(x', y') dx dy \\
 & \quad + \int_{\mathbb{R}^4} C(|x - x'|^2 + |y - y'|^2) \Theta_\varepsilon(x - x', y - y') d\Pi_0(x', y') dx dy \\
 & \leq \int_{\mathbb{R}^2} V d\Pi_0 + C \int_{\mathbb{R}^4} (|x - x'|^2 + |y - y'|^2) \Theta_\varepsilon(x - x', y - y') d\Pi_0(x', y') dx dy,
 \end{aligned}$$

because the integral in the second line vanishes (it's the integral of a linear function in x, y with respect to a symmetric kernel). The terms involving $|x - x'|^2$ and $|y - y'|^2$ in the last integral can be both computed as

$$\begin{aligned}
 & \int_{\mathbb{R}^4} |y - y'|^2 \Theta_\varepsilon(x - x', y - y') d\Pi_0(x', y') dx dy \\
 & = \int_{\mathbb{R}^3} |y - y'|^2 \theta_\varepsilon(y - y') d\Pi_0(x', y') dy \\
 & = \int_{\mathbb{R}^2} |y - y'|^2 \theta_\varepsilon(y - y') dy d\sigma^2(y') \\
 & = C\sqrt{\varepsilon} \|\sigma^2\|_1 = C\sqrt{\varepsilon} \|\Pi_0\|_1.
 \end{aligned}$$

Therefore

$$\int V d\Pi_\varepsilon \leq \int V d\Pi_0 + C\sqrt{\varepsilon} \|\Pi_0\|_1.$$

Using again the Taylor expansion of V centered at (x', y') we can proceed to estimate that

$$\begin{aligned}
 \int V(x, y) d\tilde{\pi}_\varepsilon(x, y) & \leq \int_{\mathbb{R}^4} V(x', y') P(x, y, x', y') dx dy dx' dy' \\
 & \quad + \int_{\mathbb{R}^4} \nabla V(x', y') \cdot (x - x', y - y') P(x, y, x', y') dx dy dx' dy' \\
 & \quad + \int_{\mathbb{R}^4} C(|x - x'|^2 + |y - y'|^2) P(x, y, x', y') dx dy dx' dy'.
 \end{aligned} \tag{5.6}$$

By the definition of P and integrating first the variable x and secondly the variable y , the first integral in the right-hand side is

$$\begin{aligned}
 \int_{\mathbb{R}^4} V(x', y') P dx dy dx' dy' & = \int_{\mathbb{R}^4} V(x', y') \frac{\cancel{\sigma_\varepsilon^1(x')}}{\cancel{\sigma_\varepsilon^1(x')}} \frac{\sigma_\varepsilon^2(y) \theta_\varepsilon(y' - y)}{\sigma_\varepsilon^2(y')} \Pi_\varepsilon(x', y') dy dx' dy' \\
 & = \int_{\mathbb{R}^4} V(x', y') \frac{\sigma_\varepsilon^2(y')}{\cancel{\sigma_\varepsilon^2(y')}} \Pi_\varepsilon(x', y') dx' dy' = \int_{\mathbb{R}^2} V d\Pi_\varepsilon.
 \end{aligned}$$

Given $x \in \Omega'_H$ and y such that $d((x, y), \text{graph } T) \leq c_{0,H}$, let $(x_0, y_0) \in \text{graph}(T)$ be a projection, i.e. a point realising the minimum of the distance, and let $v = (x - x_0, y - y_0)$ be the difference vector and $d = \mathbf{d}((x, y), \text{graph}(T)) = |v|$ the distance. By looking at the Taylor expansion of V at (x_0, y_0) , recalling that $\nabla V(x_0, y_0) = 0$, we have that

$$|V(x, y) - v^T \nabla^2 V(x_0, y_0) v| \leq \|\nabla^3 V\|_\infty d^3 \leq C_H d^3,$$

where the L^∞ norm is taken over the set of points (x, y) such that $x \in \Omega'_H$ and y such that $d((x, y), \text{graph } T) \leq c_{0,H}$. This L^∞ norm is in turn estimated by C_H thanks to the explicit computation of the derivatives of V (see Lemma 4.1) and thanks to the quantities that we have under control at H fixed in (4.7). The difference vector v is orthogonal to $\text{graph}(T)$ at (x_0, y_0) , therefore it is in the direction of the eigenvector of $\sqrt{\nabla^2 V(x_0, y_0)}$ associated to the non-zero eigenvalue $q(x_0, y_0)$. Therefore

$$V(x, y) \geq v^T \nabla^2 V(x_0, y_0) v - \|\nabla^3 V\|_\infty d^3 \geq q^2 d^2 - C_H d^3 \geq \frac{1}{2} \left(\min_{\Omega'_H} q \right)^2 d^2$$

as soon as $d \leq \frac{1}{2} C_H^{-1} (\min_{\Omega'_H} q)^2$. Therefore, for

$$\mathbf{d}((x, y), \text{graph}(T)) < \min \{c_{0,H}, \frac{1}{2} C_H^{-1} (\min_{\Omega'_H} q)^2\} =: c_L, \quad (5.7)$$

we have

$$|\nabla V(x, y)|^2 \leq \|\nabla^2 V\|_\infty^2 d^2 \leq 2 \|\nabla^2 V\|_\infty^2 \|1/q\|_{L^\infty(\Omega'_H)}^2 V(x, y) \leq C_H V(x, y). \quad (5.8)$$

Using Cauchy-Schwarz, the second integral in (5.6) can be bounded by

$$\begin{aligned} & \int_{\mathbb{R}^4} \nabla V(x', y') \cdot (x - x', y - y') P(x, y, x', y') dx dy dx' dy' \\ & \leq \int_{\mathbb{R}^4} (|\nabla V(x', y')|^2 + (|x - x'|^2 + |y - y'|^2)) P(x, y, x', y') dx dy dx' dy'. \end{aligned}$$

Using (5.8) in the point (x', y') , which for ε sufficiently small satisfies (5.7) thanks to (5.4) and (5.2), the first term can be estimated as

$$\begin{aligned} & \int_{\mathbb{R}^4} |\nabla V(x', y')|^2 P(x, y, x', y') dx dy dx' dy' = \int_{\mathbb{R}^2} |\nabla V(x', y')|^2 \Pi_\varepsilon(x', y') dx' dy' \\ & = \int_{\mathbb{R}^4} |\nabla V(x, y)|^2 \Theta_\varepsilon(x - x', y - y') d\Pi_0(x', y') dx dy \\ & \leq 2 \int_{\mathbb{R}^4} |\nabla V(x', y')|^2 \Theta_\varepsilon(x - x', y - y') d\Pi_0(x', y') dx dy \\ & \quad + 2 \int_{\mathbb{R}^4} |\nabla V(x, y) - \nabla V(x', y')|^2 \Theta_\varepsilon(x - x', y - y') d\Pi_0(x', y') dx dy \\ & \leq 2 \int_{\mathbb{R}^2} |\nabla V(x', y')|^2 d\Pi_0(x', y') + 2(\text{Lip}(\nabla V))^2 \end{aligned}$$

$$\begin{aligned} & \times \int_{\mathbb{R}^4} (|x - x'|^2 + |y - y'|^2) \Theta_\varepsilon(x - x', y - y') d\Pi_0(x', y') dx dy \\ & \leq C_H \int_{\mathbb{R}^2} V d\Pi_0 + C\sqrt{\varepsilon} \|\Pi_0\|_1. \end{aligned}$$

The third integral in (5.6) can be computed exploiting (5.1) with $F(x, x') = |x - x'|^2$ to get

$$\begin{aligned} \int_{\mathbb{R}^4} |x - x'|^2 P dx dy dx' dy' &= \int_{\mathbb{R}^2} |x - x'|^2 \theta_\varepsilon(x' - x) \sigma^1(x) dx dx' \\ &= \|\sigma^1\|_1 \int_{\mathbb{R}} |z|^2 \theta_\varepsilon(z) dz = C\sqrt{\varepsilon} \|\sigma^1\|_1 = C\sqrt{\varepsilon} \|\Pi_0\|_1 \end{aligned}$$

and analogously for the term with $|y - y'|^2$. Putting the pieces together, we obtain

$$\int_{\mathbb{R}^2} V d\tilde{\Pi}_\varepsilon \leq C_H \int_{\mathbb{R}^2} V d\Pi_\varepsilon + C\sqrt{\varepsilon} \|\Pi_0\|_1.$$

□

6. Potential cost of the remaining mass and kinetic energy of its marginals

In Sect. 4 we described how to build the main portion of the recovery sequence $\tilde{\gamma}_\varepsilon$. As stated in (4.18)–(4.19), this procedure leaves out a small positive mass that we still have to account for. The goal of this section is to build a transport plan for the remaining mass which satisfies suitable energy bounds and allows to apply the deconvolution procedure of Sect. 5, namely Proposition 5.1. To this end we need to build a transport plan $\pi \in \Pi(\rho - \rho_\varepsilon^1, \rho - \rho_\varepsilon^2)$ with potential energy smaller than $\sqrt{\varepsilon}$ and to have a suitable control on how the kinetic energy of its marginals explodes in ε . This is the content of the following proposition and this section is dedicated to its proof.

Proposition 6.1. *Let $\rho \in C^1(\mathbb{R})$, $\gamma \in \Pi(\rho, \rho)$, $u \in W^{2,\infty}$ and $V(x, y) = |x - y|^{-1} - u(x) - u(y)$ be as in Theorem 1.3. Let $\tilde{\gamma}_\varepsilon$ be the plan introduced in (4.14) and (4.15) and let $\rho_\varepsilon^1 = \text{proj}_\#^1 \tilde{\gamma}_\varepsilon$ and $\rho_\varepsilon^2 = \text{proj}_\#^2 \tilde{\gamma}_\varepsilon$ be its two marginals introduced in (4.16). Then for every $c > 0$ and for some constant $C_H > 1$ depending only on H*

i) (construction of a plan with $o_L(\sqrt{\varepsilon})$ potential energy) for every ε sufficiently small there is $\pi_\varepsilon \in \Pi(\rho - \rho_\varepsilon^1, \rho - \rho_\varepsilon^2)$ such that

$$\int V d\pi_\varepsilon \leq C_H \left(\frac{\delta^4}{\tau} + \frac{\varepsilon N^2}{\beta \tau} + \frac{\beta \delta^2}{\tau} \right) \quad (6.1)$$

and

$$d((x, y), \text{graph}(T)) \leq c \mathbf{1}_{\Omega'_H}(x) \quad \text{for every } (x, y) \in \text{supp } \Pi_0; \quad (6.2)$$

ii) (bound on the kinetic energy of the marginals) we have

$$\text{KE}(\rho - \rho_\varepsilon^1) + \text{KE}(\rho - \rho_\varepsilon^2) \leq \frac{C_H}{\tau} \left(\text{KE}(\rho) + \frac{1}{\beta} \right), \quad (6.3)$$

Notice that the right hand side of (6.1) is $o_H(\varepsilon^{1/2})$ and the right hand side of (6.3) is $o_H(\varepsilon^{-1/2})$ as $\varepsilon \rightarrow 0$ because of (4.9)-(4.11).

To build the plan π_ε , we will first estimate the Wasserstein distance between two suitable measures. We reduce to this estimate since an essential tool of its proof is the Benamou-Brenier formulation of optimal transport, which is not available with a general potential V . As a matter of fact, even a slightly more precise estimate than the bound on the Wasserstein distance by $o_L(\varepsilon^{1/2})$ is needed: namely, we need a map with $o_L(\varepsilon^{1/2})$ cost and with a control on the support. Since this second condition is more technical and comes naturally from the proof of (6.4), the reader may decide to skip it at a first reading and focus on (6.4).

Proposition 6.2. *Let Ω_H'' be the convex hull of Ω_H' . Under the same assumptions as in Proposition 6.1, there exists a constant C_H depending only on H such that*

$$W_2^2 \left(\rho - \rho_\varepsilon^2, T_{\delta\#}(\rho - \rho_\varepsilon^1) \right) \leq C_H \left(\frac{\delta^4}{\tau} + \frac{\varepsilon N^2}{\beta\tau} + \frac{\beta\delta^2}{\tau} \right) \quad (6.4)$$

Moreover, the optimal map S_ε between $\rho - \rho_\varepsilon^2$ and $T_{\delta\#}(\rho - \rho_\varepsilon^1)$ satisfies a pointwise bound

$$|S_\varepsilon(x) - x| \leq C_H \left(\frac{\delta^4}{\tau^2} + \frac{\varepsilon N^2}{\beta\tau^2} + \frac{\beta\delta^2}{\tau^2} \right)^{1/3} \mathbf{1}_{\Omega_H'}(x) \quad \text{for a.e. } x \in \mathbb{R}. \quad (6.5)$$

In particular, the right-hand sides of (6.4) and (6.5) are $o_H(\sqrt{\varepsilon})$ and $O_H(1)$ respectively as soon as (4.9) and (4.11) are satisfied.

We immediately show how Proposition 6.1(i) can be deduced from this estimate.

Proof of Proposition 6.1(i) from Proposition 6.2. Let $\sigma_1 = \rho - \rho_\varepsilon^1$, $\sigma_2 = \rho - \rho_\varepsilon^2$ and let S_ε be the map given by Proposition 6.2. Let then $\pi_\varepsilon = (\text{Id}, S_\varepsilon \circ T_\delta)_\# \sigma_1 \in \Pi(\sigma_1, \sigma_2)$. We observe that (recalling that T_δ maps Ω_H' in itself and coincides with T outside, and that $S_\varepsilon(x) = x$ outside Ω_H')

$$\begin{aligned} |S_\varepsilon(T_\delta(x)) - T(x)| &\leq |S_\varepsilon(T_\delta(x)) - T_\delta(x)| + |T_\delta(x) - T(x)| \\ &\leq \left[C_H \left(\frac{\delta^4}{\tau^2} + \frac{\varepsilon N^2}{\beta\tau^2} + \frac{\beta\delta^2}{\tau^2} \right) + L\delta^2 \right] \mathbf{1}_{\Omega_H'}(x) \end{aligned} \quad (6.6)$$

and the right-hand side converges to 0 as $\varepsilon \rightarrow 0$, therefore proving (6.2) for ε sufficiently small. Under the assumptions of Proposition 6.1 we have that (4.4) of Lemma 4.1 holds, that is $V(x, y) \leq C_H d((x, y), \text{graph}(T))^2$ in a neighborhood (independent of ε) of $\text{graph}(T)$. For ε sufficiently small, the plan π_ε is supported

in a neighborhood of $\text{graph}(T)$ which converges to 0 as $\varepsilon \rightarrow 0$ thanks to (6.5), so in particular the quadratic upper bound for V applies. Therefore,

$$\begin{aligned}
 \int V(x, y) d\pi_\varepsilon(x, y) &= \int_{\Omega'_H} V(x, S_\varepsilon(T_\delta(x))) d\sigma_1(x) \\
 &\leq C_H \int_{\Omega'_H} |S_\varepsilon(T_\delta(x)) - T_\delta(x) - T(x) + T_\delta(x)|^2 d\sigma_1(x) \\
 &\leq 2C_H \int_{\Omega'_H} |S(T_\delta(x)) - T_\delta(x)|^2 d\sigma_1(x) + 2C_H \|T - T_\delta\|_\infty^2 \\
 &\leq 2C_H \int_{\Omega'_H} |S(x) - x|^2 d(T_{\delta\#}\sigma_1)(x) + 2C_H (L\delta^2)^2 \\
 &\leq C_H \left(\frac{\delta^4}{\tau} + \frac{\varepsilon N^2}{\beta\tau} + \frac{\beta\delta^2}{\tau} \right).
 \end{aligned}$$

□

The estimate (6.4) relies on the particular structure that the marginals of the convolved plan have in the model case when T is (locally) a linear map, which is the main reason why we introduce T_δ in the construction. We carry out this surprising computation in Sect. 6.1 below. This computation could be interpreted by seeing the convolutions with Gaussian measures as the action of the heat semigroup. However, in our context the Gaussians need to be truncated to have a finite-range interaction, since the control of the potential is only local around its 0 level set rather than global, and hence this analogy remains only formal and we proceed in a different way.

It can be seen that ρ and $T_\delta\rho$ are very close in Wasserstein distance, as well as ρ_ε^2 and $T_{\delta\#}\rho_\varepsilon^1$. This sole property does not guarantee in general that they remain close when we subtract each other, namely when we consider the measures $\rho - \rho_\varepsilon^2$ and $T_{\delta\#}(\rho - \rho_\varepsilon^1)$: for instance, when one considers the three measures $\nu_1 = 1_{[0,1]}\mathcal{L}$, $\nu_2 = 1_{[\alpha,1+\alpha]}\mathcal{L}$ and $\nu_0 = 1_{[\alpha,1]}\mathcal{L}$, then $W_2(\nu_1, \nu_2) = \alpha$ but $W_2(\nu_1 - \nu_0, \nu_2 - \nu_0) = \sqrt{\alpha} \gg \alpha$ as $\alpha \rightarrow 0$. However, in our particular situation we can control the distance (6.4) via the Benamou-Brenier formula, since we can provide an interpolating curve between $\rho - \rho_\varepsilon^2$ and $T_{\delta\#}(\rho - \rho_\varepsilon^1)$ which enjoys some extra properties. For instance, we estimate how close the density of the interpolating curve is to that of the endpoints, providing then a lower bound on this density along the entire curve.

The proof of Proposition 6.2 in Sect. 6.2 deals then with this Benamou-Brenier estimate and combines it with a linearization argument, which allows to pass from T to its piecewise linear approximation T_δ , and with a suitable control of the errors generated at the interface between two consecutive intervals where T_δ is linear. A crucial ingredient of the proof is the particular way the two marginals of a truncated Gaussian can be mapped one onto the other at a low cost, which is explained in Sect. 6.1.

Finally, we dedicate Sect. 6.3 to the kinetic energy bounds of Proposition 6.1(ii).

6.1. Marginals of kernels of product type along a linear map

In this section we study the properties of the marginals of the rectangularly truncated Gaussian (3.1), which is a convolution kernel of product type

$$\Gamma_{M_{\varepsilon,\beta},N}(\mathbf{x}) = \frac{\left(e^{-\left(\frac{q}{\varepsilon^{1/2}} + \frac{1}{\beta}\right)w^2/2} - e^{-N/2}\right)_+^2}{G_{\frac{q}{\varepsilon^{1/2}} + \frac{1}{\beta},N}} \cdot \frac{\left(e^{-\frac{1}{\beta}z^2/2} - e^{-N/2}\right)_+^2}{G_{\frac{1}{\beta},N}} = h_{\varepsilon}^1(w)h_{\varepsilon}^2(z)$$

where $M_{\varepsilon,\beta} = A/\varepsilon^{1/2} + I/\beta$ as usual, q is the positive eigenvalue of A which is assumed to lie in a certain interval $[1/L, L]$, z is the coordinate in the direction of $\ker(A)$ and w is the transversal coordinate. The change of variables is given by the rotation

$$(w, z) = R(x, y) = \left(\frac{y - ax}{\sqrt{1+a^2}}, \frac{ay+x}{\sqrt{1+a^2}} \right) = (\cos \theta y - \sin \theta x, \sin \theta y + \cos \theta x).$$

where $(1, a) \in \ker(A)$, hence $\ker(A) = \text{graph}(a\text{Id} + b)$, and $\theta = \arctan(a)$. Observe that $c_L < \theta < \pi/2 - c_L$ for some constant $c_L > 0$ depending only on L .

In fact, for the computations of this section the precise form of h_{ε}^i does not play a role: we only need h_{ε}^1 and h_{ε}^2 to be positive, symmetric L^1 functions supported in an interval of length $C_H \varepsilon^{1/4} N^{1/2}$ and $\beta^{1/2} N^{1/2}$ respectively, and with

$$\beta^{1/2} \|\partial_z h_{\varepsilon}^2(z)\|_{L^1} + \beta \|\partial_z^2 h_{\varepsilon}^2(z)\|_{L^1} \leq C. \quad (6.7)$$

This property is satisfied by our kernel, since

$$\begin{aligned} (h_{\varepsilon}^2)'(z) &= -2G_{\frac{1}{\beta},N}^{-1} \frac{z}{\beta} e^{-\frac{1}{\beta}z^2/2} \left(e^{-\frac{1}{\beta}z^2/2} - e^{-N/2} \right)_+, \\ (h_{\varepsilon}^2)''(z) &= 2G_{\frac{1}{\beta},N}^{-1} e^{-\frac{1}{\beta}z^2/2} \left[\left(-\frac{1}{\beta} + \frac{z^2}{\beta^2} \right) \left(e^{-\frac{1}{\beta}z^2/2} - e^{-N/2} \right)_+ \right. \\ &\quad \left. + \frac{z^2}{\beta^2} \mathbf{1}_{[-\sqrt{\beta N}, \sqrt{\beta N}]}(z) \right], \end{aligned}$$

from which

$$\beta^{1/2} |(h_{\varepsilon}^2)'(z)| + \beta |(h_{\varepsilon}^2)''(z)| \leq 2G_{\frac{1}{\beta},N}^{-1} \left(\frac{|z|}{\beta^{1/2}} + 1 + 2\frac{z^2}{\beta} \right) e^{-\frac{1}{\beta}z^2} \mathbf{1}_{[-\sqrt{\beta N}, \sqrt{\beta N}]}(z);$$

integrating this inequality and disregarding the last factor $\mathbf{1}_{[-\sqrt{\beta N}, \sqrt{\beta N}]}(z)$ we get (6.7).

We adopt the following notation to denote rescaled functions by putting a subscript between curly braces

$$\varphi_{\{\zeta\}}(x) = \frac{1}{\zeta} \varphi\left(\frac{x}{\zeta}\right)$$

and we observe that $\|\varphi_{\{\zeta\}}\|_{L^1} = \|\varphi\|_{L^1}$, $(f * g)_{\{\zeta\}} = f_{\{\zeta\}} * g_{\{\zeta\}}$ and $(a\text{Id} + b)_{\#}\rho(y) = \rho_{\{a\}}(y - b)$.

Let $\eta_{\varepsilon}^i = \text{proj}_{\#}^i \Gamma_{M_{\varepsilon,\beta},N}$ be the two marginals of the Gaussian, namely,

$$\begin{aligned}\eta_{\varepsilon}^1(x) &= \int_{\mathbb{R}} h_{\varepsilon}^1(w) h_{\varepsilon}^2(z) dy = \int_{\mathbb{R}} h_{\varepsilon}^1(\cos \theta y - \sin \theta x) h_{\varepsilon}^2(\sin \theta y + \cos \theta x) dy, \\ \eta_{\varepsilon}^2(y) &= \int_{\mathbb{R}} h_{\varepsilon}^1(w) h_{\varepsilon}^2(z) dx = \int_{\mathbb{R}} h_{\varepsilon}^1(\cos \theta y - \sin \theta x) h_{\varepsilon}^2(\sin \theta y + \cos \theta x) dx,\end{aligned}$$

and $\tilde{\eta}_{\varepsilon}^2 = (a\text{Id})_{\#}\eta_{\varepsilon}^1$. We claim that we can write them as a convolution of rescalings of h_{ε}^1 and h_{ε}^2 , where the only difference between η_{ε}^2 and $\tilde{\eta}_{\varepsilon}^2$ lies in the parameter of rescaling of the first function

$$\begin{aligned}\eta_{\varepsilon}^2(y) &= (h_{\varepsilon}^1)_{\{\sin \theta \cot \theta\}} * (h_{\varepsilon}^2)_{\{\sin \theta\}}, \\ \tilde{\eta}_{\varepsilon}^2(y) &= (h_{\varepsilon}^1)_{\{\sin \theta \tan \theta\}} * (h_{\varepsilon}^2)_{\{\sin \theta\}}.\end{aligned}$$

Indeed, with the change of variable $\cos \theta y - \sin \theta x = \frac{t-x}{\sin \theta}$, which rewrites also as $\sin \theta y + \cos \theta x = \frac{t}{\cos \theta}$ and $dy = \frac{dt}{\sin \theta \cos \theta}$, and using the fact that h_{ε}^1 is symmetric, we can compute

$$\begin{aligned}\eta_{\varepsilon}^1(x) &= \int_{\mathbb{R}} \frac{1}{\sin \theta} h_{\varepsilon}^1\left(\frac{t-x}{\sin \theta}\right) \frac{1}{\cos \theta} h_{\varepsilon}^2\left(\frac{t}{\cos \theta}\right) dt \\ &= \int_{\mathbb{R}} (h_{\varepsilon}^1)_{\{\sin \theta\}}(x-t) \cdot (h_{\varepsilon}^2)_{\{\cos \theta\}}(t) dt = [(h_{\varepsilon}^1)_{\{\sin \theta\}} * (h_{\varepsilon}^2)_{\{\cos \theta\}}](x).\end{aligned}\tag{6.8}$$

In a similar fashion, with another substitution $\cos \theta y - \sin \theta x = \frac{t}{\cos \theta}$, which implies $\sin \theta y + \cos \theta x = \frac{y-t}{\sin \theta}$ and $dx = \frac{dt}{\sin \theta \cos \theta}$, we find that

$$\begin{aligned}\eta_{\varepsilon}^2(y) &= \int_{\mathbb{R}} \Gamma_{M_{\varepsilon,\beta},N}(x, y) dx = \int_{\mathbb{R}} h_{\varepsilon}^1(w) h_{\varepsilon}^2(z) dx \\ &= \int_{\mathbb{R}} \frac{1}{\cos \theta} h_{\varepsilon}^1\left(\frac{t}{\cos \theta}\right) \frac{1}{\sin \theta} h_{\varepsilon}^2\left(\frac{y-t}{\sin \theta}\right) dt \\ &= \int_{\mathbb{R}} (h_{\varepsilon}^1)_{\{\cos \theta\}}(t) \cdot (h_{\varepsilon}^2)_{\{\sin \theta\}}(y-t) dt = [(h_{\varepsilon}^1)_{\{\cos \theta\}} * (h_{\varepsilon}^2)_{\{\sin \theta\}}](y).\end{aligned}\tag{6.9}$$

Finally, by the properties of the rescalings, we have that

$$\tilde{\eta}_{\varepsilon}^2 = (\tan \theta \text{Id})_{\#}\eta_{\varepsilon}^1 = (\eta_{\varepsilon}^1)_{\{\tan \theta\}} = (h_{\varepsilon}^1)_{\{\sin \theta \tan \theta\}} * (h_{\varepsilon}^2)_{\{\sin \theta\}}.$$

A simple consequence of these representations of η_{ε}^i is for instance an L^{∞} bound which will be useful later: from (6.8), since $\|(h_{\varepsilon}^1)_{\{\sin \theta\}}\|_{L^1} = \|h_{\varepsilon}^1\|_{L^1} = 1$ and by (7.2) we have

$$\|\eta_{\varepsilon}^1\|_{L^{\infty}} \leq \|(h_{\varepsilon}^1)_{\{\sin \theta\}}\|_{L^1} \|(h_{\varepsilon}^2)_{\{\cos \theta\}}\|_{L^{\infty}} \leq C_H \|h_{\varepsilon}^2\|_{L^{\infty}} \leq \frac{C_H}{G_{\frac{1}{\beta},N}} \leq \frac{C_H}{G_{\frac{1}{\beta},\infty}} = \frac{C_H}{\beta^{1/2}}.\tag{6.10}$$

The next proposition considers a plan λ supported on the graph of $(a\text{Id} + b)$ and introduces the convolved plan $\lambda * \Gamma_{M_{\varepsilon,\beta},N}$, its marginals ρ_{ε}^1 and ρ_{ε}^2 and the pushforward $\tilde{\rho}_{\varepsilon}^2$ of the first marginal through the linear map $(a\text{Id} + b)$; it uses the representation of the marginals in terms of convolutions of η_{ε}^i with the marginal of λ to provide estimates on their distance and an interpolating curve between ρ_{ε}^2 and $\tilde{\rho}_{\varepsilon}^2$.

Proposition 6.3. *Let $a \in (L^{-1}, L)$, $\theta = \arctan(a)$, $b \in \mathbb{R}$, A a degenerate positive-semidefinite symmetric 2×2 matrix with a positive eigenvalue $q \in (L^{-1}, L)$, $\ker(A) = \text{graph}(a\text{Id})$ and $M_{\varepsilon,\beta} = A/\varepsilon^{1/2} + I/\beta$.*

*Given $\lambda \in \mathcal{M}_+(\mathbb{R}^2)$ with $\text{supp } \lambda \subset \text{graph}(a\text{Id} + b)$, define $\lambda_{\varepsilon} = \lambda * \Gamma_{M_{\varepsilon,\beta},N}$, let $\rho^i = \text{proj}_{\#}^i \lambda$, $\rho_{\varepsilon}^i = \text{proj}_{\#}^i \lambda_{\varepsilon}$ for $i = 1, 2$ and define $\tilde{\rho}_{\varepsilon}^2 = (a\text{Id} + b)_{\#} \rho_{\varepsilon}^1$. Then we have that*

$$\mu_t = (h_{\varepsilon}^1)_{\{\sin \theta (\tan \theta)^{2t-1}\}} * (h_{\varepsilon}^2)_{\{\sin \theta\}} * \rho^2, \quad t \in [0, 1],$$

is a curve interpolating between ρ_{ε}^2 and $\tilde{\rho}_{\varepsilon}^2$ satisfying $\partial_t \mu_t = \text{div}(m_t)$ with

$$\|\mu_t - \rho_{\varepsilon}^2\|_{\infty} \leq CL \frac{\varepsilon^{1/2} N}{\beta} \|\rho^2\|_{\infty}, \quad (6.11)$$

μ_t and m_t supported on the convex hull of $\text{supp } \tilde{\rho}_{\varepsilon}^2 \cup \text{supp } \rho_{\varepsilon}^2$ and

$$|m_t| \leq C_H \frac{\varepsilon^{1/2} N}{\beta^{1/2}} \|\rho^2\|_{\infty}. \quad (6.12)$$

Proof. Up to a translation, we may assume that $b = 0$. We have that $\rho_{\varepsilon}^i = \eta_{\varepsilon}^i * \rho^i$ and, recalling that the kernel h_{ε}^2 is symmetric,

$$\begin{aligned} \mu_t(y) - \rho_{\varepsilon}^2(y) &= \int_{\mathbb{R}} [(h_{\varepsilon}^1)_{\{\sin \theta (\tan \theta)^{2t-1}\}} - (h_{\varepsilon}^1)_{\{\sin \theta \cot \theta\}}](y - z) \\ &\quad \cdot ((h_{\varepsilon}^2)_{\{\sin \theta\}} * \rho^2)(z) dz \\ &= \int_{\mathbb{R}} [(h_{\varepsilon}^1)_{\{\sin \theta (\tan \theta)^{2t-1}\}} - (h_{\varepsilon}^1)_{\{\sin \theta \cot \theta\}}](y - z) \\ &\quad \cdot \left[((h_{\varepsilon}^2)_{\{\sin \theta\}} * \rho^2)(y) + ((h_{\varepsilon}^2)_{\{\sin \theta\}} * \rho^2)'(y) \cdot (z - y) \right. \\ &\quad \left. + \frac{1}{2} ((h_{\varepsilon}^2)_{\{\sin \theta\}} * \rho^2)''(y_z) \cdot (z - y)^2 \right] dz \\ &= \int_{\mathbb{R}} [(h_{\varepsilon}^1)_{\{\sin \theta (\tan \theta)^{2t-1}\}} - (h_{\varepsilon}^1)_{\{\sin \theta \cot \theta\}}](y - z) \\ &\quad \cdot \frac{((h_{\varepsilon}^2)_{\{\sin \theta\}} * \rho^2)''(y_z)}{2} \cdot (z - y)^2 dz, \end{aligned}$$

where y_z is a point in the segment between y and z . Therefore, since the integrand is nonzero only for $|y - z|^2 \leq C_H \varepsilon^{1/2} N$ and thanks to (6.7), we obtain (6.11)

$$\begin{aligned} \|\mu_t - \rho_\varepsilon^2\|_\infty &\leq C_H \left\| (h_\varepsilon^1)_{\{\sin \theta (\tan \theta)^{2t-1}\}} - (h_\varepsilon^1)_{\{\sin \theta \cot \theta\}} \right\|_1 \\ &\quad \cdot \left\| [(h_\varepsilon^2)_{\{\sin \theta\}} * \rho^2]'' \right\|_\infty \varepsilon^{1/2} N \\ &\leq C_H \varepsilon^{1/2} N \|[(h_\varepsilon^2)_{\{\sin \theta\}}]''\|_1 \cdot \|\rho^2\|_\infty \leq C_H \frac{\varepsilon^{1/2} N}{\beta} \|\rho^2\|_\infty. \end{aligned}$$

Define $H_\varepsilon^1(x) = \int_{-\infty}^x r h_\varepsilon^1(r) dr$, so that $(H_\varepsilon^1)_{\{\zeta\}}(x)$ is the primitive of $\frac{x}{\zeta^2} (h_\varepsilon^1)_{\{\zeta\}}(x)$ with the same compact support.

This allows us to express the derivative of $(h_\varepsilon^1)_{\{\zeta\}}(x)$ with respect to the parameter ζ as

$$\begin{aligned} \frac{\partial (h_\varepsilon^1)_{\{\zeta\}}(x)}{\partial \zeta} &= -\frac{1}{\zeta^2} h_\varepsilon^1 \left(\frac{x}{\zeta} \right) - \frac{x}{\zeta^3} (h_\varepsilon^1)' \left(\frac{x}{\zeta} \right) = -\frac{\partial}{\partial x} \left(\frac{x}{\zeta^2} h_\varepsilon^1 \left(\frac{x}{\zeta} \right) \right) \\ &= -\frac{\partial}{\partial x} \left(\zeta \frac{x}{\zeta^2} (h_\varepsilon^1)_{\{\zeta\}}(x) \right) = -\zeta \frac{\partial^2}{\partial x^2} \left((H_\varepsilon^1)_{\{\zeta\}}(x) \right), \end{aligned}$$

Let now $\zeta(t) = \sin \theta (\tan \theta)^{2t-1}$. We deduce that

$$\begin{aligned} \partial_t \mu_t &= \frac{\partial (h_\varepsilon^1)_{\{\zeta(t)\}}}{\partial t} * \tilde{h}_\varepsilon^2 * \rho^2 \\ &= \zeta'(t) \frac{\partial (h_\varepsilon^1)_{\{\zeta(t)\}}}{\partial \zeta} * \tilde{h}_\varepsilon^2 * \rho^2 = -\zeta'(t) \zeta(t) [(H_\varepsilon^1)_{\{\zeta(t)\}}]'' * \tilde{h}_\varepsilon^2 * \rho^2 \\ &= -\zeta'(t) \zeta(t) \operatorname{div} \left((H_\varepsilon^1)_{\{\zeta(t)\}} * [(h_\varepsilon^2)_{\{\sin \theta\}}]' * \rho^2 \right) = \operatorname{div}(m_t). \end{aligned}$$

We can bound by (6.7)

$$\|m_t\|_\infty \leq C_H \|(H_\varepsilon^1)_{\{\zeta(t)\}}\|_1 \cdot \|[(h_\varepsilon^2)_{\{\sin \theta\}}]'\|_1 \cdot \|\rho^2\|_\infty \leq C_H \|H_\varepsilon^1\|_1 \cdot \frac{C}{\beta^{1/2}} \cdot \|\rho^2\|_\infty$$

and estimate the first factor in the right-hand side in terms of powers of ε using that h_ε^1 is supported in a set of size $C_H \varepsilon^{1/4} N^{1/2}$

$$\begin{aligned} \|H_\varepsilon^1\|_1 &\leq -2 \int_{-\infty}^0 \int_{-\infty}^x r h_\varepsilon^1(r) dr dx = -2 \int_{-\infty}^0 \int_r^0 r h_\varepsilon^1(r) dx dr \\ &= 2 \int_{-\infty}^0 r^2 h_\varepsilon^1(r) dr \leq C_H \varepsilon^{1/2} N \|h_\varepsilon^1\|_1 \leq C_H \varepsilon^{1/2} N. \end{aligned}$$

Hence we obtain (6.12). $\square$

6.2. Wasserstein estimate of Proposition 6.2

Proof of Proposition 6.2. The proof is organized as follows: in steps 1 through 3 we prove the W_2 estimate (6.4); whereas in step 4 we show how to deduce from this the L^∞ estimate (6.5) for the optimal map, which is our true objective.

Define the map $T_t = (1-t)T + tT_\delta$ and define T_δ^i to be the affine extension of $T_\delta|_{I_i^1}$. Define also the marginals $\rho_{\varepsilon,i}^1$ and $\rho_{\varepsilon,i}^2$ of $\bar{\gamma}_{\varepsilon,i}$. We introduce the following three curves of measures:

- $\rho_t = T_{t\#}\rho$ from $T_{\#}\rho = \rho$ to $T_{\delta\#}\rho$;
- ξ_t^i from $\rho_{\varepsilon,i}^2$ to $T_{\delta\#}^i\rho_{\varepsilon,i}^1$ using the construction presented in Proposition 6.3
- ζ_t^i from $T_{\delta\#}^i\rho_{\varepsilon,i}^1$ to $T_{\delta\#}\rho_{\varepsilon,i}^1$ defined by linearly stretching $\rho_{\varepsilon,i}^1 \llcorner I_{i+1}^1$ from T_δ^i to T_δ^{i+1} and similarly for $\rho_{\varepsilon,i}^1 \llcorner I_{i-1}^1$ with Lemma 6.4.

We transform the remaining mass in three steps:

$$\begin{aligned} \rho - \rho_\varepsilon^2 &\xrightarrow{\rho_t - \rho_\varepsilon^2} T_{\delta\#}\rho - \rho_\varepsilon^2 \xrightarrow{T_{\delta\#}\rho - \sum_i \xi_t^i} T_{\delta\#}\rho - \sum_i T_{\delta\#}^i\rho_{\varepsilon,i}^1 \\ &\xrightarrow{T_{\delta\#}\rho - \sum_i \zeta_t^i} T_{\delta\#}\rho - \sum_i T_{\delta\#}\rho_{\varepsilon,i}^1. \end{aligned}$$

We estimate the W_2 distance of each step with Benamou-Brenier.

Step 1. For every $x \in \mathbb{R}$ we observe that the map $t \rightarrow T_t(x)$ solves the ODE

$$\partial_t X(t) = (T_\delta - T) \circ T_t^{-1} \circ X(t)$$

with initial datum $X(0) = x$. Notice that $(T_\delta - T) \circ T_t^{-1}$ is a locally Lipschitz vector field, therefore $T_t(x)$ represents its flow. As a consequence $\rho_t = (T_t)_\#\rho$ solves

$$\partial_t \rho_t = -\operatorname{div}((T_\delta - T) \circ T_t^{-1} \rho_t).$$

From the definition of ρ_t we get that $\rho_t(T_t x) = \rho(x)/T'_t(x)$, whereas $\rho(Tx) = \rho(x)/T'(x)$, since $T_{\#}\rho = \rho$. These can be rewritten as

$$\rho(y) = \frac{\rho(T^{-1}(y))}{T'(T^{-1}(y))}, \quad \rho_t(y) = \frac{\rho(T_t^{-1}(y))}{T'_t(T_t^{-1}(y))}.$$

Our goal is to show that these two densities are close to one another for $y \in \Omega_H$. We start by comparing the denominators. Let $x \in I_i^1$ and $y = T(x)$; we have $T_t^{-1}(y) \in I_i^1$. Then $T'(T^{-1}(y)) = T'(x) \in T'(I_i^1)$; but also $T'_t(T_t^{-1}(y)) = (1-t)T'(T_t^{-1}(y)) + tT'_\delta(T_t^{-1}(y)) \in \operatorname{conv}(T'(I_i^1)) = T'(I_i^1)$. Therefore we have

$$\begin{aligned} \left| T'_t(T_t^{-1}(y)) - T'(T^{-1}(y)) \right| &\leq \operatorname{Lip}(T') \operatorname{diam}(I_i^1) \leq L\delta, \\ \min\{T'_t(T_t^{-1}(y)) : y \in \Omega_H\} &\geq \min\{T'(T^{-1}(y)) : y \in \Omega_H\} \geq \frac{1}{L}. \end{aligned} \tag{6.13}$$

Let us now estimate the numerators of the densities. Differentiating with respect to t the identity $y = T_t(T_t^{-1}(y))$ and using the fact that $dT_t/dt = T_\delta - T$ we get

$$0 = \frac{dT_t}{dt}(T_t^{-1}(y)) + T'_t(T_t^{-1}(y)) \frac{dT_t^{-1}}{dt}(y),$$

from which we deduce, by (4.13) and (6.13), that, for $y \in \Omega_H$,

$$\left| \frac{dT_t^{-1}}{dt}(y) \right| = \left| \frac{(T - T_\delta)(T_t^{-1}(y))}{T'_t(T_t^{-1}(y))} \right| \leq L^2 \delta^2.$$

This allows us to compute for $y \in \Omega_H$ (notice that for $y \in (\Omega_H)^c$ the quantity below is 0) that

$$\left| T_t^{-1}(y) - T^{-1}(y) \right| = \left| T_t^{-1}(y) - T_0^{-1}(y) \right| \leq \int_0^t \left| \frac{dT_t^{-1}}{dt}(y) \right| dt \leq L^2 \delta^2,$$

and therefore

$$\left| \rho(T_t^{-1}(y)) - \rho(T^{-1}(y)) \right| \leq \text{Lip}(\rho) L^2 \delta^2 \leq L^3 \delta^2.$$

From this we conclude that

$$\begin{aligned} |\rho_t(x) - \rho(x)| &\leq \frac{|\rho(T^{-1}(y)) - \rho(T_t^{-1}(y))|}{T'(T^{-1}(y))} + \rho(T_t^{-1}(y)) \\ &\quad \times \frac{|T'_t(T_t^{-1}(y)) - T'(T^{-1}(y))|}{T'_t(T_t^{-1}(y))T'(T^{-1}(y))} \\ &\leq L^4 \delta^2 + L^4 \delta \leq C_H \delta. \end{aligned}$$

On the other hand, by (4.18) of Lemma 4.2 we have that $\rho_\varepsilon^2 \leq \rho - c_H \tau$.

Combined together, these two density estimates imply that

$$C_H \geq \rho_t(x) \geq \rho_\varepsilon^2(x) + c_H \tau / 2 \quad \text{for every } x \in \Omega_H.$$

Therefore, with Benamou-Brenier and noticing that $T_\delta(y) - T(y) = 0$ for $y \in (\Omega_H)^c$, we can compute that

$$\begin{aligned} W_2^2(\rho - \rho_\varepsilon^2, T_{\delta\#}\rho - \rho_\varepsilon^2) &\leq \int_0^1 \int_{\Omega'_H} \frac{1}{\rho_t(x) - \rho_\varepsilon^2(x)} \left| (T_\delta - T) \circ T_t^{-1}(x) \rho_t(x) \right|^2 dx dt \\ &\leq \int_0^1 \int_{\Omega'_H} \frac{C_H}{\tau} \|T_\delta - T\|_\infty^2 \|\rho\|_\infty^2 dx dt \leq \frac{C_H \delta^4}{\tau}, \end{aligned}$$

where in the last step we use the estimate (4.13).

Step 2. By Proposition 6.3 applied with $\lambda = (\text{Id}, T_\delta)_\#((\rho - \tau)\mathbf{1}_{I^1})$ (recall that by (4.15) with this definition λ_ε has marginals $\rho_{\varepsilon,i}^1$ and $\rho_{\varepsilon,i}^2$) for every i there exists a curve ξ_t^i connecting $\rho_{\varepsilon,i}^1$ and $\rho_{\varepsilon,i}^2$ such that $\partial_t \xi_t^i = \text{div}(m_t^i)$ and

$$\|\xi_t^i - \rho_{\varepsilon,i}^2\|_\infty = \|\xi_t^i - \xi_0^i\|_\infty \leq C_H \frac{\varepsilon^{1/2} N}{\beta},$$

$$|m_t^i| \leq C_H \frac{\varepsilon^{1/2} N}{\beta^{1/2}}.$$

The second curve to consider is $T_{\delta\#}\rho - \sum_i \xi_t^i$, which solves $\partial_t(T_{\delta\#}\rho - \sum_i \xi_t^i) = \operatorname{div}(\sum_i m_t^i)$ and connects $T_{\delta\#}\rho - \rho_\varepsilon^2$ and $T_{\delta\#}\rho - \sum_i T_{\delta\#\varepsilon,i}^i \rho_{\varepsilon,i}^1$. Since both ξ_t^i and m_t^i are supported in a small neighborhood of I_δ^i , at each point x at most two of each of these objects can overlap. Therefore for a.e. $x \in \Omega_H$

$$\left| \sum_i \xi_t^i - \rho_\varepsilon^2 \right| + \left| \sum_i m_t^i \right| \leq 4C_H \frac{\varepsilon^{1/2} N}{\beta^{1/2}}$$

because the terms overlap at most twice. This means again that the density of $T_{\delta\#}\rho - \sum_i \xi_t^i$ is at least $C_H \tau/2$ and

$$\begin{aligned} W_2^2 \left(T_{\delta\#}\rho - \rho_\varepsilon^2, T_{\delta\#}\rho - \sum_i T_{\delta\#\varepsilon,i}^i \rho_{\varepsilon,i}^1 \right) &\leq \int_0^1 \int_{\Omega'_H} \frac{1}{T_{\delta\#}\rho - \sum_i \xi_t^i} \left| \sum_i m_t^i \right|^2 dx dt \\ &\leq \int_0^1 \int_{\Omega'_H} C_H \frac{\varepsilon N^2}{\beta \tau} dx dt \leq C_H \frac{\varepsilon N^2}{\beta \tau}. \end{aligned}$$

Step 3. Finally, we deal with the curve $T_{\delta\#}\rho - \sum_i \zeta_t^i$ which connects $T_{\delta\#}\rho - \sum_i T_{\delta\#\varepsilon,i}^i \rho_{\varepsilon,i}^1$ to $T_{\delta\#}\rho - \sum_i T_{\delta\#}\rho_{\varepsilon,i}^1$. We split each term $\rho_{\varepsilon,i}^1$ as the sum of three contributions

$$\rho_{\varepsilon,i}^1 = \rho_{\varepsilon,i}^1 \mathbf{1}_{I_{i-1}^1} + \rho_{\varepsilon,i}^1 \mathbf{1}_{I_i^1} + \rho_{\varepsilon,i}^1 \mathbf{1}_{I_{i+1}^1}.$$

On the interval I_i^1 the maps T_δ^i and T_δ coincide, therefore we leave the central mass $\rho_{\varepsilon,i}^1 \mathbf{1}_{I_i^1}$ still. On the interval I_{i+1}^1 , on the other hand, the map T_δ coincides with T_δ^{i+1} , so the mass $\rho_{\varepsilon,i}^1 \mathbf{1}_{I_{i+1}^1}$ must be transformed from $T_{\delta\#}^i(\rho_{\varepsilon,i}^1 \mathbf{1}_{I_{i+1}^1})$ to $T_{\delta\#}^{i+1}(\rho_{\varepsilon,i}^1 \mathbf{1}_{I_{i+1}^1})$. We perform this transformation by applying Lemma 6.4 to $\mu = \rho_{\varepsilon,i}^1 \mathbf{1}_{I_{i+1}^1}$, x_0 equal to the point separating I_i^1 and I_{i+1}^1 , $\ell = \sqrt{\beta N}$, $S_{\lambda_0} = T_\delta^i$ and $S_{\lambda_1} = T_\delta^{i+1}$. Similarly, we need to transform the mass in I_{i-1}^1 from $T_{\delta\#}^i(\rho_{\varepsilon,i}^1 \mathbf{1}_{I_{i-1}^1})$ to $T_{\delta\#}^{i-1}(\rho_{\varepsilon,i}^1 \mathbf{1}_{I_{i-1}^1})$, and we do so with a mirrored version of Lemma 6.4. In summary, the curve ζ_t^i is composed of three terms: a central mass $\rho_{\varepsilon,i}^1 \mathbf{1}_{I_i^1}$ which does not move, and two masses one on each side which move according to the construction in Lemma 6.4. Notice that, when summing over i , all the contributions in $\sum_i \zeta_t^i$ deriving from the application of Lemma 6.4 are disjoint, because there are two terms in every I_i^1 but they don't overlap (since $\sqrt{\beta N} \ll \delta$).

The lower bound on the density of $T_{\delta\#}\rho - \sum_i \zeta_t^i$ follows from the same estimate as in the first step and the estimate of the density in Lemma 6.4. Moreover, Lemma 6.4 also gives us the estimate on the momentum

$$|m_t^i| \leq \beta^{1/2} \delta \frac{\max T'}{(\min T')^2} \leq L^3 \beta^{1/2} \delta,$$

which again do not overlap. Therefore we can compute that

$$\begin{aligned} W_2^2 \left(T_{\delta\#} \rho - \sum_i T_{\delta\#}^i \rho_{\varepsilon,i}^1, T_{\delta\#} \rho - \sum_i T_{\delta\#}^i \rho_{\varepsilon,i}^1 \right) \\ \leq \int_0^1 \int_{\Omega'_H} \frac{1}{T_{\delta\#} \rho - \sum_i \zeta_t^i} \left| \sum_i m_t^i \right|^2 dx dt \\ \leq \int_0^1 \int_{\Omega'_H} \frac{C_H}{\tau} \beta \delta^2 dx dt \leq C_H \frac{\beta \delta^2}{\tau}. \end{aligned}$$

Step 4: proof of (6.5). We apply the main estimate in [5, Proposition 1.2]: if μ_0, μ_1 are probability measures on a bounded interval I_0 , with μ_0 absolutely continuous and with density bounded below by τ_0 , then the optimal transport map S between μ_0 and μ_1 satisfies the estimate

$$\|S(x) - x\|_{L^\infty}^3 \leq \frac{C_\ell}{\tau_0} W_2^2(\mu_0, \mu_1). \quad (6.14)$$

Let S_ε be the optimal map from

$$\rho - \rho_\varepsilon^2 = \rho \mathbf{1}_{\Omega'_H} - \rho_\varepsilon^2 + \rho \mathbf{1}_{\Omega''_H \setminus \Omega_H} + \rho \mathbf{1}_{\mathbb{R} \setminus \Omega''_H}$$

to

$$T_{\delta\#}(\rho - \rho_\varepsilon^1) = T_{\delta\#}(\rho \mathbf{1}_{\Omega'_H} - \rho_\varepsilon^1) + \rho \mathbf{1}_{\Omega''_H \setminus \Omega_H} + \rho \mathbf{1}_{\mathbb{R} \setminus \Omega''_H}.$$

Since the tails outside Ω''_H of both the marginals coincide with $\rho \mathbf{1}_{\mathbb{R} \setminus \Omega''_H}$, we claim that $S_\varepsilon(x) = x$ for $x \in \mathbb{R} \setminus \Omega''_H$ and that S_ε is also an optimal map from

$$\mu_0 = (\rho - \rho_\varepsilon^2) \mathbf{1}_{\Omega''_H} = (\rho \mathbf{1}_{\Omega'_H} - \rho_\varepsilon^2) + \rho \mathbf{1}_{\Omega''_H \setminus \Omega_H}$$

to

$$\mu_1 = T_{\delta\#}(\rho - \rho_\varepsilon^1) \mathbf{1}_{\Omega''_H} = T_{\delta\#}(\rho \mathbf{1}_{\Omega'_H} - \rho_\varepsilon^1) + \rho \mathbf{1}_{\Omega''_H \setminus \Omega_H}.$$

Applying (6.14) with $I_0 = \Omega''_H$ and $\tau_0 = c_H \tau$ (notice that in the gap $\Omega''_H \setminus \Omega'_H$ we have $\mu_0 = \rho \gg \tau$) we deduce that

$$\begin{aligned} \|S_\varepsilon(x) - x\|_{L^\infty(\Omega''_H)}^3 &\leq \frac{C_H}{\tau} W_2^2 \left((\rho - \rho_\varepsilon^2) \mathbf{1}_{\Omega''_H}, T_{\delta\#}(\rho - \rho_\varepsilon^1) \mathbf{1}_{\Omega''_H} \right) \\ &= \frac{C_H}{\tau} \int_{\Omega''_H} |S_\varepsilon(x) - x|^2 (\rho(x) - \rho_\varepsilon^2(x)) dx \\ &\leq \frac{C_H}{\tau} \int_{\mathbb{R}} |S_\varepsilon(x) - x|^2 (\rho(x) - \rho_\varepsilon^2(x)) dx \\ &= \frac{C_H}{\tau} W_2^2 \left(\rho - \rho_\varepsilon^2, T_{\delta\#}(\rho - \rho_\varepsilon^1) \right) \\ &\leq C_H \left(\frac{\delta^4}{\tau^2} + \frac{\varepsilon N^2}{\beta \tau^2} + \frac{\varepsilon^{1/2} N}{\beta \tau} \right). \end{aligned}$$

As a final remark, notice that even without invoking (6.14) one could directly verify that the map between $\rho - \rho_\varepsilon^2$ and $T_{\delta\#}(\rho - \rho_\varepsilon^1)$ implicitly constructed above (via the curves of measures proposed in Steps 1,2,3) satisfies an L^∞ estimate of the type (6.5). The proof would meet more technical difficulties than this Step 4, but would however be sufficient. $\square$

Lemma 6.4. *Let $\mu \in \mathcal{M}_+([x_0, x_0 + \ell])$, $\mu \ll \mathcal{L}^1$, $\lambda_0, \lambda_1 \in \mathbb{R}_+$ and $S_\lambda : \mathbb{R} \rightarrow \mathbb{R} : x \mapsto x_0 + \lambda(x - x_0)$. Then $\mu_t = (S_{(1-t)\lambda_0 + t\lambda_1})_\# \mu$ solves $\partial_t \mu_t = \operatorname{div}(m_t)$ with*

$$|m_t| \leq \ell \frac{|\lambda_1 - \lambda_0| \max\{\lambda_0, \lambda_1\}}{\min\{\lambda_0, \lambda_1\}^2} \|\mu\|_\infty.$$

Moreover,

$$\|\mu_t - \mu_s\|_{L^\infty(0,\infty)} \leq C|\lambda_t - \lambda_s| + \ell \operatorname{Lip}(\mu) \max \left\{ \frac{\lambda_t}{\lambda_s}, \frac{\lambda_s}{\lambda_t} \right\}.$$

Proof. Without loss of generality, up to translation we may assume $x_0 = 0$. Let $\lambda_t = (1-t)\lambda_0 + t\lambda_1$. Then $\mu_t(x) = \frac{1}{\lambda_t} \mu\left(\frac{x}{\lambda_t}\right)$. The measure μ_t is advected by the vector field $v_t(x) = \frac{\lambda'_t}{\lambda_t} x = \frac{\lambda_1 - \lambda_0}{\lambda_t} x$: in fact

$$\partial_t \mu_t + \operatorname{div}(v_t \mu_t) = \frac{\partial}{\partial t} \left[\frac{1}{\lambda_t} \mu\left(\frac{x}{\lambda_t}\right) \right] + \frac{\partial}{\partial x} \left[\frac{\lambda'_t}{\lambda_t^2} x \mu\left(\frac{x}{\lambda_t}\right) \right] = 0.$$

Therefore $m_t = -v_t \mu_t$ and

$$|m_t| \leq \ell \frac{|\lambda_1 - \lambda_0| \max\{\lambda_0, \lambda_1\}}{\min\{\lambda_0, \lambda_1\}^2} \|\mu\|_\infty$$

because $|x| < \max \lambda_0, \lambda_1 \ell$ where $\mu(x/\lambda_t) > 0$.

The L^∞ estimate comes from the explicit formula for the density of μ_t . $\square$

6.3. Kinetic energy of the remaining mass

Proof of Proposition 6.1(ii). We start by showing the bound for $\operatorname{KE}(\rho - \rho_\varepsilon^1)$. We split the kinetic energy in the domain $S = \operatorname{supp} \rho_\varepsilon^1 \subset \Omega'_H$ and the complement, making use of (4.18) of Lemma 4.2 and the fact that in $S^c = (\operatorname{supp} \rho_\varepsilon^1)^c$ we have

$\rho - \rho_\varepsilon^1 = \rho$. What we obtain is

$$\begin{aligned}
 \text{KE}(\rho - \rho_\varepsilon^1) &= \int_S \frac{|\partial_x[\rho(x) - \rho_\varepsilon^1(x)]|^2}{8[\rho(x) - \rho_\varepsilon^1(x)]} dx \\
 &= \int_S \frac{|\partial_x[\rho(x) - \rho_\varepsilon^1(x)]|^2}{8[\rho(x) - \rho_\varepsilon^1(x)]} dx + \int_{S^c} \frac{|\partial_x[\rho(x) - \rho_\varepsilon^1(x)]|^2}{8[\rho(x) - \rho_\varepsilon^1(x)]} dx \\
 &\leq \frac{C_H}{\tau} \int_S (|\partial_x \rho(x)|^2 + |\partial_x \rho_\varepsilon^1(x)|^2) dx + \int_{S^c} \frac{|\partial_x \rho(x)|^2}{4\rho(x)} dx \\
 &\leq \frac{C_H \|\rho\|_{L^\infty(\Omega'_H)}}{\tau} \int_S \frac{|\partial_x \rho(x)|^2}{4\rho(x)} dx + \frac{C_H}{\tau} \int_S |\partial_x \rho_\varepsilon^1(x)|^2 dx \\
 &\quad + \int_{S^c} \frac{|\partial_x \rho(x)|^2}{4\rho(x)} dx \\
 &\leq \frac{C_H L}{\tau} \text{KE}(\rho) + \frac{C_H}{\tau} \int_S |\partial_x \rho_\varepsilon^1(x)|^2 dx.
 \end{aligned}$$

Let us recall that the domain Ω_H defined in (4.5) has two connected components which are both covered by an essentially disjoint family of intervals $I_i^1 = [a_i, b_i]$ of length $\delta/2 < b_i - a_i < \delta$ with

$$\begin{aligned}
 T(r_H) &= a_1 < b_1 = a_2 < \dots < b_{k_0} = T(H), \\
 r_H &= a_{k_0+1} < b_{k_0+1} = a_{k_0+2} < \dots < b_{k_1} = H.
 \end{aligned}$$

Notice that the cardinality of the intervals is $k_1 \leq C_H/\delta$. As already mentioned in (4.20), we have that

$$\rho_\varepsilon^1(x) = \sum_{i=1}^{k_1} \left([(\rho - \tau)\mathbf{1}_{I_i^1}] * \eta_{\varepsilon,i}^1 \right)(x),$$

where $\eta_{\varepsilon,i}^1 = \text{proj}_{\#}^1 \Gamma_{M_{\varepsilon,\beta}(x_i),N}$ were first introduced in the proof of Lemma 4.2. As a consequence, using the fact that the terms in the sum overlap at most twice, we have

$$\begin{aligned}
 \|\partial_x \rho_\varepsilon^1\|_2^2 &= \left\| \sum_{i=1}^{k_1} [(\rho - \tau)\mathbf{1}_{I_i^1}] * \partial_x \eta_{\varepsilon,i}^1 \right\|_2^2 \\
 &\leq 2 \sum_{i=1}^{k_1} \left\| [(\rho - \tau)\mathbf{1}_{I_i^1}] * \partial_x \eta_{\varepsilon,i}^1 \right\|_2^2 \leq 2 \sum_{i=1}^{k_1} \left\| (\rho - \tau)\mathbf{1}_{I_i^1} \right\|_2^2 \|\partial_x \eta_{\varepsilon,i}^1\|_1^2.
 \end{aligned}$$

Using the representation (6.8) we see that the function $\eta_{\varepsilon,i}^1$ is increasing in $(-\infty, 0]$, decreasing in $[0, \infty)$ and vanishes at infinity, hence

$$\begin{aligned}
 \|\partial_x \eta_{\varepsilon,i}^1\|_1 &= 2\|\eta_{\varepsilon,i}^1\|_\infty \leq 2\|(h_\varepsilon^1)_{\{\sin \theta\}}\|_1 \|(h_\varepsilon^2)_{\{\cos \theta\}}\|_\infty \\
 &= 2(\cos \theta)^{-1} \|h_\varepsilon^2\|_\infty = 2(\cos \theta)^{-1} G_{\frac{1}{\beta},N}^{-1} \leq C_H \beta^{-1/2}.
 \end{aligned}$$

As a consequence,

$$\begin{aligned}\|\partial_x \rho_\varepsilon^1\|_2^2 &\leq \frac{C_H}{\beta} \sum_{i=1}^{k_1} \left\| (\rho - \tau) \mathbf{1}_{I_i^1} \right\|_2^2 = \frac{C_H}{\beta} \|\rho - \tau\|_{L^2(\Omega_H)}^2 \\ &\leq \frac{C_H}{\beta} \|\rho\|_1 \|\rho\|_{L^\infty(\Omega'_H)} \leq \frac{C_H}{\beta}.\end{aligned}$$

Using this inequality in the original estimate of the kinetic energy we finally get

$$\text{KE}(\rho - \rho_\varepsilon^1) \leq \frac{C_H}{\tau} \text{KE}(\rho) + \frac{C_H}{\tau\beta}.$$

Let us now move on to the analogous estimate for $\text{KE}(\rho - \rho_\varepsilon^2)$. With the same argument as we did at the beginning, splitting the kinetic energy in the domain $S = \text{supp } \rho_\varepsilon^2 \subset \Omega'_H$ and the complement we get

$$\text{KE}(\rho - \rho_\varepsilon^2) \leq \frac{C_H}{\tau} \text{KE}(\rho) + \frac{C_H}{\tau} \int_{\mathbb{R}} |\partial_x \rho_\varepsilon^2(x)|^2 dx.$$

As already mentioned in the proof of (4.18) of Lemma 4.2, we have that

$$\rho_\varepsilon^2(y) = \sum_{i=1}^{k_1} \left([T_{\delta\#}(\rho - \tau) \mathbf{1}_{I_i^2}] * \eta_{\varepsilon,i}^2 \right)(y),$$

where $\eta_{\varepsilon,i}^2 = \text{proj}_{\#}^2 \Gamma_{M_{\varepsilon,\beta}(x_i), N}$ are the second marginals of the Gaussians. The intervals $[c_i, d_i] = I_i^2 = T_\delta(I_i^1) = T(I_i^1) = [T(a_i), T(b_i)]$ have lengths $\delta/(2L) \leq \text{diam}(I_i^2) \leq L\delta$. If $y \in I_i^2$ we have that

$$T_{\delta\#}(\rho - \tau)(y) = [\rho(T_\delta^{-1}(y)) - \tau]/T'(x_i) \leq L\|\rho\|_{L^\infty(\Omega'_H)};$$

on the other hand, from (6.9), we have

$$\|\eta_{\varepsilon,i}^2\|_\infty \leq \|(h_\varepsilon^1)_{\{\cos\theta\}}\|_1 \|(h_\varepsilon^2)_{\{\sin\theta\}}\|_\infty \leq (\sin\theta)^{-1} \|h_\varepsilon^2\|_\infty \leq C_H \beta^{-1/2},$$

and therefore

$$\|\partial_y \eta_{\varepsilon,i}^2\|_1 = 2\|\eta_{\varepsilon,i}^2\|_\infty \leq C_H \beta^{-1/2}.$$

This implies that

$$\begin{aligned}\|\partial_y \rho_\varepsilon^2\|_2^2 &= \left\| \sum_{i=1}^{k_1} [T_{\delta\#}(\rho - \tau) \mathbf{1}_{I_i^2}] * \partial_y \eta_{\varepsilon,i}^2 \right\|_2^2 \leq 2 \sum_{i=1}^{k_1} \left\| [T_{\delta\#}(\rho - \tau) \mathbf{1}_{I_i^2}] * \partial_y \eta_{\varepsilon,i}^2 \right\|_2^2 \\ &\leq 2 \sum_{i=1}^{k_1} \left\| T_{\delta\#}(\rho - \tau) \mathbf{1}_{I_i^2} \right\|_2^2 \|\partial_y \eta_{\varepsilon,i}^2\|_1^2 \leq \frac{C_H}{\beta} \sum_{i=1}^{k_1} \left\| T_{\delta\#} \rho \mathbf{1}_{I_i^2} \right\|_1 \left\| T_{\delta\#} \rho \mathbf{1}_{I_i^2} \right\|_\infty \\ &\leq \frac{C_H}{\beta} L \|\rho\|_{L^\infty(\Omega'_H)} \sum_{i=1}^{k_1} \left\| \rho \mathbf{1}_{I_i^2} \right\|_1 \leq \frac{C_H}{\beta} L \|\rho\|_{L^\infty(\Omega'_H)} \|\rho\|_{L^1(\Omega'_H)} \leq \frac{C_H}{\beta},\end{aligned}$$

which, inserted into the original estimate of the kinetic energy, leads to

$$\text{KE}(\rho - \rho_\varepsilon^2) \leq \frac{C_H}{\tau} \text{KE}(\rho) + \frac{C_H}{\tau\beta}.$$

□

7. Proof of Theorems 1.1 and 1.3

Proof of Theorem 1.3. Given $H > 1$ and $L = L(H)$ defined by (4.7), let $\bar{\gamma}_\varepsilon$ be as defined in (4.14), built with admissible parameters N, β, δ, τ satisfying (4.8)–(4.11), for instance the explicit choice (4.12). By Lemma 4.2 we have that

$$\limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\sqrt{\bar{\gamma}_\varepsilon}) \leq \frac{1}{2} \int_{\Omega_H} \operatorname{tr} \left(\sqrt{\nabla^2 V(x, T(x))} \right) \rho(x) dx \leq \frac{1}{2} \int_{\mathbb{R}^2} \operatorname{tr}(\sqrt{D^2 V}) d\gamma.$$

The problem is that $\bar{\gamma}_\varepsilon$ is not a recovery sequence because it has the wrong mass and marginals, so we have to use the remaining mass to fix the marginals.

For $i = 1, 2$, let $v_\varepsilon^i = \operatorname{proj}_\#^i(\gamma) - \operatorname{proj}_\#^i(\bar{\gamma}_\varepsilon) = \rho - \rho_\varepsilon^i$, which is a positive measure as remarked in (4.18) of Lemma 4.2.

Let $\pi_\varepsilon \in \Pi(v_\varepsilon^1, v_\varepsilon^2)$ be the plan given by Proposition 6.1(i). Notice that, by (4.19), $\|v_\varepsilon^i\|_1 = \|\pi_\varepsilon\|_1 \leq C_H \tau + \rho(\Omega_H^c)$. With H fixed the first term goes to 0 when $\varepsilon \rightarrow 0$, whereas the second term goes to zero as $H \rightarrow \infty$.

Let $\tilde{\pi}_\varepsilon \in \Pi(v_\varepsilon^1, v_\varepsilon^2)$ be the deconvolved plan defined in Sect. 5 starting from $\Pi_0 = \pi_\varepsilon$ (whose marginals are $\sigma^1 = v_\varepsilon^1$ and $\sigma^2 = v_\varepsilon^2$). By Proposition 5.1 (notice that the assumption (5.4) is satisfied because of (6.2)) and the bound on the mass of π_ε we have

$$\begin{aligned} E_\varepsilon(\sqrt{\tilde{\pi}_\varepsilon}) &= \varepsilon^{1/2} \operatorname{KE}(\tilde{\pi}_\varepsilon) + \varepsilon^{-1/2} \int_{\mathbb{R}^2} V d\tilde{\pi}_\varepsilon \\ &\leq \varepsilon^{1/2} [\operatorname{KE}(v_\varepsilon^1) + \operatorname{KE}(v_\varepsilon^2)] + C_H \varepsilon^{-1/2} \int_{\mathbb{R}^2} V d\pi_\varepsilon + C \|\pi_\varepsilon\|_1 \\ &\leq \varepsilon^{1/2} [\operatorname{KE}(v_\varepsilon^1) + \operatorname{KE}(v_\varepsilon^2)] + C_H \varepsilon^{-1/2} \int_{\mathbb{R}^2} V d\pi_\varepsilon + C_H \tau + C \rho(\Omega_H^c). \end{aligned}$$

By Proposition 6.1, we have $\lim_{\varepsilon \rightarrow 0} \varepsilon^{-1/2} \int_{\mathbb{R}^2} V d\pi_\varepsilon = 0 = \lim_{\varepsilon \rightarrow 0} \varepsilon^{1/2} \operatorname{KE}(v_\varepsilon^i)$, and therefore

$$\lim_{\varepsilon \rightarrow 0} E_\varepsilon(\sqrt{\tilde{\pi}_\varepsilon}) \leq C \rho(\Omega_H^c). \quad (7.1)$$

Finally, let $\psi_\varepsilon = \sqrt{\bar{\gamma}_\varepsilon + \tilde{\pi}_\varepsilon}$; then ψ_ε has the correct marginals to be a recovery sequence for γ . Moreover, by the subadditivity of $E_\varepsilon(\sqrt{\gamma})$ with respect to γ and (7.1), we have

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\psi_\varepsilon) &\leq \limsup_{\varepsilon \rightarrow 0} [E_\varepsilon(\sqrt{\bar{\gamma}_\varepsilon}) + E_\varepsilon(\sqrt{\tilde{\pi}_\varepsilon})] \\ &\leq \limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\sqrt{\bar{\gamma}_\varepsilon}) + C \rho(\Omega_H^c) \\ &\leq \frac{1}{2} \int_{\mathbb{R}^2} \operatorname{tr}(\sqrt{D^2 V}) d\gamma + C \rho(\Omega_H^c). \end{aligned}$$

We now use a diagonal argument to conclude. For every $n \in \mathbb{N}_+$ we can find H_n large enough such that $C \rho(\Omega_{H_n}^c) < 1/n$. We can then find ε_n such that the ψ_{ε_n} constructed as above with parameters $H_n, \varepsilon_n, \beta_n, \delta_n, \tau_n, N_n$ satisfies

$$E_{\varepsilon_n}(\psi_{\varepsilon_n}) \leq \frac{1}{2} \int_{\mathbb{R}^2} \operatorname{tr}(\sqrt{D^2 V}) d\gamma + C \rho(\Omega_{H_n}^c) + 1/n$$

$$\leq \frac{1}{2} \int_{\mathbb{R}^2} \operatorname{tr}(\sqrt{D^2 V}) d\gamma + 2/n,$$

which is precisely the thesis of the theorem. $\square$

Proof of Theorem 1.1. To prove (1.4), we take a sequence ψ_ε which almost realizes the minimum in (1.6) up to a vanishing error. Up to subsequence, $|\psi_\varepsilon|^2 \rightharpoonup \gamma$ where $\gamma \in \Pi_0(\rho)$ with $\gamma(\{V > 0\}) = 0$. Applying Theorem 1.2 to this sequence we get

$$\liminf_{\varepsilon \rightarrow 0} \frac{F^\varepsilon(\rho) - F_{OT}(\rho)}{\sqrt{\varepsilon}} \geq \liminf_{\varepsilon \rightarrow 0} E_\varepsilon(\psi_\varepsilon) \geq \frac{1}{2} \int_{\mathbb{R}^{dN}} \operatorname{tr}(\sqrt{D^2 V}) d\gamma = F_{ZPO}(\rho).$$

To prove (1.5), we take $\gamma \in \Pi_0(\rho)$ with $\gamma(\{V > 0\}) = 0$ and $F_{ZPO}(\rho) = \frac{1}{2} \int_{\mathbb{R}^{dN}} \operatorname{tr}(\sqrt{D^2 V}) d\gamma$. If ψ_ε is a recovery sequence given by Theorem 1.3, we get

$$\limsup_{\varepsilon \rightarrow 0} \frac{F^\varepsilon(\rho) - F_{OT}(\rho)}{\sqrt{\varepsilon}} \leq \limsup_{\varepsilon \rightarrow 0} E_\varepsilon(\psi_\varepsilon) \leq F_{ZPO}(\rho).$$

$\square$

Appendix: proof of Lemmas 3.1 and 3.2

Proof of Lemma 3.1. Notice that $G_{M,N} < G_{M,\infty}$ by definition, so we only need to prove the second inequality of (3.2). We have $G_{\alpha,N} < G_{\alpha,\infty}$ and we claim that

$$G_{\alpha,\infty} - G_{\alpha,N} \leq 3e^{-N/2} G_{\alpha,\infty}. \quad (7.2)$$

By a change of variable we reduce to prove it only for $\alpha = 1$. We apply the pointwise inequality

$$f^2 \leq (f - e^{-N/2})^2 + 2fe^{-N/2} \quad (7.3)$$

to $f = e^{-t^2/2}$ for $t < \sqrt{N}$ and use $N \geq 1$ to get

$$\begin{aligned} G_{1,\infty} &= \int_{\{|t| \leq \sqrt{N}\}} e^{-t^2} dt + \int_{\{|t| > \sqrt{N}\}} e^{-t^2} dt \\ &\leq \int_{\{|t| \leq \sqrt{N}\}} \left[\left(e^{-t^2/2} - e^{-N/2} \right)^2 + 2e^{-t^2/2} e^{-N/2} \right] dt + \int_{\{|t| > \sqrt{N}\}} te^{-t^2} dt \\ &\leq G_{1,N} + 2\sqrt{2\pi} e^{-N/2} + e^{-N} = G_{1,N} + \left(2\sqrt{2} + \frac{e^{-N/2}}{\sqrt{\pi}} \right) e^{-N/2} G_{1,\infty}, \end{aligned}$$

which proves the claim (7.2) for $N \geq 3$, because the factor inside parentheses is less than 3. From this, (3.2) follows because

$$\begin{aligned} G_{M,\infty} - G_{M,N} &= G_{a,\infty}(G_{b,\infty} - G_{b,N}) + G_{b,N}(G_{a,\infty} - G_{a,N}) \\ &\leq G_{a,\infty} 3e^{-N/2} G_{b,\infty} + G_{b,N} 3e^{-N/2} G_{a,\infty} \\ &\leq 6e^{-N/2} G_{a,\infty} G_{b,\infty} = 6e^{-N/2} G_{M,\infty}. \end{aligned}$$

As a consequence, we also get that

$$\frac{1}{G_{M,N}} - \frac{1}{G_{M,\infty}} = \frac{G_{M,\infty} - G_{M,N}}{G_{M,\infty}G_{M,N}} \leq \frac{6e^{-N/2}}{G_{M,N}} \leq \frac{6e^{-N/2}}{(1 - 6e^{-N/2})G_{M,\infty}} \leq \frac{7e^{-N/2}}{G_{M,\infty}}. \quad (7.4)$$

Now let $f = e^{-aw^2/2}$ and $g = e^{-bz^2/2}$. Notice that $f, g \leq 1$ and that we can apply (7.3) to f and g . Therefore,

$$\begin{aligned} \left| \tilde{\Gamma}_{M,\infty}(\mathbf{x}) - \tilde{\Gamma}_{M,N}(\mathbf{x}) \right| &= f^2 g^2 - (f - e^{-N/2})_+^2 (g - e^{-N/2})_+^2 \\ &= f^2 \left(g^2 - (g - e^{-N/2})_+^2 \right) + \left(f^2 - (f - e^{-N/2})_+^2 \right) \\ &\quad (g - e^{-N/2})_+^2 \\ &\leq 2(f^2 g + f g^2) e^{-N/2} \leq 4e^{-N/2}. \end{aligned}$$

We can now use this information and (7.4) to estimate the difference of the two normalized Gaussians as

$$\begin{aligned} \|\Gamma_{M,\infty} - \Gamma_{M,N}\|_\infty &= \left\| \frac{\tilde{\Gamma}_{M,\infty} - \tilde{\Gamma}_{M,N}}{G_{M,\infty}} - \tilde{\Gamma}_{M,N} \left(\frac{1}{G_{M,N}} - \frac{1}{G_{M,\infty}} \right) \right\|_\infty \\ &= \frac{\|\tilde{\Gamma}_{M,\infty} - \tilde{\Gamma}_{M,N}\|_\infty}{G_{M,\infty}} + \|\tilde{\Gamma}_{M,N}\|_\infty \left(\frac{1}{G_{M,N}} - \frac{1}{G_{M,\infty}} \right) \\ &\leq \frac{4e^{-N/2}}{G_{M,\infty}} + \frac{7e^{-N/2}}{G_{M,\infty}} = \frac{11e^{-N/2}}{G_{M,\infty}} = \frac{11}{\pi} \sqrt{\det M} e^{-N/2}, \end{aligned}$$

proving (3.3). To prove the L^1 estimate (3.4), we divide the integral on $\text{supp } \Gamma_{M,N}$ and the complement. In the first, thanks to (3.3), we obtain

$$\begin{aligned} \|\Gamma_{M,N} - \Gamma_{M,\infty}\|_{L^1(\text{supp } \Gamma_{M,N})} &\leq \|\Gamma_{M,N} - \Gamma_{M,\infty}\|_\infty \cdot |\text{supp } \Gamma_{M,N}| \\ &\leq C \sqrt{\det M} e^{-N/2} \frac{4N}{\sqrt{ab}} \leq C e^{-N/2} N. \end{aligned}$$

In the complement, only $\Gamma_{M,\infty}$ is nonzero, and by (3.2) we have that

$$\begin{aligned} \|\Gamma_{M,N} - \Gamma_{M,\infty}\|_{L^1((\text{supp } \Gamma_{M,N})^c)} &= \int_{(\text{supp } \Gamma_{M,N})^c} \Gamma_{M,\infty}(\mathbf{x}) d\mathbf{x} = 1 - \int_{\text{supp } \Gamma_{M,N}} \Gamma_{M,\infty}(\mathbf{x}) d\mathbf{x} \\ &\leq 1 - \int_{\text{supp } \Gamma_{M,N}} \frac{\tilde{\Gamma}_{M,N}(\mathbf{x})}{G_{M,\infty}} d\mathbf{x} = 1 - \frac{G_{M,N}}{G_{M,\infty}} \leq C e^{-N/2} N. \end{aligned}$$

Let us now prove (3.5). Define R_x to be the section above x of the rectangle $\text{supp } \Gamma_{M,N}$, i.e.

$$R_x = \{y\} \mid (x, y) \in \text{supp } \Gamma_{M,N} = \{y\} \mid aw^2 \leq N, \quad bz^2 \leq N.$$

Since, $by^2 \leq b(w^2 + z^2) \leq aw^2 + bz^2 \leq a(w^2 + z^2) = a(x^2 + y^2)$, we have that

$$\{(x, y) \in \mathbb{R}^2 \mid a(x^2 + y^2) \leq N\} \subset \text{supp } \Gamma_{M,N}$$

and

$$R_x \subseteq \{y\}by^2 \leq 2N, \quad R_x^c \subseteq \{y\}ay^2 > N - ax^2.$$

Therefore, using (3.3), we can estimate that

$$\begin{aligned} |\eta_{M,\infty}(x) - \eta_{M,N}(x)| &\leq \int_{\mathbb{R}} |\Gamma_{M,\infty}(x, y) - \Gamma_{M,N}(x, y)| dy \\ &= \int_{R_x} |\Gamma_{M,\infty}(x, y) - \Gamma_{M,N}(x, y)| dy + \int_{R_x^c} \Gamma_{M,\infty}(x, y) dy. \end{aligned}$$

The first integral can be estimated with (3.3) as

$$\begin{aligned} \int_{R_x} |\Gamma_{M,\infty}(x, y) - \Gamma_{M,N}(x, y)| dy &\leq |R_x| \cdot \|\Gamma_{M,\infty} - \Gamma_{M,N}\|_{\infty} \\ &\leq \left(2\sqrt{2} \frac{\sqrt{N}}{\sqrt{b}}\right) (C\sqrt{ab}e^{-N/2}) \\ &\leq C\sqrt{a}\sqrt{N}e^{-N/2}. \end{aligned}$$

For the second integral we proceed as follows. Let $B = B(0, \sqrt{N/b}) = \{b(w^2 + z^2) < N\} = \{b(x^2 + y^2) < N\}$. Note that in R_x^c we have $e^{-aw^2 - bz^2} < e^{-N}$ because at least one among $aw^2 > N$ and $bz^2 > N$ is true. As a consequence, we have that

$$\int_{\mathbb{R}_x^c \cap B} e^{-aw^2 - bz^2} dy \leq e^{-N} \text{diam}(B) = \frac{2}{\sqrt{b}} e^{-N} \sqrt{N},$$

while on the other hand,

$$\begin{aligned} \int_{\mathbb{R}_x^c \setminus B} e^{-aw^2 - bz^2} dy &\leq \int_{\mathbb{R}_x^c \setminus B} e^{-bw^2 - bz^2} dy = \int_{\{by^2 > N - bx^2\}} e^{-bx^2 - by^2} dy \\ &= \frac{\sqrt{\pi}}{\sqrt{b}} e^{-bx^2} \text{erfc}\left(\sqrt{N - bx^2}\right) \\ &\leq \frac{\sqrt{\pi}}{\sqrt{b}} e^{-bx^2} e^{-N + bx^2} = \frac{\sqrt{\pi}}{\sqrt{b}} e^{-N}, \end{aligned}$$

and therefore

$$\int_{\mathbb{R}_x^c} e^{-aw^2 - bz^2} dy = \int_{\mathbb{R}_x^c \cap B} e^{-aw^2 - bz^2} dy + \int_{\mathbb{R}_x^c \setminus B} e^{-aw^2 - bz^2} dy \leq \frac{C}{\sqrt{b}} e^{-N} \sqrt{N}.$$

Hence,

$$\begin{aligned} \int_{R_x^c} \Gamma_{M,\infty}(x, y) dy &= G_{M,\infty}^{-1} \int_{R_x^c} e^{-(aw^2 + bz^2)} dy \leq \frac{\sqrt{ab}}{\pi} \frac{C}{\sqrt{b}} e^{-N} \sqrt{N} \\ &\leq C\sqrt{a}e^{-N}\sqrt{N}. \end{aligned}$$

In conclusion, putting the two estimates together, we have

$$|\eta_{M,\infty}(x) - \eta_{M,N}(x)| \leq C\sqrt{a}\sqrt{N}e^{-N/2}.$$

□

Proof of Lemma 3.2. In some computations of this proof we use the standard error function and the complementary error function

$$\operatorname{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt, \quad \operatorname{erfc}(z) = \frac{2}{\sqrt{\pi}} \int_z^\infty e^{-t^2} dt = 1 - \operatorname{erf}(z).$$

The crucial property that we will need is the fast decay at infinity of erfc implied by the bound

$$\operatorname{erfc}(z) \leq \frac{2}{\sqrt{\pi}} \int_z^\infty \frac{t}{z} e^{-t^2} dt \leq \frac{e^{-z^2}}{\sqrt{\pi} z}.$$

Since

$$(\operatorname{supp} \Gamma_{M,N})^c \subset \{|w| > \sqrt{N/a}\} \cup \{|z| > \sqrt{N/b}\},$$

with the aid of (3.4) we can estimate that

$$\begin{aligned} & \int_{\mathbb{R}^2} |M\mathbf{x}|^2 |\Gamma_{M,N}(\mathbf{x}) - \Gamma_{M,\infty}(\mathbf{x})| d\mathbf{x} \\ & \leq \int_{\operatorname{supp} \Gamma_{M,N}} |M\mathbf{x}|^2 |\Gamma_{M,N}(\mathbf{x}) - \Gamma_{M,\infty}(\mathbf{x})| d\mathbf{x} \\ & \quad + \int_{(\operatorname{supp} \Gamma_{M,N})^c} |M\mathbf{x}|^2 \Gamma_{M,\infty}(\mathbf{x}) d\mathbf{x} \\ & \leq \left(\sup_{\operatorname{supp} \Gamma_{M,N}} |M\mathbf{x}|^2 \right) \|\Gamma_{M,N} - \Gamma_{M,\infty}\|_1 \\ & \quad + G_{M,\infty}^{-1} 2 \int_{\sqrt{N/a}}^\infty \int_{\mathbb{R}} (a^2 w^2 + b^2 z^2) e^{-aw^2 - bz^2} dz dw \\ & \quad + G_{M,\infty}^{-1} 2 \int_{\sqrt{N/b}}^\infty \int_{\mathbb{R}} (a^2 w^2 + b^2 z^2) e^{-aw^2 - bz^2} dw dz \\ & \leq (a+b) C N e^{-N/2} + (a+b) \left(\operatorname{erfc}(\sqrt{\pi}) + \frac{\sqrt{N}}{\sqrt{\pi}} e^{-N} \right) \\ & \leq C \operatorname{tr}(M) N e^{-N/2}. \end{aligned}$$

For the kinetic energy, we can compute that

$$\nabla_{w,z} \sqrt{\Gamma_{M,N}(\mathbf{x})} = G_{M,N}^{-1/2} \begin{pmatrix} -aw e^{-aw^2/2} (e^{-bz^2/2} - e^{-N/2}) \\ -bz e^{-bz^2/2} (e^{-aw^2/2} - e^{-N/2}) \end{pmatrix}_+ \mathbf{1}_{\operatorname{supp} \Gamma_{M,N}}, \quad (7.5)$$

and hence,

$$\begin{aligned} \left| \nabla \sqrt{\Gamma_{M,N}(\mathbf{x})} \right|^2 &= G_{M,N}^{-1} \left[a^2 w^2 e^{-aw^2} (e^{-bz^2/2} - e^{-N/2})^2 \right. \\ & \quad \left. + b^2 z^2 e^{-bz^2} (e^{-aw^2/2} - e^{-N/2})^2 \right]_+ \mathbf{1}_{aw^2 \leq N, bz^2 \leq N}, \end{aligned}$$

and the kinetic energy of the truncated Gaussian ends up being

$$\begin{aligned}
 \text{KE}(\Gamma_{M,N}) &= \frac{1}{2} G_{M,N}^{-1} \left(\int_{\{aw^2 \leq N\}} a^2 w^2 e^{-aw^2} dw \right) G_{b,N} \\
 &\quad + \frac{1}{2} G_{M,N}^{-1} \left(\int_{\{bz^2 \leq N\}} b^2 z^2 e^{-bz^2} dz \right) G_{a,N} \\
 &= \frac{1}{2} G_{a,N}^{-1} \sqrt{a} \left(\frac{\sqrt{\pi}}{2} \text{erf}(\sqrt{N}) - e^{-N} \sqrt{N} \right) \\
 &\quad + \frac{1}{2} G_{b,N}^{-1} \sqrt{b} \left(\frac{\sqrt{\pi}}{2} \text{erf}(\sqrt{N}) - e^{-N} \sqrt{N} \right) \\
 &= (G_{a,N}^{-1} \sqrt{a} + G_{b,N}^{-1} \sqrt{b}) \frac{\sqrt{\pi}}{4} \left(\text{erf}(\sqrt{N}) - \frac{2}{\sqrt{\pi}} e^{-N} \sqrt{N} \right),
 \end{aligned}$$

where to pass from the first to the second line we integrated by parts and changed variables. Recalling that $G_{\alpha,\infty}^{-1} = \sqrt{\alpha}/\sqrt{\pi}$, for $N = \infty$, this gives

$$\text{KE}(\Gamma_{M,\infty}) = \left(\frac{\sqrt{a}}{\sqrt{\pi}} \sqrt{a} + \frac{\sqrt{b}}{\sqrt{\pi}} \sqrt{b} \right) \frac{\sqrt{\pi}}{4} = \frac{a+b}{4} = \frac{\text{tr } M}{4},$$

as already stated in (3.7). In order to compare the two energies, we estimate separately the difference of the terms involving a and b . Using the fact that, thanks to (7.2),

$$\frac{1}{G_{\alpha,N}} - \frac{1}{G_{\alpha,\infty}} = \frac{G_{\alpha,\infty} - G_{\alpha,N}}{G_{\alpha,\infty} G_{\alpha,N}} \leq \frac{3e^{-N/2}}{G_{\alpha,N}} \leq \frac{3e^{-N/2}}{(1 - 3e^{-N/2}) G_{\alpha,\infty}} \leq \frac{4e^{-N/2}}{G_{\alpha,\infty}},$$

we can estimate that

$$\begin{aligned}
 &\left| \frac{a}{4} - G_{a,N}^{-1} \sqrt{a} \frac{\sqrt{\pi}}{4} \left(\text{erf}(\sqrt{N}) - \frac{2}{\sqrt{\pi}} e^{-N} \sqrt{N} \right) \right| \\
 &\leq \left| \frac{a}{4} - G_{a,\infty}^{-1} \sqrt{a} \frac{\sqrt{\pi}}{4} \left(\text{erf}(\sqrt{N}) - \frac{2}{\sqrt{\pi}} e^{-N} \sqrt{N} \right) \right| \\
 &\quad + \left| (G_{a,N}^{-1} - G_{a,\infty}^{-1}) \sqrt{a} \frac{\sqrt{\pi}}{4} \left(\text{erf}(\sqrt{N}) - \frac{2}{\sqrt{\pi}} e^{-N} \sqrt{N} \right) \right| \\
 &\leq \frac{a}{4} \left| \text{erfc}(\sqrt{N}) + \frac{2}{\sqrt{\pi}} e^{-N} \sqrt{N} \right| + 4e^{-N/2} G_{a,\infty}^{-1} \sqrt{a} \frac{\sqrt{\pi}}{4} \\
 &\quad \times \left| \text{erf}(\sqrt{N}) - \frac{2}{\sqrt{\pi}} e^{-N} \sqrt{N} \right| \\
 &\leq C \frac{a}{4} e^{-N/2},
 \end{aligned}$$

and similarly for b , and therefore

$$|\text{KE}(\Gamma_{M,\infty}) - \text{KE}(\Gamma_{M,N})| \leq C \frac{a+b}{4} e^{-N/2} = C \text{KE}(\Gamma_{M,\infty}) e^{-N/2}.$$

□

Acknowledgements M.C. and F.S. have been supported by the SNSF Grant 182565.

Funding Open access funding provided by EPFL Lausanne This project has received funding from Ministero dell'Università e della Ricerca (PRIN project 202244A7YL of S.D.M. and MIUR Excellence Department Project 2023-2027 awarded to the DIMA of the University of Genova, CUP D33C23001110001), Air Force (AFOSR project FA8655-22-1-7034 of S.D.M.) and Istituto Nazionale di Alta Matematica (S.D.M. is a member of GNAMPA)

Declarations

Conflict of Interest The authors have no relevant financial or non-financial interests to disclose.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. BINDINI, UGO: Smoothing operators in multi-marginal optimal transport. *Math. Phys. Anal. Geometry* **23**(2), 21, 2020
2. BINDINI, UGO, DE PASCALE, LUIGI: Optimal transport with Coulomb cost and the semi-classical limit of density functional theory. *J. Éc. polytech. Math.* **4**, 909–934, 2017
3. BINDINI, UGO, PASCALE, DE.: Luigi, Kausamo, Anna: On deterministic solutions for multi-marginal optimal transport with Coulomb cost. *Commun. Pure Appl. Anal.* **21**(4), 1189–1208, 2022
4. BLIEDTNER, J., LOEB, P.: A reduction technique for limit theorems in analysis and probability theory. *Arkiv för Matematik* **30**(1), 25–43, 1992
5. BOUCHITTÉ, G., JIMENEZ, C., RAJESH, M.: A new L^∞ estimate in optimal mass transport. *Proc. Amer. Math. Soc.* **135**(11), 3525–3535, 2007
6. BOUCHITTÉ, GUY, BUTTAZZO, GIUSEPPE, CHAMPION, THIERRY, DE PASCALE, LUIGI: Dissociating limit in Density Functional Theory with Coulomb optimal transport cost, 2018.
7. BUTTAZZO, GIUSEPPE, CHAMPION, THIERRY, DE PASCALE, LUIGI: Continuity and estimates for multimarginal optimal transportation problems with singular costs. *Appl. Math. Optim.* **78**(1), 185–200, 2018
8. BUTTAZZO, GIUSEPPE, DE PASCALE, LUIGI, GORI-GIORGI, PAOLA: Optimal-transport formulation of electronic density-functional theory. *Phys. Rev. A* **85**, 062502, 2012
9. COLOMBO, M., DE PASCALE, L., DI MARINO, S.: Multimarginal optimal transport maps for 1-dimensional repulsive costs. *Canad. J. Math.* **67**, 350–368, 2015
10. COLOMBO, MARIA, DI MARINO, SIMONE, STRA, FEDERICO: Continuity of multimarginal optimal transport with repulsive cost. *SIAM J. Math. Anal.* **51**(4), 2019
11. COLOMBO, MARIA, STRA, FEDERICO: Counterexamples in multimarginal optimal transport with Coulomb cost and spherically symmetric data. *Math. Models Methods Appl. Sci.* **26**(6), 1025–1049, 2016

12. CONFORTI, GIOVANNI, TAMANINI, LUCA: A formula for the time derivative of the entropic cost and applications. *J. Funct. Anal.* 108964, 2021
13. COTAR, CODINA, FRIESECKE, GERO, KLÜPPELBERG, CLAUDIA: Density functional theory and optimal transportation with Coulomb cost. *Comm. Pure Appl. Math.* **66**(4), 548–599, 2013
14. COTAR, CODINA, FRIESECKE, GERO, KLÜPPELBERG, CLAUDIA: Smoothing of transport plans with fixed marginals and rigorous semiclassical limit of the Hohenberg–Kohn functional. *Arch. Rat. Mech. Ana.* **228**(3), 891–922, 2018
15. COTAR, CODINA, FRIESECKE, GERO, PASS, BRENDAN: Infinite-body optimal transport with Coulomb cost. *Calculus Variat. Part. Diff. Equ.* **54**(1), 717–742, 2015
16. COTAR, CODINA, PETRACHE, MIRCEA: Equality of the jellium and uniform electron gas next-order asymptotic terms for Coulomb and Riesz potentials, 2019.
17. COTAR, CODINA, PETRACHE, MIRCEA: Next-order asymptotic expansion for n-marginal optimal transport with Coulomb and Riesz costs. *Adv. Math.* **344**, 137–233, 2019
18. DE PASCALE, L.: Optimal transport with Coulomb cost: approximation and duality. *ESAIM: M2AN* **49**, 1643–1657, 2015
19. DI MARINO, SIMONE, LOUET, JEAN: The entropic regularization of the Monge problem on the real line. *SIAM J. Math. Anal.* **50**, 2017
20. FRIESECKE, GERO, GEROLIN, AUGUSTO, GORI-GIORGI, PAOLA: The strong-interaction limit of density functional theory. *Density Functional Theory: Modeling, Mathematical Analysis, Computational Methods, and Applications*, 183–266, 2022.
21. GEROLIN, AUGUSTO, KAUSAMO, ANNA, RAJALA, TAPIO: Duality theory for multi-marginal optimal transport with repulsive costs in metric spaces. *ESAIM Control Optim. Calc. Var.* **25**, 62, 21, 2019
22. GORI-GIORGI, P., VIGNALE, G., SEIDL, M.: Electronic zero-point oscillations in the strong-interaction limit of density functional theory. *J. Chem. Theory Comput.* **5**(4), 743–753, 2009
23. GORI-GIORGI, P., SEIDL, M., VIGNALE, G.: Density-functional theory for strongly interacting electrons. *Phys. Rev. Lett.* **103**, 166402, 2009
24. HOHENBERG, PIERRE, KOHN, WALTER: Inhomogeneous electron gas. *Phys. Rev.* **136**(3B), B864, 1964
25. LEWIN, MATHIEU: Semi-classical limit of the Levy-Lieb functional in density functional theory. *Comptes Rendus Mathématique* **356**(4), 449–455, 2018
26. LEWIN, MATHIEU, LIEB, ELLIOTT H., SEIRINGER, ROBERT: Statistical mechanics of the uniform electron gas. *Journal de l'École polytechnique — Mathématiques* **5**, 79–116, 2018
27. LEWIN, MATHIEU, LIEB, ELLIOTT H., SEIRINGER, ROBERT: The local density approximation in density functional theory. *Pure Appl. Anal.* **2**(1), 35–73, 2020
28. LEWIN, M., LIEB, E.H., SEIRINGER, R.: Universal functionals in density functional theory. In: *Density Functional Theory: Modeling, Mathematical Analysis, Computational Methods, and Applications*, pp. 115–182. Springer, 2022.
29. LIEB, ELLIOTT H.: A lower bound for Coulomb energies. *Phys. Lett. A* **70**(5), 444–446, 1979
30. LIEB, E.H.: Density functionals for Coulomb systems. *Int. J. Quantum Chem.* **24**(3), 243–277, 1983
31. LIEB, ELLIOTT H., OXFORD, STEPHEN: Improved lower bound on the indirect Coulomb energy. *Int. J. Quantum Chem.* **19**(3), 427–439, 1981
32. MALET, F., MIRTSCHINK, A., MENDEL, C.B., BJERLIN, J., KARABULUT, E.Ö., REIMANN, S.M., GORI-GIORGI, P.: Density-functional theory for strongly correlated bosonic and fermionic ultracold dipolar and ionic gases. *Phys. Rev. Lett.* **115**(3), 033006, 2015
33. MALET, F., MIRTSCHINK, A., GIESBERTZ, K.J.H., WAGNER, L.O., GORI-GIORGI, P.: Exchange-correlation functionals from the strong interaction limit of DFT: applications to model chemical systems. *Phys. Chem. Chem. Phys.* **16**(28), 14551–14558, 2014
34. MORSE, A.P.: Perfect blankets. *Trans. Am. Math. Soc.* **61**(3), 418–442, 1947

35. PAL, SOUMIK: On the difference between entropic cost and the optimal transport cost. *Ann. Appl. Probab.* **34**(1B), 1003–1028, 2024
36. PETRACHE, MIRCEA, SERFATY, SYLVIA: Next order asymptotics and renormalized energy for Riesz interactions. *J. Inst. Math. Jussieu* **16**(3), 501–569, 2017
37. SANDIER, ÉTIENNE, SERFATY, SYLVIA, et al.: 2d Coulomb gases and the renormalized energy. *Ann. Probab.* **43**(4), 2026–2083, 2015
38. SEIDL, M., DI MARINO, S., GEROLIN, A., NENNA, L., GIESBERTZ, K., GORI-GIORGI, P.: The strictly-correlated electron functional for spherically symmetric systems revisited.
39. SEIDL, MICHAEL: Strong-interaction limit of density-functional theory. *Phys. Rev. A* **60**(6), 4387, 1999
40. SEIDL, MICHAEL, GORI-GIORGI, PAOLA, SAVIN, ANDREAS: Strictly correlated electrons in density-functional theory: a general formulation with applications to spherical densities. *Phys. Rev. A* **75**(4), 042511, 2007
41. SIMON, BARRY: Semiclassical analysis of low lying eigenvalues. i. non-degenerate minima: asymptotic expansions. *Annales de l'IHP Physique théorique* **38**, 295–308, 1983
42. SIMON, BARRY: Semiclassical analysis of low lying eigenvalues, ii. tunneling. *Ann. Math.* 89–118, 1984
43. VUCKOVIC, STEFAN, GORI-GIORGI, PAOLA: Simple fully nonlocal density functionals for electronic repulsion energy. *J. Phys. Chem. Lett.* **8**(13), 2799–2805, 2017

MARIA COLOMBO

EPFL, AMCV,

Lausanne

Switzerland.

e-mail: maria.colombo@epfl.ch

and

SIMONE DI MARINO

Università di Genova (DIMA),

MaLGa,

Genova

Italy.

e-mail: simone.dimarino@unige.it

and

FEDERICO STRA

Politecnico di Torino (DISMA),

Torino

Italy.

e-mail: federico.stra@polito.it

(Received September 20, 2022 / Accepted October 28, 2025)

© The Author(s) (2025)