

Jacobi Processes with Jumps as Neuronal Models: A First Passage Time Analysis

Original

Jacobi Processes with Jumps as Neuronal Models: A First Passage Time Analysis / D'Onofrio, Giuseppe; Patie, Pierre; Sacerdote, Laura. - In: SIAM JOURNAL ON APPLIED MATHEMATICS. - ISSN 0036-1399. - 84:1(2024), pp. 189-214. [10.1137/22M1516877]

Availability:

This version is available at: 11583/2985732 since: 2024-04-04T08:34:00Z

Publisher:

SIAM Society for industrial and applied mathematics

Published

DOI:10.1137/22M1516877

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

JACOBI PROCESSES WITH JUMPS AS NEURONAL MODELS : A FIRST PASSAGE TIME ANALYSIS

GIUSEPPE D'ONOFRIO, PIERRE PATIE, AND LAURA SACERDOTE

ABSTRACT. To overcome some limits of classical neuronal models, we propose a Markovian generalization of the classical model based on Jacobi processes by introducing downwards jumps to describe the activity of a single neuron. The statistical analysis of inter-spike intervals is performed by studying the first-passage times of the proposed Markovian Jacobi process with jumps through a constant boundary. In particular, we characterize its Laplace transform which is expressed in terms of some generalization of hypergeometric functions that we introduce, and deduce a closed-form expression for its expectation. Our approach, which is original in the context of first passage time problems, relies on intertwining relations between the semigroups of the classical Jacobi process and its generalization, which have been recently established in [11]. A numerical investigation of the firing rate of the considered neuron is performed for some choices of the involved parameters and of the jumps distributions.

1. INTRODUCTION

Among the models used for the description of single neuron's activity the leaky integrate-and-fire (LIF) model is still an extremely useful tool, despite its age and simplicity [16, 53]. The LIF model describes the time evolution of the voltage across the membrane of the neuron until it reaches a certain threshold. This event is called action potential (or spike) and it is believed that the distribution of these spikes encodes the information that the neurons transfer. It is assumed that the neuron under study is point-like and receives inputs from the surrounding network of neurons that are summed up (*integrate*) producing a change in the voltage value. The term *leaky* indicates that, in the absence of input, the membrane potential decays exponentially to its resting value. In accordance with the model, the spikes are instantaneous events that are generated as soon as the voltage reaches a certain value for the first time (*fire*). After that, the process is reset to its starting value and the evolution starts over again. Sometimes a refractory period is added to the model, i.e. there is a time interval after a spike in which a nerve cell is unable to fire an action potential.

Since for some types of neurons the incoming inputs are frequent and relatively small, a diffusion limit over the discrete process, see [56], describing the membrane potential evolution is performed to gain the higher mathematical tractability of the Ornstein-Uhlenbeck process [50]. The latter has been widely used for decades, although it presents some drawbacks. The Ornstein-Uhlenbeck process, indeed, allows unlimited values for the neuronal potential, it does not include that the changes in the potential of a nerve cell depend on its actual value and it does not take into account the geometry of the neuron. Some models with multiplicative noise have been proposed to overcome the first two unrealistic features of the classical LIF model [10],[37]. Among them, recently, a Jacobi process has been proposed for the description of the activity of a neuronal membrane [18]. The Jacobi process has a bounded state space, that is the value of the membrane potential is confined below and above by two fixed values that, for physiological reasons, are called the inhibitory and excitatory reversal potentials. Moreover, the change in the membrane potential determined by an

2010 *Mathematics Subject Classification.* 37A30, 47D06, 47G20, 60J75, 60J70.

Key words and phrases. Jacobi process, first passage time, non-local Wright-Fisher process, intertwining relations, Markov semigroups, infinitesimal generators, neuronal modeling.

incoming input depends on the distance between its actual state and the two reversal potentials, fact that is well established about the physiology of the synapses, see [21] or [30] for classical references.

However, the pure-diffusion models do not account for the spatial geometry of the neurons and do not discriminate among different sources of incoming inputs. In fact, more realistic models should assign different weights for the synaptic contribution impinging the neuron in different points of the membrane depending on whether they are more or less close to the trigger zone, as a first attempt to overcome the point-size assumption, in the spirit of multi-compartmental models, see for instance [52]. In order to include these features in the model, and since not all the inputs are infinitesimal and their frequencies may prevent a diffusion limit, jump-diffusion models have been proposed [23, 55]. These models have been proven to describe the activity of motor neurons and pyramidal neurons [32, 42]. Here, we investigate the features of a neuronal model in which the membrane potential evolves, between two consecutive spikes, according to a Jacobi type process with state-dependent downward jumps.

To develop the analysis on its firing activity, we investigate the first-passage-time (FPT) problem for the Jacobi process with jumps, that is the mathematical counterpart of the time of generation of the action potential. To this end, we propose an original approach in this context, which is based on intertwining relationships between the semigroups of the generalized Jacobi processes and the one of the classical Jacobi diffusion process which were identified recently in [11], see also [44] for further analytical results on these semigroups. Intertwining relations are a type of commutation relations that form a classification scheme for linear operators. They have proved to be a natural and powerful concept in a variety of contexts in mathematics ranging from the construction of new Markov semigroups to the spectral and ergodicity theory of non-self-adjoint semigroups, see [11, 12, 33, 49, 47] and the references therein. This paper provides an additional application of such concept in the potential theory of Markov processes by transferring q -invariant functions from a reference semigroup to semigroups that are in its (intertwining) orbit. This device enables us to characterize the Laplace transform of the first passage time of the process through a boundary (or in the modeling framework the firing time of the neuron) in terms of some generalized hypergeometric functions that we introduce, thanks to the relationship with the classical Jacobi model. We point out that the intertwining approach could also be used to map q -invariant functions, or more generally q -excessive functions, between semigroups associated to Markov processes with arbitrary jumps.

The strengths of the presented model rely on the improved adherence to phenomenological reality: the jumps are state-dependent and are able to reduce the firing rate and introduce saturation. The latter feature is observed in other models only if a non-zero refractory period is introduced. Otherwise the firing rate can generally grow unbounded, and this is clearly unrealistic. Moreover the high degree of freedom in the choice of the jump distribution allows the description of different situations.

Finally we stress that, despite the application in the context of mathematical neurosciences, the results on the Jacobi process with jumps and its first passage time through a constant boundary are novel and of a general nature. We mention that Jacobi processes have been popular in applications such as population genetics, under the name Wright-Fisher diffusion, see e.g. Griffiths et al. [28, 27], Huillet [31], and Pal [46], and in finance, see e.g. Delbaen and Shirikawa [17] and Gourieroux and Jasiak [25].

The paper is organized as follows. In Section 2, we introduce the neuronal model based on the jump-diffusion Jacobi process through its infinitesimal generator and a qualitative description of the dynamics together with the involved parameters. Mathematical results on the non-local Jacobi operator, necessary for the analysis of the model, are obtained in Section 3 using intertwining relations between the classical Jacobi semigroup and the non-local one. In particular we show that the process under study has only downward jumps and we provide an explicit form of the Laplace transform of the related first-passage time. Using these results, in Section 4, an analysis of the firing

activity of the neuron described by a Jacobi process with jumps is carried out focusing on some illustrative examples. In particular, we show that the jumps reduce the firing rate and introduce a saturation effect, despite the absence of a refractory period. We point out that the readers with stronger interest in the biophysical aspect of this work may skip the mathematical details provided in Section 3.

2. A JACOBI PROCESS WITH JUMPS AS A NEURONAL MODEL

We describe the evolution of the neuronal membrane potential between two consecutive spikes of a single neuron as the Markovian realization $X = (X_t)_{t \geq 0}$ of a non-local perturbation of the generator \mathbf{J}_V of the classical Jacobi neuronal model $V = (V_t)_{t \geq 0}$. We recall that the latter is obtained as a Kurtz-type diffusion approximation of a Stein's model with reversal potentials [56]. In that model two independent homogeneous Poisson processes represent the excitatory and inhibitory neuronal inputs, with intensities $\nu_E > 0$ and $\nu_I > 0$, respectively. They describe the arrival of excitatory and inhibitory potentials and are such that the input parameters are

$$(2.1) \quad \mu_e = e\nu_E \quad \text{and} \quad \mu_i = i\nu_I$$

where i and e are constants such that $-1 < i < 0 < e < 1$. Denoting by $V_I < 0 < V_E$ the inhibitory and excitatory reversal potentials, respectively, we recall that V is a diffusion on $E_V = [V_I, V_E]$ solving the following stochastic differential equation

$$(2.2) \quad dV_t = \left(-\frac{1}{\tau}V_t + \mu_e(V_E - V_t) + \mu_i(V_t - V_I) \right) dt + \sigma \sqrt{(V_E - V_t)(V_t - V_I)} dW_t.$$

where the diffusion coefficient $\sigma > 0$ controls the amplitude of the noise, $W = (W_t)_{t \geq 0}$ is a standard Wiener process and $\tau > 0$ is the membrane time constant taking into account the spontaneous voltage decay (*leak*) toward the resting potential (set equal to zero here) in the absence of inputs, μ_e and μ_i . Finally, the refractory period is assumed equal to zero. The model $V = (V_t)_{t \geq 0}$ often goes under the name of *leaky integrate-and-fire* with reversal potentials [37].

Alternatively V can be described through its infinitesimal generator taking the form, for a smooth function f on E_V ,

$$(2.3) \quad \mathbf{J}_V f(x) = \frac{\sigma^2}{2}(V_E - x)(x - V_I)f''(x) - \left(\frac{1}{\tau}x - \mu_e(V_E - x) - \mu_i(x - V_I) \right) f'(x),$$

throughout the paper this way of writing will be more convenient.

A limit of neuronal models based on diffusion processes is that all the inputs which affect the membrane potential are summed together and homogenized disregarding their origin or strength, with the advantage of a continuous trajectory for the dynamics. In [55], [23], [57] there are first attempts to introduce jumps but occurring at exponential times. Moreover the mathematical tractability requested the use of jumps of constant amplitude or the use of numerical simulations. Taking advantage of the intertwining approach here we introduce and study mathematically the case of jumps that are state-dependent both in frequency and amplitude. The dynamics of the voltage X between two consecutive spikes of the new class of neuronal models that we propose here is described as a Markov (in fact, a Feller) process on E_V with càdlàg trajectories whose infinitesimal generator, for a smooth function f on E_V , is given as the following non-local perturbation of the generator of the classical Jacobi process

$$(2.4) \quad \mathbf{J}_X f(x) = \mathbf{J}_V f(x) + \int_{V_I}^{V_E} (f(r) - f(x)) N_V(x, dr)$$

where the kernel $N_V(x, dr) = \frac{V_E - V_I}{x - V_I} \Pi_V(x, dr) \mathbb{1}_{\{r < x\}}$ with Π_V the measure image, by the mapping $r \mapsto \ln(\frac{x - V_I}{r - V_I}) \mathbb{1}_{\{r < x\}}$, of Π a finite non-negative Radon measure on \mathbb{R}_+ with $\int_0^\infty r \Pi(dr) < \infty$. We shall

show in Proposition 3.1 that the family (indexed by Π) of linear operators \mathbb{J}_X is the infinitesimal generator of a Feller process admitting an unique stationary measure. This will be achieved by identifying a homeomorphism, i.e. an intertwining relation à la Dynkin, between these semigroups and the one of Jacobi processes with jumps on $(0, 1)$ recently introduced in [11], where the process jumps from state x to state $e^{-r}x$ at a frequency given by $\Pi(dr)/x$, that is inversely proportional to the achieved state.

We point out that the assumptions on the measure Π ensure that the operator defined in (2.4), endowed with its domain, is the generator of a Markov process. We also remark that the jumps are only downwards, but, both the amplitude and the intensity of the jumps are state-dependent. In fact if the voltage approaches the inhibitory reversal potential, the number of jumps is high but the corresponding depolarization is small. Conversely for higher values of the voltage the frequency of jumps decreases whereas their amplitude depends on Π . See Fig. 1 for an example of a possible path of the proposed model.

If the jump kernel is a finite measure, which is the case of Π_V here, there is a nice and more formal path interpretation of the Markov process that can be read off from its generator, see e.g. Bass [6]. Indeed, one has the following description of dynamics of the voltage X : the potential starts by undergoing the same dynamics than the classical Jacobi neuronal model V until being killed at a random time \mathcal{T} whose survival probability up to time t is given by $e^{-\frac{V_E - V_I}{X_t - V_I} \Pi(\mathbb{R}^+)}$, where we used the fact that $N_V(x, E_V) = \frac{V_E - V_I}{x - V_I} \Pi(\mathbb{R}^+)$. At the time of death \mathcal{T} , restart it (in a sense made precise through for instance the work of Meyer [43]) with distribution $\frac{N_V(X_{\mathcal{T}-}, dr)}{N_V(X_{\mathcal{T}}, E_V)}$, where $X_{t-} = \lim_{s \uparrow t} X_s$ stands for the left-limit, and, repeat the procedure. In other words, the neuronal model X behaves like the classical Jacobi neuronal model but at some random times performs downwards jumps (as we have, by definition, the support of the kernel $N_V(x, dr)$ is $V_I < r < x$) according to the distribution given above. In particular, the closer x gets to the inhibitory reversal potential V_I , the larger $N_V(x, E_V)$ is, that is the number of jumps becomes more frequent but the corresponding hyperpolarization is small, as the support of the distribution of the amplitude of jumps is $[0, x - V_I]$. Also, different choices of Π allow different sizes of the jumps: the more mass Π concentrates around zero, the smaller is the amplitude of jumps, if Π admits large values with high probability, the voltage can be almost reset after the jump. We stress that the latter scenario could be the result of a rare event for the type of measures Π considered in the following. We consider the case when $\Pi(dr) = e^{-\alpha r} dr, \alpha, r > 0$. Then, easy computation yields that $N_V(x, dr) = (V_E - V_I) \frac{(r - V_I)^{\alpha-1}}{(x - V_I)^{\alpha+1}} \mathbb{I}_{\{r < x\}} dr$ and thus $N_V(x, E_V) = \frac{V_E - V_I}{\alpha(x - V_I)}$. In this case, the probability that there is a jump of amplitude lower than $y \in (0, x - V_I)$ is given by $\left(1 - \frac{y}{x - V_I}\right)^{\alpha+1}$.

At first sight, one may be surprised that the dynamics of the neuronal stochastic model is described in terms of the generator compared to the usual path definition of diffusions as solution to a stochastic differential equation. However, this is probably the most natural way when one is dealing with state dependent jumps processes. Indeed, the theory of stochastic differential equations for Markov processes with jumps is still incomplete regarding for instance the existence and uniqueness of a solution, and when available for state dependent jumps processes, it involves integral with respect to some Poisson random measures which makes its interpretation scarcely intuitive.

Remark 2.1. For a better understanding of the dynamics and the interpretation of the involved parameters, we draw a parallel between the model (2.4) and the equation of a conductance-based neuronal model. The evolution in time of the potential difference $\tilde{V} = (\tilde{V}_t)_{t \geq 0}$ across the membrane of a neuron is given by

$$(2.5) \quad C \frac{d\tilde{V}_t}{dt} = -g_L \tilde{V}_t + I_{syn} + I = -g_L \tilde{V}_t - g_E(t)(\tilde{V}_t - V_E) + g_I(t)(\tilde{V}_t - V_I) + I,$$




FIGURE 1. A realization of a Jacobi process with jumps (bottom figure) and corresponding number of jumps (top figure). The jump frequency increases if the depolarization V_t approaches V_I (in the figure $V_I = -10$, $V_E = 100$), as it can be seen from the two plots for $t > 0.6$, while the amplitude of the jumps decreases.

where $C > 0$ is the membrane capacitance, $g_L > 0$ is the conductance of the leak current, while $g_E(t) > 0$ and $g_I(t) > 0$ are the conductances of the excitatory and inhibitory components of the synaptic current I_{syn} , see e.g. [51]. The current I accounts for large or external inhibitory inputs that cannot be considered in the diffusion limit. Differently from classical external currents considered in the literature, that are constant or periodic functions, here I has random components. The dynamics (2.5) with $I = 0$ is analogous to the deterministic version of the leaky integrate and fire model (2.2), with C/g_L playing the role of the membrane constant τ . Moreover at time t_k , the time of the arrival of the k -th incoming excitatory pulse, distributed according to a Poisson process with parameter ν_e , the conductance $g_E(t)$ increases by a factor of Ce . Consequently the increase in the voltage is $\Delta V = e(V_E - V_t)$, where e is a dimensionless constant measuring the strength of the synapse (similarly for $g_I(t)$ and ν_i). This increment corresponds to the one of the LIF model before taking the diffusion limit. Analogously the current I (the jump part in the proposed model) describes a possibly external (or strong) inhibitory current that at random times τ_k , determined by a measure Π , decreases instantaneously the voltage of a quantity that depends on N_V as described above, but in any case of a quantity smaller than $(V_{\tau_k} - V_I)$.

The motivation for considering the Jacobi process with state dependent jumps as a model of neuron's activity are several folds. On the one hand, the inhibition is well known to be regulatory of neuronal excitability and has a role in information transmission. The study, for state-dependent inputs, of the effect of inhibition on output indicators like signal to noise ratio, effective diffusion coefficient of the spike count and degree of coherence demonstrates that inhibitory input acts to decrease membrane potential fluctuations increasing spike regularity, see for example [4, 5, 19, 55]. Moreover the state dependence of the jumps preserves the fundamental improvement with respect to the Ornstein-Uhlenbeck model that the changes in the potential depends on its actual value. In addition it allows the possible description of the sites in which the neuron receives the inputs giving the chance to relax the assumption that the cell is point-like.

We prove in Lemma 3.1 that, under assumptions motivated by realistic interpretation of the involved parameters, X has only downward jumps. This property suggests to apply, possibly, the model to the probabilistic study of the effect of anti-epileptic drugs on a neuron whose firing activity is too intense, see [58].

3. JACOBI PROCESSES WITH JUMPS AND THEIR FIRST PASSAGE TIME PROBLEMS

In this section, we start by providing a homeomorphism between the neuronal model X defined in the previous section and generalized Jacobi processes with jumps that have been introduced in [11]. Then, we proceed by characterizing the Laplace transform of the first passage time to a fixed level by these generalized Jacobi processes.

3.1. Jacobi processes and neuronal models with jumps. Let us denote by $Y = (Y_t)_{t \geq 0}$ the generalized Jacobi process with jumps defined in [11] as follows. It is a Feller process on $[0, 1]$ whose infinitesimal generator is given, for a smooth function f on $[0, 1]$, by

$$(3.1) \quad \mathbb{J}_Y f(y) = \mathbf{J}_\mu f(y) + \int_0^\infty (f(e^{-r}y) - f(y)) \frac{\Pi(dr)}{y}$$

where \mathbf{J}_μ is the classical Jacobi operator

$$\mathbf{J}_\mu f(y) = \frac{\sigma^2}{2} y(1-y) f''(y) - (\lambda y - \mu) f'(y)$$

with $\sigma^2 > 0$, Π is, as in (2.4), a finite, non-negative Radon measure on \mathbb{R}_+ with $\hbar = \int_0^\infty r \Pi(dr) < \infty$, and, we have set, to simplify the notation,

$$(3.2) \quad \lambda = \frac{1}{\tau} + \mu_e - \mu_i \text{ and } \mu = \mu_e - \frac{V_I}{\tau(V_E - V_I)},$$

where these parameters were introduced in (2.1) and (2.3). Throughout, we impose the following assumption that guarantees that V_I is an entrance boundary

$$(3.3) \quad \mu_e > \hbar + \frac{\sigma^2}{2} + \frac{V_I}{\tau(V_E - V_I)} \text{ or, equivalently, } \lambda > \mu > \hbar + \frac{\sigma^2}{2},$$

since $\mu_i < 0$. Note that in [11], $\sigma^2 = 2$, and we shall explain in Lemma 3.4 below how to relate our generator \mathbb{J}_Y to the one of [11]. In what follows, we recall some basic results from [11] that will be useful for our analysis by adapting them, in an obvious way, for any $\sigma^2 > 0$. For instance, the condition (3.3) is the appropriate modification of the standing assumption in [11]. Next, let us write, for $u \geq 0$,

$$(3.4) \quad \phi(u) = u + \frac{2}{\sigma^2} \left(\mu - \hbar - \frac{\sigma^2}{2} + \int_0^\infty (1 - e^{-ur}) \bar{\Pi}(r) dr \right)$$

where $\bar{\Pi}(r) = \int_r^\infty \Pi(du)$. We observe that the condition (3.3), i.e. $\mu > \hbar + \frac{\sigma^2}{2}$, is also equivalent to

$$(3.5) \quad \phi(0) = \frac{2}{\sigma^2} \left(\mu - \hbar - \frac{\sigma^2}{2} \right) > 0.$$

Under this condition, it is not difficult to check that ϕ is a Bernstein function, i.e. $\phi : [0, \infty) \rightarrow [0, \infty)$ is infinitely differentiable on \mathbb{R}_+ and $(-1)^{n+1} \frac{d^n}{du^n} \phi(u) \geq 0$, for all $n = 1, 2, \dots$ and $u \geq 0$, see Schilling et al. [54] for a thorough exposition on Bernstein functions and subordinators. We denote throughout by \mathbf{B}_J the subset of Bernstein functions of the form (3.4) which satisfies the condition $\phi(0) > 0$.

We also observe that \mathbb{J}_Y (resp. ϕ) is uniquely determined by σ^2, Π, μ and λ (resp. σ^2, Π, μ) so that, for a fixed λ , there is a one-to-one correspondence between ϕ and \mathbb{J}_Y .

Next, we set, $W_\phi(1) = 1$ and for any $n \geq 1$,

$$(3.6) \quad W_\phi(n+1) = \prod_{k=1}^n \phi(k)$$

Note that W_ϕ is solution to the recurrence equation $W_\phi(n+1) = \phi(n)W_\phi(n)$, with $W_\phi(0) = 1$, and we refer to Patie and Savov [48] for a thorough account on this set of functions that generalizes the gamma function, which appears as a special case when $\phi(n) = n$. Then, it is shown, in [11, Theorem 2.1], that there exists an absolutely continuous probability measure whose support is $[0, 1]$, with a continuous density denoted by β that is positive on $(0, 1)$. Being of compact support, its law is moment determinate, and, more specifically, one has, for any $n \in \mathbb{N}$,

$$(3.7) \quad \int_0^1 y^n \beta(y) dy = \frac{W_\phi(n+1)\Gamma(\frac{2\lambda}{\sigma^2})}{\Gamma(\frac{2\lambda}{\sigma^2} + n)}.$$

Note that, in particular, using (3.4), one gets the following expression for the first moment of β

$$(3.8) \quad \int_0^1 y\beta(y)dy = \sigma^2 \frac{\phi(1)}{2\lambda} = \sigma^2 \frac{(\frac{2}{\sigma^2}\mu - \frac{2}{\sigma^2}\hbar) + \frac{2}{\sigma^2} \int_0^\infty (1 - e^{-r})\bar{\Pi}(r)dr}{2\lambda} = \frac{\mu - \int_0^\infty e^{-r}\bar{\Pi}(r)dr}{\lambda}.$$

We also point out that when $\phi(u) = u$, β boils down to the Beta distribution, which is easily identified from the expression of its moments as in this case $W_\phi(n+1) = n!$. Other examples will be provided in Section 4.1. β turns out to be the stationary measure of the Feller semigroup \mathbb{Q} , that is for all $f \in C([0, 1])$, the Banach space of continuous functions on $[0, 1]$ equipped with the sup-norm $\|\cdot\|_\infty$, and $t \geq 0$,

$$(3.9) \quad \beta[\mathbb{Q}_t f] = \beta[f] = \int_0^1 f(y)\beta(dy)$$

where the last equality serves as a definition for the notation $\beta[f]$. The extension of \mathbb{J}_Y to an operator on $L^2(\beta)$, still denoted by \mathbb{J}_Y , is the infinitesimal generator, having \mathcal{P} , the algebra of polynomials, as a core, of an ergodic Markov semigroup $\mathbb{Q} = (\mathbb{Q}_t)_{t \geq 0}$ on $L^2(\beta)$ whose unique invariant measure is β .

It is then classical, see either Bakry et al. [7] or Da Prato [14], that given a Markov semigroup on $C([0, 1])$ with invariant probability measure β one may extend it to a Markov semigroup on $L^2(\beta)$, the weighted Hilbert space being defined as

$$L^2(\beta) = \{f : [0, 1] \rightarrow \mathbb{R} \text{ measurable with } \beta[f^2] < \infty\}.$$

Such a semigroup is said to be ergodic if, for every $f \in L^2(\beta)$, $\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \mathbb{Q}_t f dt = \beta[f]$ in the $L^2(\beta)$ -norm.

Proposition 3.1. *Let $X = (X_t)_{t \geq 0}$ where, for any $t \geq 0$, $X_t = g(Y_t)$ with $g(x) = (V_E - V_I)x + V_I$. Then X is a Feller process on $E_V = [V_I, V_E]$ which admits the measure $(V_E - V_I)^{-1} \beta\left(\frac{x - V_I}{V_E - V_I}\right) dx$ as the unique stationary measure. Its infinitesimal generator is the closure of $(\mathbb{J}_X, \mathcal{P})$, where \mathbb{J}_X is defined in (3.1) and \mathcal{P} is a core. Moreover, we have, on \mathcal{P} ,*

$$(3.10) \quad \mathbb{J}_X Gf = G\mathbb{J}_Y f$$

where $Gf(x) = f \circ g(x)$ is a homeomorphism from $[0, 1]$ onto $[V_I, V_E]$.

Proof. Since g is a homeomorphism from $[0, 1]$ onto $[V_I, V_E]$ with inverse function $h(x) = \frac{x - V_I}{V_E - V_I}$ and, from [11, Lemma 3.10 and its proof], Y is a Feller process on $[0, 1]$, we deduce that X is also

a Feller process on E_V . Next, using (2.4) and the notation (3.2), simple algebra yields, for any $f \in \mathcal{P}$,

$$\begin{aligned}
G^{-1}\mathbb{J}_X Gf(y) &= \sigma^2(V_E - V_I)^2 y(1-y)f''(y) \frac{1}{(V_E - V_I)^2} \\
&- \left(\left(\frac{1}{\tau} + \mu_e - \mu_i \right) (V_E - V_I)y - \mu_e(V_E - V_I) + \frac{V_I}{\tau} \right) f'(y) \frac{1}{(V_E - V_I)} \\
&+ \int_0^\infty (f(e^{-r}y) - f(y)) \frac{\Pi(dr)}{y} \\
&= \sigma^2 y(1-y)f''(y) - (\lambda y - \mu) f'(y) + \int_0^\infty (f(e^{-r}y) - f(y)) \frac{\Pi(dr)}{y} \\
&= \mathbb{J}_Y f(y)
\end{aligned}$$

which completes the proof of the intertwining relation. Since \mathcal{P} is core for \mathbb{J}_Y , see [11, Theorem 2.1] we deduce, by the homeomorphism G , that \mathcal{P} is also a core for \mathbb{J}_X . Next, by taking the inverse of G , from the left and from the right, in the relation (3.10), one gets that $G^{-1}\mathbb{J}_X f = \mathbb{J}_Y G^{-1}f$. Since from [11], we have that β is the unique stationary measure for Y , that is the unique measure β such that $\beta\mathbb{J}_Y f = 0$, $f \in \mathcal{D}_Y$, we deduce that the measure on $[V_I, V_E]$, defined by $\beta_G = \beta G^{-1}$, is the unique one such that $\beta_G \mathbb{J}_X Gf = 0$, which completes the proof. \blacksquare

Remark 3.1. We point out that the requirement that the support of the measure of the Lévy kernel is the negative half-line (or, equivalently, the process has only negative jumps) comes from Proposition 3.1. Indeed, it ensures that the corresponding integro-differential operator satisfies the maximum principle, and, hence it is the generator of a Markov semigroup. The interested reader can consult [11] for a detailed discussion on this technical aspect.

3.2. Laplace transform of first passage times. Let us write

$$(3.11) \quad T_a = \inf\{t > 0; Y_t \geq a\}$$

for the first passage time to the level $0 < a < 1$ of the generalized Jacobi process Y . Note that, from Proposition 3.1, one gets, when Y is issued from $y \in (0, 1)$, the identity in distribution, with the obvious notation, $T_{y \rightarrow a}(Y) \stackrel{d}{=} T_{g(y) \rightarrow g(a)}(X)$. To characterize the Laplace transform of T_a , we introduce the mapping

$$(3.12) \quad {}_2F_1(a, b, \phi; y) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{n!} \frac{y^n}{W_\phi(n+1)}$$

with $(a)_n = \frac{\Gamma(a+n)}{\Gamma(a)}$, $n \in \mathbb{N}$, $a \in \mathbb{C}$. Note that when $\Pi \equiv 0$, we have, from (3.4), $W_\phi(n+1) = (\phi(0) + 1)_n$, and, thus, in this case

$${}_2F_1(a, b; \phi; y) = {}_2F_1(a, b; \phi(0) + 1; y)$$

which is the Gauss hypergeometric function, explaining the notation. There are several representations of this function which provides an analytical continuation to the entire complex plane cut along $[1, \infty]$ with $\lim_{x \uparrow 1} {}_2F_1(a, b, 1; z) = \frac{\Gamma(1-a-b)}{\Gamma(1-a)\Gamma(1-b)}$, $\Re(a+b) < 1$, see [39, Chap. 9]. We are now ready to state the following.

Theorem 3.1. *Let $\phi \in \mathbf{B}_J$. Then, for any $a, b \in \mathbb{C}$, the mapping $z \mapsto {}_2F_1(a, b; \phi; z)$ defines an analytic function on the unit disc. Moreover, for any $0 < y < a < 1$ and $q > 0$, we have*

$$(3.13) \quad \mathbb{E}_y [e^{-qT_a}] = \frac{{}_2F_1(\kappa(q), \theta(q); \phi; y)}{{}_2F_1(\kappa(q), \theta(q); \phi; a)},$$

where $\kappa(q)$ and $\theta(q)$ are solution to the system

$$(3.14) \quad \kappa(q)\theta(q) = \frac{2q}{\sigma^2} \text{ and } \kappa(q) + \theta(q) + 1 = \frac{2\lambda}{\sigma^2}.$$

3.3. Proof of Theorem 3.1. The proof is split into several intermediate results. We start with the following result that shows that the dynamics of Y has discontinuities which are due to negative jumps only.

Lemma 3.1. *We have, for all $y \in [0, 1], t \geq 0$ and \tilde{f} a positive borelian function on $[0, 1] \times [0, 1]$,*

$$\mathbb{E}_y \left[\sum_{s \leq t} \tilde{f}(Y_{s-}, Y_s) \mathbb{I}_{\{Y_{s-} \neq Y_s\}} \right] = \mathbb{E}_y \left[\int_0^t ds \int_0^1 \tilde{f}(Y_{s-}, r) \mathbb{I}_{\{r \leq Y_{s-}\}} \tilde{\Pi}(Y_{s-}, dr) \right]$$

where $\tilde{\Pi}(y, \cdot)$ is the image measure of $\frac{\Pi(\cdot)}{y}$ by the mapping $r \mapsto -\ln(r/y)$. Consequently, for all $y \in [0, 1], \mathbb{P}_y(Y_{t-} \geq Y_t \text{ for all } t \geq 0) = 1$, i.e. Y has only downward jumps.

Proof. First, by [11, Lemma 3.1], we know that Y is a Feller process, and hence, from [8], we have that Y admits a Lévy kernel, say N , that we now characterize. To this end, one observes, from (3.1), that for a smooth function f that vanishes in the neighborhood of $y \in [0, 1]$, we have

$$(3.15) \quad \mathbf{J}f(y) = \int_0^\infty f(e^{-s}y) \frac{\Pi(ds)}{y} = \int_0^1 f(s) \mathbb{I}_{\{r \leq y\}} \tilde{\Pi}(y, dr)$$

where $\tilde{\Pi}(y, \cdot)$ is the measure defined in the claim. Hence, the Lévy kernel $N(y, dr) = \mathbb{I}_{\{r \leq y\}} \tilde{\Pi}(y, dr)$, see e.g. [43]. The first claim follows from the definition of the Lévy kernel whereas the second one is deduced from the first one by choosing the function $\tilde{f}(y, r) = \mathbb{I}_{\{y \leq r\}}$. ■

We proceed with the following.

Lemma 3.2. *Let us write $F_q(y) = {}_2F_1(\kappa(q), \theta(q); 1; y)$, $q > 0$, then F_q is positive increasing on $(0, 1)$ and we have*

$$\mathbf{J}F_q(y) = qF_q(y), \quad y \in [0, 1],$$

where, simplifying the notation, $\mathbf{J} = \mathbf{J}_{\frac{\sigma^2}{2}}$, that is $\mathbf{J}f(y) = \frac{\sigma^2}{2}y(1-y)f''(y) - \left(\lambda y - \frac{\sigma^2}{2}\right)f'(y)$.

Consequently, for any $t, q \geq 0$ and $y \in [0, 1]$,

$$(3.16) \quad e^{-qt} \mathbf{Q}_t F_q(y) = F_q(y)$$

that is, F_q is a q -invariant function for $\mathbf{Q} = (\mathbf{Q}_t)_{t \geq 0}$ the semigroup associated to \mathbf{J} .

Proof. The first part is classical, see Appendix A. Note that $\kappa(q) + \theta(q) = 1 - 2\lambda/\sigma^2 < 1$ which ensures that $\lim_{y \uparrow 1} {}_2F_1(\kappa(q), \theta(q), 1; y)$ exists. Next, using the fact that, in addition, the mapping $x \mapsto F_q(y)$ is twice continuously differentiable on $[0, 1]$, one can apply Itô's formula to get

$$\begin{aligned} e^{-qt} F_q(Y_t) &= F_q(y) + \int_0^t \mathbf{J}F_q(Y_s) - qF_q(Y_s) ds + \sqrt{2}\sigma \int_0^t \sqrt{Y_s(1-Y_s)} F'_q(Y_s) dB_s \\ &= F_q(y) + \sqrt{2}\sigma \int_0^t \sqrt{Y_s(1-Y_s)} F'_q(Y_s) dB_s. \end{aligned}$$

Since the last term has a squared integrable integrant, it defines a martingale. Then, taking the expectation on both sides of the previous identity yields the second claim. ■

Let us now denote by $\varrho = (\varrho_t)_{t \geq 0}$ a subordinator that is a positive valued stochastic process with stationary and independent increments, and recall that its law is uniquely determined by a Bernstein function ϕ . More specifically, one has, for any $t, u \geq 0$,

$$(3.17) \quad \mathbb{E}[e^{-u\varrho t}] = e^{-\phi(u)t}.$$

Next, for each subordinator ϱ associated to $\phi \in \mathbf{B}_J$, we define the random variable

$$I_\phi = \int_0^\infty e^{-\varrho t} dt$$

which is the so-called exponential functional of the subordinator ϱ . We point out that this random variable has been studied intensively over the last two decades see e.g. [48] and the references therein.

Lemma 3.3. *Let $\phi \in \mathbf{B}_J$ and write $F_{a,b}(y) = {}_2F_1(a, b; 1; y)$. Then, we have, for any $y \in [0, 1]$,*

$$(3.18) \quad \Lambda_\phi F_{a,b}(y) = {}_2F_1(a, b; \phi; y)$$

where $\Lambda_\phi : C([0, 1]) \mapsto C([0, 1])$ is the Markov multiplicative operator associated to the random variable I_ϕ , that is

$$(3.19) \quad \Lambda_\phi f(y) = \mathbb{E}[f(yI_\phi)].$$

Moreover, $z \mapsto {}_2F_1(a, b; \phi; z)$ defines a function which is analytic on the unit disc.

Proof. First, we recall, from e.g. [11, Lemma 3.3], that Λ_ϕ is a Markov bounded operator from $C([0, 1])$ into itself, and, with $p_n(y) = y^n$, $n \in \mathbb{N}$,

$$(3.20) \quad \Lambda_\phi p_n(y) = \frac{n!}{W_\phi(n+1)} p_n(y).$$

Then, an application of Tonelli Theorem and (3.20) yield, for any $0 \leq y \leq 1$,

$$\Lambda_\phi F_{a,b}(y) = \mathbb{E}[F_{a,b}(yI_\phi)] = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{n!} \frac{\Lambda_\phi p_n(y)}{n!} = {}_2F_1(a, b; \phi; y).$$

Moreover, as $W_\phi(n+2) = \phi(n+1)W_\phi(n+1)$ and $\lim_{n \rightarrow \infty} \frac{\phi(n+1)}{n+1} = 1$, we easily get that the power series ${}_2F_1(a, b; \phi; \cdot)$ defines an analytic function on the unit disc. \blacksquare

Lemma 3.4. *Writing $F_q^{(\phi)}(y) = {}_2F_1(\kappa(q), \theta(q); \phi; y)$, we have, for any $t, q \geq 0$,*

$$e^{-qt} \mathbf{Q}_t F_q^{(\phi)}(y) = F_q^{(\phi)}(y)$$

that is $F_q^{(\phi)}$ is a q -invariant function for \mathbf{Q} , which is positive and increasing on $[0, 1]$.

Proof. First, let us denote by $(\tilde{\mathbf{Q}}_t)_{t \geq 0}$ the semigroup associated to the non-local Jacobi generator given, for any $\sigma^2 > 0$ and $y \in (0, 1)$, by

$$(3.21) \quad \tilde{\mathbf{J}}^Y f(y) = y(1-y)f''(y) - \left(\frac{\lambda}{\sigma^2} y - \frac{\mu}{\sigma^2} \right) f'(y) + \int_0^\infty (f(e^{-r}y) - f(y)) \frac{\Pi(dr)}{y\sigma^2}.$$

Then, observes that, for any $c > 0$,

$$(3.22) \quad \mathbf{J}^Y f(y) = \lim_{t \rightarrow 0} \frac{\mathbf{Q}_t f(y) - f(y)}{t} = c \lim_{t \rightarrow 0} \frac{\tilde{\mathbf{Q}}_{ct} f(y) - f(y)}{ct} = c \tilde{\mathbf{J}}^Y f(y)$$

and note that the same relationship holds between $(\mathbf{Q}_t)_{t \geq 0}$ and $(\tilde{\mathbf{Q}}_t)_{t \geq 0}$ the semigroups of the classical Jacobi processes with generator $\mathbf{J}f(y) = \frac{\sigma^2}{2} y(1-y)f''(y) - \left(\lambda y - \frac{\sigma^2}{2} \right) f'(y)$ and $\tilde{\mathbf{J}}f(y) = y(1-y)f''(y) - (2\lambda\sigma^{-2}y - 1)f'(y)$ respectively. Now, we recall from [11, Proposition 3.3], taking with the notation thereout $\epsilon = d_\phi$ and $r_1 = 1$, that the following intertwining relation

$$(3.23) \quad \tilde{\mathbf{Q}}_t \Lambda_\phi = \Lambda_\phi \tilde{\mathbf{Q}}_t$$

holds on the weighted Hilbert space $L^2(\beta_{\lambda_1})$, where $\lambda_1 = 2\lambda\sigma^{-2} > 1$, by the assumption (3.3), and, $\beta_{\lambda_1}(dy) = (\lambda_1 - 1)(1-y)^{\lambda_1-2} dy$, $y \in (0, 1)$. Hence, for any $t \geq 0$, on $L^2(\beta_{\lambda_1})$,

$$(3.24) \quad \mathbf{Q}_t \Lambda_\phi = \tilde{\mathbf{Q}}_{\sigma^2/2t} \Lambda_\phi = \Lambda_\phi \tilde{\mathbf{Q}}_{\sigma^2/2t} = \Lambda_\phi \mathbf{Q}_t.$$

Thus, using successively that $F_q \in L^2(\beta_{\lambda_1})$, (3.18), (3.24) and (3.16), one gets that

$$e^{-qt} \mathbf{Q}_t F_q^{(\phi)}(y) = e^{-qt} \mathbf{Q}_t^\phi \Lambda_\phi F_q(y) = e^{-qt} \Lambda_\phi \mathbf{Q}_t F_q(y) = \Lambda_\phi F_q(y) = F_q^{(\phi)}(y)$$

which proves the first claim. Next, since Λ_ϕ is clearly a Markov operator, i.e. $\Lambda_\phi f \geq 0$ for any $f \geq 0$ and $\Lambda_\phi p_0(y) = 1$, we get that ${}_2F_1(a, b; \phi; \cdot) \geq 0$ on $[0, 1]$. Finally, $F_q^{(\phi)}$ being a power series with non-negative coefficients, we deduce the monotonicity property. \blacksquare

End of the proof of Theorem 3.1. First, one invokes the previous lemma and Dynkin's theorem to the bounded stopping time $T_a^t = T_a \wedge t$, to get, for any $t, q > 0$ and $0 < y < a < 1$,

$$\mathbb{E}_y \left[e^{-qT_a^t} F_q^{(\phi)}(Y_{T_a^t}) \right] = F_q^{(\phi)}(y).$$

Then, letting $t \rightarrow \infty$, using the fact that $F_q^{(\phi)}$ is increasing on $[0, 1]$ and by absence of positive jumps, see Lemma (3.1), $\mathbb{P}_y(X_{T_a} = a) = 1$, combined with a dominated convergence argument yield

$$\mathbb{E}_y \left[e^{-qT_a} \mathbb{I}_{\{T_a < \infty\}} \right] = \frac{F_q^{(\phi)}(y)}{F_q^{(\phi)}(a)}.$$

Next, observe that, if $\bar{\lambda} = \frac{\lambda}{\sigma^2} - \frac{1}{2} \geq 0$ (resp. < 0) then, by Taylor's expansion, one gets that $\lim_{q \rightarrow 0} \sigma^2 \bar{\lambda} \frac{\theta(q)}{q} = 1$ (resp. $\lim_{q \rightarrow 0} \theta(q) = 2\bar{\lambda}$), and thus $\lim_{q \rightarrow 0} \kappa(q) = 2\bar{\lambda}$ (resp. $= 0$). It is not difficult to check that in both cases, one has for all $y \in [0, 1]$, $\lim_{q \rightarrow 0} F_q^{(\phi)}(y) = 1$ and hence $\mathbb{P}_y(T_a < \infty) = 1$, which completes the proof of Theorem 3.1.

3.4. Mean of the first passage times. We proceed by deriving the expression of the first moment of the first passage times of our family of Markov processes whose proof is split into several intermediate results.

Theorem 3.2. *Let $\phi \in \mathbf{B}_J$. Then, for any $0 < y < a < 1$,*

$$(3.25) \quad \mathbb{E}_y[T_a] = \frac{2}{\sigma^2} \sum_{n=0}^{\infty} \frac{(2\lambda/\sigma^2)_n a^{n+1} - y^{n+1}}{n+1} \frac{1}{W_\phi(n+2)}.$$

Remark 3.2. We note, from (3.25), that when $\Pi = 0$, we recover the expression of $\mathbb{E}_y[T_a]$ for the classical Jacobi process given in (A.5). Indeed, in this case,

$$W_\phi(n+2) = \prod_{k=1}^{n+1} \phi(k) = \left(\frac{2\mu}{\sigma^2} \right)_{n+1} = \frac{2\mu}{\sigma^2} \left(\frac{2\mu}{\sigma^2} + 1 \right)_n.$$

As a by-product of this Theorem, we state and prove the following comparison result between the first moment of the first passage times of our class of Jacobi processes with jumps.

Corollary 3.1. *Let $\phi, \phi_1 \in \mathbf{B}_J$ be such that $\phi \leq \phi_1$ on \mathbb{R}_+ . Then, we have, for any $0 < y < a < 1$, with the obvious notation,*

$$(3.26) \quad \mathbb{E}_y[T_a^{\phi_1}] \leq \mathbb{E}_y[T_a^\phi]$$

The conditions hold, for instance, when $\phi, \phi_1 \in \mathbf{B}_J$ and $\bar{\Pi} \geq \bar{\Pi}_1$ on \mathbb{R}_+ , or when $\phi \in \mathbf{B}_J$ and $\phi_1(u) = \phi(u) + \frac{2}{\sigma^2} \int_0^\infty (1 - e^{-ur}) \bar{\Pi}_1(r) dr, u \geq 0$, where $\bar{\Pi}_1$ satisfies the same conditions than $\bar{\Pi}$ in (3.1). Note that, with the notation of (3.5), in the former instance, we have $\mu = \frac{\sigma^2}{2}(\phi(0) + 1) + \hbar = \frac{\sigma^2}{2}(\phi_1(0) + 1) + \hbar_1 = \mu_1$ whereas, in the latter case, $\mu_1 = \mu + \hbar_1$.

Remark 3.3. It is interesting to note that, in the second example, i.e. when $\mu_1 = \mu + \hbar_1$, although the intensity of jumps is larger for the Jacobi process associated to ϕ_1 , the first moment of its first passage times above the starting point is smaller, meaning that adding the mean of the jump measure to the drift term compensates the presence of additional downwards jumps.

Proof. We first observe, that with $\phi, \phi_1 \in \mathbf{B}_J$ such that $\phi \leq \phi_1$ on \mathbb{R}_+ , we have, for all $n \geq 0$, $W_\phi(n+1) \leq W_{\phi_1}(n+1)$, the inequality between the first moments of the first passage times is deduced easily from the identity (3.25) as the other parameters in the two expressions are identical. Finally, on the one hand, since $\hbar = \int_0^\infty \bar{\Pi}(r)dr$ we get, from (3.4), that

$$(3.27) \quad \phi(u) = u + \frac{2}{\sigma^2} \left(\mu - \frac{\sigma^2}{2} - \int_0^\infty e^{-ur} \bar{\Pi}(r) dr \right)$$

which provides the first inequality between the Bernstein functions. On the other hand, writing $\hbar_1 = \int_0^\infty \bar{\Pi}_1(r)dr$, we get that

$$\phi_1(u) = u + \frac{2}{\sigma^2} \left(\mu + \hbar_1 - (\hbar + \hbar_1) - \frac{\sigma^2}{2} + \int_0^\infty (1 - e^{-ur})(\bar{\Pi} + \bar{\Pi}_1)(r) dr \right) \in \mathbf{B}_J$$

Then, plainly $\phi \leq \phi_1$ on \mathbb{R}_+ , and the last remarks follow readily. \blacksquare

We now turn to the proof of Theorem 3.2 which relies on taking the derivative of the Laplace transform (3.13) which is given in the following lemma. This extends the result of [2] on the derivative of the Gauss hypergeometric function ${}_2F_1$.

Lemma 3.5. *Let $\phi \in \mathbf{B}_J$. Then, for any $|z| < 1$,*

$$(3.28) \quad \frac{\partial}{\partial a} {}_2F_1(a, b, \phi, z)|_{a=0} = b \sum_{n=0}^{\infty} \frac{(b+1)_n}{n+1} \frac{z^{n+1}}{W_\phi(n+2)}$$

$$(3.29) \quad \frac{\partial}{\partial b} {}_2F_1(a, b, \phi, z)|_{a=0} = 0.$$

Proof. Using that $\frac{\partial}{\partial a} (a)_n = (a)_n[\Psi(a+n) - \Psi(a)]$, where Ψ is the Digamma function, we get that

$$(3.30) \quad \frac{\partial}{\partial a} {}_2F_1(a, b, \phi, z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{n!} (\Psi(a+n) - \Psi(a)) \frac{z^n}{W_\phi(n+1)}.$$

From [1, Formulas 6.3.5 and 6.3.6], one has

$$(3.31) \quad \Psi(a+n) - \Psi(a) = \sum_{k=0}^{n-1} \frac{1}{k+a}.$$

Moreover, observing that

$$(3.32) \quad \frac{1}{k+a} = \frac{1}{a} \frac{(a)_k}{(a+1)_k}$$

one gets

$$(3.33) \quad \begin{aligned} \frac{\partial}{\partial a} {}_2F_1(a, b, \phi, z) &= \frac{1}{a} \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{(a)_{n+1} (a)_k (b)_{n+1}}{(a+1)_k (n+1)!} \frac{z^{n+1}}{W_\phi(n+2)} \\ &= \frac{1}{a} \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{(a)_{n+k+1} (a)_k (b)_{n+k+1}}{(a+1)_k (n+k+1)!} \frac{z^{n+k+1}}{W_\phi(n+k+2)} \\ &= bz \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{(a+1)_{n+k} (a)_k (b+1)_{n+k}}{(a+1)_k (n+k+1)!} \frac{z^{n+k}}{W_\phi(n+k+2)} \end{aligned}$$

where in the last equality we have used that $(a)_{n+k+1} = a(a+1)_{n+k}$. From (3.33) with $a = 0$, noting that we have non-zero terms only for $k = 0$, the identity (3.28) follows. The expression of $\frac{\partial}{\partial b} {}_2F_1(a, b, \phi, z)$ can be obtained directly by interchanging a with b in (3.33), i.e

$$(3.34) \quad \frac{\partial}{\partial b} {}_2F_1(a, b, \phi, z) = az \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{(b+1)_{n+k} (b)_k (a+1)_{n+k}}{(b+1)_k (n+k+1)!} \frac{z^{n+k}}{W_\phi(n+k+2)}$$

that is always equal to zero for $a = 0$. ■

End of the proof of Theorem 3.2. For $\phi \in \mathbf{B}_J$, that is $\lambda > \mu > \hbar + \frac{\sigma^2}{2}$, we have, writing $\Phi_y(q) := \mathbb{E}_y[e^{-qT_a} \mathbb{1}_{\{T_a < \infty\}}]$,

$$(3.35) \quad \begin{aligned} \mathbb{E}_y[T_a] &= -\frac{\partial \Phi_y(q)}{\partial q} \Big|_{q=0} = -\Phi_y(0) \frac{\partial \ln \Phi_y(q)}{\partial q} \Big|_{q=0} = -\frac{\partial \ln \Phi_y(q)}{\partial q} \Big|_{q=0} \\ &= \frac{\partial}{\partial q} ({}_2F_1(\kappa(q), \theta(q); \phi; a) - {}_2F_1(\kappa(q), \theta(q); \phi; y)) \Big|_{q=0}. \end{aligned}$$

We have

$$(3.36) \quad \begin{aligned} \frac{\partial}{\partial q} {}_2F_1(\kappa(q), \theta(q); \phi; y) \Big|_{q=0} &= \frac{\partial}{\partial \kappa(q)} {}_2F_1(\kappa(q), \theta(q); \phi; y) \Big|_{q=0} \frac{\partial \kappa(q)}{\partial q} \Big|_{q=0} \\ &+ \frac{\partial}{\partial \theta(q)} {}_2F_1(\kappa(q), \theta(q); \phi; y) \Big|_{q=0} \frac{\partial \theta(q)}{\partial q} \Big|_{q=0}. \end{aligned}$$

and, with $\bar{\lambda} := \frac{\lambda}{\sigma^2} - \frac{1}{2} \geq 0$, it is easy to check, from the system (3.14), that

$$(3.37) \quad \frac{\partial \kappa(q)}{\partial q} \Big|_{q=0} = \frac{1}{\bar{\lambda} \sigma^2}, \quad \kappa(0) = 0, \quad \theta(0) = 2\bar{\lambda}.$$

Moreover, one gets, by Lemma 3.5, that

$$(3.38) \quad \begin{aligned} \frac{\partial}{\partial \kappa(q)} {}_2F_1(\kappa(q), \theta(q); \phi; y) \Big|_{q=0} &= \frac{\partial}{\partial \kappa(q)} {}_2F_1(\kappa(q), \theta(q); \phi; y) \Big|_{\kappa(q)=0} \\ &= 2\bar{\lambda} \sum_{n=0}^{\infty} \frac{(1)_n (2\bar{\lambda} + 1)_n}{(2)_n} \frac{y^{n+1}}{W_\phi(n+2)}, \end{aligned}$$

and

$$(3.39) \quad \frac{\partial}{\partial \theta(q)} {}_2F_1(\kappa(q), \theta(q); \phi; y) \Big|_{\kappa(q)=0} = 0.$$

Finally, combining (3.35) and (3.36)-(3.39), we obtain the expression of $\mathbb{E}_y[T_a]$, which completes the proof of the Theorem.

4. FIRING ACTIVITY OF THE JACOBI PROCESS WITH JUMPS

Let X be the Jacobi process with jumps with state space $E_V = [V_I, V_E]$ defined in (2.4). As mentioned above, according to the model, the spikes are generated when the process X crosses a voltage threshold $V_I < S < V_E$ for the first time, that is at time $T_S = T_S(X)$. After the spike, the process is reset instantaneously to the starting position $V_I < x < V_E$, ready to start its evolution over again. This renewal condition guarantees that the inter-spike intervals, i.e the time between two consecutive spikes, are independent and all identically distributed as the first inter-spike interval T_S . Proposition 3.1 guarantees that we can consider equivalently the Jacobi process with jumps Y with state space $[0, 1]$ starting at $y = h(x) = (x - V_I)/(V_E - V_I)$ in the presence of the threshold $a = h(S) = (S - V_I)/(V_E - V_I)$ and other parameters defined in (3.2). This enables us to use the mathematical results obtained in Section 3. For these reasons the quantity of interest in the

mathematical analysis of the neuronal activity is the first passage time $T_a = T_a(Y)$. The probability of firing (i.e. the probability that the process Y crosses the boundary a within a finite time) is given by (3.13) for $q = 0$. Under hypothesis (3.3), the crossing of a occurs almost surely in finite time.

Furthermore, it is interesting, for the analysis of the firing activity, to study the first moment of T_a . In fact, it is assumed that neurons express information about their input mainly by means of the average frequency of spikes described by the neuronal firing rate. It can be mathematically defined in several different ways [35], here we choose the classical definition of the instantaneous firing rate as the reciprocal of the mean first passage time.

We distinguish between three possible regimes to characterize the neuronal activity. If the asymptotic mean membrane potential is larger than the firing threshold a , then the process is in the so-called suprathreshold regime. In the classical case, in this regime, the spikes are regular and the dynamics is driven mainly by the drift part. If the asymptotic mean membrane potential is smaller than a , then the process is said to be in the subthreshold regime, and the noise plays a prominent role for the crossing of the threshold. Finally, if the asymptotic mean is equal to a , the process is said to be in the threshold regime. The classical Jacobi process is in the suprathreshold regime for

$$\mu > a\lambda$$

whereas the analogous condition for the Jacobi with jumps, using (3.8), is

$$(4.1) \quad \mu > a\lambda + \int_0^\infty e^{-r} \bar{\Pi}(r) dr.$$

We observe that in (4.1) the asymptotic mean μ/λ of the classical Jacobi has to exceed the threshold plus a term given by the downward jumps.

The dependence of $\mathbb{E}_y[T_a]$ on a and y is the same as the classical Jacobi (and all other classical single neuron models). $\mathbb{E}_y[T_a]$ decreases with the difference $a - y$ as it can be easily seen from the expression (3.25).

However, the dependence of $\mathbb{E}_y[T_a]$ on the inputs parameters ν_e , ν_i , \hbar is non-trivial since the contribution of μ is hidden in the function W_ϕ that merges the contribution of the drift and the diffusion component. To investigate it, we consider the following examples in which we choose a special form of the measure Π .

4.1. Example. We consider a parametric family of non-local Jacobi operators for which $\bar{\Pi}(r) = \int_r^\infty \Pi(du) = e^{-\alpha r}$, $r > 0$, is of exponential type, that is $\Pi(dr) = \alpha e^{-\alpha r} dr$. In particular, let $\alpha \geq 1$ and consider the integro-differential operator \mathbb{J}_α given, from (3.1), by

$$(4.2) \quad \mathbb{J}_\alpha f(y) = \frac{\sigma^2}{2} y(1-y) f''(y) - (\lambda y - \mu) f'(y) - \int_0^1 (f(r) - f(y)) \frac{r^\alpha}{y^{\alpha+1}} dr,$$

Then \mathbb{J}_α is a non-local Jacobi operator with $\hbar = \int_1^\infty \bar{\Pi}(r) dr = 1/\alpha$ and

$$(4.3) \quad \phi(u) = u + \frac{2}{\sigma^2} \left(\mu - \frac{1}{u + \alpha} \right) - 1.$$

Assumption (3.3) is satisfied whenever

$$(4.4) \quad \frac{\sigma^2}{2} < \mu - \frac{1}{\alpha},$$

suggesting that the noise amplitude has to be smaller than in the classical case. The more is the contribution of the downward jumps (smaller values of α) the higher is the risk that a large value of σ can lead the process across the lower boundary, a condition that we want to avoid. Under

assumption (4.4), the first moment of T_a for the Jacobi process with jumps with generator (4.2) is (see Appendix B)

$$(4.5) \quad \mathbb{E}_y[T_a] = \frac{2(\alpha + 1)}{\sigma^2(k_+ + 1)(k_- + 1)} \left({}_4F_3\left(1, 1, \alpha + 2, \frac{2\lambda}{\sigma^2}; 2, k_+ + 2, k_- + 2; a\right) a - {}_4F_3\left(1, 1, \alpha + 2, \frac{2\lambda}{\sigma^2}; 2, k_+ + 2, k_- + 2; y\right) y \right)$$

where


$$(4.6) \quad k_{\pm} = \frac{1}{2} \left(\alpha + 2\mu/\sigma^2 - 1 \pm \sqrt{(\alpha + 2\mu/\sigma^2 - 1)^2 - 4(2\alpha\mu/\sigma^2 - \alpha - 2/\sigma^2)} \right).$$

We want to investigate the sensitivity of the mean FPT to a change in the input parameters μ , λ , σ^2 and \hbar . The precise analysis requires the derivative of generalized hypergeometric functions with respect to the relevant parameters. To avoid lengthy calculation, we only show by plots the qualitative behavior using numerical evaluations in correspondence of physiologically realistic parameters chosen as in [37]. In this case the firing regime is suprathreshold if


$$(4.7) \quad \mu > a\lambda + \frac{1}{1 + \alpha},$$

we observe that the asymptotic mean μ/λ of the classical Jacobi is decreased by a term given by the downward jumps. The result is that the asymptotic mean of the Jacobi process with jumps increases with α . The reason lies in the shape of the distribution Π , see Fig.2-left. For small values of α there is a higher probability that r takes large values with corresponding large jumps. Conversely for large values of α the probability mass is concentrated around zero favoring small jumps. The consequence is shown in Fig.2-right: the mean FPT decreases as α increases. In Fig.2-right we also use a discretization scheme for simulating sample paths developed recently in [20]. At each time-step of the algorithm, a value for r is sampled from the distribution $\Pi(dr)$ and according to the survival probability of \mathcal{T} described in Section 2, a jump may occur. Then the trajectory moves according to the diffusion, from state e^{-rY_t} if there was a jump, otherwise from state Y_t . Between the jump epochs the dynamics of the constructed process are purely diffusive and are simulated using the Milstein's discretization method. The curve is obtained simulating $2 \cdot 10^4$ sample paths of the Jacobi process with jumps for each of the 100 values of α considered. For each simulation the FPT is recorded and for every value of α a value of the mean FPT is obtained. The simulation results are of course subject to numerical errors, mainly due to the choice of the time-discretization step (here $5 \cdot 10^{-4}$) and the relative low number of FPTs considered. On the one hand, this confirms the importance of an analytic result, and, on the other hand, supports the validity of formula (3.25). In fact, it is known that discretization schemes overestimate the mean FPT since undetected threshold crossings may occur inside each discretization interval (see for instance [24]). We stress, anyway, that one must be careful in the evaluation of Eq.(3.25) or Eq.(4.5) since the involved functions become soon very large or very small as n increases. To be sure of our evaluations, we applied different numerical approaches obtaining the same results as a guarantee of the correctness of our computations. In Fig.2-right, one also observes that the mean first passage time, as a function of α , is nonincreasing, which is an illustration of the comparison result provided in (3.26). Indeed, one uses the fact that, for any $0 < \alpha \leq \alpha_1$, the function $e^{-\alpha_1 r} = \bar{\Pi}_1(r) \leq \bar{\Pi}(r) = e^{-\alpha r}$, $r > 0$.

Let us now investigate how sensitive is $\mathbb{E}_y[T_a]$ to a change in the incoming input rates. As expected we find that the mean FPT decreases for stronger excitatory inputs and increases with the inhibitory inputs. This dependence is clearly visible in the color change in the heatmap in Fig.3 where the excitatory and inhibitory inputs are tuned simultaneously. The blue lines are the contour plots, i.e., the couples (ν_e, ν_i) that produce the same mean FPT. The values of ν_e are chosen to



(A) Function $\alpha e^{-\alpha r}$ for three values of α given in the legend.



(B) Mean FPT $\mathbb{E}_y[T_a]$ for the Jacobi process with jumps from (3.25) with $\Pi(dr) = \alpha e^{-\alpha r} dr$, as function of α . The series is truncated after 150 terms, but it has been checked numerically that the quick convergence of the series guarantees a correct evaluation of the whole sum. The other parameters are $V_I = -10\text{mV}$, $V_E = 100\text{mV}$, $S = 10 \text{ mV}$, $x = 0 \text{ mV}$, $\tau = 15 \text{ ms}$, $a = 0.18 \text{ mV}$, $y = 0.09 \text{ mV}$, $e = 0.5$, $i = -1$, $\nu_i = 1 \text{ ms}^{-1}$, $\nu_e = 2.8 \text{ ms}^{-1}$, $\sigma^2 = 0.5 \text{ ms}^{-1}$. In grey, $\mathbb{E}_y[T_a]$ obtained from simulations of $2 \cdot 10^4$ FPTs for each of the 100 values of α with time step $dt = 5 \cdot 10^{-4}$.

FIGURE 2

meet condition (4.4) or equivalently

$$(4.8) \quad \nu_e > \frac{1}{e} \left(\frac{V_I}{\tau(V_E - V_I)} + \frac{\sigma^2}{2} + \frac{1}{\alpha} \right).$$

The heatmaps are obtained from (3.25) with $\Pi(dr) = \alpha e^{-\alpha r} dr$ and $\alpha = 3$ (Fig.3 - left) and from (A.5) (Fig.3 - right). Alternatively one can evaluate (4.5) with the package `hypergeo` [29] for the software environment `R`.

We observe three main differences between the mean FPT of the two processes:

- in the non-local case, due to the presence of the term $1/\alpha$ in (4.8), we need a larger excitatory input rate to guarantee a finite FPT,
- for the same choices of parameters, the waiting time before the first spike in the classical case is shorter than in the non-local case,
- the shape of the contour plots changes.

Regarding the third item, in the classical case, if we increase the inhibitory input rate ν_i , then we have to increase linearly the excitatory input rate ν_e to get the same mean FPT. In the non-local case the jump part comes into play breaking this tight coupling.

Fig.4 plots the mean FPT of the Jacobi process with jumps with infinitesimal generator (4.2) as a function of σ^2 . As in the classical Jacobi model, $\mathbb{E}_y[T_a]$ decreases as σ^2 increases. This result is generally explained noting that an increase of variability facilitates the boundary crossing.

Since a closed form formula for the variance of T_a is not available, it is natural to look at the asymptotic variance of the process Y to study the role of σ^2 . From (3.7) and (3.8) we calculate the





FIGURE 3. Mean FPT, $\mathbb{E}_y[T_a]$, for the non-local (left) and classical (right) Jacobi processes as a function of the excitatory and inhibitory input rates ν_e and ν_i . The heatmaps are obtained from (3.25) with $\Pi(dr) = \alpha e^{-\alpha r} dr$, $\alpha = 3$ (left) and from (A.5) (right). The other parameters are chosen as in Fig.2.



(A) $\mathbb{E}_y[T_a]$ as a function of σ^2 for the Jacobi process with jumps with infinitesimal generator (4.2) for different values of the ratio λ/μ given in the legend, using (3.25). In the plot $\alpha = 3$, $\tau = 15$ ms, $a = 0.18$ mV, $y = 0.09$ mV, $\nu_e = 2.1$ ms $^{-1}$.

(B) We consider the effect of α in the case $\lambda \sim \mu$. All curves are plotted as function of σ^2 to meet assumption (4.4), and this is the reason why some lines stop before others.

FIGURE 4

asymptotic variance of Y , $\text{Var}(Y_\infty)$, as

$$\begin{aligned} \text{Var}(Y_\infty) &= \beta[p_2] - \beta[p_1]^2 \\ &= \frac{\left(\mu - \frac{1}{1+\alpha}\right) \left(\sigma^4 + \sigma^2 \left(\mu - \frac{1}{2+\alpha}\right)\right) - \mu^2 - \frac{1}{(1+\alpha)^2} + \frac{2\mu\lambda^2}{1+\alpha} - \frac{\mu^2\sigma^2}{\lambda} - \frac{\sigma^2}{\lambda(1+\alpha)^2} + \frac{2\lambda\mu\sigma^2}{1+\alpha}}{\lambda^2 + \lambda\sigma^2} \end{aligned}$$

and one can get that the derivative with respect to σ^2 is positive. Then $\text{Var}(Y_\infty)$ increases with σ^2 and the variability usually favors the crossing of the threshold, explaining the result of Fig.4.

As a final remark, we look at the blue solid curve in Fig.4 (A). All the curves are obtained keeping fixed $\nu_e = 2.1$ ms $^{-1}$ and changing ν_i (0.1; 1.5; 1.9 ms $^{-1}$) to get different ratios λ/μ , as it is usually done in the classical case. The blue solid curve is obtained in the case of a very weak inhibition $\nu_i = 0.1$ ms $^{-1}$, that is the reason why the mean FPT is smaller and the behavior is different from the other two cases. We observe in Fig.4 (B) that the presence of the jump part compensates the absence of the inhibitory inputs, increasing the waiting time before the neuronal spike.

In Fig.5, we compare the firing rate, that is here the reciprocal of $\mathbb{E}_y[T_a]$, for the classical and the non-local Jacobi processes for the same choices of the common parameters. In the classical case a strong excitation rate ν_e produces an intense activity of the neuron that grows linearly with ν_e . In the non-local case the value of the firing rate is almost halved and shows a sub-linear growth with respect to ν_e .

In Fig.5 the vertical lines indicate the threshold regimes for the two dynamics, separating the subthreshold on the left from the suprathreshold on the right. We have used three colors to highlight the intervals of sub and suprathreshold for the two processes. We observe that the difference between the two firing rates is smaller in the subthreshold regime, whereas the gap increases in the suprathreshold regime where the dynamics of the classical Jacobi is mainly driven by the drift component, especially being ν_i much smaller than ν_e . Moreover, numerical evidences suggest that the firing rate for the Jacobi process with jumps saturates, differently from the classical one (at least for this range of parameters). A similar kind of saturation is observed in the classical case, but only in the presence of a non-zero refractory period, see for instance Fig.2 of [41].




FIGURE 5. Firing rate $1/\mathbb{E}_y[T_a]$ for classical and non-local Jacobi processes as function of the excitatory input rate ν_e . Curves are obtained from (3.25) and (A.5) for $y = 0.09$ mV, $a = (S - V_I)/(V_E - V_I) = 0.75$ mV, $\alpha = 3$, $\nu_i = 0.2$ ms $^{-1}$, $\tau = 5$ ms, $e = 0.2$, $i = -0.2$, $\mu_e = e\nu_e$, $\mu_i = i\nu_i$, $\sigma^2 = 0.1$ ms $^{-1}$. The firing rate is reduced by the downward jumps. The vertical lines indicate the threshold regimes for the two dynamics, separating the subthreshold and the suprathreshold regimes. The blue region corresponds to subthreshold regime for both processes, the red one corresponds to subthreshold for the non-local and suprathreshold for the classical Jacobi process and finally the yellow area represents suprathreshold regime for both processes.

Then, in the case of a strong excitatory input, the presence of the jump part can contribute to reduce the firing rate. We stress that we choose incoming input parameters that are up to 10 times stronger than those of an healthy neuron, see for instance physiological parameter values chosen in [37], to illustrate instances in which anomalous behaviors arise. We speculate, that one can refine this model to describe a pharmacological treatment of neurons whose activity is too intense, like in epileptic seizures or eventually to model the effect of drug consumption.

4.1.1. *Example: A special case.* Let us consider the previous example in the special case of an input dependent distribution $\bar{\Pi}$, in particular let, with $\delta = \mu - 1$, $\bar{\Pi}(r) = e^{-\delta r}$, $r > 0$, and, $\sigma^2 = 1$.

Let $\delta > 1$ and consider the integro-differential operator \mathbb{J}_δ given by (4.2) for $\alpha = \delta$. One gets that in this case $\hbar = 1/\delta$ and

$$(4.9) \quad \phi(u) = u + 2 \left(\mu - \frac{1}{u + \mu - 1} \right) - 1 = \frac{(u + \mu)(u + \mu - 2)}{u + \mu - 1}.$$

Since $\mu = \delta + 1 > 2$, $\phi(0) = \frac{\mu(\mu-2)}{\mu-1} > 0$, the required assumption (3.5) is satisfied.

In this case, the distribution $\bar{\Pi}$ being dependent on the incoming excitatory inputs, we have that the contribution of the jump part reduces as ν_e increases. This means that if the excitatory input is strong then the neuron fires with a weak contrast of the jump part, whereas if the input is weak and the potential is far from the threshold then the jump component tends to make the neuron silent. This behavior avoids unnecessary spikes and enhances the information transmission. This case may describe the situation in which inhibitory neurons inside the network, that are regulatory for the neuron activity, are not able to oppose to an increment in the excitatory inputs that may lead to an excessive spiking activity of the neuron under study.

On the contrary, if one wants to extend the model to address a pharmacological treatment of neurons whose activity is too intense we suggest to choose some jump distribution that depends on the inverse of μ or in general a heavy-tailed distribution that favors the large jumps reducing consistently the firing activity.

As a future work we plan to investigate the effects of other distributions of the jumps, with special attention to heavy-tailed distributions that favors large jumps. Moreover it would be interesting to add also upward jumps to the model and investigate the case of a signal dependent noise as in [26],[38] and [41], to study the possible role of the inhibitory jumps in improving the information transmission through a coherence resonance between the input and the output.

5. CONCLUSIONS

The contribution of this paper is twofold. On the one hand, it allows to advance in the LIF modeling in the mathematical neuroscience context. On the other hand, it also contains an original methodology, based on intertwining relationship, to study the classical first passage time problem of a Markov process with possible jumps.

Starting from the idea of endowing the classical LIF model with features that are more in keeping with the phenomenological reality we introduce a diffusion process with jumps for the description of the activity of a single neuron. Among the strengths of the presented model we have that on one hand the *good* properties of the classical Jacobi process are preserved: the state space is limited and the frequency and the amplitude of the jumps are state-dependent. On the other hand it also accounts for inputs that prevent the diffusion limit due to their amplitudes and/or to their frequencies. In this way one can assign different weights of the incoming inputs depending on whether they arrive more or less close to the trigger zone, as a first attempt to consider the neuron not only as a point. Moreover, these downward jumps are able to reduce the firing rate and introduce saturation even in the absence of a refractory period. This constitutes a novelty compared to other LIF models for which the firing rate increases linearly (and unbounded). The feature that the model accounts possibly large downward jumps suggests its use for describing the contribution of a strong internal inhibitory input or the effect of an external factor, like a pharmacological treatment or the intake of drugs and alcohol that interfere with the standard activity of the neuron. The tuning of this quantity can also help the investigation of the role of inhibition in the information transmission. Moreover, the fact that the jumps are more frequent close to the inhibitory reversal potential could describe the phenomenon of neuronal accommodation (see for instance [3]). In this framework after a spike, if a current which rises sufficiently slowly is applied, it will never evoke an action potential until the inactive phase is over. Finally, the high degree of freedom in the choice of the jump

distribution and the relatively easy numerical implementation permit the description of multiple different situations.

As far as the mathematical novelty is concerned, we note that for the study of the firing rate, we had to develop the study of the first-passage times of the proposed Markovian Jacobi process with jumps through a constant boundary. We are able to inherit some results from the classical Jacobi process to the process with jumps, thanks to a general strategy, original in the context of FPT problems, that relies on intertwining relations between the semigroups of the classical Jacobi process and its generalization. Therefore, this paper provides an additional application of such a concept in the theory of Markov processes by transferring q -invariant functions from a reference semigroup to semigroups that are in its intertwining orbit. This new approach enables us to characterize the Laplace transform of the FPT, expressed in terms of a generalization of the Gauss hypergeometric function that we introduce. As by-product, we obtain a closed-form expression for its expectation. This result appears of particular interest since an exact simulation method for the paths of the process considered is not yet available. We also mention that, relying on the recent works [12, 40], where a comprehensive fluctuation theory for skip-free Markov chains is established, one could exploit the intertwining relation to identify the Laplace transform of the first exit time from an interval. More generally, intertwining relations enable to relate the set of q -invariant functions (martingales), and, more generally, the convex cone of q -excessive functions (supermartingales) between semigroups. In potential theory, q -excessive functions are well-known to characterize the Laplace transform of the first passage time of a set for the processes. For this standpoint, the intertwining approach seems to be a natural and promising way to deal with the first exit time problem for a Markov process with two sided-jumps. These will be the subject of future investigations.

Finally, we stress that, despite our application in the context of mathematical neuroscience, the results on the Jacobi process with jumps and its first passage time through a constant boundary are novel and of a general nature. In particular, we mention the use of the Jacobi process in the context of population genetics and mathematical finance, where it often goes under the name of Wright-Fisher diffusion.

APPENDIX A. FIRST PASSAGE TIMES OF THE CLASSICAL JACOBI PROCESS

Let $Y = (Y_t)_{t \geq 0}$ be the Jacobi process with infinitesimal generator given, for a smooth function f on $[0, 1]$, by

$$(A.1) \quad \mathbf{J}_\mu f(y) = \frac{\sigma^2}{2} y(1-y) f''(y) - (\lambda y - \mu) f'(y)$$

with $\mu > \sigma^2/2$ to ensure that 0 is an entrance boundary [22]. We recall that, from (3.4), $W_\phi(n+1) = (\phi(0) + 1)_n = \left(\frac{2\mu}{\sigma^2}\right)_n$, and, thus, in this case

$${}_2F_1(a, b; \phi; y) = {}_2F_1(a, b; \phi(0) + 1; y)$$

the latter being the Gauss hypergeometric function. In what follows, we recall the expression of the Laplace transform and the first moment of its first passage time, which can be found in [37], see also [18].

Proposition A.1. *Let $0 < y < a < 1$, the Laplace transform of the first passage time*

$$(A.2) \quad T_a = \inf\{t > 0; Y_t \geq a\}$$

of the Jacobi process (A.1) is given by

$$(A.3) \quad \mathbb{E}_y[e^{-qT_a} \mathbb{I}_{\{T_a < \infty\}}] = \frac{{}_2F_1\left(\kappa(q), \theta(q); \frac{2\mu}{\sigma^2}; y\right)}{{}_2F_1\left(\kappa(q), \theta(q), \frac{2\mu}{\sigma^2}; a\right)}$$

where $\kappa(q)$ and $\theta(q)$ are solution of the system

$$(A.4) \quad \kappa(q) + \theta(q) + 1 = \frac{2\lambda}{\sigma^2}, \quad \kappa(q)\theta(q) = \frac{2q}{\sigma^2}$$

More specifically,

$$\kappa(q) = \frac{2q}{\theta(q)\sigma^2},$$

and

$$\theta(q) = \frac{2\lambda - \sigma^2 \pm \sqrt{(\sigma^2 - 2\lambda)^2 - 8q\sigma^2}}{2\sigma^2} = \bar{\lambda} \pm \sqrt{\left|\bar{\lambda}^2 - \frac{2q}{\sigma^2}\right|} \left(\mathbb{I}_{\{q \leq \frac{\sigma^2 \bar{\lambda}^2}{2}\}} + i \mathbb{I}_{\{q > \frac{\sigma^2 \bar{\lambda}^2}{2}\}} \right).$$

where $\bar{\lambda} = \frac{\lambda}{\sigma^2} - \frac{1}{2} \geq 0$, as, by condition (3.3), $\lambda > \mu > \sigma^2/2$. Finally, the first moment of T_a is

$$(A.5) \quad \mathbb{E}_y[T_a] = \frac{1}{\mu} \sum_{n=0}^{\infty} \frac{\left(\frac{2\lambda}{\sigma^2}\right)_n}{\left(\frac{2\mu}{\sigma^2} + 1\right)_n} \frac{a^{n+1} - y^{n+1}}{n+1}$$

$$(A.6) \quad = \frac{1}{\mu} \left({}_3F_2\left(1, 1, \frac{2\lambda}{\sigma^2}; 2, \frac{2\mu}{\sigma^2}; a\right) a - {}_3F_2\left(1, 1, \frac{2\lambda}{\sigma^2}; 2, \frac{2\mu}{\sigma^2}; y\right) y \right).$$

APPENDIX B. MEAN OF THE FIRST PASSAGE TIME OF THE JACOBI PROCESS WITH EXPONENTIAL JUMPS TYPE

Proposition B.1. *Under the condition $\frac{\sigma^2}{2} < \mu - \frac{1}{\alpha}$, the first moment of T_a for the Jacobi process with jumps with generator (4.2) is, for any $0 < y < a$,*

$$(B.1) \quad \mathbb{E}_y[T_a] = \frac{2(\alpha + 1)}{\sigma^2(k_+ + 1)(k_- + 1)} \left({}_4F_3\left(1, 1, \alpha + 2, \frac{2\lambda}{\sigma^2}; 2, k_+ + 2, k_- + 2; a\right) - {}_4F_3\left(1, 1, \alpha + 2, \frac{2\lambda}{\sigma^2}; 2, k_+ + 2, k_- + 2; y\right) y \right)$$

where

$$(B.2) \quad k_{\pm} = \frac{1}{2} \left(\alpha + 2\mu/\sigma^2 - 1 \pm \sqrt{(\alpha + 2\mu/\sigma^2 - 1)^2 - 4(2\alpha\mu/\sigma^2 - \alpha - 2/\sigma^2)} \right).$$

Proof. For $\hbar = 1/\alpha$, $\alpha \geq 1$, the Bernstein function ϕ defined in (4.3) can be written as

$$(B.3) \quad \phi(u) = \frac{(u + k_+)(u + k_-)}{(u + \alpha)}$$

with k_+ and k_- are defined in (B.2). This implies that, for any $n \geq 0$,

$$(B.4) \quad W_{\phi}(n+2) = \frac{(k_+ + 1)_{n+1}(k_- + 1)_{n+1}}{(\alpha + 1)_{n+1}}.$$

Using that $(k_{\pm} + 1)_{n+1} = (k_{\pm} + 1)(k_{\pm} + 2)_n$ and (3.25), one gets that

$$(B.5) \quad \mathbb{E}_y[T_a] = \frac{2(\alpha + 1)}{\sigma^2(k_+ + 1)(k_- + 1)} \sum_{n=0}^{\infty} \frac{(1)_n(1)_n\left(\frac{2\lambda}{\sigma^2}\right)_n(\alpha + 2)_n}{(2)_n(k_+ + 2)_n(k_- + 2)_n} \frac{a^{n+1} - y^{n+1}}{n!}.$$

Finally, the definition of the generalized hypergeometric function proves (B.1). ■

ACKNOWLEDGMENTS

The authors are indebted to the associated editors and an anonymous referee for valuable and constructive comments that improved the presentation of the paper. G.D. and L.S. have been partially supported by the MIUR-PRIN 2022 project “Non-Markovian dynamics and non-local equations”, no. 202277N5H9.

REFERENCES

- [1] M. Abramowitz and I.A. Stegun. *Handbook of Mathematical Functions: With Formulas, Graphs, and Mathematical Tables*. Applied mathematics series, Dover Publications, 1965.
- [2] L.U. Ancarani and G. Gasaneo. Derivatives of any order of the Gaussian hypergeometric function ${}_2F_1(a, b, c; z)$ with respect to the parameters a, b and c . *J. Phys. A: Math. Theor.*, 42(39):395208, 2009.
- [3] M. Baker and H. Bostock. Depolarization changes the mechanism of accommodation in rat and human motor axons. *J Physiol.*, 411:545–561, 1989.
- [4] T. Barta and L. Kostal. The effect of inhibition on rate code efficiency indicators. *PLOS Computational Biology*, 15(12):e1007545, 2019.
- [5] T. Barta and L. Kostal. Regular spiking in high-conductance states: The essential role of inhibition. *Phys. Rev. E*, 103(2):022408, 2021.
- [6] R. F. Bass. Adding and subtracting jumps from Markov processes. *Trans. Amer. Math. Soc.* 255, 363-376, 1979.
- [7] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and geometry of Markov diffusion operators*, volume 348 of *Grundlehren der Mathematischen Wissenschaften*. Springer, Cham, 2014.
- [8] A. Benveniste, and J. Jacod. Systèmes de Lévy des processus de Markov. *Inventiones mathematicae*, 21: 183–198, 1973.
- [9] J. Bertoin. Subordinators: examples and applications. In *Lectures on probability theory and statistics (Saint-Flour, 1997)*, volume 1717 of *Lecture Notes in Math.*, pages 1–91. Springer, Berlin, 1999.
- [10] J. Bauermann and B. Lindner. Multiplicative noise is beneficial for the transmission of sensory signals in simple neuron models. *Biosystems*, 178:25–31, 2019.
- [11] P. Cheridito, P. Patie, A. Srapionyan, and A. Vaidyanathan. On non-local ergodic jacobi semigroups: spectral theory, convergence-to-equilibrium and contractivity. *Journal de l'Ecole Polytechnique-Mathématiques*, 8:331–378, 2021.
- [12] M.C.H. Choi and P. Patie, Skip-free Markov chains, *Trans. Amer. Math. Soc.*, 371(10), 7301-7342, 2019.
- [13] C. Cuchiero, M. Larsson, and S. Svaluto-Ferro. Polynomial jump-diffusions on the unit simplex. *Ann. Appl. Probab.*, 28(4):2451–2500, 2018.
- [14] G. Da Prato. *An introduction to infinite-dimensional analysis*. Universitext. Springer-Verlag, Berlin, 2006.
- [15] D.A. Darling and A.J.F. Siegert. The First Passage Problem for a Continuous Markov Process. *Ann. Math. Statist.*, 24(4):624–639, 1953.
- [16] P. Dayan, and L.F- Abbott. *Theoretical Neuroscience: Computational And Mathematical Modeling of Neural Systems*. Massachusetts Institute of Technology Press, 2005.
- [17] F. Delbaen and H. Shirakawa. An interest rate model with upper and lower bounds. *Asia-Pacific Financial Markets*, 9(3):191–209, 2002.
- [18] G. D’Onofrio, M. Tamborrino and P. Lansky. The Jacobi diffusion process as a neuronal model. *Chaos*, 28(10):103119, 2018.
- [19] G. D’Onofrio, P. Lansky and M. Tamborrino. Inhibition enhances the coherence in the Jacobi neuronal model. *Chaos, Solitons & Fractals*, 128:108–113, 2019.
- [20] G. D’Onofrio and A. Lanteri. Approximating the First Passage Time Density of Diffusion Processes with State-Dependent Jumps. *Fractal and Fractional*, 7(1):30, 2023.
- [21] J.C. Eccles *The physiology of synapses*. Academic Press, 1964.
- [22] J. Forman and M. Sorensen. The Pearson Diffusions: A Class of Statistically Tractable Diffusion Processes. *Scandinavian Journal of Statistics*, 35(3):438–465, 2008.
- [23] M.T. Giraudo and L. Sacerdote. Jump-Diffusion processes as models for neuronal activity. *Biosystems*, 40(1-2):75–82, 1997.
- [24] M.T. Giraudo, L. Sacerdote and C. Zucca. A Monte Carlo method for the simulation of first passage times of diffusion processes. *Methodology and computing in applied probability*, 3: 215–231, 2001.
- [25] C. Gourieroux and J. Jasiak. Multivariate Jacobi process with application to smooth transitions. *J. Econometrics*, 131(1-2):475–505, 2006.
- [26] P.E. Greenwood and P. Lansky. Optimum signal in a simple neuronal model with signal-dependent noise. *Biol Cybern*, 92(3):199-205, 2005.

- [27] R. C. Griffiths, P. A. Jenkins, and D. Spanò. Wright–Fisher diffusion bridges. *Theoretical Population Biology*, 122:67 – 77, 2018.
- [28] R. C. Griffiths and D. Spanó. Diffusion processes and coalescent trees. In *Probability and mathematical genetics*, volume 378 of *London Math. Soc. Lecture Note Ser.*, pages 358–379. Cambridge Univ. Press, Cambridge, 2010.
- [29] K. S. Hankin. *Hypergeo: The Gauss Hypergeometric Function*, R Package Version 1.2-13, available at <https://CRAN.R-project.org/package=hypergeo>, 2016.
- [30] F.B. Hanson and H.C. Tuckwell. Diffusion approximations for neuronal activity including synaptic reversal potentials. *J Theor Neurobiol*, 2:127–153, 1983.
- [31] T. Huillet. On Wright–Fisher diffusion and its relatives. *Journal of Statistical Mechanics: Theory and Experiment*, 2007(11):P11006, 2007.
- [32] P. Jahn, R.W. Berg, J. Hounsgaard and S. Ditlevsen. Motoneuron membrane potentials follow a time inhomogeneous jump diffusion process. *J Comput Neurosci.*, 31(3):563–579, 2011.
- [33] R.A. Jarrow, P. Patie, A. Srapionyan and Y. Zhao, Risk-neutral pricing techniques and example, *Math. Finance*, 31: 857–884, 2021.
- [34] R. Koekoek and R. F. Swarttouw. The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue, 1998.
- [35] L. Kostal, P. Lansky and M. Stiber. Statistics of inverse interspike intervals: The instantaneous firing rate revisited. *Chaos*, 28:106305, 2018.
- [36] A. E. Kyprianou. *Fluctuations of Lévy processes with applications*. Universitext. Springer, Heidelberg, second edition, 2014. Introductory lectures.
- [37] V. Lanska, P. Lansky and C.E. Smith. Synaptic Transmission in a Diffusion Model for Neural Activity. *J. Theor. Biol.*, 166(4):393–406, 1994.
- [38] P. Lansky, L. Sacerdote. The Ornstein–Uhlenbeck neuronal model with signal-dependent noise. *Physics Letters A*, 285(3-4):132–140, 2001.
- [39] N. N. Lebedev, *Special functions and their applications*, Prentice-Hall, Englewood Cliffs, NJ, 1965.
- [40] R.L. Loeffen, P. Patie and J. Wang. Fluctuation theory of continuous-time skip-free downward Markov chains with applications to branching processes with immigration, [arXiv:2208.14425](https://arxiv.org/abs/2208.14425), 2022.
- [41] A. Longtin, B. Doiron and A. R. Bulsara. Noise-induced divisive gain control in neuron models. *Biosystems*, 67(1):147–156, 2002.
- [42] A. Melanson and A. Longtin. Data-driven inference for stationary jump-diffusion processes with application to membrane voltage fluctuations in pyramidal neurons. *J. Math. Neurosc.*, 9(6), 30p, 2019.
- [43] P.-A. Meyer. Intégrales stochastiques IV. *Séminaire de probabilités de Strasbourg*, Tome 1, pp. 142-162, 1967.
- [44] L. Miclo and P. Patie. On interweaving relations. *Journal of Functional Analysis*, 280(3):108816, 2021.
- [45] B. Øksendal and A. Sulem. *Applied Stochastic Control of Jump Diffusions*. Springer Berlin Heidelberg, 2007.
- [46] S. Pal. Wright–Fisher diffusion with negative mutation rates. *Ann. Probab.*, 41(2):503–526, 2013.
- [47] P. Patie and R. Sarkar, Weak similarity orbit of the (log)-Bessel semigroup on the Euclidean space, *Proceedings of London Math. Soc.*, 126(5), 1522-1584, 2023.
- [48] P. Patie and M. Savov. Bernstein-gamma functions and exponential functionals of Lévy processes. *Electron. J. Probab.*, 23:Paper No. 75, 101p., 2018.
- [49] P. Patie and M. Savov. Spectral expansion of non-self-adjoint generalized Laguerre semigroups. *Mem. Amer. Math. Soc.*, 272, no. 1336, vii+182 pp, 2021.
- [50] L.M. Ricciardi and L. Sacerdote. The Ornstein-Uhlenbeck process as a model for neuronal activity. *Biol. Cybern.*, 35:1–9, 1979.
- [51] M.J. Richardson. Effects of synaptic conductance on the voltage distribution and firing rate of spiking neurons. *Physical Review E*, 69(5): 051918, 2004.
- [52] R. Rodriguez and P. Lansky. A simple stochastic model of spatially complex neurons. *Biosystems*, 58(1-3), 49–58, 2000.
- [53] L. Sacerdote and M.T. Giraudo. Stochastic Integrate and Fire Models: A Review on Mathematical Methods and Their Applications. In: *Stochastic Biomathematical Models. Lecture Notes in Mathematics*, vol 2058, pages 99–148, Springer, Berlin, 2013.
- [54] R. L. Schilling, R. Song, and Z. Vondraček. *Bernstein functions*, volume 37 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 2010.
- [55] R. Sirovich, L. Sacerdote and A.E.P. Villa. Cooperative behavior in a jump diffusion model for a simple network of spiking neurons. *Math. Biosc. for Engineering*, 11(2):385–401, 2013.
- [56] R.B. Stein. A theoretical analysis of neuronal variability. *Biophys J.*, 5(2):173–94, 1965.
- [57] M. Tamborrino, L. Sacerdote and M. Jacobsen Weak convergence of marked point processes generated by crossings of multivariate jump processes. Applications to neural network modeling *Physica D*, 288:45–52, 2014.

- [58] A. J. Trevelyan; S. F. Muldoon; E. M. Merricks; C. Racca; K. J. Staley. The Role of Inhibition in Epileptic Networks. *Journal of Clinical Neurophysiology*, 32(3):227–234, 2015.
- [59] G. Szegő. *Orthogonal polynomials*. American Mathematical Society, Providence, R.I., fourth edition, 1975. American Mathematical Society, Colloquium Publications, Vol. XXIII.

DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO, 10129 TURIN, ITALY
Email address: `giuseppe.donofrio@polito.it`

SCHOOL OF OPERATIONS RESEARCH AND INFORMATION ENGINEERING, CORNELL UNIVERSITY, ITHACA, NY 14853.
Email address: `pp396@cornell.edu`

DIPARTIMENTO DI MATEMATICA ‘G. PEANO’, UNIVERSITÀ DEGLI STUDI DI TORINO, VIA CARLO ALBERTO 10, 10123 TORINO, ITALY
Email address: `laura.sacerdote@unito.it`