

Homogenization and 3D-2D dimension reduction of a functional on manifold valued BV space

Original

Homogenization and 3D-2D dimension reduction of a functional on manifold valued BV space / Lussardi, Luca; Torricelli, Andrea; Zappale, Elvira. - In: MATHEMATICS IN ENGINEERING. - ISSN 2640-3501. - 8:3(2026), pp. 368-401. [10.3934/mine.2026012]

Availability:

This version is available at: 11583/3010685 since: 2026-05-08T11:59:40Z

Publisher:

AIMS

Published

DOI:10.3934/mine.2026012

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Research article

Homogenization and 3D-2D dimension reduction of a functional on manifold valued BV space

Luca Lussardi^{1,*}, Andrea Torricelli² and Elvira Zappale³

¹ Dipartimento di Scienze Matematiche G.L. Lagrange, Politecnico di Torino, C.so Duca degli Abruzzi 24, 10129 Torino, Italy

² Dipartimento di Matematica, Università di Trento, Via Sommarive 14, 38123 Povo, Italy

³ Dipartimento di Scienze di base e Applicate per l'Ingegneria, Sapienza - Università di Roma, Via Antonio Scarpa 16, 00161 Roma, Italy and CIMA, Universidade de Évora, Portugal

* **Correspondence:** Email: luca.lussardi@polito.it.

Abstract: We study the simultaneous homogenization and dimension reduction of an energy functional with linear growth defined on the space of manifold valued Sobolev functions. The study is carried out by Γ -convergence, providing an integral representation result in the space of manifold constrained functions with bounded variation.

Keywords: homogenization; dimensional reduction; manifold-valued spaces of functions with bounded variation; Γ -convergence; micromagnetics; linear growth

1. Introduction and main results

The study of variational problems constrained to manifold-valued functions has attracted increasing attention due to its relevance in various fields such as micromagnetics and liquid crystals (see, for instance, [1–4]). In this paper, we investigate the simultaneous process of homogenization and dimension reduction for a class of energy functionals with linear growth, defined on the space of functions taking values in a smooth manifold without boundary. Specifically, we consider a one-parameter family of functionals defined on $W^{1,1}$ functions with values in a compact, connected manifold $\mathcal{M} \subset \mathbb{R}^3$, and study their behavior under the joint asymptotic regime of periodic homogenization and thin domain limit.

The joining of this two processes is a classical field of study, indeed the simultaneous homogenization and dimension reduction of an integral functional defined on real valued Sobolev functions has been studied in [5] in the case $p > 1$, while the case $p = 1$ can be covered by the global method [6], but without explicit formulae which connect the limiting energy density with the

original ones. In the application framework, regarding either the solely study of homogenization or the study of thin structures we refer to [7–23] among a wider literature, Having in mind applications in the superlinear setting we refer, for instance, to [24–28]. As far as we know, there are no rigorous derivation on dimensional reduction for functionals with linear growth defined on manifold-valued Sobolev functions.

Our analysis is carried out in the framework of Γ -convergence (see [29]), which provides a rigorous tool for deriving effective models and capturing the asymptotic behavior of infimizers. A key feature of our approach is the use of blow-up techniques and relaxation arguments adapted to the manifold setting.

The main result of the paper establishes the Γ -limit of the rescaled energy functional in the space $BV(\omega; \mathcal{M})$, where $\omega \subset \mathbb{R}^2$ represents the cross-section of the thin domain.

We assume that the domain is an inhomogeneous cylinder, whose microstructure is distributed with periodicity within the material described by the small parameter h comparable with the height of the domain. The equilibrium configurations are detected as minimizers of an integral functional of the form

$$\int_{\omega \times (-\frac{h}{2}, \frac{h}{2})} f\left(\frac{x}{h}, \nabla u\right) dx \quad u : \omega \times \left(-\frac{h}{2}, \frac{h}{2}\right) \rightarrow \mathbb{R}^3,$$

under suitable boundary conditions, where $\omega \subset \mathbb{R}^2$ is a bounded open set, and $f : \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}$ is a periodic integrand with respect to the first variable, and u is a manifold-valued Sobolev field, that will be specialized in the sequel. Due to the many applications, it is worth to recall that solely the homogenization of integral functionals depending on x and ∇u and defined on manifold-valued Sobolev fields has been studied in [12] for $u \in W^{1,p}$ and in [11, 30] for $u \in W^{1,1}$, we also refer to [2] for other related models. Analogously the dimensional reduction of micromagnetic and ferromagnetic energy has been studied, in several contexts, we recall [31–36] among the others.

Given a Carathéodory function $f : \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}$ we consider the functional

$$\frac{1}{h} \int_{\omega_h} f\left(\frac{x}{h}, \nabla u\right) dx, \quad u \in W^{1,1}(\omega_h; \mathcal{M}),$$

with $\omega_h := \omega \times \left(-\frac{h}{2}, \frac{h}{2}\right)$, $h > 0$, $\omega \subset \mathbb{R}^2$ open and bounded, and \mathcal{M} a smooth submanifold of \mathbb{R}^3 without boundary. In particular, here and in the sequel, when not stated otherwise, we always assume that f has the following properties:

(H1) $f(\cdot, x_3, \xi)$ is 1-periodic, i.e., for every $(x_\alpha, x_3) \in \mathbb{R}^3$ and $\xi \in \mathbb{R}^{3 \times 3}$, it holds

$$f(x_\alpha + \mathbf{e}_i, x_3, \xi) = f(x_\alpha, x_3, \xi), \quad \forall i = 1, 2,$$

where $\{\mathbf{e}_1, \mathbf{e}_2\}$ is the canonical basis of \mathbb{R}^2 ;

(H2) f has linear growth: there exists $\alpha, \beta > 0$ such that

$$\alpha|\xi| \leq f(x, \xi) \leq \beta(1 + |\xi|), \quad (1.1)$$

for a.e. $x \in \mathbb{R}^3$ and for every $\xi \in \mathbb{R}^{3 \times 3}$;

(H3) f is a Lipschitz continuous function with respect to the second variable uniformly in the first variable, i.e., there exists $L > 0$ such that it holds

$$|f(x, \xi) - f(x, \eta)| \leq L|\xi - \eta|$$

for a.e. $x \in \mathbb{R}^3$ and $\xi, \eta \in \mathbb{R}^{3 \times 3}$;

(H4) there exist $C > 0$ and $0 < q < 1$ such that

$$|f(x, \xi) - f^\infty(x, \xi)| \leq C(1 + |\xi|^{1-q}),$$

for a.e. $x \in \mathbb{R}^3$ and $\xi \in \mathbb{R}^{3 \times 3}$ and where $f^\infty : \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}$ is the (strong) recession function of f and it is defined as

$$f^\infty(x, \xi) := \limsup_{t \rightarrow +\infty} \frac{f(x, t\xi)}{t}. \tag{1.2}$$

We observe that by (1.1) it holds

$$\alpha|\xi| \leq f^\infty(x, \xi) \leq \beta|\xi|, \tag{1.3}$$

for a.e. $x \in \mathbb{R}^3$ and every $\xi \in \mathbb{R}^{3 \times 3}$.

We consider the functional $\tilde{I}^h : L^1(\omega_h; \mathcal{M}) \rightarrow \overline{\mathbb{R}}$,

$$\tilde{I}^h(u) := \begin{cases} \frac{1}{h} \int_{\omega_h} f\left(\frac{x}{h}, \nabla u\right) dx, & \text{if } u \in W^{1,1}(\omega_h; \mathcal{M}), \\ +\infty, & \text{elsewhere.} \end{cases}$$

By a change of variable, to study the Γ -limit of \tilde{I}^h is equivalent to study the Γ -limit of the rescaled functional I^h defined as

$$I^h(u) := \begin{cases} \int_{\Omega} f\left(\frac{x_\alpha}{h}, x_3, \nabla_h u\right) dx, & \text{if } u \in W^{1,1}(\Omega; \mathcal{M}), \\ +\infty, & \text{elsewhere} \end{cases} \tag{1.4}$$

with

$$\Omega := \omega \times \left(-\frac{1}{2}, \frac{1}{2}\right) = \omega_{,1} \quad \text{and} \quad \nabla_h := \left[\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \frac{1}{h} \frac{\partial}{\partial x_3}\right].$$

From now on, given an appropriate matrix A and vector b we denote by $(A|b)$ the matrix that has b as the last column and the columns of A in the remaining columns. It follows that if $\nabla_\alpha := [\frac{\partial}{\partial x_1} | \frac{\partial}{\partial x_2}]$ and $\nabla_3 := \frac{\partial}{\partial x_3}$, then $\nabla_h = [\nabla_\alpha | \frac{1}{h} \nabla_3]$. Moreover, we denote with ξ_α an element of $\mathbb{R}^{3 \times 2}$ and with ξ an element of $\mathbb{R}^{3 \times 3}$.

Theorem 1.1. *Assume \mathcal{M} compact and connected, and let $f : \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}$ be a Carathéodory function satisfying (H1)–(H4). Then, the Γ -limit of I^h with respect to the L^1 -topology is the functional I defined as*

$$I(u) := \int_{\omega} T f_{\text{hom}}^0(u, \nabla_\alpha u) dx_\alpha + \int_{\omega} T f_{\text{hom}}^{0,\infty}\left(u, \frac{dD_\alpha^c u}{d|D_\alpha^c u|}\right) d|D_\alpha^c u| + \int_{\omega \cap S_u} \theta(u^+, u^-, \nu_u) d\mathcal{H}^1, \tag{1.5}$$

if $u \in BV(\omega; \mathcal{M})$ and $I(u) = +\infty$ elsewhere, with $Tf_{\text{hom}}^0 : \mathbb{R}^3 \times \mathbb{R}^{3 \times 2} \rightarrow \mathbb{R}$ defined as

$$Tf_{\text{hom}}^0(s, \xi_\alpha) := \liminf_{t \rightarrow +\infty} \inf_{\varphi} \left\{ \frac{1}{t^2} \int_{(tQ')_{,1}} f(x_\alpha, x_3, \xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi) dx_\alpha dx_3 : \right. \\ \left. \varphi \in W^{1,\infty}((tQ')_{,1}; T_s(\mathcal{M})), \varphi(x_\alpha, x_3) = 0 \text{ for every } (x_\alpha, x_3) \in \partial(tQ') \times \left(-\frac{1}{2}, \frac{1}{2}\right) \right\}, \quad (1.6)$$

where $Q' := \left(-\frac{1}{2}, \frac{1}{2}\right)^2$, $(tQ')_{,1} := tQ' \times \left(-\frac{1}{2}, \frac{1}{2}\right)$, and $T_s(\mathcal{M})$ denotes the tangent space to \mathcal{M} in s . Moreover, for every $s \in \mathbb{R}^3$, $Tf_{\text{hom}}^{0,\infty}(s, \cdot)$ is the (strong) recession function of $Tf_{\text{hom}}^0(s, \cdot)$, i.e., for every $(s, \xi_\alpha) \in \mathbb{R}^3 \times \mathbb{R}^{3 \times 2}$:

$$Tf_{\text{hom}}^{0,\infty}(s, \xi_\alpha) = (Tf_{\text{hom}}^0)^{\infty}(s, \xi_\alpha) = \limsup_{t \rightarrow +\infty} \frac{Tf_{\text{hom}}^0(s, t\xi_\alpha)}{t}, \quad (1.7)$$

while $\theta : \mathcal{M} \times \mathcal{M} \times \mathbb{S}^1 \rightarrow \mathbb{R}$, is defined as

$$\theta(a, b, \nu) := \lim_{t \rightarrow +\infty} \inf_{\phi} \left\{ \frac{1}{t} \int_{(tQ_\nu)_{,1}} f^\infty(x_\alpha, x_3, \nabla \phi) dx_\alpha dx_3 : \right. \\ \left. \phi \in W^{1,1}((tQ_\nu)_{,1}, \mathcal{M}), \phi = a \text{ on } \partial(tQ_\nu) \times \left(-\frac{1}{2}, \frac{1}{2}\right) \cap \{x_\alpha \cdot \nu > 0\} \text{ and} \right. \\ \left. \phi = b \text{ on } \partial(tQ_\nu) \times \left(-\frac{1}{2}, \frac{1}{2}\right) \cap \{x_\alpha \cdot \nu < 0\} \right\}, \quad (1.8)$$

where Q_ν is an open cube of \mathbb{R}^2 centered at the origin and with two of its faces parallel to ν , $Q_{\nu,1} := Q_\nu \times \left(-\frac{1}{2}, \frac{1}{2}\right)$, and \mathbb{S}^1 is the unit sphere in \mathbb{R}^2 .

2. Notation and preliminaries

This section is devoted to fix notation, recall previous results that will be useful in the sequel and establish properties of the energy densities appearing in the main results. In what follows $\Omega := \omega \times \left(-\frac{1}{2}, \frac{1}{2}\right)$, $\omega \subset \mathbb{R}^2$ is open and bounded, and \mathcal{M} is a smooth submanifold of \mathbb{R}^3 without boundary, further assumptions on \mathcal{M} will be stated explicitly if needed. Given $s \in \mathcal{M}$, we write $T_s(\mathcal{M})$ for the tangent space to \mathcal{M} in s . Given $a, b \in \mathcal{M}$, we introduce the family of geodesic curves between a and b by

$$\mathcal{G}(a, b) := \left\{ \gamma \in C^\infty(\mathbb{R}; \mathcal{M}) : \gamma(t) = a, \text{ if } t \geq 1/2, \right. \\ \left. \gamma(t) = b, \text{ if } t \leq -1/2, \int_{\mathbb{R}} |\dot{\gamma}| dt = \mathbf{d}_{\mathcal{M}}(a, b) \right\}, \quad (2.1)$$

where $\mathbf{d}_{\mathcal{M}}$ denotes the geodesic distance on \mathcal{M} .

We denote by $\mathcal{A}(\Omega)$ the family of all open subsets of Ω and with $\mathcal{A}(\omega)$ the family of all open subset of ω . We write $B^k(s, r)$ for the closed ball in \mathbb{R}^k , $k \in \mathbb{N}$, of center $s \in \mathbb{R}^k$ and radius $r > 0$. Moreover, we denote by Q the cube $\left(-\frac{1}{2}, \frac{1}{2}\right)^3$ and with $Q(x_0, \rho)$ the rescaled and translated cube $x_0 + \rho Q$, with $x_0 \in \mathbb{R}^3$, $\rho > 0$. In a similar way, given $\nu \in \mathbb{S}^1$, Q_ν stands for the open unit square in \mathbb{R}^2 centered at the origin, with the two sides orthogonal to ν ; we set $Q_\nu(x_0, \rho) := x_0 + \rho Q_\nu$.

Furthermore, given a set $A \subset \omega$ we denote by A_h , with $h > 0$, the set $A \times (-\frac{h}{2}, \frac{h}{2})$ and so $A_{,1} := A \times (-\frac{1}{2}, \frac{1}{2})$. It follows that, in particular, $\Omega = \omega_{,1}$. We also denote by Q' the cube $(-\frac{1}{2}, \frac{1}{2})^2$, so $Q'_{,h} = (-\frac{1}{2}, \frac{1}{2})^2 \times (-\frac{h}{2}, \frac{h}{2})$ for $h > 0$, while $Q'_{,1} = (-\frac{1}{2}, \frac{1}{2})^3 = Q$ and $Q_{v,1} = Q_v \times (-\frac{1}{2}, \frac{1}{2})$. $\mathcal{M}(\Omega)$ is the space of real valued Radon measure in Ω with finite total variation, \mathcal{L}^k , is the k -dimensional Lebesgue measure, with $k \in \mathbb{N}$. Finally, given $\lambda, \mu \in \mathcal{M}(\Omega)$ we denote by $\frac{d\lambda}{d\mu}$ the Radon-Nykodým derivative of λ with respect to μ . By a generalization of Besicovitch differentiation theorem, see for instance [37, Proposition 2.2], there exists a Borel set E such that $\mu(E) = 0$ and $\frac{d\lambda}{d\mu}(x) = 0$ for every $x \in \text{supp } \lambda \setminus E$.

Given $s \in \mathcal{M}$, we consider the orthogonal projection

$$P_s : \mathbb{R}^3 \rightarrow T_s(\mathcal{M}), \tag{2.2}$$

and we define the function $\mathbf{P}_s : \mathbb{R}^{3 \times 3} \rightarrow [T_s(\mathcal{M})]^3$ as

$$\mathbf{P}_s(\xi) := [P_s(\xi_1)|P_s(\xi_2)|P_s(\xi_3)], \tag{2.3}$$

for every $\xi = [\xi_1|\xi_2|\xi_3] \in \mathbb{R}^{3 \times 3}$. For a Carathéodory function $f : \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}$, we set

$$\bar{f}(x, s, \xi) := f(x, \mathbf{P}_s(\xi)) + |\xi - \mathbf{P}_s(\xi)|. \tag{2.4}$$

This function will play a crucial role in our subsequent analysis, since it will appear in the formulas to detect our limiting energy densities. By construction the function $\bar{f} : \mathbb{R}^3 \times \mathcal{M} \times \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}$ is Carathéodory, i.e., it is measurable with respect to the first variable and continuous with respect to the last two variables. Moreover, if conditions (H1) and (H2) are satisfied then \bar{f} also satisfies (H1) and (H2), i.e., \bar{f} is 1-periodic in the first variable and there exists $C > 0$ such that

$$\frac{1}{C}|\xi| \leq \bar{f}(x, s, \xi) \leq C(1 + |\xi|), \tag{2.5}$$

for every $(s, \xi) \in \mathcal{M} \times \mathbb{R}^{3 \times 3}$ and for a.e. $x \in \mathbb{R}^3$. Now we extend the function \bar{f} in (2.4) to the whole space $\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^{3 \times 3}$. This can be done as in [11, Lemma 3.1]. For $\delta_0 > 0$ fixed, let $\mathcal{U} := \{s \in \mathbb{R}^3 : \text{dist}(s, \mathcal{M}) < \delta_0\}$ be the δ_0 -neighborhood of \mathcal{M} . Choosing $\delta_0 > 0$ small enough, we may assume that the nearest point projection $\Pi : \mathcal{U} \rightarrow \mathcal{M}$ is a well defined Lipschitz mapping. Then the map $s \in \mathcal{U} \mapsto P_{\Pi(s)}$ is Lipschitz, with $P_{\Pi(s)}$ as in (2.2). We denote by χ a cut-off function in $C_c^\infty(\mathbb{R}^3; [0, 1])$ such that $\chi(t) = 1$ if $\text{dist}(t, \mathcal{M}) \leq \delta_0/2$, and $\chi(t) = 0$ if $\text{dist}(t, \mathcal{M}) \geq \frac{3\delta_0}{4}$. We define $\mathbb{P}_s(\xi) := \chi(s)\mathbf{P}_{\Pi(s)}(\xi)$ for $(s, \xi) \in \mathbb{R}^3 \times \mathbb{R}^{3 \times 3}$, with $\mathbf{P}_{\Pi(s)}$ as in (2.3). In the sequel we will consider the integrand $g : \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow [0, +\infty)$ given by

$$g(y, s, \xi) = f(y, \mathbb{P}_s(\xi)) + |\xi - \mathbb{P}_s(\xi)|. \tag{2.6}$$

We also recall an important property of manifold-valued Sobolev functions. The following theorem has been proved in [38, 39] and we will use it to prove Theorem 1.1.

Theorem 2.1. *Let $\Omega \subset \mathbb{R}^3$ be open and bounded, let \mathcal{S} be the family of all finite unions of 1-dimensional manifolds in \mathbb{R}^3 and let $\pi_1(\mathcal{M})$ be the fundamental group of \mathcal{M} . Denote by $\mathcal{D}(\Omega, \mathcal{M}) \subset W^{1,1}(\Omega, \mathcal{M})$ the set*

$$\mathcal{D}(\Omega, \mathcal{M}) := \begin{cases} W^{1,1}(\Omega, \mathcal{M}) \cap C^\infty(\Omega, \mathcal{M}), & \text{if } \pi_1(\mathcal{M}) = 0, \\ \{u \in W^{1,1}(\Omega, \mathcal{M}) \cap C^\infty(\Omega \setminus \Sigma, \mathcal{M}) \text{ for some } \Sigma \in \mathcal{S}\}, & \text{otherwise.} \end{cases}$$

Then $\mathcal{D}(\Omega, \mathcal{M})$ is dense in $W^{1,1}(\Omega, \mathcal{M})$ with respect to the strong topology of $W^{1,1}(\Omega, \mathbb{R}^3)$.

Remark 2.2. Given $A \subset \mathcal{A}(\omega)$, the set

$$V(A) := \left\{ v \in W^{1,p}(A,1; \mathcal{M}) : \frac{\partial v}{\partial x_3} = 0 \text{ a.e. on } A,1 \right\},$$

is isomorphic to the Sobolev space $W^{1,p}(A; \mathcal{M})$.

2.1. Bulk part density

Now we characterize the density of the bulk part of the Γ -limit. First, we recall that the extended function g defined in (2.6) has the following properties.

Lemma 2.3. [11, Lemma 3.1] Assume that \mathcal{M} is compact. Let $f : \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow [0, +\infty)$ be a Carathéodory function satisfying (H1)–(H3). Then the function $g : \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow [0, +\infty)$ in (2.6), turns out to be a Carathéodory function such that

$$g(y, s, \xi) = f(y, \xi), \quad g^\infty(y, s, \xi) = f^\infty(y, \xi), \quad (2.7)$$

for a.e. $y \in \mathbb{R}^3$, for every $s \in \mathcal{M}$ and $\xi \in [T_s(\mathcal{M})]^3$. Furthermore it satisfies

- (i) $g(y_\alpha, y_3, s, \xi)$ is 1-periodic in the first variable, with $(y_\alpha, y_3) \in \mathbb{R}^3$;
- (ii) there exist $0 < \alpha' \leq \beta'$ such that

$$\alpha' |\xi| \leq g(y, s, \xi) \leq \beta' (1 + |\xi|), \quad (2.8)$$

for a.e. $y \in \mathbb{R}^3$ and for every $(s, \xi) \in \mathbb{R}^3 \times \mathbb{R}^{3 \times 3}$;

- (iii) there exist constants $C > 0$ and $C' > 0$ such that

$$|g(y, s, \xi) - g(y, s', \xi)| \leq C |s - s'| |\xi|, \quad (2.9)$$

and

$$|g(y, s, \xi) - g(y, s, \xi')| \leq C' |\xi - \xi'|, \quad (2.10)$$

for a.e. $y \in \mathbb{R}^3$, for every $s, s' \in \mathbb{R}^3$, and for every $\xi \in \mathbb{R}^{3 \times 3}$;

- (iv) if (H4) holds, there exists $0 < q < 1$ and C'' such that

$$|g(y, s, \xi) - g^\infty(y, s, \xi)| \leq C'' (1 + |\xi|^{1-q}), \quad (2.11)$$

for a.e. $y \in \mathbb{R}^3$, for every $(s, \xi) \in \mathbb{R}^3 \times \mathbb{R}^{3 \times 3}$, and where $g^\infty : \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}$ is the (strong) recession function of g and it is defined as

$$g^\infty(y, s, \xi) := \limsup_{t \rightarrow +\infty} \frac{g(y, s, t\xi)}{t}.$$

Now we prove some properties of the bulk part $T f_{hom}^0$ (see (1.6)) of I (see (1.5)).

Proposition 2.4. Given a Carathéodory function $f : \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}$ satisfying conditions (H1) and (H2), let $g : \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow [0, +\infty)$ be the functions defined as in (2.6), then following properties hold.

(i) For every $s \in \mathcal{M}$ and $\xi_\alpha \in [T_s(\mathcal{M})]^2$

$$Tf_{\text{hom}}^0(s, \xi_\alpha) = g_{\text{hom}}^0(s, \xi_\alpha), \quad (2.12)$$

where Tf_{hom}^0 is defined as in (1.6) and

$$g_{\text{hom}}^0(s, \xi_\alpha) = \liminf_{t \rightarrow +\infty} \inf_{\varphi} \left\{ \frac{1}{t^2} \int_{(tQ')_{,1}} g(x_\alpha, x_3, s, \xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi) dx_\alpha dx_3 : \right. \\ \left. \varphi \in W^{1,\infty}((tQ')_{,1}; \mathbb{R}^3), \varphi(x_\alpha, x_3) = 0 \text{ for every } (x_\alpha, x_3) \in \partial(Q') \times \left(-\frac{1}{2}, \frac{1}{2}\right) \right\}. \quad (2.13)$$

(ii) The function Tf_{hom}^0 is tangentially quasiconvex in the second variable, i.e.,

$$Tf_{\text{hom}}^0(s, \xi_\alpha) \leq \int_{Q'} Tf_{\text{hom}}^0(s, \xi_\alpha + \nabla_\alpha \psi) dx_\alpha,$$

for every $s \in \mathcal{M}$, $\xi_\alpha \in [T_s(\mathcal{M})]^2$, and $\psi \in W_0^{1,\infty}(Q'; T_s(\mathcal{M}))$. In particular, $Tf_{\text{hom}}^0(s, \cdot)$ is rank one convex.

(iii) Tf_{hom}^0 is uniformly 1-coercive and has uniform linear growth in the second variable (i.e., it satisfies inequalities as in (2.5), uniformly with respect to s). Moreover, there exists $C > 0$ such that for every $s \in \mathcal{M}$ and $\xi_\alpha, \xi'_\alpha \in [T_s(\mathcal{M})]^2$

$$|Tf_{\text{hom}}^0(s, \xi_\alpha) - Tf_{\text{hom}}^0(s, \xi'_\alpha)| \leq C|\xi_\alpha - \xi'_\alpha|.$$

(iv) Finally, if hypothesis (H4) is also satisfied, then

$$|Tf_{\text{hom}}^0(s, \xi_\alpha) - Tf_{\text{hom}}^{0,\infty}(s, \xi_\alpha)| \leq C(1 + |\xi_\alpha|^{1-q}),$$

for some positive constant C and for every $s \in \mathcal{M}$ and $\xi_\alpha \in [T_s(\mathcal{M})]^2$, where $Tf_{\text{hom}}^{0,\infty}$ is the function appearing in (1.7).

Proof. The proof of (i)–(iii) follows from [28, Proposition 2.1]. Now we prove (iv) following the technique of [11, Proposition 3.1-(v)]. Consider a sequence $(t_n)_n \subset \mathbb{R}$ such that $t_n \rightarrow +\infty$ as $n \rightarrow +\infty$, a sequence $(k_n)_n \subset \mathbb{N}$, and $(\varphi_n)_n \subset W^{1,\infty}((0, k_n)_{,1}^2; T_s(\mathcal{M}))$ such that $\varphi_n(x_\alpha, x_3) = 0$ for every $(x_\alpha, x_3) \in \partial(0, k_n)^2 \times \left(-\frac{1}{2}, \frac{1}{2}\right)$ and such that for every $s \in \mathcal{M}$ and $\xi_\alpha \in \mathbb{R}^{3 \times 2}$

$$Tf_{\text{hom}}^{0,\infty}(s, \xi_\alpha) = \lim_{n \rightarrow +\infty} \frac{Tf_{\text{hom}}^0(s, t_n \xi_\alpha)}{t_n}, \quad (2.14)$$

and

$$\int_{(0, k_n)_{,1}^2} f(y_\alpha, y_3, t_n \xi_\alpha + t_n \nabla_\alpha \varphi_n | \nabla_3 \varphi_n) dy \leq Tf_{\text{hom}}^0(s, t_n \xi_\alpha) + \frac{1}{n}.$$

By (H2) and since Tf_{hom}^0 has linear growth, we have that

$$\int_{(0, k_n)_{,1}^2} |\nabla_{t_n} \varphi_n| dx \leq C(1 + |\xi_\alpha|), \quad (2.15)$$

for some positive $C = C(\alpha, \beta)$. Since

$$\lim_{n \rightarrow +\infty} \frac{1}{t_n} \int_{(0, k_n)_1^2} f(y_\alpha, y_3, t_n \xi_\alpha + t_n \nabla_\alpha \varphi_n | \nabla_3 \varphi_n) dy = T f_{hom}^0(s, \xi_\alpha),$$

then by (2.14) we have

$$\begin{aligned} & T f_{hom}^0(s, \xi_\alpha) - T f_{hom}^{0, \infty}(s, \xi_\alpha) \\ & \leq \liminf_{n \rightarrow \infty} \left\{ \int_{(0, k_n)_1^2} \left| f(y_\alpha, y_3, \xi_\alpha + \nabla_\alpha \varphi_n | \nabla_3 \varphi_n) - \frac{f(y_\alpha, y_3, t_n \xi_\alpha + t_n \nabla_\alpha \varphi_n | \nabla_3 \varphi_n)}{t_n} \right| dy \right\}. \end{aligned}$$

Summing and subtracting $f^\infty(y_\alpha, y_3, \xi_\alpha + \nabla_\alpha \varphi_n | \nabla_3 \varphi_n)$, we get

$$\begin{aligned} & T f_{hom}^0(s, \xi_\alpha) - T f_{hom}^{0, \infty}(s, \xi_\alpha) \\ & \leq \liminf_{n \rightarrow \infty} \left\{ \int_{(0, k_n)_1^2} |f(y_\alpha, y_3, \xi_\alpha + \nabla_\alpha \varphi_n | \nabla_3 \varphi_n) - f^\infty(y_\alpha, y_3, \xi_\alpha + \nabla_\alpha \varphi_n | \nabla_3 \varphi_n)| dy \right. \\ & \quad \left. + \int_{(0, k_n)_1^2} \left| f^\infty(y_\alpha, y_3, \xi_\alpha + \nabla_\alpha \varphi_n | \nabla_3 \varphi_n) - \frac{f(y_\alpha, y_3, t_n \xi_\alpha + t_n \nabla_\alpha \varphi_n | \nabla_3 \varphi_n)}{t_n} \right| dy \right\}. \end{aligned}$$

By (H4), (2.14), and the 1-homogeneity of f^∞ , we get

$$\begin{aligned} & T f_{hom}^0(s, \xi_\alpha) - T f_{hom}^{0, \infty}(s, \xi_\alpha) \\ & \leq \liminf_{n \rightarrow \infty} C \left\{ \int_{(0, k_n)_1^2} 1 + |(\xi_\alpha + \nabla_\alpha \varphi_n | \nabla_3 \varphi_n)|^{1-q} dy \right. \\ & \quad \left. + \int_{(0, k_n)_1^2} \frac{1 + t_n^{1-q} |(\xi_\alpha + \nabla_\alpha \varphi_n | \nabla_3 \varphi_n)|^{1-q}}{t_n} dy \right\}. \end{aligned}$$

Using Hölder inequality and (2.15), we get

$$T f_{hom}^0(s, \xi_\alpha) - T f_{hom}^{0, \infty}(s, \xi_\alpha) \leq C(1 + |\xi_\alpha|^{1-q}).$$

Now we have to prove the inverse inequality. First, fix $k \in \mathbb{N}$ and $\varphi \in W^{1, \infty}((0, k)_1^2; T_s(\mathcal{M}))$ such that $\varphi(x_\alpha, x_3) = 0$ for every $(x_\alpha, x_3) \in \partial(0, k)^2 \times (-\frac{1}{2}, \frac{1}{2})$, then from (H2), we have that for every $x \in (0, k)_1^2$ and for every $t > 0$

$$\frac{f(x, t(\xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi))}{t} \leq \beta(1 + |\xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi|),$$

so that

$$\frac{f(\cdot, t(\xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi))}{t} \in L^1((0, k)_1^2).$$

By the dominated convergence theorem, we get

$$T f_{hom}^{0, \infty}(s, \xi_\alpha) \leq \limsup_{t \rightarrow \infty} \int_{(0, k)_1} \frac{f(y, t(\xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi))}{t} dy \leq \int_{(0, k)_1} f^\infty(y, \xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi) dy,$$

which implies

$$Tf_{hom}^{0,\infty}(s, \xi_\alpha) \leq T(f^\infty)_{hom}^0(s, \xi_\alpha). \tag{2.16}$$

Now, fix $\eta > 0$ and let $\varphi \in W^{1,\infty}((0, k)_{,1}^2; T_s(\mathcal{M}))$ such that $\varphi(x_\alpha, x_3) = 0$ for every $(x_\alpha, x_3) \in \partial(0, k)^2 \times (-\frac{1}{2}, \frac{1}{2})$ such that

$$\int_{(0,k)_{,1}^2} f(y_\alpha, y_3, \xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi) dy \leq Tf_{hom}^0(s, \xi_\alpha) + \eta.$$

As before, by (H2), we have

$$\int_{(0,k)_{,1}^2} |\nabla \varphi| \leq C(1 + |\xi_\alpha|),$$

so from (2.16) and (H4), we obtain

$$\begin{aligned} Tf_{hom}^{0,\infty}(s, \xi_\alpha) - Tf_{hom}^0(s, \xi_\alpha) &\leq T(f^\infty)_{hom}^0(s, \xi_\alpha) - Tf_{hom}^0(s, \xi_\alpha) \\ &\leq \int_{(0,k)_{,1}^2} |f^\infty(y_\alpha, y_3, \xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi) - f(y_\alpha, y_3, \xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi)| dy + \eta \\ &\leq C \int_{(0,k)_{,1}^2} 1 + |(\xi_\alpha + \nabla_\alpha \varphi | \nabla_3 \varphi)|^{1-q} dy + \eta. \end{aligned}$$

By Hölder inequality, we get

$$Tf_{hom}^{0,\infty}(s, \xi_\alpha) - Tf_{hom}^0(s, \xi_\alpha) \leq C(1 + |\xi|^{1-q}) + \eta.$$

By the arbitrariness of η this gives the desired inequality. □

2.2. Jump part density

Now we prove some properties for the density θ defined in (1.8). To this aim, we introduce some further notations. We remark the fact that the results below are valid for any dimension while reducing any number of variables. We present here the main steps of the proof in dimension 3 reducing the third variable. We also observe that when the number of “surviving” dimensions is 2 (as in our case) the proof could be significantly simplified. In the remainder of this section there is an abuse of notation which will not generate contradiction with (1.8) in view of the results we are going to show.

Given $\nu = \{\nu_1, \nu_2\}$ an orthonormal basis of \mathbb{R}^2 and $(a, b) \in \mathcal{M} \times \mathcal{M}$, we denote by

$$Q_\nu := \{\lambda_1 \nu_1 + \lambda_2 \nu_2 : \lambda_1, \lambda_2 \in (-1/2, 1/2)\}$$

and, for $x_\alpha \in \mathbb{R}^2$, we set

$$\|x_\alpha\|_{\nu, \infty} := \sup_{i=1,2} |x_\alpha \cdot \nu_i|, \quad x_\nu := x_\alpha \cdot \nu_1, \quad x' := x_\alpha \cdot \nu_2, \tag{2.17}$$

so that x_α can be identified with the pair (x_ν, x') . We also point out that this last definition of Q_ν is a slight abuse of notation, indeed at the beginning of this section Q_ν has been introduced as the unit cube of \mathbb{R}^2 , centered at the origin, and with two sides orthogonal to ν , (with $\nu \in \mathbb{S}^1$). Here $\nu = \{\nu_1, \nu_2\}$ is an orthonormal basis of \mathbb{R}^2 , but the notation Q_ν is still consistent since by definition $Q_{\nu_1} = Q_{\nu_2} = Q_\nu$.

Let $u_{a,b,\nu} : \mathbb{R}^2 \times (-1/2, 1/2) \rightarrow \mathcal{M}$ be the function defined by

$$u_{a,b,\nu} := \begin{cases} a, & \text{if } x_\nu > 0, \\ b, & \text{if } x_\nu \leq 0. \end{cases}$$

For every $t, h > 0$, we also introduce the following classes of functions

$$\begin{aligned} \mathcal{A}_t(a, b, \nu) &:= \{ \phi \in W^{1,1}((tQ_\nu)_{,1}; \mathcal{M}) : \phi = u_{a,b,\nu} \text{ on } \partial(tQ_\nu) \times (-1/2, 1/2) \}, \\ \mathcal{B}_h(a, b, \nu) &:= \left\{ u \in W^{1,1}(Q_{\nu,1}; \mathcal{M}) : u(x) = \gamma \left(\frac{x_\nu}{h} \right) \text{ on } \partial Q_\nu \times (-1/2, 1/2) \text{ for some } \gamma \in \mathcal{G}(a, b) \right\}, \end{aligned} \tag{2.18}$$

where \mathcal{G} is the set introduced in (2.1).

Proposition 2.5. *Let $f : \mathbb{R}^3 \times \mathbb{R}^{3 \times 3} \rightarrow \mathbb{R}$ be a Carathéodory function satisfying (H1)–(H4). For every $(a, b, \nu_1) \in \mathcal{M} \times \mathcal{M} \times \mathbb{S}^1$, there exists*

$$\theta(a, b, \nu_1) := \liminf_{t \rightarrow +\infty} \inf_{\phi} \left\{ \frac{1}{t} \int_{(tQ_\nu)_{,1}} f^\infty(y, \nabla \phi) dy : \phi \in \mathcal{A}_t(a, b, \nu) \right\} \tag{2.19}$$

$$= \liminf_{h \rightarrow 0} \inf_u \left\{ \int_{Q_{\nu,1}} f^\infty \left(\frac{x_\alpha}{h}, x_3, \nabla_h u \right) dx : u \in \mathcal{B}_h(a, b, \nu) \right\}, \tag{2.20}$$

where $\nu = (\nu_1, \nu_2)$ is any orthonormal base of \mathbb{R}^2 with first element ν_1 , i.e., the limit is dependent only on ν_1 .

Proof. The proof of Proposition 2.5 follows in a similar way as in [11], which in turn is very much inspired by [13, Proposition 2.2] passing through the introduction of an appropriate rescaled surface energy density. We present the main steps. First we prove that, for every $(a, b) \in \mathcal{M} \times \mathcal{M}$ and any orthonormal basis ν , the limit in (2.20) exists (see [11, Proposition 3.3]), then we prove it also coincides with (2.19), i.e., a result analogous to [11, Proposition 3.2] holds. It follows that the limit in (2.20) exists and it is the same for any two ν, ν' are orthonormal rational bases (i.e., there exist scalars $\gamma_1, \gamma'_1, \gamma_2, \gamma'_2 \in \mathbb{R} \setminus \{0\}$ such that $\gamma_1 \nu_1, \gamma'_1 \nu'_1, \gamma_2 \nu_2$, and $\gamma'_2 \nu'_2 \in \mathbb{Z}^2$), with equal first vector $\nu_1 = \nu'_1$. Then, we claim that for every $\sigma > 0$ there exists $\delta > 0$ (independent of a and b) such that if ν and ν' are two orthonormal bases of \mathbb{R}^2 with $|\nu_i - \nu'_i| < \delta$ for every $i = 1, 2$, then

$$\liminf_{h \rightarrow 0} I_h(\nu) - K\sigma \leq \liminf_{h \rightarrow 0} I_h(\nu') \leq \limsup_{h \rightarrow 0} I_h(\nu') \leq \limsup_{h \rightarrow 0} I_h(\nu) + K\sigma, \tag{2.21}$$

where K is a positive constant which only depends on \mathcal{M}, β (see (1.1)) and

$$I_h(\nu) := I_h(a, b, \nu) := \inf \left\{ \int_{Q_{\nu,1}} f^\infty \left(\frac{x_\alpha}{h}, x_3, \nabla_h u \right) dx : u \in \mathcal{B}_h(a, b, \nu) \right\}.$$

Let $Q_{\nu,\eta} := (1 - \eta)Q_\nu$ where $0 < \eta < 1$. Let $\sigma > 0$ be fixed and let $0 < \eta < 1$ be such that $\eta < \frac{1}{34}$ and

$$\max \left\{ \eta, \frac{(1 - \eta)(1 - 2\eta)}{(1 - 3\eta)} - (1 - 2\eta) \right\} < \sigma. \tag{2.22}$$

Consider $\delta_0 > 0$ (that may be chosen so that $\delta_0 \leq \frac{\eta}{2\sqrt{2}}$). Then take any $0 < \delta \leq \delta_0$ and any pair v and v' of orthonormal basis of \mathbb{R}^2 satisfying $|v_i - v'_i| \leq \delta$ for $i = 1, 2$ such that one has

$$Q_{v,3\eta} \subset Q_{v',2\eta} \subset Q_{v,\eta} \tag{2.23}$$

and $\{x \cdot v'_1 = 0\} \cap \partial Q_{v,\eta} \subset \{|x \cdot v_1| \leq 1/8\}$. Given $h > 0$ small, we consider $u_h \in \mathcal{B}_h(a, b, v')$ such that

$$\int_{Q_{v',1}} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h u_h\right) dx \leq I_h(v') + \sigma,$$

where $u_h(x) = u_h(x_\alpha, x_3) = \gamma_h\left(\frac{x_{v'}}{h}\right)$ for some $\gamma_h \in \mathcal{G}(a, b)$ and for $x \in \partial Q_{v'} \times (-\frac{1}{2}, \frac{1}{2})$. Then one can construct $v_h \in \mathcal{B}_{(1-2\eta)h}(a, b, v)$ satisfying the boundary condition $v_h(x) = \gamma_h\left(\frac{x_v}{(1-2\eta)h}\right)$ for $x \in \partial Q_v \times (-1/2, 1/2)$. Consider $F_\eta : \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$F_\eta(x_\alpha) := \left(\frac{1 - 2\|x'\|_{v,\infty}}{\eta}\right) \frac{x_{v'}}{1 - 2\eta} + \left(\frac{\eta - 1 + 2\|x'\|_{v,\infty}}{\eta}\right) \frac{x_v}{1 - 2\eta},$$

and define

$$v_h(x) = v_h(x_\alpha, x_3) := \begin{cases} u_h\left(\frac{x_\alpha}{1-2\eta}, x_3\right), & \text{if } x \in Q_{v',2\eta} \times (-1/2, 1/2), \\ \gamma_h\left(\frac{x_{v'}}{(1-2\eta)h}\right), & \text{if } x \in (Q_{v,\eta} \setminus Q_{v',2\eta}) \times (-1/2, 1/2), \\ a, & \text{if } x \in (Q_v \setminus Q_{v,\eta}) \times (-1/2, 1/2) \text{ and } x_v \geq \frac{1}{4}, \\ \gamma_h\left(\frac{F_\eta(x_\alpha)}{h}\right), & \text{if } x \in A_\eta \cap (Q_v \setminus Q_{v,\eta}) \times (-1/2, 1/2), \\ b, & \text{if } x \in (Q_v \setminus Q_{v,\eta}) \times (-1/2, 1/2) \text{ and } x_v \leq \frac{-1}{4}, \end{cases}$$

and $A_\eta := \{x_\alpha : |x_v| \leq 1/4\}$.

One can check that v_h is well defined for h small enough and that $v_h \in \mathcal{B}_{(1-2\eta)h}(a, b, v)$. Therefore

$$\begin{aligned} I_{(1-2\eta)h}(v) &\leq \int_{Q_{v,1}} f^\infty\left(\frac{x_\alpha}{(1-2\eta)h}, x_3, \nabla_h v_h\right) dx \\ &= \int_{Q_{v',2\eta,1}} f^\infty\left(\frac{x_\alpha}{(1-2\eta)h}, x_3, \nabla_h v_h\right) dx \\ &\quad + \int_{(Q_{v,\eta} \setminus Q_{v',2\eta}) \times (-1/2, 1/2)} f^\infty\left(\frac{x_\alpha}{(1-2\eta)h}, x_3, \nabla_h v_h\right) dx \\ &\quad + \int_{A_\eta \times (-1/2, 1/2)} f^\infty\left(\frac{x_\alpha}{(1-2\eta)h}, x_3, \nabla_h v_h\right) dx \\ &=: I_1 + I_2 + I_3. \end{aligned} \tag{2.24}$$

We now estimate each of the three integrals in (2.24). First, we easily get that

$$I_1 = (1 - 2\eta) \int_{Q_{v',1}} f^\infty\left(\frac{y_\alpha}{h}, x_3, \nabla_h u_h\right) dy_\alpha dx_3 \leq I_h(v') + \sigma. \tag{2.25}$$

In view of (2.23), we have

$$Q_{v,\eta} \subset (1 - \eta)(1 - 2\eta)(1 - 3\eta)^{-1} Q_{v'} =: D_\eta.$$

Then we infer from (1.3) and Fubini's theorem, that

$$\begin{aligned}
 I_2 &\leq \beta \int_{(D_\eta \setminus \mathcal{Q}_{v', 2\eta}) \times (-1/2, 1/2)} |\nabla_h v_h| dx & (2.26) \\
 &= \frac{\beta}{(1-2\eta)h} \int_{((D_\eta \setminus \mathcal{Q}_{v', 2\eta}) \cap \{|x_v| \leq (1-2\eta)h/2\}) \times (-1/2, 1/2)} \left| \dot{\gamma}_h \left(\frac{x_v}{(1-2\eta)h} \right) \right| dx_\alpha dx_3 \\
 &= \beta \mathcal{H}^1((D_\eta \setminus \mathcal{Q}_{v', 2\eta}) \cap \{x_{v'} = 0\}) \frac{1}{(1-2\eta)h} \int_{-(1-2\eta)h/2}^{(1-2\eta)h/2} \left| \dot{\gamma}_h \left(\frac{t}{(1-2\eta)h} \right) \right| dt \\
 &= \beta \text{dist}_{\mathcal{M}}(a, b) \left(\frac{(1-\eta)(1-2\eta)}{(1-3\eta)} - (1-2\eta) \right).
 \end{aligned}$$

Now we estimate I_3 . In order to do so, we observe that from (2.23), we get

$$\|\nabla_\alpha F_\eta(x_\alpha)\|_{L^\infty(A_\eta, \mathbb{R}^2)} \leq C \quad (2.27)$$

for some absolute constant $C > 0$, and

$$|\nabla_\alpha F_\eta(x_\alpha) \cdot v_1| \geq 1, \quad (2.28)$$

for a.e. $x_\alpha \in A_\eta$. Combining (1.3), (2.27), and (2.28), we get

$$\begin{aligned}
 I_3 &\leq \beta \int_{A_\eta} |\nabla_h v_h| dx_\alpha \leq \frac{C\beta}{h} \int_{A_\eta} \left| \dot{\gamma}_h \left(\frac{F_\eta(x_\alpha)}{h} \right) \right| dx_\alpha \\
 &\leq \frac{C\beta}{h} \int_{A_\eta} \left| \dot{\gamma}_h \left(\frac{F_\eta(x_\alpha)}{h} \right) \right| |\nabla_\alpha F_\eta(x_\alpha) \cdot v_1| dx_\alpha \\
 &= C\beta \int_{A'_\eta} \frac{1}{h} \int_{-1/4}^{1/4} \left| \dot{\gamma}_h \left(\frac{F_\eta(tv_1 + x')}{h} \right) \right| |\nabla_\alpha F_\eta(tv_1 + x') \cdot v_1| dt d\mathcal{H}^1(x'),
 \end{aligned}$$

where we used Fubini's theorem in the last equality and $A'_\eta := A_\eta \cap \{x_v = 0\}$. By the change of variable $s = (1/h)F_\eta(tv_1 + x')$ it follows that for \mathcal{H}^1 -a.e. $x' \in A_\eta$

$$\int_{-1/4}^{1/4} \left| \dot{\gamma}_h \left(\frac{F_\eta(tv_1 + x')}{h} \right) \right| |\nabla_\alpha F_\eta(tv_1 + x') \cdot v_1| dt \leq \int_{\mathbb{R}} |\dot{\gamma}_h(s)| ds = \text{dist}_{\mathcal{M}}(a, b).$$

Combining the previous two inequalities we get

$$I_3 \leq C\beta \mathcal{H}^1(A'_\eta) \text{dist}_{\mathcal{M}}(a, b) = C\beta \eta \text{dist}_{\mathcal{M}}(a, b) \quad (2.29)$$

By (2.22), (2.25), (2.26), and (2.29), we get

$$I_{(1-2\eta)h}(v) \leq I_h(v') + K\sigma,$$

with $K = 1 + \beta\delta(1 + C)$, δ diameter of \mathcal{M} and C the constant from (2.27). Passing to the limit $h \rightarrow 0$ and by the arbitrariness of v and v' , we get (2.21). Arguing as in [13, Proposition 2.2, Step 4], we get that for any orthonormal basis v and v' with first element $v_1 \in \mathbb{S}^1$, we get the existence of the limit of I_h for every v and

$$\lim_{h \rightarrow 0} I_h(v) = \lim_{h \rightarrow 0} I_h(v'). \quad (2.30)$$

Now we prove the equality between (2.19) and (2.20). Fix $h > 0$ and an orthonormal basis $\nu = (\nu_1, \nu_2)$ of \mathbb{R}^2 . Let

$$J_h(\nu) = J_h(a, b, \nu) := \inf \left\{ \int_{Q_\nu \times (-1/2, 1/2)} f^\infty \left(\frac{x_\alpha}{h}, x_3, \nabla_h u \right) dx : u \in \mathcal{A}_1(a, b, \nu) \right\},$$

by a change of variable we have also that

$$J_h(\nu) = \inf \left\{ h \int_{\frac{1}{h} Q_\nu \times (-1/2, 1/2)} f^\infty(x_\alpha, x_3, \nabla u) dx : u \in \mathcal{A}_{1/h}(a, b, \nu) \right\}.$$

We want to prove that

$$\lim_{h \rightarrow 0} J_h(\nu) = \lim_{h \rightarrow 0} I_h(\nu). \tag{2.31}$$

For h small enough, set $\tilde{h} := h/(1 - h)$ and consider $u_{\tilde{h}} \in \mathcal{B}_{\tilde{h}}(a, b, \nu)$ such that

$$\int_{Q_\nu \times (-1/2, 1/2)} f^\infty \left(\frac{x_\alpha}{\tilde{h}}, x_3, \nabla_{\tilde{h}} u_{\tilde{h}} \right) dx \leq I_{\tilde{h}}(\nu) + h,$$

with $u_{\tilde{h}}(x) = \gamma_{\tilde{h}}(x_\nu/\tilde{h})$ in $\partial Q_\nu \times (-1/2, 1/2)$, for some $\gamma_{\tilde{h}} \in \mathcal{G}(a, b)$. Now, for every $x \in Q_\nu \times (-1/2, 1/2)$, we define

$$v_h(x) := \begin{cases} u_{\tilde{h}} \left(\frac{x_\alpha}{1-h}, x_3 \right), & \text{if } x \in Q_{\nu, h}, \\ \gamma_{\tilde{h}} \left(\frac{x_\nu}{1-2\|x'\|_{\nu, \infty}} \right), & \text{otherwise.} \end{cases}$$

It follows that

$$\begin{aligned} J_h(\nu) &\leq \int_{Q_\nu \times (-1/2, 1/2)} f^\infty \left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h \right) dx \\ &\leq \int_{Q_{\nu, h} \times (-1/2, 1/2)} f^\infty \left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h \right) dx + \int_{(Q_\nu \setminus Q_{\nu, h}) \times (-1/2, 1/2)} f^\infty \left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h \right) dx \\ &= I_1 + I_2. \end{aligned} \tag{2.32}$$

Now we estimate I_1 and I_2 . For I_1 we have that by construction

$$I_1 = (1 - h) \int_{Q_\nu \times (-1/2, 1/2)} f^\infty \left(\frac{x_\alpha}{\tilde{h}}, x_3, \nabla_{\tilde{h}} u_{\tilde{h}} \right) dx \leq (1 - h)(I_{\tilde{h}}(\nu) + h). \tag{2.33}$$

Now we turn to I_2 .

$$\begin{aligned} I_2 &\leq \beta \int_{(Q_\nu \setminus Q_{\nu, h}) \times (-1/2, 1/2)} \left| \dot{\gamma}_{\tilde{h}} \left(\frac{x_\nu}{1-2\|x'\|_{\nu, \infty}} \right) \right| \left(\frac{1}{1-2\|x'\|_{\nu, \infty}} + \frac{|x_\nu| \|\nabla_\alpha(\|x'\|_{\nu, \infty})\|}{(1-2\|x'\|_{\nu, \infty})^2} \right) dx \\ &\leq 2\beta \int_{(Q_\nu \setminus Q_{\nu, h}) \cap \{x_\nu \leq (1-2\|x'\|_{\nu, \infty})/2\}} \left| \dot{\gamma}_{\tilde{h}} \left(\frac{x_\nu}{1-2\|x'\|_{\nu, \infty}} \right) \right| \left(\frac{1}{1-2\|x'\|_{\nu, \infty}} \right) dx_\alpha \end{aligned}$$

where, in order, we used the growth condition of f^∞ , the fact that $\dot{\gamma}_{\tilde{h}} \left(\frac{x_\nu}{1-2\|x'\|_{\nu, \infty}} \right) = 0$ on the complementary set of $\{x_\nu \leq (1 - 2\|x'\|_{\nu, \infty})/2\}$, and the fact that

$$\|\nabla_\alpha(\|x'\|_{\nu, \infty})\|_{L^\infty(Q_\nu, \mathbb{R}^2)} \leq 1.$$

Denoting $Q'_v := Q_v \cap \{x_v = 0\}$ and $Q'_{v,h} := Q_{v,h} \cap \{x_v = 0\}$, then by Fubini's theorem, we get

$$\begin{aligned} I_2 &\leq 2\beta \int_{(Q'_v \setminus Q'_{v,h})} \int_{-(1-2\|x'\|_{v,\infty})/2}^{(1-2\|x'\|_{v,\infty})/2} \left| \dot{\gamma}_{\tilde{h}} \left(\frac{t}{1-2\|x'\|_{v,\infty}} \right) \right| \left(\frac{1}{1-2\|x'\|_{v,\infty}} \right) dt dx' \\ &\leq 2\beta \text{dist}_{\mathcal{M}}(a, b) \mathcal{H}^1(Q'_v \setminus Q'_{v,h}) \\ &\leq 2\beta \text{dist}_{\mathcal{M}}(a, b) h, \end{aligned} \quad (2.34)$$

and so from (2.32)–(2.34) and (2.30), we infer that

$$\limsup_{h \rightarrow 0} J_h(v) \leq \lim_{h \rightarrow 0} I_h(v).$$

On the other hand, if we fix $h > 0$ small enough and $\tilde{u}_h \in \mathcal{A}_1(a, b, v)$ such that

$$\int_{Q_v \times (-1/2, 1/2)} f^\infty \left(\frac{x_\alpha}{h}, x_3, \nabla_h \tilde{u}_h \right) dx \leq J_h(v) + h,$$

and we denote for $x \in Q_v \times (-1/2, 1/2)$,

$$w_h(x) := \begin{cases} \tilde{u}_h \left(\frac{x_\alpha}{1-h}, x_3 \right), & \text{if } x_\alpha \in Q_{v,h}, \\ \gamma \left(\frac{x_v}{(2\|x'\|_{v,\infty} - 1 + h)(1-h)} \right), & \text{otherwise,} \end{cases}$$

with $\gamma \in \mathcal{G}(a, b)$, then it is easy to check that $w_h \in \mathcal{B}_{(1-h)h}(a, b, v)$. Arguing as in the previous estimates we get that

$$I_{(1-h)h}(v) \leq (1-h)(J_h(v) + h) + 2\beta h \text{dist}_{\mathcal{M}}(a, b),$$

which leads to

$$\liminf_{h \rightarrow 0} J_h(v) \geq \lim_{h \rightarrow 0} I_h(v),$$

and this concludes the proof. \square

Proposition 2.6. *The function θ defined in (1.8) is continuous on $\mathcal{M} \times \mathcal{M} \times \mathbb{S}^1$ and there exist constants $C_1, C_2 > 0$ such that*

$$|\theta(a_1, b_1, v_1) - \theta(a_2, b_2, v_1)| \leq C_1(|a_1 - a_2| + |b_1 - b_2|), \quad (2.35)$$

$$\theta(a_1, b_1, v_1) \leq C_2|a_1 - b_1|, \quad (2.36)$$

for every $a_1, b_1, a_2, b_2 \in \mathcal{M}$ and $v_1 \in \mathbb{S}^1$.

Proof. It follows in a similar way to [11, Proposition 3.4], taking into account similar modifications as we did in Proposition 2.5. For the reader's convenience, we present a short proof. For shortness sake, we use the same notation as in the previous proof. From the previous proof it is clear that $\theta(a, b, \cdot)$ is uniformly continuous on \mathbb{S}^1 so it is enough to prove (2.35) in order to prove the continuity of θ on $\mathcal{M} \times \mathcal{M} \times \mathbb{S}^1$, and thus we start from there. Fix $v_1 \in \mathbb{S}^1$ and let $v = (v_1, v_2)$ be an orthonormal base of

\mathbb{R}^2 . For $h > 0$, we denote $\tilde{h} := h/(1 - h)$ and we consider $\gamma_{\tilde{h}} \in \mathcal{G}(a_1, b_1)$ and $u_{\tilde{h}} \in \mathcal{B}_{\tilde{h}}(a_1, b_1, \nu)$ such that $u_{\tilde{h}}(x) = \gamma_{\tilde{h}}(\frac{x_\nu}{\tilde{h}})$ on $(\partial Q_\nu)_1$ and

$$\int_{Q_{\nu,1}} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_{\tilde{h}} u_{\tilde{h}}\right) dx \leq I_{\tilde{h}}(a_1, b_1, \nu) + h.$$

As in the previous proof, now we construct a function $v_h \in \mathcal{A}_1(a_2, b_2, \nu)$ starting from $u_{\tilde{h}}$. Fix $\gamma_a \in \mathcal{G}(a_2, a_1)$, $\gamma_b \in \mathcal{G}(b_2, b_1)$, and define

$$v_h(x) := \begin{cases} u_{\tilde{h}}\left(\frac{x_\alpha}{1-h}, x_3\right), & x \in Q_{\nu,h} \times (-1/2, 1/2), \\ \gamma_{\tilde{h}}\left(\frac{x_\nu}{1-2\|x'\|_{\nu,\infty}}\right), & x \in A_1 \times (-1/2, 1/2), \\ \gamma_a\left(\frac{2\|x_\alpha\|_{\nu,\infty}-1}{h} + \frac{1}{2}\right), & x \in A_2 \times (-1/2, 1/2), \\ \gamma_b\left(\frac{2\|x_\alpha\|_{\nu,\infty}-1}{h} + \frac{1}{2}\right), & x \in A_3 \times (-1/2, 1/2), \\ \gamma_a\left(\frac{2\|x'\|_{\nu,\infty}-1}{2x_\nu} + \frac{1}{2}\right) & x \in A_4 \times (-1/2, 1/2), \\ \gamma_b\left(\frac{1-2\|x'\|_{\nu,\infty}}{2x_\nu} + \frac{1}{2}\right), & x \in A_5 \times (-1/2, 1/2), \end{cases}$$

with

$$\begin{aligned} A_1 &:= \left\{ \frac{1-h}{2} \leq \|x'\|_{\nu,\infty} < \frac{1}{2} \text{ and } |x_\nu| \leq -\|x'\|_{\nu,\infty} + \frac{1}{2} \right\}, \\ A_2 &:= (Q_\nu \setminus Q_{\nu,h}) \cap \{x_\nu \geq h/2\}, \\ A_3 &:= (Q_\nu \setminus Q_{\nu,h}) \cap \{x_\nu \leq -h/2\}, \\ A_4 &:= \left\{ 0 < x_\nu \leq \frac{h}{2}, \frac{1}{2} - x_\nu \leq \|x'\|_{\nu,\infty} < \frac{1}{2} \right\}, \\ A_5 &:= \left\{ -\frac{h}{2} < x_\nu \leq 0, \frac{1}{2} + x_\nu \leq \|x'\|_{\nu,\infty} < \frac{1}{2} \right\}. \end{aligned}$$

Since by construction $v_h \in \mathcal{A}_1(a_2, b_2, \nu)$, then

$$J_h(a_2, b_2, \nu) \leq \int_{Q_{\nu,1}} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h\right) dx.$$

By the same argument as in the proof of Proposition 2.5, we get

$$\int_{Q_{\nu,1}} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h\right) dx \leq I_{\tilde{h}}(a_1, b_1, \nu) + h,$$

and

$$\int_{A_1 \times (-1/2, 1/2)} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h\right) dx \leq C \text{dist}_{\mathcal{M}}(a_1, b_1)h,$$

where $\text{dist}_{\mathcal{M}}$ denotes the geodesic distance on \mathcal{M} . Now we estimate the integrals over A_2 and A_4 . The integrals on A_3 and A_5 follow in a similar way. We start with A_2 . First, we define

$$F_h(x_\alpha) := \frac{2\|x_\alpha\|_{\nu,\infty} - 1}{h} + \frac{1}{2}.$$

From the linear growth of f^∞ , Fubini's theorem, the fact that $A_2 \subset F_h^{-1}([-1/2, 1/2])$, the Coarea formula, and the fact that v_h does not depend on x_3 on A_2 , we get

$$\begin{aligned} \int_{A_2 \times (-1/2, 1/2)} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h\right) dx &\leq \beta \int_{A_2} |\dot{\gamma}_a(F_h(x_\alpha))| |\nabla_\alpha F_h(x_\alpha)| dx_\alpha \\ &\leq \beta \int_{F_h^{-1}([-1/2, 1/2])} |\dot{\gamma}_a(F_h(x_\alpha))| |\nabla_\alpha F_h(x_\alpha)| dx_\alpha \\ &\leq \beta \int_{(-1/2, 1/2)} |\dot{\gamma}_a(t)| \mathcal{H}^1(F_h^{-1}(t)) dt. \end{aligned}$$

Since $F_h^{-1}(t) = \partial Q_{v, \frac{h(1-2t)}{2}}$ for every $t \in (-1/2, 1/2)$, then it holds $\mathcal{H}^1(F_h^{-1}(t)) \leq \mathcal{H}^1(\partial Q_v) = 2$, which gives us

$$\int_{A_2 \times (-1/2, 1/2)} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h\right) dx \leq 2\beta \text{dist}_M(a_1, a_2).$$

Now we estimate the integral on A_4 . We define

$$G_h(x_\alpha) := \frac{2\|x'\|_{v, \infty} - 1}{2x_v} + \frac{1}{2}.$$

As before, by the linear growth of f^∞ and Fubini's theorem, we get

$$\begin{aligned} \int_{A_4 \times (-1/2, 1/2)} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h\right) dx \\ \leq \beta \int_{(0, h/2)} \int_{G^{-1}(\cdot, x_v)([-1/2, 1/2])} |\dot{\gamma}_a(G(x', x_v))| |\nabla_\alpha G(x', x_v)| dx' dx_v. \end{aligned}$$

Since $|\nabla_{x'} G(x_\alpha)| = 1/x_v$ and $|\nabla_{x_v} G(x_\alpha)| \leq 1/x_v$ for a.e. $x_\alpha \in A_4$, we get that $|\nabla_\alpha G(x_\alpha)| \leq 2|\nabla_{x'} G(x_\alpha)|$ and thus

$$\begin{aligned} \int_{A_4 \times (-1/2, 1/2)} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h\right) dx \\ \leq 2\beta \int_{(0, h/2)} \int_{G^{-1}(\cdot, x_v)([-1/2, 1/2])} |\dot{\gamma}_a(G(x', x_v))| |\nabla_{x'} G(x', x_v)| dx' dx_v. \end{aligned}$$

By construction, $G(\cdot, x_v)$ is Lipschitz for every $x_v \in (0, h/2)$ and so by the Coarea formula we get again that

$$\begin{aligned} \int_{A_4 \times (-1/2, 1/2)} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h\right) dx \\ \leq 2\beta \int_{(0, h/2)} \int_{(-1/2, 1/2)} |\dot{\gamma}_a(t)| \mathcal{H}^0(\{x' : G(x', x_v) = t\}) dt dx_v \\ \leq C h \text{dist}_M(a_1, a_2), \end{aligned}$$

where we used the fact that $\mathcal{H}^0(\{x' : G(x', x_v) = t\}) = 1$ for every $t \in (-1/2, 1/2)$. Estimating the integrals over A_3 and A_5 in a similar way we get

$$J_h(a_2, b_2, v) \leq \int_{Q_{v,1}} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h v_h\right) dx \leq I_{\bar{h}}(a_1, b_1, v) + C(h + \text{dist}_M(a_1, a_2) + \text{dist}_M(b_1, b_2)).$$

Passing to the limit for $h \rightarrow 0$ we get

$$\theta(a_1, b_1, \nu) \leq \theta(a_2, b_2, \nu) + C(\text{dist}_{\mathcal{M}}(a_1, a_2) + \text{dist}_{\mathcal{M}}(b_1, b_2)),$$

since $\text{dist}_{\mathcal{M}}$ is equivalent to the Euclidean distance, and by the arbitrariness of a_1, a_2, b_1 , and b_2 in \mathcal{M} we get (2.35).

Now we prove (2.36). Fix an orthonormal basis $\nu = (\nu_1, \nu_2)$ of \mathbb{R}^2 and let $\gamma \in \mathcal{G}(a_1, b_1)$. Consider $u_h(x) = \gamma(x_\nu/h)$. By construction, $u_h \in \mathcal{B}_h(a_1, b_1, \nu)$. Moreover, since $\lim_{h \rightarrow 0} J_h(\nu) = \lim_{h \rightarrow 0} I_h(\nu)$ it holds

$$\begin{aligned} \theta(a_1, b_1, \nu) &\leq \liminf_{h \rightarrow 0} \int_{Q_{\nu,1}} f^\infty\left(\frac{x_\alpha}{h}, x_3, \nabla_h u_h\right) dx \\ &\leq \liminf_{h \rightarrow 0} \frac{\beta}{h} \int_{Q_\nu} \left| \dot{\gamma}\left(\frac{x_\alpha \cdot \nu}{h}\right) \right| dx_\alpha \\ &\leq \beta \text{dist}_{\mathcal{M}}(a_1, b_1). \end{aligned}$$

Again, by the equivalence of $\text{dist}_{\mathcal{M}}$ with the Euclidean distance we get (2.36). \square

In order to prove Theorem 1.1, we need to introduce the auxiliary functional I^0 defined as

$$I^0(u, A) := \inf \left\{ \liminf_{n \rightarrow +\infty} I^{h_n}(u_n, A) : u_n \rightarrow u \text{ in } L^1(A_{,1}; \mathcal{M}) \right\}, \quad (2.37)$$

for any $u \in BV(\omega; \mathcal{M})$ and $A \in \mathcal{A}(\omega)$.

Proposition 2.7. *There exists a function $K : \mathcal{M} \times \mathcal{M} \times \mathbb{S}^1 \rightarrow [0, +\infty)$ continuous in the last variable and such that*

- (i) $K(a, b, \nu) = K(b, a, -\nu)$ for every $(a, b, \nu) \in \mathcal{M} \times \mathcal{M} \times \mathbb{S}^1$,
- (ii) for every finite subset T of \mathcal{M} ,

$$I^0(u, S) = \int_S K(u^+, u^-, \nu_u) d\mathcal{H}^1, \quad (2.38)$$

for every $u \in BV(\omega; T)$ and every Borel subset S of $\omega \cap S_u$.

Proof. Let T be a finite subset of \mathcal{M} and $\omega \subset \mathbb{R}^2$ open and bounded. For every $A \in \mathcal{A}(\omega)$ and $u \in BV(\omega; T)$ we define

$$G_T(u, A) := I^0(u, S_u \cap A),$$

where $I^0(u, S_u \cap A)$ is defined as in (2.37).

We aim to show that the assumptions of [40, Theorem 3.1] are satisfied by $G_T : BV(\omega; T) \times \mathcal{A}(\omega) \rightarrow [0, +\infty)$, namely:

- (1) $0 \leq G_T(u, A) \leq \Lambda \mathcal{H}^1(A \cap S_u)$ with a fixed constant $\Lambda \in \mathbb{R}$;
- (2) $G_T(u, \cdot)$ is the restriction to $\mathcal{A}(\omega)$ of a Borel measure on ω ;
- (3) $G_T(u, A) = G_T(v, A)$ if $u = v$ a.e. in A ;
- (4) If $u_h \rightarrow u$ a.e. in A , then

$$G_T(u, A) \leq \liminf_{h \rightarrow +\infty} G_T(u_h, A);$$

(5) For every $y \in \mathbb{R}^2$, we have

$$G_T(\tau_y u, \tau_y A) = G_T(u, A),$$

where $(\tau_y u)(x) = u(x - y)$ and $\tau_y A = A + y$, provided $\tau_y A \subset \Omega$.

Property (1) and (3) follows immediately from the definition of I^0 . Property (2), follows from the fact the Borel measure $I^0(u, \cdot)$ on ω (see 3.1), restricted to S_u , is an extension of $G_T(u, \cdot)$.

For (4), if $u_h \rightarrow u$ a.e. on A , then the sequence (u_h) is equibounded in A . It follows that $(u_h)_h$ converges to u in $L^1(A; \mathcal{M})$. It not difficult to see that both u_h and u can be extended to Ω by taking the extension constant in the third variable, and so that $u_h \rightarrow u$ a.e. on $A_{,1}$ and $(u_h)_h$ converges to u in $L^1(A_{,1}; \mathcal{M})$. For every open set $E \subset A$ with $S_u \cap A \subset E$, we have

$$I^0(u, E) \leq \liminf_{h \rightarrow \infty} I^0(u_h, E).$$

Moreover, from (3.1), (H2), and the equiboundedness of $(u_h)_h$, we get

$$I^0(u_h, E) = I^0(u_h, S_{u_h} \cap E) + I^0(u_h, E \setminus S_{u_h}) \leq I^0(u_h, S_{u_h} \cap A) + C|E|.$$

Taking the infimum over $E \supset S_u \cap A$, we obtain

$$I^0(u, S_u \cap A) \leq \liminf_{h \rightarrow \infty} I^0(u_h, S_{u_h} \cap A).$$

Finally, property (5) follows from the translation invariance of I^h (see (H1) and (1.4)) with respect to the x_α -variable. We can now apply Theorem 3.1 in [40], from which follows the existence of a continuous function $K_\omega^T : \omega \times T \times T \times \mathbb{S}^1 \rightarrow [0, +\infty[$ such that $K_\omega^T(x_\alpha, a, b, \nu) = K_\omega^T(x_\alpha, b, a, -\nu)$ and

$$I^0(u, S_u \cap A) = G_T(u, A) = \int_{S_u \cap A} K_\omega^T(x_\alpha, u^+, u^-, \nu_u) d\mathcal{H}^1$$

for every $u \in BV(\omega; T)$ and $A \in \mathcal{A}(\omega)$. For $x_\alpha, x_0 \in \mathbb{R}^2$, $a, b \in \mathcal{M}$ and $\nu \in \mathbb{S}^1$ define

$$u_{x_0, \nu}^{a, b}(x) := \begin{cases} a & \text{if } (x_\alpha - x_0) \cdot \nu_1 \geq 0, \\ b & \text{if } (x_\alpha - x_0) \cdot \nu_1 < 0, \end{cases} \quad \Pi_{x_0, \nu_1} := \{x_\alpha \in \mathbb{R}^2 : (x_\alpha - x_0) \cdot \nu_1 = 0\}.$$

The continuity of K_ω^T implies that

$$K_\omega^T(x_0, a, b, \nu) = \lim_{\rho \rightarrow 0} \frac{I^0(u_{x_0, \nu_1}^{a, b}(x), B_\rho(x_0) \cap \Pi_{x_0, \nu_1})}{\mathcal{H}^1(B_\rho(x_0) \cap \Pi_{x_0, \nu_1})},$$

for every $(x_0, a, b, \nu) \in \omega \times T \times T \times \mathbb{S}^1$. By the arbitrariness of T and ω it follows that K_ω^T is independent from T and ω , i.e., $K_\omega^T = K$. Exploiting the translation invariance of I_0 and by a blow-up argument it is possible to show that K does not depend from the x_α -variable (for details see [13, Proposition 4.1]). Since $I^0(u, \cdot)$ restricted to S_u is a regular Borel measure, it follows the integral representation on the Borel subsets of S_u and this concludes the proof. \square

3. Proof of Theorem 1.1

In this section we prove Theorem 1.1. In particular, we assume f to satisfy hypothesis (H1)–(H4). The first step in order to prove Theorem 1.1 is to localize I^0 .

Lemma 3.1 (Localization). *Given $u \in BV(\omega; \mathcal{M})$ the set function $I^0(u, \cdot) : \mathcal{A}(\omega) \rightarrow \mathbb{R}$ is the restriction to $\mathcal{A}(\omega)$ of a Radon measure absolutely continuous with respect to $\mathcal{L}^2 + |Du|$.*

Proof. Fix $u \in BV(\omega; \mathcal{M})$ and $A \in \mathcal{A}(\omega)$. By [41, Theorem 3.9] there exists a sequence $(u'_n)_n \subset W^{1,1}(A; \mathbb{R}^3) \cap C^\infty(A; \mathbb{R}^3)$ such that $u'_n \rightarrow u$ in $L^1(A; \mathbb{R}^3)$ and $\|\nabla_\alpha u'_n\|_{L^1(A; \mathbb{R}^3)} \rightarrow |D_\alpha u|(A)$. So, if we consider $u_n(x) := u'_n(x_\alpha)$, we have that $(u_n)_n \subset W^{1,1}(A_{,1}; \mathbb{R}^3) \cap C^\infty(A_{,1}; \mathbb{R}^3)$ and $\|\nabla_{h_n} u_n\|_{L^1(A_{,1}; \mathbb{R}^3)} \rightarrow |D_\alpha u|(A)$. By construction, $u_n(x) \in co(\mathcal{M})$ for a.e. $x \in A_{,1}$ and every $n \in \mathbb{N}$, where $co(\mathcal{M})$ stands for the convex envelope of \mathcal{M} . Following the argument of [11, Proposition 2.1], there exists a sequence $(w_n)_n \subset W^{1,1}(A_{,1}; \mathcal{M})$ satisfying

$$\int_{A_{,1}} |\nabla_{h_n} w_n| dx \leq C^* \int_{A_{,1}} |\nabla_{h_n} u_n| dx,$$

for some constant C^* depending only on \mathcal{M} . Moreover, we have $w_n \rightarrow u$ in $L^1(A_{,1}; \mathbb{R}^3)$ and by the growth condition (H2) we have

$$I^0(u, A) \leq \beta(\mathcal{L}^2(A) + C^*|Du|(A)).$$

Now we just have to prove that

$$I^0(u, A) \leq I^0(u, B) + I^0(u, A \setminus \bar{C}),$$

for any $A, B, C \in \mathcal{A}(\omega)$ with $\bar{C} \subset B \subset A$. The proof of the previous inequality follows from a standard slicing argument similar to [28, Lemma 3.2]. \square

Now we prove the limsup inequality.

Proposition 3.2. *For every $u \in BV(\omega; \mathcal{M})$ it holds*

$$I^0(u) \leq I(u),$$

with I defined in (1.5) and I^0 defined in (2.37).

Proof. For every $u \in BV(\omega; \mathcal{M})$, we can write

$$I^0(u, \omega) = I^0(u, \omega \setminus S_u) + I^0(u, S_u), \quad (3.1)$$

where S_u denotes the jump set of u . It follows that in order to prove our claim we can separately estimate the two terms on the right hand side of (3.1). First, by [28, Proposition 3.3], we have for every $A \in \mathcal{A}(\omega)$

$$I^0(u, A) \leq \begin{cases} \int_A T f_{\text{hom}}^0(u, \nabla_\alpha u) dx, & \text{if } u \in W^{1,1}(\omega; \mathcal{M}), \\ +\infty, & \text{otherwise in } L^1(\omega; \mathbb{R}^3). \end{cases}$$

By means of a relaxation argument (see [11, Theorem 7.1] which holds by Proposition 2.4) and exploiting the coercivity of f (see H2), a standard diagonalization argument, and the lower semicontinuity of I^0 with respect to $L^1(\Omega, \mathbb{R}^3)$ strong convergence we get

$$I^0(u, A) \leq \int_A T f_{\text{hom}}^0(u, \nabla_\alpha u) dx_\alpha + \int_{S_u \cap A} H(u^+, u^-, \nu_u) d\mathcal{H}^1 + \int_A T f_{\text{hom}}^{0, \infty} \left(\tilde{u}, \frac{dD_\alpha^c u}{d|D_\alpha^c u|} \right) d|D_\alpha^c u| \quad (3.2)$$

for some function $H : \mathcal{M} \times \mathcal{M} \times \mathbb{S}^1 \rightarrow [0 + \infty)$. By outer regularity, in (3.2) we can choose $A = \omega \setminus S_u$, which leads to

$$I^0(u, \omega \setminus S_u) \leq \int_A T f_{\text{hom}}^0(u, \nabla_\alpha u) dx_\alpha + \int_A T f_{\text{hom}}^{0, \infty} \left(\tilde{u}, \frac{dD_\alpha^c u}{d|D_\alpha^c u|} \right) d|D_\alpha^c u|.$$

Now we estimate $I^0(u, S_u)$. By Proposition 2.7, we have

$$I^0(u, S) = \int_S K(u^+, u^-, \nu_u) d\mathcal{H}^1,$$

for every $u \in BV(\omega; T)$ and every Borel subset S of $\omega \cap S_u$, so to conclude it is enough to prove that

$$K(a, b, \nu) \leq \theta(a, b, \nu)$$

for every $(a, b, \nu) \in \mathcal{M} \times \mathcal{M} \times \mathbb{S}^1$.

Since the proof is similar to Proposition 2.5 we refer to it for the notation. Let $\nu = (\nu_1, \nu_2)$ be an orthonormal basis of \mathbb{R}^2 . Since K and θ are continuous functions in the \mathbb{S}^1 variable, we can assume that ν is a rational basis, i.e., for all $i \in \{1, 2\}$, there exists $\gamma_i \in \mathbb{R} \setminus \{0\}$ such that $\nu_i := \gamma_i \nu_i \in \mathbb{Z}^2$. The general case, then, follows by a density argument. Given an arbitrary $0 < \eta < 1$, by Proposition 2.5 and (2.31), there exists $h_0 > 0$, $u_0 \in \mathcal{B}_{h_0}(a, b, \nu)$ and $\gamma_{h_0} \in \mathcal{G}(a, b)$ such that

$$u_0(x) = \gamma_{h_0} \left(\frac{x_\alpha \cdot \nu_1}{h_0} \right) \quad \text{and} \quad \int_{Q_{\nu,1}} f^\infty \left(\frac{x_\alpha}{h_0}, x_3, \nabla_{h_0} u_0 \right) dx \leq \theta(a, b, \nu_1) + \eta.$$

Given $\lambda \in \mathbb{Z}$, let $(h_n)_n$ be a vanishing positive sequence and set

$$x_n^{(\lambda)} := h_n \lambda \nu_2, \quad Q_{\nu, n}^{(\lambda)} := x_n^{(\lambda)} + \left(\frac{h_n}{h_0} \right) Q_{\nu,1}.$$

Next, we define

$$\Lambda_n := \left\{ \lambda \in \mathbb{Z} : Q_{\nu, n}^{(\lambda)} \subset Q_{\nu,1} \text{ and } \frac{x_n^{(\lambda)}}{h_n} \in l \left(\frac{1}{h_0} + \gamma_2 \right) \nu_2 + P \text{ for some } l \in \mathbb{Z} \right\},$$

with

$$P := \{\alpha \nu_2 : \alpha \in [-1/2, 1/2)\}.$$

Now we consider the sequence of functions

$$u_n(x_\alpha, x_3) := \begin{cases} u_0 \left(\frac{h_0}{h_n} (x_\alpha - x_n^{(\lambda)}) \cdot \nu_1 \right), & \text{if } x \in Q_{\nu, n}^{(\lambda)} \text{ for some } \lambda \in \Lambda_n, \\ \gamma_{h_0} \left(\frac{x_\alpha \cdot \nu_1}{h_0} \right), & \text{otherwise.} \end{cases}$$

By construction, $u_n \in W^{1,1}(Q_{v_1}; \mathcal{M})$, $(\nabla_{h_n} u_n)_n$ is bounded in $L^1(Q_{v_1}; \mathbb{R}^{3 \times 2})$, and $u_n \rightarrow u_{v_1}^{a,b}$ in $L^1(Q_{v_1}; \mathbb{R}^3)$ as $n \rightarrow +\infty$ with

$$u_{v_1}^{a,b}(x) := \begin{cases} a, & \text{if } x_\alpha \cdot v_1 \geq 0, \\ b, & \text{if } x_\alpha \cdot v_1 < 0. \end{cases}$$

By a similar argument as in Step 1 of the proof of [13, Proposition 2.2], we get that

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \int_{Q_{v_1}} f^\infty\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} u_n\right) dx &\leq \int_{Q_{v_1}} f^\infty\left(\frac{x_\alpha}{h_0}, x_3, \nabla_{h_0} u_0\right) dx \\ &\leq \theta(a, b, v_1) + \eta. \end{aligned} \tag{3.3}$$

For $\rho > 0$, let $A_\rho := Q_{v_1} \cap \{|x_\alpha \cdot v_1| < \rho\}_1$ and set $\Pi_{v_1} := \{x_\alpha \in \mathbb{R}^2 : x_\alpha \cdot v_1 = 0\}$. By construction, the sequence $(u_n)_n$ is admissible for $I^0(u_{v_1}^{a,b}, A_\eta)$, so that

$$\begin{aligned} I^0(u_{v_1}^{a,b}, A_\eta \cap \Pi_{v_1}) &\leq I^0(u_{v_1}^{a,b}, A_\eta) \leq \liminf_{n \rightarrow +\infty} \int_{A_{\eta,1}} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} u_n\right) dx \\ &\leq \beta \mathcal{L}^2(A_\eta) + \liminf_{n \rightarrow +\infty} \int_{A_{h_n,1}} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} u_n\right) dx \\ &\leq \liminf_{n \rightarrow +\infty} \int_{A_{h_n,1}} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} u_n\right) dx + \beta \eta. \end{aligned} \tag{3.4}$$

Here we used assumption (H2) and the fact that $\nabla_{h_n} u_n = 0$ outside $A_{h_n,1}$. On the other hand, by (2.38) we have

$$I^0(u_{v_1}^{a,b}, A_\eta \cap \Pi_{v_1}) = \int_{A_\eta \cap \Pi_{v_1}} K(a, b, v_1) d\mathcal{H}^1 = K(a, b, v_1). \tag{3.5}$$

Using (H4), the boundedness of $(\nabla_{h_n} u_n)_n$ in $L^1(Q_{v_1}; \mathbb{R}^{3 \times 2})$, the fact that $f^\infty(\cdot, 0) \equiv 0$, and Hölder’s inequality, we derive

$$\begin{aligned} &\left| \int_{A_{h_n,1}} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} u_n\right) dx - \int_{Q_{v_1}} f^\infty\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} u_n\right) dx \right| \\ &\leq C \int_{A_{h_n,1}} (1 + |\nabla_{h_n} u_n|^{1-q}) dx \leq C \left(h_n + h_n^q \|\nabla_{h_n} u_n\|_{L^1(Q_{v_1}; \mathbb{R}^{3 \times 2})}^{1-q} \right) \rightarrow 0. \end{aligned} \tag{3.6}$$

From (3.3)–(3.6), we get

$$K(a, b, v_1) \leq \theta(a, b, v_1) + (1 + \beta)\eta,$$

and from the arbitrariness of η we get the desired inequality. By exploiting Propositions 2.6, 2.7, and Lemma 3.1 we can argue as in [11, Corollary 5.1], which, in turn rely on the approximation arguments in [42, Lemma 4.7 and Proposition 4.8], and together with (3.1) we get

$$\begin{aligned} I^0(u) &\leq \int_\omega T f_{\text{hom}}^0(u, \nabla_\alpha u) dx_\alpha \\ &\quad + \int_{S_u \cap \omega} \theta(u^+, u^-, v_u) d\mathcal{H}^1 + \int_\omega T f_{\text{hom}}^{0,\infty}\left(\tilde{u}, \frac{dD_\alpha^c u}{d|D_\alpha^c u|}\right) d|D_\alpha^c u| = I(u), \end{aligned}$$

where I is defined in (1.5), which yields the conclusion. □

This concludes the proof of the upper bound. We now prove the lower bound using the blow up method.

Proposition 3.3. *For every $u \in BV(\omega, \mathcal{M})$ it holds*

$$I^0(u) \geq I(u),$$

with I defined in (1.5) and I^0 defined in (2.37).

Proof. Consider $u \in BV(\omega; \mathcal{M})$ and let $(u_n)_n \subset W^{1,1}(\Omega; \mathcal{M})$ be such that

$$I^0(u, \omega) = \lim_{n \rightarrow +\infty} \int_{\Omega} f\left(\frac{x_{\alpha}}{h_n}, x_3, \nabla_{h_n} u_n\right) dx.$$

Define the sequence of nonnegative Radon measures

$$\mu_n := \left(\int_{(-\frac{1}{2}, \frac{1}{2})} f\left(\frac{(\cdot)_{\alpha}}{h_n}, x_3, \nabla_{h_n} u_n((\cdot)_{\alpha}, x_3)\right) dx_3 \right) \mathcal{L}^2 \llcorner \omega.$$

Possibly extracting a subsequence, we can infer the existence of a nonnegative Radon measure $\mu \in \mathcal{M}(\omega)$ such that $\mu_n \xrightarrow{*} \mu$ in $\mathcal{M}(\omega)$. Exploiting the Besicovitch differentiation theorem, we are able to split μ into the sum of three nonnegative measure which are mutually singular, namely

$$\mu = \mu^a + \mu^j + \mu^c,$$

where

$$\mu^a \ll \mathcal{L}^2, \quad \mu^j \ll \mathcal{H}^1 \llcorner S_u \quad \mu^c \ll |D_{\alpha}^c u|.$$

As long as $\mu(\omega) \leq I^0(u, \omega)$, it is sufficient to check that

$$\frac{d\mu}{d\mathcal{L}^2}(x_0) \geq T_{f_{\text{hom}}^0}(u(x_0), \nabla_{\alpha} u(x_0)) \quad \text{for } \mathcal{L}^2\text{-a.e. } x_0 \in \omega, \tag{3.7}$$

$$\frac{d\mu}{d|D_{\alpha}^c u|}(x_0) \geq T_{f_{\text{hom}}^{0,\infty}}\left(\tilde{u}(x_0), \frac{dD_{\alpha}^c u}{d|D_{\alpha}^c u|}(x_0)\right) \quad \text{for } |D_{\alpha}^c u|\text{-a.e. } x_0 \in \omega, \tag{3.8}$$

and

$$\frac{d\mu}{d\mathcal{H}^1 \llcorner S_u}(x_0) \geq \theta(u^+(x_0), u^-(x_0), \nu_u(x_0)) \quad \text{for } \mathcal{H}^1\text{-a.e. } x_0 \in S_u. \tag{3.9}$$

Proof of (3.7). Before proving the result we recall that all the operations of sum and difference between the functions $u_n, v_n, w_k, w_{n,k}$ and u must be intended in the sense of Remark 2.2.

Step 1. Consider $x_0 \in \omega$ to be a Lebesgue point of u and $\nabla_{\alpha} u$, a point of approximate differentiability of u , in a way that $u(x_0) \in \mathcal{M}, \nabla_{\alpha} u(x_0) \in [T_{u(x_0)}(\mathcal{M})]^2$ and such that the Radon-Nykodým derivative of μ with respect to the Lebesgue measure \mathcal{L}^2 exists and it is finite. We observe that \mathcal{L}^2 -almost every points $x_0 \in \omega$ satisfy these properties. We set

$$s_0 := u(x_0) \quad \xi_0 := \nabla_{\alpha} u(x_0).$$

We may assume that, up to a subsequence, there exists a nonnegative Radon measure $\lambda \in \mathcal{M}(\omega)$ such that

$$\left(\int_{-\frac{1}{2}}^{\frac{1}{2}} (1 + |\nabla_{h_n} u_n|) dx_3 \right) \mathcal{L}^2 \llcorner \omega \xrightarrow{*} \lambda \quad \text{in } \mathcal{M}(\omega).$$

Let us consider a sequence $(\rho_k)_k \rightarrow 0^+$ such that $Q'(x_0, 2\rho_k) \subset \Omega$ and $\mu(\partial Q'(x_0, \rho_k)) = \lambda(\partial Q(x_0, \rho_k)) = 0$ for each $k \in \mathbb{N}$. Since μ is the limit of μ_n , we have that

$$\mu(Q'(x_0, \rho_k)) = \lim_{n \rightarrow +\infty} \int_{Q'(x_0, \rho_k)_1} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} u_n\right) dx. \quad (3.10)$$

Since

$$\tau_n := h_n \left\lfloor \frac{x_0}{h_n} \right\rfloor \in h_n \mathbb{Z}^2 \rightarrow x_0,$$

given $r \in (0, 1/2)$ we have that $Q'(\tau_n, \rho_k) \subset Q'(x_0, r\rho_k)$ for sufficiently large values of k ; therefore we can define, for $x \in Q'(0, \rho_k)_1$

$$v_n(x) := u_n(x_\alpha + \tau_n, x_3). \quad (3.11)$$

The continuity of the translation in L^1 allows us to obtain the following estimate, holding for $n \rightarrow +\infty$

$$\begin{aligned} \int_{Q'(0, \rho_k)_1} |v_n(x) - u(x_\alpha + x_0)| dx &= \int_{Q'(\tau_n, \rho_k)_1} |u_n(x) - u(x_\alpha + x_0 - \tau_n)| dx \\ &\leq \int_{Q'(x_0, r\rho_k)_1} |u_n(x) - u(x_\alpha + x_0 - \tau_n)| dx \rightarrow 0. \end{aligned} \quad (3.12)$$

A change of variable in (3.10) together with the periodicity condition (H1) of $f(\cdot, x_3, \xi)$, the growth condition (H2), and (3.11) lead to

$$\begin{aligned} \mu(Q'(x_0, \rho_k)) &= \lim_{n \rightarrow +\infty} \int_{Q'(x_0 - \tau_n, \rho_k)_1} f\left(\frac{x_\alpha + \tau_n}{h_n}, x_3, \nabla_{h_n} u_n(x_\alpha + \tau_n, x_3)\right) dx \\ &= \lim_{n \rightarrow +\infty} \int_{Q'(x_0 - \tau_n, \rho_k)_1} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} v_n\right) dx \\ &\geq \limsup_{n \rightarrow +\infty} \int_{Q'(0, \rho_k)_1} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} v_n\right) dx \\ &\quad - \beta \limsup_{n \rightarrow +\infty} \int_{Q_1(\tau_n, \rho_k)_1 \setminus Q'(x_0, \rho_k)_1} (1 + |\nabla_{h_n} u_n|) dx. \end{aligned} \quad (3.13)$$

The last term in (3.13) vanishes since, by the choice of ρ_k

$$\begin{aligned} &\limsup_{n \rightarrow +\infty} \int_{Q'(\tau_n, \rho_k)_1 \setminus Q'(x_0, \rho_k)_1} (1 + |\nabla_{h_n} u_n|) dx \\ &\leq \limsup_{r \rightarrow 1^+} \limsup_{n \rightarrow +\infty} \int_{Q'(x_0, r\rho_k)_1 \setminus Q'(x_0, \rho_k)_1} (1 + |\nabla_{h_n} u_n|) dx \\ &\leq \limsup_{r \rightarrow 1^+} \lambda\left(\overline{Q'(x_0, r\rho_k)_1} \setminus Q'(x_0, \rho_k)_1\right) \\ &\leq \lambda(\partial Q'(x_0, \rho_k)_1) = 0. \end{aligned}$$

Summing up, this entails

$$\mu(Q'(x_0, \rho_k)) \geq \limsup_{n \rightarrow +\infty} \int_{Q'(0, \rho_k)_1} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} v_n\right) dx,$$

where $(v_n)_n \subset W^{1,1}(Q'(0, \rho_k), 1; \mathcal{M})$ is such that $v_n \rightarrow u(x_0 + (\cdot)_\alpha)$ in $L^1(Q'(0, \rho_k), 1; \mathbb{R}^3)$ thanks to (3.12). At this point, let us consider, for every n a sequence $(v_{n,j})_j \subset C^\infty(Q'(0, \rho_k), 1; \mathbb{R}^3)$ (not necessarily taking values in \mathcal{M} !) such that $v_{n,j} \rightarrow v_n$ in $W^{1,1}(Q'(0, \rho_k), 1; \mathbb{R}^3)$, $v_{n,j} \rightarrow v_n$ and $\nabla v_{n,j} \rightarrow \nabla v_n$ a.e. in $Q'(0, \rho_k), 1$ as $j \rightarrow +\infty$ and consider the function g introduced in (2.7). It is possible to prove that

$$\begin{aligned} & \lim_{j \rightarrow +\infty} \int_{Q'(0, \rho_k), 1} g\left(\frac{x_\alpha}{h_n}, x_3, v_{n,j}, \nabla_{h_n} v_{n,j}\right) dx \\ &= \int_{Q'(0, \rho_k), 1} g\left(\frac{x}{h_n}, x_3, v_n, \nabla_{h_n} v_n\right) dx \\ &= \int_{Q'(0, \rho_k), 1} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} v_n\right) dx, \end{aligned}$$

therefore, by possibly passing to a diagonal sequence $\bar{v}_n := v_{n,j_n}$, such that $\bar{v}_n \rightarrow u(x_0 + (\cdot)_\alpha)$ in $L^1(Q'(0, \rho_k), 1; \mathbb{R}^3)$, we may deduce

$$\mu(Q'(x_0, \rho_k)) \geq \limsup_{n \rightarrow +\infty} \int_{Q'(0, \rho_k), 1} g\left(\frac{x_\alpha}{h_n}, x_3, \bar{v}_n, \nabla_{h_n} \bar{v}_n\right) dx.$$

If we perform a change of variable in the previous inequality and set

$$w_{n,k}(x) := \frac{\bar{v}(\rho_k x_\alpha, x_3) - s_0}{\rho_k},$$

we may deduce

$$\begin{aligned} \frac{d\mu}{d\mathcal{L}^2}(x_0) &\geq \limsup_{k \rightarrow +\infty} \limsup_{n \rightarrow +\infty} \int_Q g\left(\frac{\rho_k x_\alpha}{h_n}, x_3, \bar{v}_n(\rho_k x_\alpha, x_3), \nabla_{\frac{\rho_k}{h_n}} \bar{v}_n(\rho_k x_\alpha, x_3)\right) dx \\ &= \limsup_{k \rightarrow +\infty} \limsup_{n \rightarrow +\infty} \int_Q g\left(\frac{\rho_k x_\alpha}{h_n}, x_3, s_0 + \rho_k w_{n,k}, \nabla_{\frac{\rho_k}{h_n}} w_{n,k}\right) dx. \end{aligned} \tag{3.14}$$

Since x_0 is a point of approximate differentiability of u and $\bar{v}_n \rightarrow u(x_0 + (\cdot)_\alpha)$ in $L^1(Q'(0, \rho_k), 1; \mathbb{R}^3)$, we have

$$\lim_{k \rightarrow +\infty} \lim_{n \rightarrow +\infty} \int_Q [w_{n,k}(x) - \xi_0 x_\alpha] dx = \lim_{k \rightarrow +\infty} \int_{Q'(x_0, \rho_k)} \frac{|u(x_\alpha) - s_0 - \xi_0(x_\alpha - x_0)|}{\rho_k^3} dx_\alpha = 0. \tag{3.15}$$

At this point, (3.14) and (3.15) allow us to take a diagonal sequence $h_{n_k} < \rho_k^2$ in such a way that $w_k := w_{n_k, k} \rightarrow w_0$ in $L^1(Q; \mathbb{R}^d)$ with $w_0(x) := \xi_0 x_\alpha$ and

$$\frac{d\mu}{d\mathcal{L}^2}(x_0) \geq \limsup_{k \rightarrow +\infty} \int_Q g\left(\frac{x}{\delta_k}, x_3, s_0 + \rho_k w_k, \nabla_{\delta_k} w_k\right) dx,$$

where $\delta_k := \frac{h_{n_k}}{\rho_k} \rightarrow 0$.

Step 2. This second step can be performed following the lines of [12, Lemma 5.2], which, in turn, rely on [43, Theorem 2.1]. In fact we observe that the rescaled gradient can be seen as the gradient of a composed function. Moreover, the dependence on x_3 and the 'homogenization procedure' in the first

two variables and the dimension reduction with respect to the third one do not influence the strategy and the estimates, which rely only on the growth of f .

Thus we can show the existence of a sequence $(\bar{w}_k)_k \subset W^{1,\infty}(Q; \mathbb{R}^3)$ such that $\bar{w}_k \rightarrow w_0$ in $L^\infty(Q; \mathbb{R}^3)$, \bar{w}_k is uniformly bounded in $W^{1,1}(Q; \mathbb{R}^3)$ and $\nabla_{\delta_k} \bar{w}_k$ is bounded in $L^1(Q; \mathbb{R}^{3 \times 3})$.

$$\frac{d\mu}{d\mathcal{L}^2}(x_0) \geq \limsup_{k \rightarrow +\infty} \int_Q g\left(\frac{x_\alpha}{\delta_k}, x_3, s_0 + \rho_k \bar{w}_k, \nabla_{\delta_k} \bar{w}_k\right) dx. \quad (3.16)$$

Step 3. Thanks to the fact that $(\bar{w}_k)_k \subset L^\infty(Q; \mathbb{R}^3)$ and $(\nabla_{\delta_k} \bar{w}_k)_k \subset L^\infty(Q; \mathbb{R}^{3 \times 3})$ are uniformly bounded in L^1 , (2.9) gives

$$\lim_{k \rightarrow +\infty} \int_Q \left| g\left(\frac{x_\alpha}{\delta_k}, x_3, s_0 + \rho_k \bar{w}_k, \nabla_{\delta_k} \bar{w}_k\right) - g\left(\frac{x_\alpha}{\delta_k}, x_3, s_0, \nabla_{\delta_k} \bar{w}_k\right) \right| dx = 0,$$

which in turn entails, together with (3.16)

$$\frac{d\mu}{d\mathcal{L}^2}(x_0) \geq \limsup_{k \rightarrow +\infty} \int_Q g\left(\frac{x_\alpha}{\delta_k}, x_3, s_0, \nabla_{\delta_k} \bar{w}_k\right) dx.$$

Now we claim that

$$\limsup_{k \rightarrow +\infty} \int_Q g\left(\frac{x_\alpha}{\delta_k}, x_3, s_0, \nabla_{\delta_k} \bar{w}_k\right) dx = g_{\text{hom}}^0(s_0, \xi_0), \quad (3.17)$$

with the latter defined in (2.13). This concludes the proof since by (2.12) we have

$$\frac{d\mu}{d\mathcal{L}^2}(x_0) \geq g_{\text{hom}}^0(s_0, \xi_0) = T f_{\text{hom}}^0(s_0, \xi_0).$$

The proof of (3.17) is the following. Consider the family of integral functionals

$$(w, A) \in W^{1,1}(Q'; \mathbb{R}^3) \times \mathcal{A}(Q') \mapsto G^{\delta_k}(w, A) := \int_A \int_{-\frac{1}{2}}^{\frac{1}{2}} g\left(\frac{x_\alpha}{\delta_k}, x_3, s_0, \nabla_{\delta_k} w\right) dx_3 dx_\alpha,$$

for some vanishing sequence $(\delta_k)_k$. Let $G : BV(Q', \mathbb{R}^3) \times \mathcal{A}(Q') \rightarrow \mathbb{R}$ be

$$G(w, A) := \inf \left\{ \liminf_{k \rightarrow +\infty} G^{\delta_k}(w_k, A) : w_k \rightarrow w \text{ in } L^1(A; \mathcal{M}) \right\}.$$

It follows that G is L^1 lower semicontinuous. Moreover, since g satisfies hypothesis (H1) to (H4), then by the same argument as Lemma 3.1 we get that G is the restriction on $\mathcal{A}(Q')$ of a positive Radon measure. Furthermore, by Lemma 2.3-(ii) it is easy to see that

$$\alpha' |D_\alpha u|(A) \leq G(w, A) \leq \beta' (\mathcal{L}^2(A) + |D_\alpha u|(A)),$$

for every $(w, A) \in BV(Q'; \mathbb{R}^3) \times \mathcal{A}(Q')$. Standard arguments ensure that $G(w + c, A) = G(w, A)$ for any $c \in \mathbb{R}^3$ and $(w, A) \in BV(Q'; \mathbb{R}^3) \times \mathcal{A}(Q')$.

By Lemma 2.3-(i) and classical arguments in homogenization theory, it is possible to deduce that

$$G(\tau_y u, y + A) = G(u, A),$$

for any $A \in \mathcal{A}(Q')$, $y \in \mathbb{R}^2$ such that $y + A \subset Q'$, $u \in BV(Q'; \mathcal{M})$, and with $\tau_y w(x) := w(x - y)$. Combining the last to identities we get that

$$G(\tau_y w + c, y + A) = G(w, A).$$

Now we apply [6, Theorem 3.12, and Remark 3.8 (1) and (2)] and we obtain that

$$G(w, A) = \int_A \varphi^b(s_0, \nabla_\alpha w) dx_\alpha + \int_{S_w \cap A} \varphi^j(s_0, (w^- - w^+) \otimes \nu_w) d\mathcal{H}^1(x_\alpha) + \int_A \varphi^{b,\infty} \left(s_0, \frac{dD_\alpha^c w}{d|D_\alpha^c w|} \right) d|D_\alpha^c w|,$$

for every $(w, A) \in BV(Q'; \mathbb{R}^3) \times \mathcal{A}(Q')$. In what follow, we prove that, for every $\xi_\alpha \in \mathbb{R}^{3 \times 2}$,

$$\begin{aligned} \varphi^b(s_0, \xi_\alpha) &= \liminf_{t \rightarrow +\infty} \inf_{\varphi} \left\{ \int_{(tQ')_1} g(y, s_0, \xi_\alpha + \nabla_\alpha \varphi(y) | \nabla_3 \varphi(y)) dy : \right. \\ &\quad \left. \varphi \in W^{1,\infty}((tQ')_1; \mathbb{R}^3), \varphi(x_\alpha, x_3) = 0 \text{ for every } (x_\alpha, x_3) \in \partial(tQ') \times \left(-\frac{1}{2}, \frac{1}{2}\right) \right\} \\ &=: g_{\text{hom}}^0(s_0, \xi_\alpha), \end{aligned}$$

in (2.13). The proof of this identity follows the same argument as [5, pages 1390-1391], and reasoning as in [44, Proposition 2.4] we can substitute the space $W^{1,p}$ with $W^{1,\infty}$.

Proof of (3.8). For this proof we follow the strategy of [30, Section 5.1]. Let $G : BV(\omega, \mathbb{R}^3) \rightarrow \mathbb{R}$ be defined as

$$G(u) := \int_\omega g_{\text{hom}}^0(u, \nabla_\alpha u) dx_\alpha.$$

By (3.7) we have that for every $u \in BV(\omega, \mathcal{M})$ it holds

$$I^0(u) \geq G(u).$$

So, in particular, it holds

$$I^0(u) \geq \bar{G}(u), \quad u \in BV(\omega, \mathcal{M}),$$

with

$$\bar{G}(u) := \inf \left\{ \liminf_{n \rightarrow \infty} G(u_n) : u_n \rightharpoonup^* u \text{ in } BV(\omega, \mathbb{R}^3) \right\}.$$

For $\varepsilon > 0$, consider now the function $\tilde{g}(s, \xi_\alpha) : \mathbb{R}^3 \times \mathbb{R}^{3 \times 2} \rightarrow \mathbb{R}$ defined as

$$\tilde{g}_\varepsilon(s, \xi_\alpha) := g_{\text{hom}}^0(s, \xi_\alpha) + \varepsilon |\xi_\alpha|.$$

Fixed $u \in BV(\omega, \mathcal{M})$ and a sequence $(u_n)_n \subset BV(\omega, \mathcal{M})$ such that $u_n \rightharpoonup^* u$ in $BV(\omega, \mathcal{M})$, then in particular $u_n \rightharpoonup^* u$ in $BV(\omega, \mathbb{R}^3)$ and

$$\begin{aligned} \liminf_{n \rightarrow \infty} G(u_n) &\geq \liminf_{n \rightarrow \infty} \int_\omega \tilde{g}_\varepsilon(u_n, \nabla_\alpha u_n) dx_\alpha - \varepsilon \sup_n \|\nabla_\alpha u_n\|_{L^1(\omega, \mathbb{R}^{3 \times 2})} \\ &\geq \int_\omega \tilde{g}_\varepsilon(u, \nabla_\alpha u) dx_\alpha + \int_\omega \tilde{g}_\varepsilon^\infty \left(u, D_\alpha^c \frac{dD_\alpha^c u}{d|D_\alpha^c u|} \right) d|D_\alpha^c u| - \varepsilon \sup_n \|\nabla_\alpha u_n\|_{L^1(\omega, \mathbb{R}^{3 \times 2})}, \end{aligned}$$

where in the last inequality we used [45, Theorem 2.16]. Passing to the limit for $\varepsilon \rightarrow 0$, then considering that $\tilde{g}_\varepsilon^\infty(s, \xi_\alpha) = g_{hom}^{0,\infty}(s, \xi_\alpha) + \varepsilon|\xi_\alpha|$ we get

$$\begin{aligned} I^0(u) &\geq \int_\omega g_{hom}^0(u, \nabla_\alpha u) dx_\alpha + \int_\omega g_{hom}^{0,\infty}\left(u, \frac{dD_\alpha^c u}{d|D_\alpha^c u|}\right) d|D_\alpha^c u| \\ &= \int_\omega T f_{hom}^0(u, \nabla_\alpha u) dx_\alpha + \int_\omega T f_{hom}^{0,\infty}\left(u, \frac{dD_\alpha^c u}{d|D_\alpha^c u|}\right) d|D_\alpha^c u| \end{aligned}$$

wich proves (3.8), passing to densities via Besicovitch’s differentiation.

Proof of (3.9). In this step we use a strategy similar to the one employed in [11, Lemma 6.1]. In particular, we will still rely on the blow up method together with the projection argument in [11, Proposition 2.1] which we already used in the localization (see Lemma 3.1).

Step 1. Let $x_0 \in S_u$ be such that

$$\lim_{\rho \rightarrow 0^+} \int_{Q_{\nu_u(x_0)}^\pm(x_0, \rho)} |u(x_\alpha) - u^\pm(x_0)| dx_\alpha = 0, \tag{3.18}$$

where $u^\pm(x_0) \in \mathcal{M}$,

$$\lim_{\rho \rightarrow 0^+} \frac{\mathcal{H}^1(S_u \cap Q_{\nu_u(x_0)}(x_0, \rho))}{\rho} = 1, \tag{3.19}$$

and such that the Radon-Nikodým derivative of μ with respect to $\mathcal{H}^1 \llcorner S_u$ exists and it is finite. By [41, Lemma 2.1, Theorem 3.78, Theorem 2.83 (i)] with cubes instead of balls, it turns out that \mathcal{H}^1 -a.e. $x_0 \in S_u$ satisfy these properties. Now let us set $s_0^\pm := u^\pm(x_0)$ and $\nu_0 := \nu_u(x_0)$. Given a sequence $(u_n)_n \subset W^{1,1}(\Omega, \mathbb{R}^3)$ such that $\liminf_{n \rightarrow \infty} I^{h_n}(u_n) \leq c$, then, possibly extracting a further subsequence, we may assume that

$$\left(\int_{-\frac{1}{2}}^{\frac{1}{2}} (1 + |\nabla_{h_n} u_n|) dx_3 \right) \mathcal{L}^2 \llcorner \omega \xrightarrow{*} \lambda \quad \text{in } \mathcal{M}(\omega),$$

for some nonnegative Radon measure $\lambda \in \mathcal{M}(\omega)$. At this point, let us consider a sequence $\rho_k \rightarrow 0^+$ satisfying

$$\mu(\partial Q'_{\nu_0}(x_0, \rho_k)) = \lambda(\partial Q'_{\nu_0}(x_0, \rho_k)) = 0,$$

for each $k \in \mathbb{N}$. By (3.19), we have

$$\begin{aligned} \frac{d\mu}{d\mathcal{H}^1 \llcorner S_u}(x_0) &= \lim_{k \rightarrow +\infty} \frac{\mu(Q'_{\nu_0}(x_0, \rho_k))}{\mathcal{H}^1(S_u \cap Q'_{\nu_0}(x_0, \rho_k))} \\ &= \lim_{k \rightarrow +\infty} \frac{\mu(Q'_{\nu_0}(x_0, \rho_k))}{\rho_k} \\ &= \lim_{k \rightarrow +\infty} \lim_{n \rightarrow +\infty} \frac{1}{\rho_k} \int_{Q'_{\nu_0}(x_0, \rho_k)_1} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} u_n\right) dx. \end{aligned}$$

By exploiting [11, Theorem 2.1], it is possible to assume, without loss of generality, that $u_n \in \mathcal{D}(\Omega; \mathcal{M})$ (see Theorem 2.1) for each $n \in \mathbb{N}$. Therefore, arguing as in the proof of (3.7) (with $Q'_{\nu_0}(x_0, \rho_k)$

instead of $Q'(x_0, \rho_k)$ we get a sequence $(v_n)_n \subset \mathcal{D}((Q'_{v_0}(0, \rho_k))_1; \mathcal{M})$ such that $v_n \rightarrow u(x_0 + \cdot)$ in $L^1((Q'_{v_0}(0, \rho_k))_1; \mathbb{R}^3)$ as $n \rightarrow +\infty$ and

$$\frac{d\mu}{d\mathcal{H}^1 \llcorner S_u}(x_0) \geq \limsup_{k \rightarrow +\infty} \limsup_{n \rightarrow +\infty} \frac{1}{\rho_k} \int_{(Q'_{v_0}(0, \rho_k))_1} f\left(\frac{x_\alpha}{h_n}, x_3, \nabla_{h_n} v_n\right) dx.$$

It is worth mentioning that the construction process to get v_n starting from u_n does not affect the manifold constraint.

By means of a change of variable, setting $w_{n,k} = v_n(\rho_k x_\alpha, x_3)$ we get

$$\frac{d\mu}{d\mathcal{H}^1 \llcorner S_u}(x_0) \geq \limsup_{k \rightarrow +\infty} \limsup_{n \rightarrow +\infty} \rho_k \int_{Q'_{v_0,1}} f\left(\frac{\rho_k x_\alpha}{h_n}, x_3, \frac{1}{\rho_k} \nabla_\alpha w_{n,k} \left| \frac{1}{h_n} \nabla_3 w_{n,k} \right.\right) dx.$$

At this point, let us define

$$u_0(x) := \begin{cases} s_0^+, & \text{if } x \cdot v_0 > 0, \\ s_0^-, & \text{if } x \cdot v_0 \leq 0. \end{cases}$$

By using (3.18), we obtain

$$\lim_{k \rightarrow +\infty} \lim_{n \rightarrow +\infty} \int_{Q'_{v_0,1}} |w_{n,k} - u_0| dx = 0.$$

Exploiting a standard diagonal argument, we are able to find a sequence $n_k \rightarrow +\infty$ such that $(h_k)_k := (h_{n_k})_k$, $\delta_k := h_{n_k}/\rho_k \rightarrow 0$, $w_k := w_{n_k, k} \in \mathcal{D}(Q_{v_0,1}; \mathbb{R}^3)$ converges to $u_0 \in L^1(Q_{v_0,1}; \mathbb{R}^3)$ and

$$\begin{aligned} \frac{d\mu}{d\mathcal{H}^1 \llcorner S_u}(x_0) &\geq \limsup_{k \rightarrow +\infty} \rho_k \int_{Q'_{v_0,1}} f\left(\frac{x_\alpha}{\delta_k}, x_3, \frac{1}{\rho_k} \nabla_\alpha w_k \left| \frac{1}{h_k} \nabla_3 w_k \right.\right) dx \\ &= \limsup_{k \rightarrow +\infty} \rho_k \int_{Q'_{v_0,1}} f\left(\frac{x_\alpha}{\delta_k}, x_3, \frac{1}{\rho_k} \nabla_{\delta_k} w_k\right) dx. \end{aligned} \tag{3.20}$$

Exploiting (H4), the positive 1-homogeneity of $f^\infty(y, \cdot)$ and Hölder's inequality (being $0 < q < 1$)

$$\begin{aligned} &\int_{Q'_{v_0,1}} \left| \rho_k f\left(\frac{x_\alpha}{\delta_k}, x_3, \frac{1}{\rho_k} \nabla_{\delta_k} w_k\right) - f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} w_k\right) \right| dx \\ &= \int_{Q'_{v_0,1}} \left| \rho_k f\left(\frac{x_\alpha}{\delta_k}, x_3, \frac{1}{\rho_k} \nabla_{\delta_k} w_k\right) - \rho_k f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \frac{1}{\rho_k} \nabla_{\delta_k} w_k\right) \right| dx \\ &\leq C \rho_k \int_{Q'_{v_0,1}} (1 + \rho_k^{q-1} |\nabla_{\delta_k} w_k|^{1-q}) dx \\ &\leq C \left(\rho_k + \rho_k^q \|\nabla_{\delta_k} w_k\|_{L^1(Q'_{v_0,1}; \mathbb{R}^{3 \times 3})}^{1-q} \right). \end{aligned} \tag{3.21}$$

On the other hand, from (3.20) and the coercivity condition (H2), it follows that the sequence $(\nabla_{\delta_k} w_k)_k$ is uniformly bounded in $L^1(Q'_{v_0,1}; \mathbb{R}^{3 \times 2})$. Putting together (3.20) and (3.21), we obtain

$$\frac{d\mu}{d\mathcal{H}^1 \llcorner S_u}(x_0) \geq \limsup_{k \rightarrow +\infty} \int_{Q'_{v_0,1}} f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} w_k\right) dx. \tag{3.22}$$

Step 2. In this second and last step it remains to modify the value of w_k on a neighbourhood of $\partial Q'_{v_0}$, with the aim to get an admissible test function for the surface energy density. The computations follow similarly as in [30, Lemma 5.2].

We consider $\gamma \in \mathcal{G}(s_0^+, s_0^-)$ (see (2.1) and (2.17)) and set

$$\psi_k(x) := \gamma\left(\frac{x_{v_0}}{\delta_k}\right).$$

Using a De Giorgi type slicing argument, we shall modify w_k in order to get a function which matches ψ_k to $\partial Q'_{v_0,1}$. In order to reach this purpose, we introduce

$$\begin{aligned} r_k &:= \|w_k - \psi_k\|_{L^1(Q'_{v_0,1}; \mathbb{R}^3)}^{1/2}, \\ M_k &:= k \left[1 + \|w_k\|_{W^{1,1}(Q'_{v_0,1}; \mathbb{R}^3)} + \|\psi_k\|_{W^{1,1}(Q'_{v_0,1}; \mathbb{R}^3)} \right], \\ \ell_k &:= \frac{r_k}{M_k}. \end{aligned}$$

As long as ψ_k and w_k converge to u_0 in $L^1(Q'_{v_0,1}; \mathbb{R}^d)$, we have that $r_k \rightarrow 0$ and it is possible to assume that $0 < r_k < 1$. We set

$$Q_k^{(i)} := ((1 - r_k + i\ell_k)Q'_{v_0})_{,1} \quad \text{for } i = 0, \dots, M_k.$$

For every $i \in \{1, \dots, M_k\}$, we consider a cut-off functions $\varphi_k^{(i)} \in C_c^\infty((1 - r_k + i\ell_k)Q'_{v_0}; [0, 1])$ and we extend $\varphi_k^{(i)}$ to $Q_k^{(i)}$ setting $\varphi_k^{(i)}$ constant in the third variable. For simplicity of notation we still denote $\varphi_k^{(i)}$ this extension. By construction $\varphi_k^{(i)} = 1$ on $Q_k^{(i-1)}$ and $|\nabla_{\delta_k} \varphi_k^{(i)}| \leq c/\ell_k$. We further define

$$z_k^{(i)} := \varphi_k^{(i)} w_k + (1 - \varphi_k^{(i)}) \psi_k \in W^{1,1}(Q_{v_0,1}; \mathbb{R}^3),$$

in such a way that

$$z_k^{(i)} = w_k \quad \text{in } Q_k^{(i-1)} \quad \text{and} \quad z_k^{(i)} = \psi_k \quad \text{in } Q'_{v_0,1} \setminus Q_k^{(i)}.$$

Since $z_k^{(i)}$ is smooth outside a finite union of sets contained in some 1-dimensional submanifolds and $z_k^{(i)}(x) \in co(\mathcal{M})$ for a.e. $x \in Q'_{v_0,1}$, it is possible to apply a similar argument to [11, Proposition 2.1] to obtain new functions $\hat{z}_k^{(i)} \in W^{1,1}(Q'_{v_0,1}; \mathcal{M})$ such that

$$\hat{z}_k^{(i)} = z_k^{(i)} \quad \text{on } (Q'_{v_0,1} \setminus Q_k^{(i)}) \cup Q_k^{(i-1)}$$

and

$$\begin{aligned} \int_{Q_k^{(i)} \setminus Q_k^{(i-1)}} |\nabla_{\delta_k} \hat{z}_k^{(i)}| dx &\leq C_* \int_{Q_k^{(i)} \setminus Q_k^{(i-1)}} |\nabla_{\delta_k} z_k^{(i)}| dx \\ &\leq C_* \int_{Q_k^{(i)} \setminus Q_k^{(i-1)}} \left(|\nabla_{\delta_k} w_k| + |\nabla_{\delta_k} \psi_k| + \frac{1}{\ell_k} |w_k - \psi_k| \right) dx. \end{aligned}$$

In particular $\hat{z}_k^{(i)} \in \mathcal{B}_{\delta_k}(s_0^+, s_0^-, \nu_0)$ and by the growth condition (H2),

$$\begin{aligned} \int_{Q'_{\nu_0,1}} f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} \hat{z}_k^{(i)}\right) dx &\leq \int_{Q'_{\nu_0,1}} f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} w_k\right) dx \\ &+ C \int_{Q'_{\nu_0,1} \setminus Q_k^{(i)}} |\nabla_{\delta_k} \psi_k| dx \\ &+ C \int_{Q_k^{(i)} \setminus Q_k^{(i-1)}} \left(|\nabla_{\delta_k} w_k| + |\nabla_{\delta_k} \psi_k| + \frac{1}{\ell_k} |w_k - \psi_k| \right) dx. \end{aligned}$$

Summing up over all $i = 1, \dots, M_k$ and dividing by M_k , we get that

$$\begin{aligned} &\frac{1}{M_k} \sum_{i=1}^{M_k} \int_{Q'_{\nu_0,1}} f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} \hat{z}_k^{(i)}\right) dx \\ &\leq \int_{Q'_{\nu_0,1}} f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} w_k\right) dx \\ &+ C \int_{Q'_{\nu_0,1} \setminus Q_k^{(0)}} |\nabla_{\delta_k} \psi_k| dx + \frac{C}{k} + C \|w_k - \psi_k\|_{L^1(Q'_{\nu_0,1}; \mathbb{R}^d)}^{1/2}. \end{aligned}$$

Since

$$\int_{Q'_{\nu_0,1} \setminus Q_k^{(0)}} |\nabla \psi_k| dx \leq \mathbf{d}_M(s_0^+, s_0^-) \mathcal{H}^2((Q'_{\nu_0,1} \setminus Q_k^{(0)}) \cap \{x \cdot \nu = 0\}) \rightarrow 0$$

as $k \rightarrow +\infty$, there exists a sequence $\eta_k \rightarrow 0^+$ such that

$$\frac{1}{M_k} \sum_{i=1}^{M_k} \int_{Q'_{\nu_0,1}} f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} \hat{z}_k^{(i)}\right) dx \leq \int_{Q'_{\nu_0,1}} f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} w_k\right) dx + \eta_k.$$

Hence, for each $k \in \mathbb{N}$ it is possible to find some index $i_k \in \{1, \dots, M_k\}$ satisfying

$$\int_{Q'_{\nu_0,1}} f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} \hat{z}_k^{(i_k)}\right) dx \leq \int_{Q'_{\nu_0,1}} f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} w_k\right) dx + \eta_k. \tag{3.23}$$

Putting together (3.22) and (3.23), we finally obtain

$$\frac{d\mu}{d\mathcal{H}^1 \llcorner S_u}(x_0) \geq \limsup_{k \rightarrow +\infty} \int_{Q'_{\nu_0,1}} f^\infty\left(\frac{x_\alpha}{\delta_k}, x_3, \nabla_{\delta_k} \hat{z}_k^{(i_k)}\right) dx.$$

Since $\hat{z}_k^{(i_k)} \in \mathcal{B}_{\delta_k}(s_0^+, s_0^-, \nu_0)$ (see (2.18) for definition), we infer by Proposition 2.5 that

$$\frac{d\mu}{d\mathcal{H}^1 \llcorner S_u}(x_0) \geq \theta(s_0^+, s_0^-, \nu_0)$$

which finally concludes the proof. □

From the proofs of Theorem 1.1, it should be clear that both results hold in dimensions $N \geq 3$. In particular, we can assume $N, k, d \in \mathbb{N}$ with the $1 \leq k < N$, $\omega \subset \mathbb{R}^{N-k}$, $\Omega := \omega \times (-\frac{1}{2}, \frac{1}{2})^k$, and \mathcal{M} submanifold of \mathbb{R}^d .

4. Conclusions

In this paper, we have established a Γ -convergence result describing the simultaneous homogenization and 3D–2D dimension reduction for functionals with linear growth in the setting of manifold-valued maps. The obtained integral representation highlights the interplay between oscillations at the microscale, geometric constraints, and the emergence of lower-dimensional effective energies in the BV framework.

Our results extend previous analyses in the unconstrained or superlinear setting to the more delicate case of linear growth and manifold constraints, providing explicit formulas for the limiting densities. This contributes to a better understanding of variational models arising in applications such as micromagnetics and thin structures.

Possible directions for future research include the study of more general growth conditions, non-periodic settings, or the extension to higher codimension reductions and more complex geometrical constraints.

Use of Generative-AI tools declaration

The authors declare that they have not used Artificial Intelligence (AI) tools in the creation of this article.

Acknowledgments

The authors gratefully acknowledge support from INdAM GNAMPA. A.T. and L. L. have been partially supported through the INdAM-GNAMPA 2025 project “Minimal surfaces: the Plateau problem and behind” cup E5324001950001 and the project: Geometric-Analytic Methods for PDEs and Applications (GAMPA), ref. 2022SLTHCE – cup E53D2300588 0006 - funded by European Union - Next Generation EU within the PRIN 2022 program (D.D. 104 - 02/02/2022 Ministero dell’Università e della Ricerca). This manuscript reflects only the authors’ views and opinions and the Ministry cannot be considered responsible for them. E. Z. has been supported by PRIN 2022: Mathematical Modelling of Heterogeneous Systems (MMHS) - Next Generation EU CUP B53D23009360006 and by INdAM GNAMPA Project 2024 ‘Composite materials and microstructures’.

Conflict of interest

The authors declare no conflicts of interest.

References

1. G. Gioia, R. D. James, Micromagnetics of very thin films, *Proc. A*, **453** (1997), 213–223. <https://doi.org/10.1098/rspa.1997.0013>
2. G. Pisante, Homogenization of micromagnetics large bodies, *ESAIM: COCV*, **10** (2004), 295–314. <https://doi.org/10.1051/cocv:2004008>

3. E. Sanchez-Palencia, Fluid flow in porous media, In: *Non-homogeneous media and vibration theory*, Lecture Notes in Physics, Springer, Berlin, Heidelberg, **127** (1980), 129–157. https://doi.org/10.1007/3-540-10000-8_7
4. E. G. Virga, *Variational theories for liquid crystals*, Chapman and Hall/CRC, 2018.
5. A. Braides, I. Fonseca, G. Francfort, 3D-2D asymptotic analysis for inhomogeneous thin films, *Indiana Univ. Math. J.*, **49** (2000), 1367–1404.
6. Bouchitté, I. Fonseca, L. Mascarenhas, A global method for relaxation, *Arch. Rational Mech. Anal.*, **145** (1998), 51–98. <https://doi.org/10.1007/s002050050124>
7. M. Amar, Two-scale convergence and homogenization on $BV(\Omega)$, *Asymptotic Anal.*, **16** (1998), 65–84. <https://doi.org/10.3233/ASY-1998-276>
8. M. Amar, J. Matias, M. Morandotti, E. Zappale, Periodic homogenization in the context of structured deformations, *Z. Angew. Math. Phys.*, **73** (2022), 173. <https://doi.org/10.1007/s00033-022-01817-6>
9. S. Almi, S. Tasso, Brittle fracture in linearly elastic plates, *Proc. R. Soc. Edinburgh Sect. A: Math.*, **153** (2023), 68–103. <https://doi.org/10.1017/prm.2021.71>
10. J. F. Babadjian, E. Zappale, H. Zorgati, Dimensional reduction for energies with linear growth involving the bending moment, *J. Math. Pures Appl.*, **90** (2008), 520–549. <https://doi.org/10.1016/j.matpur.2008.07.003>
11. J. F. Babadjian, V. Millot, Homogenization of variational problems in manifold valued BV -spaces, *Calc. Var.*, **36** (2009), 7–47. <https://doi.org/10.1007/s00526-008-0220-3>
12. J. F. Babadjian, V. Millot, Homogenization of variational problems in manifold valued Sobolev Spaces, *ESAIM: COCV*, **16** (2010), 833–855. <https://doi.org/10.1051/cocv/2009025>
13. A. Braides, A. Defranceschi, E. Vitali, Homogenization of free discontinuity problems, *Arch. Rat. Mech. Anal.*, **135** (1996), 297–356. <https://doi.org/10.1007/BF02198476>
14. A. Braides, I. Fonseca, Brittle thin films, *Appl. Math. Optim.*, **44** (2001), 299–323. <https://doi.org/10.1007/s00245-001-0022-x>
15. G. Carita, J. Matias, M. Morandotti, D. R. Owen, Dimension reduction in the context of structured deformations, *J. Elast.*, **133** (2018), 1–35. <https://doi.org/10.1007/s10659-018-9670-9>
16. G. Carvalho, J. Matias, E. Zappale, Asymptotic analysis of a clamped thin multidomain allowing for fractures and discontinuities, *Commun. Contemp. Math.*, **28** (2025), 2550040. <https://doi.org/10.1142/S0219199725500403>
17. E. Davoli, R. Ferreira, C. Kreisbeck, Homogenization in BV of a model for layered composites in finite crystal plasticity, *Adv. Calc. Var.*, **14** (2021), 441–473. <https://doi.org/10.1515/acv-2019-0011>
18. R. De Arcangelis, G. Gargiulo, Homogenization of integral functionals with linear growth defined on vector-valued functions, *NoDEA*, **2** (1995), 371–416. <https://doi.org/10.1007/BF01261182>
19. R. De Arcangelis, G. Gargiulo, A remark on the homogenization formula for quasiconvex functionals with linear growth, *Ric. Mat.*, **54** (2005), 31–37.

20. R. Ferreira, I. Fonseca, Reiterated homogenization in BV via multiscale convergence, *SIAM J. Math. Anal.*, **44** (2012), 2053–2098. <https://doi.org/10.1137/110826205>
21. R. Ferreira, I. Fonseca, R. Venkatraman, Homogenization of quasi-crystalline functionals via two-scale-cut-and-project convergence, *SIAM J. Math. Anal.*, **53** (2021), 1785–1817. <https://doi.org/10.1137/20M1341222>
22. J. Matias, M. Morandotti, P. M. Santos, Homogenization of functionals with linear growth in the context of A -quasiconvexity, *Appl. Math. Optim.*, **72** (2015), 523–547. <https://doi.org/10.1007/s00245-015-9289-1>
23. J. Matias, P. M. Santos, A dimension reduction result in the framework of structured deformations, *Appl. Math. Optim.*, **69** (2014), 459–485. <https://doi.org/10.1007/s00245-013-9229-x>
24. A. Chakraborty, G. Griso, J. Orlik, Dimension reduction and homogenization of composite plate with matrix pre-strain, *Asymptotic Anal.*, **138** (2024), 255–310.
25. J. Fabricius, M. Gahn, Homogenization and dimension reduction of the Stokes problem with Navier-slip condition in thin perforated layers, *Multiscale Model. Simul.*, **21** (2023), 1502–1533. <https://doi.org/10.1137/22M1528860>
26. T. Fatima, E. Ijioma, T. Ogawa, A. Muntean, Homogenization and dimension reduction of filtration combustion in heterogeneous thin layers, *Netw. Heterog. Media*, **9** (2014), 709–737. <https://doi.org/10.3934/nhm.2014.9.709>
27. C. Kreisbeck, S. Krömer, Heterogeneous thin films: combining homogenization and dimension reduction with directors, *SIAM J. Math. Anal.*, **48** (2016), 785–820. <https://doi.org/10.1137/15M1032557>
28. M. Eleuteri, L. Lussardi, A. Torricelli, E. Zappale, Homogenization and 3D-2D dimension reduction of a functional on manifold valued Sobolev space, *Nonlinear Anal.*, **91** (2026), 104579. <https://doi.org/10.1016/j.nonrwa.2025.104579>
29. G. Dal Maso, *An introduction to Γ -convergence*, Progress in nonlinear differential equations and their applications, Vol. 8, Birkhäuser Boston, 1993. <https://doi.org/10.1007/978-1-4612-0327-8>
30. R. Alicandro, A. Corbo Esposito, C. Leone, Relaxation in BV of integral functionals defined on Sobolev functions with values in the unit sphere, *J. Convex Anal.*, **14** (2007), 69–98.
31. R. Alicandro, C. Leone, 3D-2D asymptotic analysis for micromagnetic thin films, *ESAIM: COCV*, **6** (2001), 489–498. <https://doi.org/10.1051/cocv:2001119>
32. M. Baía, E. Zappale, A note on the 3D-2D dimensional reduction of a micromagnetic thin film with nonhomogeneous profile, *Appl. Anal.*, **86** (2007), 555–575. <https://doi.org/10.1080/00036810701233942>
33. L. Carbone, K. Chacouche, A. Gaudiello, Fin junction of ferroelectric thin films, *Adv. Calc. Var.*, **11** (2018), 341–371. <https://doi.org/10.1515/acv-2016-0047>
34. A. Desimone, R. Kohn, S. Müller, F. Otto, A reduced theory for thin-film micromagnetics, *Commun. Pure Appl. Math.*, **55** (2002), 1408–1460. <https://doi.org/10.1002/cpa.3028>
35. A. Gaudiello, R. Hadiji, Ferromagnetic thin multi-structures, *J. Differ. Equations*, **257** (2014), 1591–1622. <https://doi.org/10.1016/j.jde.2014.05.015>

36. A. Gaudiello, R. Hadiji, Junction of ferromagnetic thin films, *Calc. Var.*, **39** (2010), 593–619. <https://doi.org/10.1007/s00526-010-0327-1>
37. L. Ambrosio, G. Dal Maso, On the relaxation in $BV(\Omega; \mathbb{R}^m)$ of quasi-convex integrals, *J. Funct. Anal.*, **109** (1992), 76–97. [https://doi.org/10.1016/0022-1236\(92\)90012-8](https://doi.org/10.1016/0022-1236(92)90012-8)
38. F. Bethuel, The approximation problem for Sobolev maps between two manifolds, *Acta Math.*, **167** (1991), 153–206. <https://doi.org/10.1007/BF02392449>
39. F. Bethuel, X. Zheng, Density of smooth functions between two manifolds in Sobolev spaces, *J. Funct. Anal.*, **80** (1988), 60–75. [https://doi.org/10.1016/0022-1236\(88\)90065-1](https://doi.org/10.1016/0022-1236(88)90065-1)
40. L. Ambrosio, A. Braides, Functionals defined on partitions in sets of finite perimeter. I. Integral representation and G-convergence, *J. Math. Pures Appl.*, **69** (1990), 285–305.
41. L. Ambrosio, N. Fusco, D. Pallara, *Functions of bounded variation and free discontinuity problems*, New York: Oxford University Press, 2000. <https://doi.org/10.1093/oso/9780198502456.001.0001>
42. L. Ambrosio, S. Mortola, V. M. Tortorelli, Functional with linear growth defined on vector-valued BV functions, *J. Math. Pure Appl.*, **70** (1991), 269–323.
43. I. Fonseca, S. Müller, Quasi-convex integrands and lower semicontinuity in L^1 , *SIAM J. Math. Anal.*, **23** (1992), 1081–1098. <https://doi.org/10.1137/0523060>
44. J. M. Ball, F. Murat, $W^{1,p}$ -quasiconvexity and variational problems for multiple integrals, *J. Funct. Anal.*, **58** (1984), 225–253. [https://doi.org/10.1016/0022-1236\(84\)90041-7](https://doi.org/10.1016/0022-1236(84)90041-7)
45. I. Fonseca, S. Müller, Relaxation of quasiconvex functional in $BV(\Omega, \mathbb{R}^p)$ for integrands $f(x, u, \nabla u)$, *Arch. Rational Mech. Anal.*, **123** (1993), 1–49. <https://doi.org/10.1007/BF00386367>



AIMS Press

© 2026 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)