



# A CIRCULAR ECONOMY APPROACH FOR *CUPRIAVIDUS NECATOR* DSM 545 BIOSYNTHESIS OF POLY (3-HYDROXYBUTYRATE)

S. Bellini <sup>a,b</sup>, T. Tommasi <sup>a</sup> and D. Fino <sup>\* a,b</sup>



<sup>a</sup> Politecnico di Torino, Department of Applied Science and Technology, Torino, Piemonte, Italy  
<sup>b</sup> Istituto Italiano di Tecnologia, Center for Sustainable Future Technologies, Torino, Piemonte, Italy  
<sup>\*</sup>corresponding author, (presenting author e-mail: [silvia.bellini@polito.it](mailto:silvia.bellini@polito.it))



- *Cupriavidus necator* produces large amounts of the polyester poly (3-hydroxybutyric acid), or **PHB**.
- It is a **versatile bacterium**: it can grow both autotrophically and heterotrophically, and on a wide range of organic substrates, including **waste feedstocks** (Bellini et al., 2022).
- In this work, *C. necator* DSM 545 strain has been fermented to reach high yield of PHB by using **two waste substrates**.
- **Circular economy** approach for bacterial PHB biosynthesis
- This research is part of the **PRIME project** (*Processi e prodotti Innovativi di chimica verde*), promoted by Novamont® and Piedmont Region.



- ✓ 1L bioreactor Sartorius®, working volume 0,5 L, pH 6.8, 30°C, vvm 0,5.
- ✓ *C. necator* DSM 545 grown using a growth medium containing a **carbon source, phosphate, sulphate and magnesium sources, and metals** (Mozumder et al. 2014).
- ✓ **Biomass** sampled at 15, 24, 36, 40, 45 and 48 hours.
- ✓ **Two different carbon waste sources:**
  1. a **syrup (glucose and fructose in equal amount)** from PRIME project supply chain (partner Sedamyl®)
  2. the sterilized and concentrated waste medium of an acetogenic bacterium (*Acetobacterium woodii*) fermentation containing **acetate**.
- ✓ **Sugars and acetate** in the medium and **PHB concentration** (extracted by acid digestion using sulphuric acid 96%): analyzed at HPLC using a Resex18 column and a mobile phase of H<sub>2</sub>SO<sub>4</sub> 5 mM (flow rate of 0.7 mL/min).



## Results & Discussion

- The graph shows the fermentation of *C. necator* DSM 545 performed through a **“two-step”** fermentation strategy, as shown in the scheme on the left.
- **Sedamyl® syrup** (containing glucose and fructose equal concentrations) has been furnished at different concentrations during the whole **fed-batch fermentation** (“spike feeding”).
- After 24 hours, **spike feeding** of the acetogenic bacterium medium containing **acetate** (2 g L<sup>-1</sup> each feed).
- The highest PHB concentration, almost **10 g L<sup>-1</sup>**, has been reached after 48 hours of fermentation and the biomass reached about **12 g L<sup>-1</sup>** at the same hour.
- **83% of PHB content**,  $w_{PHB}/w_{biomass}$ .
- PHB is mainly accumulated in *C. necator* under **unbalanced growth conditions**, e.g. when shortage of N and P occur.

## Conclusions

- ✓ High concentration of PHB and biomass: **yield of 83% of biopolymer (w/w)**, using valuable **waste substrates**, using a **circular economy** approach.
- ✓ *C. necator* DSM 545 easily convert glucose, fructose and acetate into PHB.
- ✓ **Optimization** of fermentation operative conditions:
  - i. A **three phases C/N ratio** fermentation approach (three different concentration of C and N and relative ratio) could be used to test PHB biosynthesis improvement (Garcia-gonzalez and Wever, 2018).
  - ii. Exponential feeding and an alkali-addition monitoring strategies (Mozumder et al., 2014)
  - iii. Utilization of **carbon dioxide waste** (e.g. industrial gas-off) as carbon source for acidogenic bacterium fermentation, to improve the whole **Life Cycle Assessment analysis** of the process.

Bellini S., Tommasi T., Fino D., (2022). Poly(3-hydroxybutyrate) biosynthesis by Cupriavidus necator: a review on waste substrates utilization for a circular economy approach. Bioresource Technology Reports, Vol. 17.

Md. Salatul Islam Mozumder, Heleen De Wever, E.I.P.V., and A. L.G.-G. (2014). A robust fed-batch feeding strategy independent of the carbon source for optimal polyhydroxybutyrate production. Process Biochem. Vol. 49, Issue 3 49, 365–373.

Kourmentza, C., Pl, J., Venetsaneas, N., Burniol-figols, A., Varrone, C., Gavala, H.N., and Reis, M.A.M. (2017). Recent Advances and Challenges towards Sustainable Polyhydroxyalkanoate ( PHA ) Production. Bioingeniering 1–43.

Garcia-gonzalez, L., and Wever, H. De (2018). Acetic Acid as an Indirect Sink of CO 2 for the Synthesis of Polyhydroxyalkanoates (PHA ): Comparison with PHA Production Processes Directly Using CO 2 as Feedstock. Appl. Sci.