

Development and Validation of Prediction Models for Subtype Diagnosis of Patients With Primary Aldosteronism

Original

Development and Validation of Prediction Models for Subtype Diagnosis of Patients With Primary Aldosteronism / Burrello, Jacopo; Burrello, Alessio; Pieroni, Jacopo; Sconfienza, Elisa; Forestiero, Vittorio; Rabbia, Paola; Adolf, Christian; Reincke, Martin; Veglio, Franco; Williams, Tracy Ann; Monticone, Silvia; Mulatero, Paolo. - In: THE JOURNAL OF CLINICAL ENDOCRINOLOGY AND METABOLISM. - ISSN 0021-972X. - 105:10(2020), pp. e3706-e3717. [10.1210/clinem/dgaa379]

Availability:

This version is available at: 11583/2978534 since: 2023-05-16T12:18:17Z

Publisher:

ENDOCRINE SOC

Published

DOI:10.1210/clinem/dgaa379

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

1 **Development and validation of prediction models for subtype diagnosis of**
2 **patients with primary aldosteronism.**

3 Jacopo Burrello MD^{1*}, Alessio Burrello MS^{2*}, Jacopo Pieroni MD¹, Elisa Sconfienza MD¹, Vittorio
4 Forestiero MD¹, Paola Rabbia MD³, Christian Adolf MD⁴, Martin Reincke MD⁴, Franco Veglio
5 MD¹, Tracy Ann Williams PhD^{1,4}, Silvia Monticone PhD^{1#}, Paolo Mulatero MD^{1#}.

6

7 (1) Division of Internal Medicine and Hypertension, Department of Medical Sciences, University
8 of Torino, Italy. (2) Department of Electrical, Electronic and Information Engineering "Guglielmo
9 Marconi" (DEI), University of Bologna, Italy. (3) Division of Radiology, University of Torino, Italy.

10 (4) Medizinische Klinik und Poliklinik IV, Klinikum der Universität, Ludwig-Maximilians-
11 Universität München, Munich, Germany.

12 * Contributed equally and should be considered as joint first authors.

13 # Contributed equally and should be considered as joint last authors.

14

15 **Short title:** Diagnostic modelling for PA subtyping

16 **Keywords:** aldosterone; primary aldosteronism; adrenal venous sampling; machine learning.

17

18 **Corresponding author and person to whom reprints should be addressed:** Prof. Paolo Mulatero

19 Division of Internal Medicine and Hypertension Unit, Department of Medical Sciences, University
20 of Torino, Città della Salute e della Scienza, Via Genova 3, 10126 Torino, Italy. Telephone/Fax
21 number: 0039.011.633.6959 / 0039.011.633.6931. E-mail: paolo.mulatero@libero.it

22

23 **Source(s) of Funding:** this research did not receive any specific grant. CA and MR are supported
24 by the Else Kröner-Fresenius Stiftung (2019_A104) and by the Deutsche Forschungsgemeinschaft
25 within the CRC/Transregio 205/1 "The Adrenal: Central Relay in Health and Disease".

26 **Conflict(s) of Interest/Disclosure(s):** the authors have nothing to disclose.

27 **ABSTRACT**

28 **Context.** Primary aldosteronism (PA) comprises unilateral (lateralized, LPA) and bilateral disease
29 (BPA). The identification of LPA is important to recommend potentially curative adrenalectomy.
30 Adrenal venous sampling (AVS) is considered the gold standard for PA subtyping, but the procedure
31 is available in few referral centers.

32 **Objective.** To develop prediction models for subtype diagnosis of PA using patient clinical and
33 biochemical characteristics.

34 **Design, Patients and Setting.** Patients referred to a tertiary hypertension unit. Diagnostic
35 algorithms were built and tested in a training (N=150) and in an internal validation cohort (N=65),
36 respectively. The models were validated in an external independent cohort (N=118).

37 **Main outcome measure.** Regression analyses and supervised machine learning algorithms were
38 used to develop and validate two diagnostic models and a 20-point score to classify patients with
39 PA according to subtype diagnosis.

40 **Results.** Six parameters were associated with a diagnosis of LPA (aldosterone at screening and after
41 confirmatory testing, lowest potassium value, presence/absence of nodules, nodule diameter, and
42 computed tomography results) and were included in the diagnostic models. Machine learning
43 algorithms displayed high accuracy at training and internal validation (79.1% to 93%), whereas a
44 20-point score reached an AUC of 0.896, and a sensitivity/specificity of 91.7/79.3%. An integrated
45 flow-chart correctly addressed 96.3% of patients to surgery and would have avoided AVS in 43.7%
46 of patients. The external validation on an independent cohort confirmed a similar diagnostic
47 performance.

48 **Conclusions.** Diagnostic modelling techniques can be used for subtype diagnosis and guide surgical
49 decision in patients with PA in centers where AVS is unavailable.

50 INTRODUCTION

51 Primary aldosteronism (PA) accounts for 3-13% of primary care hypertensive patients [1-3] and is
52 associated with an increased cardio- and cerebrovascular risk compared with patients affected by
53 essential hypertension [4;5]. The two major subtypes of PA are unilateral primary aldosteronism
54 (lateralized, LPA), mainly due to an aldosterone producing adenoma (APA), and bilateral primary
55 aldosteronism (BPA). The treatments of choice are unilateral adrenalectomy, or medical therapy
56 with a mineralocorticoid receptor antagonist, respectively [6]. A timely and accurate subtype
57 diagnosis is critical to recommend the appropriate treatment and improving the outcomes of these
58 patients [7;8].

59 Over the last decades, many procedures have been proposed to differentiate LPA from BPA,
60 including posture testing, functional imaging (using 11-C-metomidate or 68-Ga-pentixafor tracers
61 [9;10] and steroid profiling [11-13]. Nevertheless, technical issues and/or the lack of sensitivity and
62 specificity hampered the introduction of these tests in the routine management of PA patients [6].

63 Adrenal venous sampling (AVS) is currently considered the gold standard for subtype diagnosis [6].
64 Nevertheless, several concerns prevent its widespread use: AVS is an invasive, time-consuming,
65 and relatively expensive procedure, requiring a high level of technical skill and is available only in
66 a limited number of referral centers [14]. Beside AVS, adrenal computed tomography (CT) scanning
67 is widely available in most centers and performed in all patients with confirmed PA [6]. Even if
68 several studies reported unreliable diagnostic performance of CT in PA subtyping [15;16], score-
69 based algorithms combining imaging findings with clinical and biochemical parameters have been
70 developed [17-23]. Küpers et al. first proposed a prediction score to bypass AVS; patients with a
71 potassium lower than 3.5 mmol/L, an estimated glomerular filtration rate (eGFR) more than 100
72 ml/min and a typical adenoma at CT imaging could avoid AVS. Sensitivity and specificity were
73 53.1% and 100%, respectively [17]. Other clinical scores were subsequently developed in the
74 attempt of differentiating LPA from BPA, with an accuracy ranging from 58.2% to 86.3% [18-23].
75 Only 2 of these scores were also validated in independent cohorts [17;22], 5 included less than 100

76 patients in the development cohort [17-20;23], and the majority of these scores was applicable only
77 in selected cohorts of patients with PA [18-21;23].

78 Considering the high prevalence of PA and the limited availability of AVS, an alternative method
79 that reduces the number of requested AVS is highly desirable. Our objective was to develop and
80 validate clinical models to discriminate LPA from BPA, which can bypass AVS for bilateral disease,
81 and indicating unilateral adrenalectomy for patients with high probability of LPA who cannot
82 undergo AVS. We propose herein two advanced diagnostic models based on supervised machine
83 learning algorithms and a flow-chart integrating our score-system (the SPACE score, Subtyping
84 Primary Aldosteronism by Clinical Evaluation) in PA patient management. Validation of previously
85 described score-based algorithms is also provided and demonstrates the superiority of our prediction
86 models.

87

88 **METHODS**

89 Data analyzed during the current study are not publicly available but are available from the
90 corresponding author on reasonable request. A detailed description of patient management and data
91 extraction, statistical analyses, and diagnostic modelling is provided as Supplemental Data [24]
92 (available at <https://github.com/ABurrello/SPACE-score>).

93 ***Study cohort and data extraction***

94 We retrospectively evaluated a cohort of 215 patients referred to the hypertension unit of the
95 University of Torino between 2008 and 2019, to train and test the diagnostic models. Eligible
96 patients were randomly assigned to the training cohort (N = 150) or to the internal validation cohort
97 (N = 65). An independent cohort of 118 consecutive patients from the Munich Klinikum der
98 Universität treated between 2008 and 2014 was used for external validation. PA was diagnosed
99 according to the Endocrine Society Guideline [6]. Inclusion criteria were: (1) confirmed diagnosis
00 of PA; (2) successful AVS for subtype diagnosis.

01 Unilateral PA mainly depends on unilateral aldosterone-producing adenoma. However, a
02 lateralization at AVS may also occur also in the presence of a dominant lesion with asymmetrical
03 autonomous aldosterone production in the context of bilateral adrenal alterations, including
04 aldosterone producing cell clusters, or diffuse/nodular hyperplasia. For this reason, the AVS-based
05 term of “lateralized PA” was used to indicate a prevalently unilateral disease throughout the present
06 study.

07 ***Statistical analysis***

08 IBM SPSS Statistics 22 (*IBM Corp., Armonk, New York, USA*) was used for statistical analyses.
09 Data distribution was assessed by the Kolmogorov–Smirnov test. Normally distributed variables
10 were analyzed by T-student test and reported as mean \pm standard deviation. Non-normally
11 distributed variables were analyzed by Mann-Whitney’s test and reported as median [interquartile
12 range]. Categorical variables were analyzed by Chi-square test and reported as absolute number and
13 proportion (%). Univariate logistic regression analysis was used to define the odds ratios (ORs) for
14 each analyzed parameter. Six selected variables were included in the multivariate logistic regression
15 analysis. An OR greater than 1 is associated with an increased likelihood of the defined outcome
16 (diagnosis of LPA), an OR less than 1, a decreased likelihood. A *P* value of less than 0.05 was
17 considered significant.

18 ***Diagnostic modelling***

19 Python 3.5 (library, scikit-learn) was used for the development and validation of diagnostic models
20 by machine learning techniques. Supervised machine learning algorithms are widely used in clinical
21 research to formulate predictions about possible outcomes based on a pre-defined set of labeled
22 paired input-output training sample data [13;25]. Supervised machine learning and in particular
23 linear discriminant analysis (LDA) and random forest (RF) algorithms were applied on the
24 combined cohort to develop diagnostic models able to discriminate patients with LPA vs. BPA. Six
25 selected variables were used in both models (aldosterone at screening and after confirmatory testing,

26 lowest potassium recorded in the absence of diuretic therapy, presence/absence of a nodule at CT
27 scanning imaging, nodule diameter, and descriptive CT scanning findings).

28 LDA maximize the separation between groups by increasing precision estimates by variance
29 reduction. The algorithm computes a set of coefficients for linear combination of each variable to
30 classify patients according to their diagnosis; a canonical plot was used to represent diagnostic
31 performance of the LDA model. The RF model was composed of 20 classification trees with a
32 maximum number of eight splits for each tree. The predicted diagnosis was defined on the basis of
33 the outcome of each classification tree of the RF: if at least 11 of 20 trees of the forest predict the
34 diagnosis of lateralized PA, the patient is classified as LPA. The RF model was integrated in a free-
35 downloadable tool which allows the application of the algorithm in clinical practice (available at
36 https://github.com/ABurrello/SPACE-score/raw/master/Random_Forest_model.zip).

37 Performance and generalizability of both LDA and RF models were evaluated by a 10K-cross
38 validation algorithm (see extended methods - Supplemental Data) [24].

39 The 6 variables were used to develop a 20-point score to predict the diagnosis of LPA. Variables
40 were categorized, points were assigned to each reference interval, and cut-offs were derived to
41 achieve the best accuracy in an automated way. The SPACE score was generated using the training
42 cohort and tested with both internal and external validation cohorts. ROC curve analysis was used
43 to assess the area under the curve and derive the best cut-off to discriminate patients with LPA by
44 evaluation of the Younden Index ($J = \text{sensitivity} + \text{specificity} - 1$). A second online tool was
45 developed to automatically calculate the score and the predicted diagnosis (available at
46 <https://github.com/ABurrello/SPACE-score/raw/master/SPACE%20Score%20Calculator.xlsm>).

47

48 **RESULTS**

49 *Patient characteristics*

50 Two hundred and fifteen patients were included in the analyses from the developmental cohort of
51 Torino, 133 with a diagnosis of LPA and 82 with BPA. Clinical and biochemical characteristics are

52 reported in Table 1. The mean age at diagnosis was 49 ± 9.5 years, mean BP was 164/99 mmHg,
53 with a duration of hypertension of 68 [27; 128] months. Patients with a diagnosis of LPA were more
54 frequently females (42.1% vs. 23.2%; $P = 0.005$), had a higher DDD (3.8 [2.2; 5.7] vs. 3.0 [1.3; 4.7]
55 $P = 0.027$) and a lower potassium level (3.1 ± 0.6 vs. 3.8 ± 0.4 ; $P < 0.001$). At the diagnostic workup,
56 patients with LPA displayed higher levels of aldosterone, both at screening (38.0 [25.7; 49.7] vs.
57 28.7 [19.8; 37.9] ng/dL; $P < 0.001$) and post-confirmatory testing (20.5 [13.3; 32.9] vs. 11.5 [8.2;
58 17.7] ng/dL; $P < 0.001$). To confirm PA diagnosis, 165 patients underwent saline infusion testing
59 (76.7%), and 50 had captopril challenge testing (23.3%). CT scanning demonstrated the presence
60 of a defined nodule in 85.7% of patients with LPA; a nodule was also detected in 41.5% of patients
61 with bilateral disease (unilateral nodule in 29 of 34 cases, bilateral in 5). In patients with BPA, CT
62 scanning was bilaterally normal in 24.4% of patients, bilaterally abnormal in 22%, and with a
63 unilateral abnormality in 53.7% of the cases (see the Supplemental Data for details on adrenal CT
64 scanning interpretation and definition of nodule) [24]. Among the 37 patients with bilateral
65 abnormalities at CT scanning, 40.6% displayed a unilateral nodule in the context of bilateral adrenal
66 thickening or contralateral thickening, 37.8% bilateral nodules, and 21.6% bilateral hyperplasia.
67 Prevalence of target organ damage (eGFR, microalbuminuria, and LVH at echocardiography) and
68 prior cardiovascular events was not significantly different between groups. As expected, the LI at
69 AVS was significantly higher in LPA than BPA patients (12.0 [6.9; 21.3] vs. 1.8 [1.3; 2.6]; $P <$
70 0.001). According to the PASO criteria [7], after a follow-up of 6-12 months from unilateral
71 adrenalectomy, patients with LPA displayed complete clinical and biochemical success in 54.1%
72 and 98.5% of cases, respectively.

73 Univariate logistic regression analysis was performed including all parameters (Table S1) [24],
74 showing a significant association with a diagnosis of LPA of female sex (OR 2.41), duration of
75 HTN (OR 1.01), DDD (Defined Daily Dose; OR 1.18), potassium (OR 0.10), aldosterone at
76 screening (OR 1.01) and after confirmatory testing (OR 1.01), presence of a nodule at CT scanning
77 (OR 8.33), nodule diameter (OR 1.12), and CT findings (OR 9.91). Six out of these 9 variables were

78 selected considering their discriminative performance and introduced in the multivariate model,
79 which confirmed a highly significant independent association with the diagnosis of LPA for all
80 parameters (Table 2).

81 ***Linear discriminant analysis model***

82 The 6 selected variables confirmed by the multivariate regression analysis were used in an LDA
83 model. The linear combination of variables included in the LDA is shown in the canonical plot
84 (Figure 1A). Each point represents a patient and the clear separation according to their subtype
85 diagnosis indicates that the model can discriminate LPA from BPA with reliable accuracy. In
86 particular, 175 of 215 patients (accuracy 81.4%) were correctly classified, with a sensitivity and
87 specificity for LPA detection of 86.5% and 73.2%, respectively (Figure 1B). To exclude overfitting
88 bias and assess how the model could generalize in an independent cohort, the LDA was validated
89 by a 10K-cross validation algorithm (see extended methods - Supplemental Data) [24]. The cross
90 validation showed a high predictive performance with an accuracy of 79.1%, compared with the
91 81.4% (at training), thus confirming a negligible overfitting bias (overfitting effect = 2.3%). In the
92 LDA model, the stronger predictor was the lowest potassium level (normalized LDA coefficient =
93 1.0), followed by presence of a defined nodule and CT findings (0.8 and 0.4, respectively; Figure
94 1C and Table S2) [24].

95 ***Random forest model***

96 Besides LDA, we also developed a non-linear classification model, exploiting RF classification
97 algorithms. The same 6 selected variables were combined in a RF model comprising 20
98 classification trees (a representative tree is reported in Figure 2A), and were able to correctly
99 discriminate 132 of 133 patients with LPA (sensitivity 99.2%), and 68 of 82 patients with BPA
:00 (specificity 82.9%), resulting in an overall accuracy of 93% at the training and of 87% after 10K-
:01 cross validation (overfitting effect 6%; Figure 2B). In this case, the stronger predictor was nodule
:02 diameter, followed by the lowest potassium level and by the presence of a nodule at CT scanning
:03 (Figure 2C).

:04 **Prediction score**

:05 Patients included in the described models (combined cohort; N = 215) were randomly assigned to a
:06 training cohort (N = 150) or internal validation cohort (N = 65). No differences were found for all
:07 evaluated parameters between the two groups (Table S3) [24]. The same 6 variables used in the
:08 LDA and RF models were then used to develop the SPACE score, a 20-point score to discriminate
:09 patients with LPA vs. BPA. The SPACE score was developed in the training cohort and then tested
:10 in the internal validation cohort. Figures 3A and 3C report the categorization of the 6 different
:11 variables and assignment of points. The analysis of the ROC curve demonstrated a high diagnostic
:12 performance (Figure 3B). The area under the curve was 0.896 (95% CI 0.852-0.940) and the cut-
:13 off with the higher accuracy was 12. At the training, a score greater than 12 correctly identified a
:14 diagnosis of LPA in 87 of 93 patients (sensitivity 93.5%), whereas a score equal or lower than 12
:15 identified a diagnosis of BPA in 47 of 57 patients (specificity 82.5%), with an overall accuracy of
:16 89.3%. Of note, the prediction score displayed a very high performance with an accuracy of 81.5%
:17 at validation, and a sensitivity and specificity of 87.5% and 72%, which was not significantly
:18 different from the machine learning models (accuracy at validation of 79.1% and 87% for the LDA
:19 and the RF model, respectively). Confusion matrix for training, internal validation, and combined
:20 developmental cohort are reported in Figure 3D. The difference between the accuracy of the
:21 prediction score in the training cohort compared with the internal validation cohort, revealed a
:22 modest bias due to an expected overfitting effect (7.8%), which did not affect the reliability of the
:23 model. A cut-off of greater than 8 or of greater than 16 optimized sensitivity or specificity,
:24 respectively (Table S4) [24]. With a cut-off of 8, sensitivity was increased to 97.8% and 95%,
:25 correctly classifying 91 of 93, and 38 of 40 patients with LPA, in the training cohort and in the
:26 validation cohorts, respectively. With a cut-off of 16, specificity was increased to 98.2% and 92%,
:27 correctly classifying 56 of 57, and 23 of 25 patients with BPA, in the training cohort and in the
:28 validation cohort, respectively.

:29 To evaluate further the diagnostic performance of the SPACE score, 7 previously published scores
:30 [17-23] were tested on our combined cohort (Table S5) [24]. The accuracy of our prediction score
:31 (89.3% and 81.5% at training and internal validation analysis, respectively) was superior to all
:32 available scores (accuracy ranging from 58.2% to 86.3% at training and from 67.3% to 78% at
:33 validation). Of note, the RF classification algorithm outperformed all other models with an accuracy
:34 of 87% at validation, higher than all score evaluated at training.

:35 ***External validation***

:36 LDA, RF model and the SPACE score were validated on an external independent cohort from
:37 Munich of 118 patients, 57 with LPA and 61 with BPA (Table S6) [24]. Compared with the
:38 developmental cohort, the prevalence of LPA was significantly lower in the external validation
:39 cohort (48.3% vs. 61.9%; $P = 0.017$) and mean BP (153/94 mmHg vs. 164/99 mmHg), DDD (2.5
:40 [1.0; 4.0] vs. 3.3 [2.0; 5.0]), potassium levels (3.1 ± 0.5 mEq/L vs. 3.4 ± 0.7 mEq/L) were also
:41 significantly lower. PRA at screening (0.29 vs 0.25 ng/mL/h) and after confirmatory testing (0.21
:42 vs 0.15 ng/mL/h) was significantly higher and aldosterone levels at screening (17.9 vs 33.4 ng/dL)
:43 and after confirmatory testing (11.2 and 16.4 ng/dL) were significantly lower ($P < 0.05$ for all
:44 comparisons) in the validation compared with the developmental cohort. The reliability of the
:45 diagnostic performance of our prediction models was confirmed at external validation. The accuracy
:46 was 78.8%, 80.5%, and 78.8%, respectively for LDA, RF, and the score system (Figure S1) [24],
:47 with a minimum overfitting bias compared with the internal validation on the developmental cohort
:48 (range between 0.3% and 6.5%).

:49 ***Management of PA patient***

:50 The SPACE score was directly correlated with the proportion of patients with a diagnosis of LPA
:51 (Table S7) [24] and with the lateralization index (LI) at the AVS (Table S8) [24]. Figure 4A clearly
:52 illustrates the stratification of patients with a diagnosis of LPA vs. BPA for the prediction score and
:53 graphically confirmed the cut-offs of 8, 12 and 16, which maximize sensitivity, accuracy, and

:54 specificity, as defined by ROC curve analysis. In addition, all patients with a score greater than 18
:55 had LPA, whereas all patients with a score lower or equal than 2 had BPA.

:56 Finally, our score was integrated in a flow chart for PA management (Figure 4B). Patients with a
:57 score less than or equal to 8 were classified as “probable BPA” and treated with MRA (N = 32),
:58 thus resulting in 28 patients with true bilateral disease correctly managed, and 4 patients with a LPA
:59 (1.9%) which missed the possibility to undergo adrenalectomy. Patients with a score greater than
:60 16 were classified as “probable LPA”, with indication to unilateral adrenalectomy (N = 62).
:61 Accordingly, 3 patients with bilateral disease would undergo inappropriate surgery (1.4%), and 1
:62 patient with LPA would have resection of the wrong adrenal (0.5%). All remaining patients (N =
:63 121), with a score comprised between 8.5 and 16 would undergo AVS with management according
:64 to the result of the procedure. Sensitivity, specificity, positive, and negative predictive values are
:65 reported in the confusion matrix (Figure 4C).

:66 We combined patients from the developmental and external validation cohorts (N=333) and
:67 stratified these patients into 3 groups according to the points of the SPACE score (score \leq 8 vs.
:68 8.5-16 vs. $>$ 16): the median LI displayed a gradual increase in the 3 groups of patients (Table S8)
:69 [24]. Moreover, clinical and biochemical outcomes in patients with LPA misclassified as BPA were
:70 worse than patients with a correct prediction of LPA (83.3% vs. 48.8% partial + absent clinical
:71 success, and 5.6% vs. 0.6% partial + absent biochemical success; Table S9) [24].

:72 After stratification for the confirmatory test performed during the diagnostic work-up (Table S10)
:73 [24], the SPACE score confirmed its applicability both for patients diagnosed by saline infusion
:74 testing (accuracy 84%) or captopril challenge testing (accuracy 84.6%).

:75 The application of the prediction score in our clinical context would result in the correct
:76 management of 207 of 215 patients (96.3%) with a reduction of 43.7% (94 of 215) of AVS
:77 procedures in the developmental combined cohorts. Notably, the accuracy of the flow chart for
:78 patient management at external validation remained high (94.9%), with a reduction of 66.1% (78 of
:79 118) of AVS procedures (Figure S1) [24].

:80

:81 **DISCUSSION**

:82 In our study, we developed and validated two different prediction models based on supervised
:83 machine learning algorithms and a clinical score for the subtype diagnosis of PA. An online tool
:84 was developed to allow the application of the RF algorithm to clinical practice. Moreover, we
:85 proposed a flow-chart for patient management which integrates our score-system in a second user-
:86 friendly downloadable tool.

:87 Küpers et al. proposed for the first time a clinical score to diagnose lateralized PA; the major
:88 advantages were easy applicability and a very high positive predictive value, resulting in the correct
:89 classification of all patients predicted as LPA [17]. However, this score displays very low
:90 sensitivity, misclassifying 43% of LPA patients, who would miss the chance of potentially curative
:91 adrenalectomy. In addition, validation on independent cohorts did not confirm its diagnostic
:92 performance with a low accuracy, between 56.0% and 72.7% [26-29]. Six other score-systems were
:93 proposed (Table S11) [24]. Two of them [18;23] used only biochemical or demographic features,
:94 thus applicable before imaging. However, these scores were useful only to detect patients that could
:95 avoid AVS due to BPA. The other scores [19;20] combined biochemical parameters with
:96 radiological findings and displayed a high negative predictive value (82.2-100%), with the
:97 identification of patients with BPA to be allocated to medical treatment. Limitations of these studies
:98 were the low number of enrolled patients, the absence of an internal or external validation, and the
:99 applicability only to patients undergoing captopril challenge [18;20] or intravenous saline loading
:00 [19;23] for confirmatory testing. The score proposed by Kamemura et. al was developed in more
:01 than 200 patients but was applicable only to patients without evidence of an adrenal mass at CT
:02 scanning, which represent a minority of patients with PA [21]. Finally, Kobayashi et al. proposed
:03 and validated a score on more than 1,000 patients, reporting a negative predictive value of 92.5%
:04 [22] but with insufficient accuracy. The application of this score in our patients resulted in an
:05 accuracy of 67.4-72.7%.

06 In our diagnostic models, the highest performance was reached by the RF algorithm, which
07 identified 132 of 133 patients with LPA and correctly classified 68 of 69 patients with BPA,
08 resulting in a sensitivity of 99.2% and a negative predictive value of 98.5%. The model accuracy
09 was 93.0% and 87.0% at training and internal validation, respectively. Our SPACE score displayed
10 an equally high performance with an overall accuracy of 89.3% in the training cohort and 81.5% in
11 the internal validation cohort (using 12 as cut-off), outperforming all previously proposed clinical
12 scores. A cut-off of 8 maximized sensitivity and negative predictive value (97% and 87.5%,
13 respectively, in the combined cohort), correctly identifying 28 of 32 patients with BPA, whereas a
14 cut-off of 16 maximized specificity and positive predictive value (96.3% and 95.2%, respectively,
15 in the combined cohort), correctly identifying 59 of 62 patients with LPA.

16 All previously proposed score-systems were applied in our cohort, to assess their generalizability.
17 The accuracy at validation was not suitable for clinical use, ranging between 67.3% and 78% and
18 suggesting a moderate overfitting bias (up to 19%). Conversely, the overfitting effect was low in
19 our models (from 2.3% to 7.8%) with a high accuracy at validation (from 81.5% to 93%).

20 To exclude selection bias and further assess the generalizability of our diagnostic models, we
21 performed an external validation on an independent cohort of patients. LDA, RF model, and the
22 SPACE score confirmed a high diagnostic performance (accuracy range 78.8-94.4%), with a
23 minimum overfitting bias.

24 We combined 3 biochemical variables with 3 imaging-related parameters associated with subtype
25 diagnosis. These parameters were selected considering the results of univariate and multivariate
26 regression analysis, and then used for the LDA, the RF model, and the score-system. Potassium
27 levels, aldosterone levels at screening and after confirmatory test are clinical criteria associated with
28 a high probability of LPA and reflect the severity of disease [30]. Imaging-related parameters
29 resulted to be crucial for subtype diagnosis; in our cohort only 5 of 133 LPA patients (3.8%)
30 displayed a bilaterally normal CT scanning, whereas 85.7% had a defined nodule.

31 The SPACE score was integrated in a flow chart for the management of patients with PA, resulting
32 in the correct classification of 96.3% of patients, potentially reducing almost half of the AVS. The
33 lower cut-off identifies patients with BPA to address to MRA treatment: 28/32 patients with BPA
34 were correctly classified, whereas 4 patients with LPA would be diagnosed as BPA and treated with
35 MRA, therefore, missing the chance of treatment by adrenalectomy. These 4 patients displayed
36 bilaterally normal adrenals at CT scanning and are thus at high risk of partial/absent clinical success
37 after surgery according with the recently proposed prognostic PASO score [7;25]. The higher cut-
38 off identifies patients with LPA, who could undergo unilateral adrenalectomy in centers where AVS
39 is not available. With this strategy, 58 patients with LPA would be correctly adrenalectomized,
40 whereas 4 patients would receive inappropriate surgery (3 patients with BPA and 1 patient with
41 lateralization on the other side). The 3 BPA patients would also be misclassified by all other
42 previously published scores.

43 The external validation resulted in similar performance, with correct management of 94.9% of
44 patients and a potential reduction of 66.1% in the number of AVS, thus excluding a significant inter-
45 center variability. The assessment of clinical and biochemical outcomes of patients with a correct
46 prediction of LPA compared with those misclassified by the SPACE score, reinforced our findings.
47 Finally, unlike previous models, our score system was applicable both to patients with PA diagnosed
48 by saline infusion testing and by captopril challenge testing, with a similar accuracy (84.0% vs.
49 84.6%, respectively).

50 The present score is expected to be of interest to hypertension and endocrine centers and in particular
51 for those that perform systematic screening of patients with hypertension, therefore having a high
52 rate of diagnosis of BPA [31]. With our score a high proportion of BPA patients can avoid
53 unnecessary AVS with a significant reduction of costs and potential complications.

54 The failure to define with certainty the side of aldosterone hypersecretion represents the main limit
55 of our score and of all others previously proposed. A second limit is the retrospective inclusion of
56 the patients with PA: a prospective validation in a large number of patients is warranted to confirm

.57 and further validate our prediction models. Moreover, our score cannot be applied to patients with
.58 PA diagnosed by the furosemide upright posture test or the oral saline loading test. Finally, it should
.59 be noted that dichotomization into LPA and BPA reflects the need to address patients to surgical vs.
.60 medical treatment and does not represent the complexity of the disease. Many patients with
.61 unilateral disease are cases with bilateral but asymmetrical aldosterone production displaying a high
.62 LI at AVS. These cases should benefit from adrenalectomy and are therefore considered as patients
.63 with unilateral or lateralized PA [32]. The strengths of our study include the reliable accuracy of
.64 our diagnostic models after internal and external validation, using both machine learning algorithms,
.65 or a simple scoring system, with a potential impact on clinical practice for centers where AVS is
.66 not available. In addition, we proposed two user-friendly downloadable tools which integrate the
.67 RF model and the flow-chart based on the SPACE score, allowing their application for the
.68 management of PA patients.

.69

.70 **CONCLUSIONS**

.71 We here developed and validated two prediction model and an easy applicable scoring system for
.72 the subtype diagnosis of PA. Our findings support the integration of clinical, biochemical, and
.73 imaging parameters by advanced computational approaches, to define PA subtype diagnosis,
.74 potentially reducing the number of AVS for patients with confirmed PA and guiding surgical
.75 decision in centers where AVS is not available.

.76

.77 **Source(s) of Funding:** This research did not receive any specific grant. CA and MR are supported
.78 by the Else Kröner-Fresenius Stiftung (2019_A104) and by the Deutsche Forschungsgemeinschaft
.79 within the CRC/Transregio 205/1 “The Adrenal: Central Relay in Health and Disease”.

.80 **Conflict(s) of Interest/Disclosure(s):** the authors have nothing to disclose.

REFERENCES

1. Käyser SC, Dekkers T, Groenewoud HJ, van der Wilt GJ, Carel Bakx J, van der Wel MC, Hermus AR, Lenders JW, Deinum J. Study Heterogeneity and Estimation of Prevalence of Primary Aldosteronism: A Systematic Review and Meta-Regression Analysis. *J Clin Endocrinol Metab.* 2016;101(7):2826-2835.
2. Buffolo F, Monticone S, Burrello J, Tetti M, Veglio F, Williams TA, Mulatero P. Is Primary Aldosteronism Still Largely Unrecognized? *Horm Metab Res.* 2017;49(12):908-914.
3. Monticone S, Burrello J, Tizzani D, Bertello C, Viola A, Buffolo F, Gabetti L, Mengozzi G, Williams TA, Rabbia F, Veglio F, Mulatero P. Prevalence and Clinical Manifestations of Primary Aldosteronism Encountered in Primary Care Practice. *J Am Coll Cardiol.* 2017;69(14):1811-1820.
4. Mulatero P, Monticone S, Bertello C, Viola A, Tizzani D, Iannaccone A, Crudo V, Burrello J, Milan A, Rabbia F, Veglio F. Long-term cardio- and cerebrovascular events in patients with primary aldosteronism. *J Clin Endocrinol Metab.* 2013;98(12):4826-4833.
5. Monticone S, D'Ascenzo F, Moretti C, Williams TA, Veglio F, Gaita F, Mulatero P. Cardiovascular events and target organ damage in primary aldosteronism compared with essential hypertension: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol.* 2018;6(1):41-50.
6. Funder JW, Carey RM, Mantero F, Murad MH, Reincke M, Shibata H, Stowasser M, Young WF Jr. The Management of Primary Aldosteronism: Case Detection, Diagnosis, and Treatment: An ES Clinical Practice Guideline. *J Clin Endocrinol Metab.* 2016;101(5):1889-1916.
7. Williams TA, Lenders JWM, Mulatero P, Burrello J, Rottenkolber M, Adolf C, Satoh F, Amar L, Quinkler M, Deinum J, Beuschlein F, Kitamoto KK, Pham U, Morimoto R, Umakoshi H, Prejbisz A, Kocjan T, Naruse M, Stowasser M, Nishikawa T, Young WF Jr, Gomez-Sanchez CE, Funder JW, Reincke M; Primary Aldosteronism Surgery Outcome (PASO) investigators. Outcomes after adrenalectomy for unilateral primary aldosteronism: an international consensus

- .07 on outcome measures and analysis of remission rates in an international cohort. *Lancet Diabetes*
.08 *Endocrinol.* 2017;5(9):689-699.
- .09 8. Hundemer GL, Curhan GC, Yozamp N, Wang M, Vaidya A. Cardiometabolic outcomes and
.10 mortality in medically treated primary aldosteronism: a retrospective cohort study. *Lancet*
.11 *Diabetes Endocrinol.* 2018;6(1):51-59.
- .12 9. Burton TJ, Mackenzie IS, Balan K, Koo B, Bird N, Soloviev DV, Azizan EA, Aigbirhio F,
.13 Gurnell M, Brown MJ. Evaluation of the sensitivity and specificity of (11)C-metomidate
.14 positron emission tomography (PET)-CT for lateralizing aldosterone secretion by Conn's
.15 adenomas. *J Clin Endocrinol Metab.* 2012;97(1):100-109.
- .16 10. Heinze B, Fuss CT, Mulatero P, Beuschlein F, Reincke M, Mustafa M, Schirbel A, Deutschbein
.17 T, Williams TA, Rhayem Y, Quinkler M, Rayes N, Monticone S, Wild V, Gomez-Sanchez CE,
.18 Reis AC, Petersenn S, Wester HJ, Kropf S, Fassnacht M, Lang K, Herrmann K, Buck AK,
.19 Bluemel C, Hahner S. Targeting CXCR4 (CXC Chemokine Receptor Type 4) for Molecular
.20 Imaging of Aldosterone-Producing Adenoma. *Hypertension.* 2018;71(2):317-325.
- .21 11. Mulatero P, di Cella SM, Monticone S, Schiavone D, Manzo M, Mengozzi G, Rabbia F, Terzolo
.22 M, Gomez-Sanchez EP, Gomez-Sanchez CE, Veglio F. 18-hydroxycorticosterone, 18-
.23 hydroxycortisol, and 18-oxocortisol in the diagnosis of primary aldosteronism and its subtypes.
.24 *J Clin Endocrinol Metab.* 2012;97(3):881-889.
- .25 12. Williams TA, Peitzsch M, Dietz AS, Dekkers T, Bidlingmaier M, Riestler A, Treitl M, Rhayem
.26 Y, Beuschlein F, Lenders JW, Deinum J, Eisenhofer G, Reincke M. Genotype-specific steroid
.27 profiles associated with aldosterone-producing adenomas. *Hypertension.* 2016;67(1):139-145
- .28 13. Yang Y, Burrello J, Burrello A, Eisenhofer G, Peitzsch M, Tetti M, Knösel T, Beuschlein F,
.29 Lenders JWM, Mulatero P, Reincke M, Williams TA. Classification of microadenomas in
.30 patients with primary aldosteronism by steroid profiling. *J Steroid Biochem Mol Biol.*
.31 2019;189:274-282.

14. Monticone S, Viola A, Rossato D, Veglio F, Reincke M, Gomez-Sanchez C, Mulatero P. Adrenal vein sampling in primary aldosteronism: towards a standardised protocol. *Lancet Diabetes Endocrinol.* 2015;3(4):296-303.
15. Kempers MJ, Lenders JW, van Outheusden L, van der Wilt GJ, Schultze Kool LJ, Hermus AR, Deinum J. Systematic review: diagnostic procedures to differentiate unilateral from bilateral adrenal abnormality in primary aldosteronism. *Ann Intern Med.* 2009;151(5):329-337.
16. Williams TA, Burrello J, Sechi LA, Fardella CE, Matrozova J, Adolf C, Baudrand R, Bernardi S, Beuschlein F, Catena C, Doumas M, Fallo F, Giacchetti G, Heinrich DA, Saint-Hilary G, Jansen PM, Januszewicz A, Kocjan T, Nishikawa T, Quinkler M, Satoh F, Umakoshi H, Widimský J Jr, Hahner S, Douma S, Stowasser M, Mulatero P, Reincke M. Computed tomography and adrenal venous sampling in the diagnosis of unilateral primary aldosteronism. *Hypertension.* 2018;72(3):641-649.
17. Küpers EM, Amar L, Raynaud A, Plouin PF, Steichen O. A clinical prediction score to diagnose unilateral primary aldosteronism. *J Clin Endocrinol Metab.* 2012;97(10):3530-3537.
18. Nanba K, Tsuiki M, Nakao K, Nanba A, Usui T, Tagami T, Hirokawa Y, Okuno H, Suzuki T, Shimbo T, Shimatsu A, Naruse M. A subtype prediction score for primary aldosteronism. *J Hum Hypertens.* 2014;28(12):716-720.
19. Kocjan T, Janez A, Stankovic M, Vidmar G, Jensterle M. A new clinical prediction criterion accurately determines a subset of patients with bilateral primary aldosteronism before adrenal venous sampling. *Endocr Pract.* 2016;22(5):587-594.
20. Kobayashi H, Haketa A, Ueno T, Ikeda Y, Hatanaka Y, Tanaka S, Otsuka H, Abe M, Fukuda N, Soma M. Scoring system for the diagnosis of bilateral primary aldosteronism in the outpatient setting before adrenal venous sampling. *Clin Endocrinol (Oxf).* 2017;86(4):467-472.
21. Kamemura K, Wada N, Ichijo T, Matsuda Y, Fujii Y, Kai T, Fukuoka T, Sakamoto R, Ogo A, Suzuki T, Umakoshi H, Tsuiki M, Naruse M. Significance of adrenal computed tomography in

- :57 predicting laterality and indicating adrenal vein sampling in primary aldosteronism. *J Hum*
:58 *Hypertens.* 2017;31(3):195-199.
- :59 22. Kobayashi H, Abe M, Soma M, Takeda Y, Kurihara I, Itoh H, Umakoshi H, Tsuiki M, Katabami
:60 T, Ichijo T, Wada N, Yoshimoto T, Ogawa Y, Kawashima J, Sone M, Inagaki N, Takahashi K,
:61 Watanabe M, Matsuda Y, Shibata H, Kamemura K, Yanase T, Otsuki M, Fujii Y, Yamamoto
:62 K, Ogo A, Nanba K, Tanabe A, Suzuki T, Naruse M; JPAS Study Group. Development and
:63 validation of subtype prediction scores for the workup of primary aldosteronism. *J Hypertens.*
:64 2018;36(11):2269-2276.
- :65 23. Leung HT, Woo YC, Fong CHY, Tan KCB, Lau EYF, Chan KW, Leung JYY. A clinical
:66 prediction score using age at diagnosis and saline infusion test parameters can predict
:67 aldosterone-producing adenoma from idiopathic adrenal hyperplasia. *J Endocrinol Invest.*
:68 2020;43(3):347-355.
- :69 24. Burrello J, Burrello A, Pieroni J, Sconfienza E, Forestiero V, Rabbia P, Adolf C, Reincke M,
:70 Veglio F, Williams TA, Monticone S, Mulatero P. Data from: Development and validation of
:71 prediction models for subtype diagnosis of patients with primary aldosteronism. *J Clin*
:72 *Endocrinol Metab* 2020. Deposited 6 May 2020. <https://github.com/ABurrello/SPACE-score>
- :73 25. Burrello J, Burrello A, Stowasser M, Nishikawa T, Quinkler M, Prejbisz A, Lenders JWM,
:74 Satoh F, Mulatero P, Reincke M, Williams TA. The Primary Aldosteronism Surgical Outcome
:75 Score for the Prediction of Clinical Outcomes After Adrenalectomy for Unilateral Primary
:76 Aldosteronism. *Ann Surg.* 2019 [Epub ahead of print]. doi: 10.1097/SLA.0000000000003200.
- :77 26. Sze WC, Soh LM, Lau JH, Reznek R, Sahdev A, Matson M, Riddoch F, Carpenter R, Berney
:78 D, Grossman AB, Chew SL, Akker SA, Druce MR, Waterhouse M, Monson JP, Drake WM.
:79 Diagnosing unilateral primary aldosteronism - comparison of a clinical prediction score,
:80 computed tomography and adrenal venous sampling. *Clin Endocrinol (Oxf).* 2014;81(1):25-30.
- :81 27. Riestler A, Fischer E, Degenhart C, Reiser MF, Bidlingmaier M, Beuschlein F, Reincke M,
:82 Quinkler M. Age below 40 or a recently proposed clinical prediction score cannot bypass

- .83 adrenal venous sampling in primary aldosteronism. *J Clin Endocrinol Metab.*
.84 2014;99(6):E1035-1039.
- .85 28. Venos ES, So B, Dias VC, Harvey A, Pasiaka JL, Kline GA. A clinical prediction score for
.86 diagnosing unilateral primary aldosteronism may not be generalizable. *BMC Endocr Disord.*
.87 2014;14:94.
- .88 29. Zhang Y, Niu W, Zheng F, Zhang H, Zhou W, Shen Z, Xu J, Tang X, Zhang J, Gao PJ, Wang
.89 JG, Zhu L. Identifying unilateral disease in Chinese patients with primary aldosteronism by
.90 using a modified prediction score. *J Hypertens.* 2017;35(12):2486-2492.
- .91 30. Young WF Jr, Klee GG. Primary aldosteronism. Diagnostic evaluation. *Endocrinol Metab Clin*
.92 *North Am.* 1988;17(2):367-395.
- .93 31. Mulatero P, Stowasser M, Loh KC, Fardella CE, Gordon RD, Mosso L, Gomez-Sanchez CE,
.94 Veglio F, Young WF Jr. Increased diagnosis of primary aldosteronism, including surgically
.95 correctable forms, in centers from five continents. *J Clin Endocrinol Metab.* 2004;89(3):1045-
.96 1050.
- .97 32. Vaidya A, Mulatero P, Baudrand R, Adler GK. The Expanding Spectrum of Primary
.98 Aldosteronism: Implications for Diagnosis, Pathogenesis, and Treatment. *Endocr Rev.*
.99 2018;39(6):1057-1088.

i00 **FIGURE LEGENDS**

i01

i02 *Legend to Figure 1 – Diagnostic Modelling: Linear Discriminant Analysis.* The LDA model
i03 included the 6 variables with the highest classification power for subtype diagnosis in the combined
i04 cohort (N = 215). **Panel A**, canonical plot representing diagnostic performance of LDA; each patient
i05 is indicated by a point and subtype diagnosis are reported by color (LPA, lateralized PA, black;
i06 BPA, bilateral PA, grey). The axes (canonical component 1 and 2) are calculated by weighted linear
i07 combination of the 6 variables included in the model to maximize the separation between groups.
i08 The crosses indicate the means of (canonical 1; canonical 2) for patients with LPA or BPA, the
i09 ellipse included patients with a linear combination coefficient that falls within the mean \pm SD. **Panel**
i10 **B**, confusion matrix reporting real and predicted diagnosis, accuracy, sensitivity, specificity, and
i11 10K-cross validation. **Panel C**, histogram representing normalized LDA coefficients for each
i12 variable included in the model. CT, Computed Tomography.

i13

i14 *Legend to Figure 2 – Diagnostic Modelling: Random Forest.* The RF algorithm included the 6
i15 variables with the highest classification power for subtype diagnosis in the combined cohort (N =
i16 215). **Panel A**, the first classification tree of the forest is shown for the prediction of LPA (lateralized
i17 PA) vs. BPA (bilateral PA). **Panel B**, confusion matrix reporting real and predicted diagnosis,
i18 accuracy, sensitivity, specificity, and 10K-cross validation. **Panel C**, histogram representing
i19 normalized predictive coefficients for each variable included in the model. CT, Computed
i20 Tomography.

i21

i22 *Legend to Figure 3 – Score development and validation.* Univariate/multivariate regression
i23 analyses and coefficients from the LDA and RF models were used to assign points to each variable
i24 according to stratification level. The score was developed in the training cohort (N = 150) and tested
i25 on the validation cohort (N = 65). **Panel A**, table showing included variables and final point system

i26 used for the score. **Panel B**, receiver operating characteristics (ROC) curve to assess AUC (area
i27 under the curve) and the best cut-off for the score in the combined cohort (N = 215). **Panel C**,
i28 representation of cut-offs and assigned points for each variable after categorization: subtype
i29 diagnosis is represented by colors (LPA, lateralized PA, black; BPA, bilateral PA, grey); the bars
i30 indicate median and interquartile range for each group. **Panel D**, confusion matrix representing real
i31 and predicted subtype diagnosis, accuracy sensitivity, specificity for the training cohort (N = 150),
i32 the validation cohort (N = 65), and the combined cohort (N = 215). CT, Computed Tomography;
i33 CI, Confidence Interval.

i34

i35 *Legend to Figure 4 – Score performance and management of PA patients.* Flow chart for PA
i36 patient management using our prediction score. **Panel A**, histogram showing the proportion of
i37 patients (y-axis, %) for each subtype diagnosis (LPA, lateralized PA, black; BPA, bilateral PA,
i38 grey), stratified by score points (x-axis) on the combined cohort (N = 215). The total number of
i39 patients (N) for each AVS score level and their proportion (%) are reported in Table S7 [24]. **Panel**
i40 **B**, PA patient management using our score; the number of patients is indicated in bold; cut-offs and
i41 misclassified patients are indicated in grey. **Panel C**, confusion matrix representing real and
i42 predicted subtype diagnosis, sensitivity, specificity, positive predictive value (PPV), and negative
i43 predictive value (NPV). AVS, Adrenal Venous Sampling; MRA, Mineral Receptor Antagonist.

i44 **Table 1. Patient Characteristics of Study Cohort**

Variable	LPA (N = 133)	BPA (N = 82)	P-value
Female sex, n (%)	56 (42.1)	19 (23.2)	0.005
Age at diagnosis (years)	49 ± 10.5	50 ± 7.7	0.248
Duration of HTN (months)	74 [27; 168]	63 [22; 123]	0.284
Systolic BP (mmHg)	165 ± 25.0	163 ± 20.5	0.613
Diastolic BP (mmHg)	99 ± 14.5	99 ± 11.7	0.873
Antihypertensive medication (DDD)	3.8 [2.2; 5.7]	3.0 [1.3; 4.7]	0.027
eGFR (mL/min)	96 [81; 109]	94 [80; 102]	0.146
Lowest Potassium (mEq/L)	3.1 ± 0.6	3.8 ± 0.4	< 0.001
PRA at screening (ng/mL/h)	0.30 [0.20; 0.40]	0.20 [0.10; 0.40]	0.554
Aldosterone at screening (ng/dL)	38.0 [25.7; 49.7]	28.7 [19.8; 37.9]	< 0.001
Confirmatory testing			
Saline infusion test, n (%)	102 (76.7)	63 (76.8)	0.982
Captopril Challenge test, n (%)	31 (23.3)	19 (23.2)	
PRA post-confirmatory test (ng/mL/h)	0.15 [0.10; 0.20]	0.15 [0.10; 0.21]	0.850
Aldosterone post-confirmatory test (ng/dL)	20.5 [13.3; 32.9]	11.5 [8.2; 17.7]	< 0.001
Microalbuminuria, n (%)	42 (31.5)	24 (29.4)	0.800
LVH at Echo, n (%)	81 (60.7)	48 (59.1)	0.831
CV events, n (%)	17 (12.6)	15 (18.1)	0.320
Presence of nodule at CT scanning, n (%)	114 (85.7)	34 (41.5)	< 0.001
Largest nodule at CT scanning (diameter, mm)	14 [10; 20]	12 [10; 19]	0.315
CT scanning findings			
Bilaterally Normal	5 (3.8)	20 (24.4)	< 0.001
Bilaterally Abnormal	19 (14.2)	18 (22.0)	
Unilateral Abnormality	109 (82.0)	44 (53.7)	
AVS protocol			
Basal, n (%)	43 (32.3)	37 (45.1)	0.051
ACTH continuous infusion, n (%)	51 (38.4)	32 (39.0)	
Both (Basal + ACTH), n (%)	39 (29.3)	13 (15.9)	
Lateralization Index at AVS	12.0 [6.9; 21.3]	1.8 [1.3; 2.6]	< 0.001
Clinical outcome: Complete, n (%)	72 (54.1)	N.A.	N.A.
[only for LPA] Partial, n (%)	55 (41.4)		
Absent, n (%)	6 (4.5)		
Biochemical outcome: Complete, n (%)	131 (98.5)	N.A.	N.A.
[only for LPA] Partial, n (%)	2 (1.5)		
Absent, n (%)	0 (0.0)		

i45

i46 Clinical characteristics of patients included in the analysis stratified for diagnosis: patients with
i47 lateralized PA (LPA; N = 133) vs. bilateral PA (BPA; N = 82). The DDD is the assumed average
i48 maintenance dose per day for a drug used for its main indication in adults. AVS, Adrenal Venous
i49 Sampling; HTN, Hypertension; BP, Blood Pressure; DDD, Defined Daily Dose; eGFR, estimated

50 Glomerular Filtration Rate; PRA, Plasma Renin Activity; LVH, Left Ventricular Hypertrophy;
 51 Echo, Echocardiography; CV, Cardiovascular; CT, Computed Tomography. Normally and non-
 52 normally distributed variables were reported as mean \pm standard deviation or median [interquartile
 53 range], respectively. Categorical variables were reported as absolute number (n) and proportion (%).
 54

55 **Table 2. Selected discriminant variables for a diagnosis of lateralized PA**

Variable (ref. LPA)	Univariate analysis		Multivariate Analysis	
	OR (CI 95%)	<i>P</i> -value	OR (CI 95%)	<i>P</i> -value
Aldosterone at screening (ng/dL)	1.04 (1.02 – 1.07)	< 0.001	1.05 (1.01 – 1.10)	0.017
Lowest Potassium (mEq/L)	0.10 (0.05 – 0.21)	< 0.001	0.09 (0.03 – 0.30)	< 0.001
Aldosterone post-confirmatory test (ng/dL)	1.09 (1.05 – 1.12)	< 0.001	1.09 (1.02 – 1.16)	0.012
Nodule at CT scanning (ref. presence)	8.33 (4.35 – 16.67)	< 0.001	12.50 (2.94 – 47.62)	0.001
Largest nodule at CT scanning (diameter, mm)	1.12 (1.07 – 1.16)	< 0.001	1.11 (1.06 – 1.16)	0.013
CT scanning findings (ref. unilateral abnormality)	9.91 (3.50 – 28.05)	< 0.001	4.44 (1.30 – 13.21)	0.016

56
 57 Logistic regression analysis was performed to assess the odds ratio (OR) and the 95% confidence
 58 interval (CI) for each variable. Univariate and multivariate analysis are shown as indicated. An OR
 59 greater than 1 indicates an increased likelihood of lateralized PA (LPA), and an OR less than 1 a
 60 decreased likelihood. Aldosterone at screening, lowest potassium, aldosterone post-confirmatory
 61 test, and largest nodule at CT were treated as continuous variables. An OR increase of 0.01
 62 represents a 1% increased likelihood of a diagnosis of LPA for each unit of the reference variable.
 63 Presence/absence of nodule at CT scanning, and CT scanning findings were treated as categorical
 64 variables. CT, Computed Tomography.