

Microphysical time scales, supersaturation fluctuations and droplet distance-neighbour statistical analysis at a warm Cloud Top Boundary

*Original*

Microphysical time scales, supersaturation fluctuations and droplet distance-neighbour statistical analysis at a warm Cloud Top Boundary / Abdunabiev, Shahbozbek; Fossa', Ludovico; GOLSHAN KOVIJI, Mina; Tordella, Daniela. - ELETTRONICO. - 75:(2022). (Intervento presentato al convegno 75th Annual Meeting of the Division of Fluid Dynamics).

*Availability:*

This version is available at: 11583/2977896 since: 2023-04-12T11:17:04Z

*Publisher:*

APS

*Published*

DOI:

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

Abstract Submitted  
for the DFD75 Meeting of  
The American Physical Society

Sorting Category: L33.00010 (experimental)

**Microphysical time scales, supersaturation fluctuations and droplet distance-neighbour statistical analysis at a warm Cloud Top Boundary** SHAHBOZBEK ABDUNABIEV, LUDOVICO FOSSA', MINA GOLSHAN KOVIJI, DANIELA TORDELLA, Dipartimento di Scienza Applicata e Tecnologia (DISAT), Politecnico di Torino, Italy — Recent results have shown that there is an acceleration in the spread of the size distribution of droplet populations in the region bordering the cloud and undersaturated ambient. We have analyzed the supersaturation balance in this region, which is typically a highly intermittent shearless turbulent mixing layer, under a condition where there is no mean updraft. We have investigated the evolution of the cloud - clear air interface and of the droplets therein via direct numerical simulations. We have compared horizontal averages of the phase relaxation, evaporation, reaction and condensation times within the cloud-clear air interface for the size distributions of the initial monodisperse and polydisperse droplet populations. For the monodisperse population, a clustering of the values of the reaction, phase and evaporation times, that is around 20-30 seconds, is observed in the central area of the mixing layer, just before the location where the maximum value of the supersaturation turbulent flux occurs. This clustering of values is similar for the polydisperse population but also includes the condensation time. The mismatch between the time derivative of the supersaturation and the condensation term in the interfacial mixing layer is correlated with the planar covariance of the horizontal longitudinal velocity derivatives of the carrier air flow and the supersaturation field, thus suggesting that a quasi-linear relationship may exist between these quantities. We will also present results on turbulence dispersion in the cloud - clear air interface based on the distance neighbor graph statistics (Richardson, 1926) obtained by tracking water droplets in the Lagrangian framework. This project has received funding from the Marie-Sklodowska Curie Actions (MSCA ITN ETN COM-LETE) under the European Union's Horizon 2020 research and innovation program. Grant agreement 675675, <http://www.complete-h2020network.eu>

Prefer Oral Session  
 Prefer Poster Session

Shahbozbek Abdunabiev  
shahbozbek.abdunabiev@polito.it  
DISAT, Politecnico di Torino, Italy

Date submitted: September 7, 2022

Electronic form version 1.2