POLITECNICO DI TORINO
Repository ISTITUZIONALE

Sea-crossing behaviour of two migratory eagles in relation to weather conditions

Original
Sea-crossing behaviour of two migratory eagles in relation to weather conditions / Agostini, N., Muñoz Arroyo, G., De La Cruz, A., Benjumea, R., Guerrero Molina, M., Pérez, B., Rollán, L., Mellone, U., Von Hardenberg, J., Chiatante, G.. - In: JOURNAL OF ORNITHOLOGY. - ISSN 2193-7206. - (2026). [10.1007/s10336-025-02310-3]

Availability:
This version is available at: 11583/3004116 since: 2025-11-24T19:38:07Z

Publisher:
Springer Nature

Published
DOI:10.1007/s10336-025-02310-3

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

12 September 2026

SEA CROSSING BEHAVIOUR OF TWO MIGRATORY EAGLES IN RELATION TO WEATHER CONDITIONS

Nicolantonio Agostini[1], Gonzalo Muñoz Arroyo[2], Andrés De La Cruz[2,3], Rafael Benjumea[4,5], Marina Guerrero Molina[6], Blanca Pérez[6], Laura Rollán[3], Ugo Mellone[1,7], Jost von Hardenberg[8], Gianpasquale Chiatante[1,9]

[1] *MEDRAPTORS (Mediterranean Raptor Migration Network), Pavia, Italy*

[2] *Department of Biology, University of Cádiz, P.O. Box 40, 11510 Puerto Real, Cádiz, Spain*

[3] *Institute of Marine Research (INMAR), International Campus of Excellence in Marine Science (CEIMAR), University of Cádiz, 11510, Puerto Real, Cádiz, Spain.*

[4] *University of Málaga, Andalucía Tech, Facultad de Ciencias, Campus de Teatinos s/n 29071 Málaga, Spain.*

[5] *Biodiversity Research Institute (IMIB), CSIC/UO/PA, Campus de Mieres, Edificio de Investigación, 33600 Mieres (Asturias), Spain.*

[6] *Vida Silvestre Ibérica, Madrid, Spain. Cl. de la Navazuela, 18, 28410 Manzanares el Real, Madrid.*

[7] *Vertebrates Zoology Research Group, University of Alicante, Alicante, Spain*

[8] *Department of Environment, Land and Infrastructure Engineering (DIATI) Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy.*

[9] *University of Tuscia, Department of Ecological and Biological Sciences, Largo dell'Università s/n, 01100, Viterbo, Italy*

ORCID

Nicolantonio Agostini: 0000-0001-9845-7794

Gonzalo Muñoz Arroyo: 0000-0002-1064-5342

Laura Rollán: 0000-0001-6502-6955

Andrés De La Cruz: 0000-0002-0363-2498

Rafael Benjumea: 0000-0002-0569-8765

Marina Guerrero Molina: 0000-0001-7803-2238.

Ugo Mellone: 0000-0001-9665-505X

Jost von Hardenberg: 0000-0002-5312-8070

Gianpasquale Chiatante: 0000-0002-4570-9350

* Corresponding author: Nicolantonio Agostini, nicolantonioagostini@gmail.com

Short title: Sea crossing behaviour of migratory eagles

Abstract

During migration, typical soaring species such as eagles and vultures primarily rely on soaring flight to minimize energy expenditure, exploiting thermal currents over land while avoiding prolonged water crossings. This study investigates the flight behaviour of the Booted Eagle (*Hieraaetus pennatus*) and the Short-toed Snake Eagle (*Circaetus gallicus*) as they approach the European coastline of the Strait of Gibraltar, a critical migration bottleneck spanning at least 14 km, during their pre-breeding migration. Visual observations were conducted from two watchpoints on the European side of the Strait to assess migration magnitude and record the birds' flight altitudes upon reaching the coastline. Both species experienced wind drift when flying in crosswinds. Under calm conditions or light easterly winds, they tended to converge at the center of the Strait, whereas westerly winds caused a shift in their migratory flow toward the eastern sector. Short-toed Snake Eagles were observed flying at lower altitudes when facing headwinds or strong crosswinds, increasing their risk of drowning. In contrast, Booted Eagles were less affected, likely due to their lower energetic cost of flapping flight, which helped mitigate drowning risks. Notably, crosswind intensity did not significantly influence the flight altitude of Booted Eagles. These findings highlight interspecific differences in response to challenging wind conditions during water crossings and provide insight into the flight strategies adopted by raptors in the Strait of Gibraltar.

Keywords *Circaetus gallicus*, *Hieraaetus pennatus*, Mediterranean, flight altitude, migration, optimization, transport costs

Statements and declarations

Acknowledgements

The spring raptor count campaigns in the Strait of Gibraltar were conducted as part of the Lifewatch ERIC INDALO project (https://www.juntadeandalucia.es/medioambiente/portal/proyecto-indalo/el-proyecto-indalo), promoted by the Regional Government of Andalusia and the Andalusian Agency for Environment and Water (AMAYA). Andrés De la Cruz was funded by the Margarita Salas Grant (021-067/PN/MS-RECUAL/CD) from the Ministry of Universities of the Government of Spain and the European Union (Funded by the European Union – NextGenerationEU. Gianpasquale Chiatante acknowledges the support of University of Tuscia, project ECS00000024 Rome Technopole (CUP B83C22002820006, PNRR Missione 4 Componente 2 Investimento 1.5), funded by the European Union – NextGenerationEU. Our colleague and friend Michele Panuccio, who sadly passed away in 2019, started the fieldwork of this project in 2011. Finally, we would like to express our sincere appreciation to the three anonymous reviewers for their insightful and constructive comments, which significantly enhanced the quality and clarity of this manuscript.

Competing interests

The authors have no competing interests to declare that are relevant to the content of this article

Data availability

The dataset generated during and/or analyzed during the current study is available from the corresponding author on a reasonable request.

Author contribution

Nicolantonio Agostini, Ugo Mellone and Gianpasquale Chiatante contributed to the study conception and design. Data were collected by Nicolantonio Agostini, Ugo Mellone, Gonzalo Muñoz Arroyo, Andrés De La Cruz, Rafael Benjumea, Marina Guerrero Molina, Blanca Pérez, and Laura Rollán. Analysis were performed by Gianpasquale Chiatante and Jost von Hardenberg. The first draft of the manuscript was written by Nicolantonio Agostini and Gianpasquale Chiatante and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

INTRODUCTION

During migration, soaring raptors take advantage of thermals and deflection updrafts to travel long distances while minimizing energy expenditure. Since vertical currents are typically weaker or absent over water surfaces (especially in temperate zones), birds cannot soar as much, therefore making them reliant on flapping flight, which is costly and risky (Agostini et al. 2015). Although recent research has shown that some facultative soaring raptors, such as the Osprey (*Pandion haliaetus*) and the Red Kite (*Milvus milvus*), can exploit thermals even over the open sea, adopting complex flight strategies that integrate wind support and available vertical uplift over marine surfaces (Duriez et al. 2018; Škrábal et al. 2023), in the case of obliged soaring raptors (e.g. eagles and vultures), landmass distribution clearly shaped their migratory strategies, particularly across the Mediterranean basin (Mellone 2020; Panuccio et al. 2021). Wind conditions play a crucial role in this phase. Favourable winds, such as tailwinds or supportive crosswinds, reduce energy consumption and travel time, facilitating crossings. In contrast, headwinds or adverse crosswinds increase the risks of overwater flights. Several studies indicate that large and medium-sized land birds face higher mortality when crossing water bodies (Zu-Aretz and Leshem 1983; Oppel et al. 2015; Newton 2023; Panuccio et al. 2021). As a result, heavier species with high energetic costs of flapping flight, such as vultures and eagles, typically circumvent the Mediterranean, concentrating their passage through isthmuses and narrow straits (e.g., the Bosphorus and Gibraltar). In contrast, lighter species, such as the European Honey Buzzard (*Pernis apivorus*), the Black Kite (*Milvus migrans*), harriers (*Circus* spp.), and falcons (*Falco* spp.), undertake longer water crossings due to their more efficient flapping flight and lower energy expenditure (Kerlinger 1989; Agostini et al. 2015).

Among European eagles, the Booted Eagle (*Hieraaetus pennatus*) is an exception. This relatively small raptor has a flight morphology associated with lower flapping flight costs, similar to facultative soarers such as the European Honey Buzzard and Black Kite, and lower than that of obligate soarers such as the Short-toed Snake Eagle (*Circaetus gallicus*) and the Lesser Spotted Eagle (*Clanga pomarina*; Agostini et al. 2015; Supplementary Materials, Fig. S1; Agostini et al. 2015). Despite its efficient flight capabilities, the Booted Eagle behaves like a typical obligate soaring species, generally avoiding long water crossings, with rare exceptions (Mellone 2021). Consequently, most Booted Eagles from Europe cross the Mediterranean Sea via the Strait of Gibraltar (14.7 km wide at its narrowest point, Fig. 1) and, to a lesser extent, via the Bosphorus (1–3 km wide; Mellone 2021), a pattern also observed in the Short-toed Snake Eagle (Agostini 2021).

In this study, we analysed the factors affecting migratory flow and flight altitude in these two eagle species as they reached the eastern and western European coasts of the Strait of Gibraltar during pre-breeding migration. Our goal was to investigate potential species-specific flight strategies shaped by differences in transport costs. We expected both species to exhibit site-specific differences in migratory flow based on wind direction. Specifically, we predicted that more individuals would be observed at the western watchsite (Cazalla) under easterly winds and at the eastern watchsite (Punta Carnero) under westerly winds due to wind-driven drift patterns. We also predicted that stronger crosswinds and headwinds would force Short-toed Snake Eagles (the larger species) to fly at lower altitudes upon reaching the European coastline, thereby increasing their risk of drowning. Conversely, Booted Eagles, owing to their lower flapping flight costs, would be less affected by wind intensity in terms of flight altitude.

Finally, we expected both species to reach the European coastline at higher altitudes under favourable thermal conditions. Under poor thermal conditions (e.g., high cloud cover and weak thermals), we predicted that individuals would compensate by increasing their flapping effort, leading to species-specific differences in crossing strategies. In particular, Booted Eagles, benefiting from a lower energetic cost of flapping flight, were expected to adopt a more risk-averse strategy by maintaining higher flight altitudes in adverse conditions (e.g., headwinds and strong crosswinds), whereas Short-toed Snake Eagles were likely to be more susceptible to altitude loss.

MATERIALS AND METHODS

Study area and data collection

The Strait of Gibraltar is the main bottleneck for soaring raptors breeding in Western Europe and wintering in Sub-Saharan Africa, funnelling them across the Mediterranean Sea during migration (De la Cruz et al. 2011a; Panuccio et al. 2021). Here, crosswinds prevail during both spring and autumn: westerly winds in spring, and easterly winds in autumn (Finlayson 2010; Hidalgo and Gallego 2019). We used a dual approach, taking advantage of two different dataset: (1) magnitude of migratory flow (from the watchsites with optimal visibility) and (2) birds' flight altitude (recorded from watchsites closer to the sea). Therefore, to quantify the magnitude of the migratory flow, we carried out systematic counts of Booted and Short-toed Snake Eagles crossing the Strait in spring. Observations were carried out daily from 11:00 to 17:00 local time between 1 March and 15 April in 2022 and 2023, using binoculars and telescopes. Data were collected from two watchpoints: Punta Carnero (-5.429 °E, 36.076 °N, elevation = 41 m a.s.l.) and Cazalla (-5.577 °E, 36.033 °N, elevation = 158 m a.s.l.) (Fig. 1).

169 Previous spring surveys in 2008 and 2009, covering six watchpoints between Bolonia
170 and Punta Carnero (Fig. 1), revealed that these two locations accounted for more than
171 70% of migrating raptors (De la Cruz et al. 2011b). Moreover, to determine flight
172 altitude over the sea, we used two observation posts along the coastline, approximately
173 15 km apart: Trafico (-5.587 °E, 36.014 °N; elevation = 52 m a.s.l.), located two km
174 south of Cazalla, and Punta Carnero (Fig. 1). We recorded flight altitude measurements
175 as soon as the raptors became visible upon reaching the coastline, while they were still
176 flying over the sea and not yet over land, between 09:00 and 19:00 local time, from 1 to
177 31 March in 2011, 2017, and 2022. This period coincides with the peak overlap of the
178 spring migration of these species at the Strait of Gibraltar (Finlayson 2010).
179 We used a Leica Rangemaster 1600-R optical range finder to determine bird
180 altitude relative to the observer, accounting for non-overhead or lower-altitude passes.
181 Using a trigonometric formula for a right triangle, we calculated bird altitude based on
182 the acute angle and distance from the observer. Finally, we derived the bird's altitude
183 above the sea was by incorporating the watchpoint's altitude (SM, Fig. S2).
184
185 **Weather data**
186 We analysed the effect of five weather variables on both species flow and flight
187 altitude, interpolated for the exact date and time of observations. These data were
188 obtained from Movebank (www.movebank.org), a free online database of animal
189 tracking and related environmental data, which compiles information from the most
190 relevant gridded global datasets (Dodge et al. 2013; Wikelski and Kays 2018). Thereby,
191 from the Movebank ECMWF ERA5 weather dataset, we obtained cloud cover (%
192 coverage) and temperature (°C), whereas from the Movebank ECMWF ERA-Interim

193 weather dataset, we extracted horizontal wind components, namely the eastward (U)
194 and the northward (V) wind speeds (m/s). Positive U values indicate flow toward the
195 east (westerly wind), while positive V values indicate flow toward the north (southerly
196 wind) (Wallace and Hobbs 2006).
197 For species flow analysis, weather data were extracted for a pressure level
198 corresponding to 110 m, approximately the mean flight altitude observed (see *Results*).
199 To explore flight altitude, data were bilinearly interpolated by Movebank for each
200 watchsite at the exact measured flight altitude. To describe wind conditions during the
201 study period, we obtained daily horizontal wind data (extracted at 12:00 AM at 100 m
202 above ground) from the Movebank ECMWF ERA5 SL dataset. Additionally, we
203 retrieved hourly ERA5 horizontal wind components from the ECMWF Climate Data
204 Store (Hersbach et al. 2023) at a 950 hPa pressure level (corresponding to
205 approximately 500 m altitude), for the observation days, matching the presumed
206 average flight altitude of raptors crossing of the Strait of Gibraltar (see Santos et al.
207 2020). Wind maps for the Gibraltar region were computed by averaging daily mean
208 values, weighted between 9:00 UTC and 17:00 UTC based on the number of
209 observations of each species at Punta Carnero and Cazalla.
210
211
212 **Statistical analysis**
213 First, we used the χ^2 test and the Kruskal-Wallis test to verify the presence of
214 significant differences in wind direction, wind speed, and the number of days with slow
215 winds (< 2 m/s) between the three watchsites (Trafico, Cazalla, Punta Carnero). In
216 addition, we compared flight altitudes between species with the non-parametric Mann-

Whitney *U* test. Then, we investigated the flow and flight altitude of both species in relation to the weather variables through regression analysis, using daily counts with ≥ 1 individual observed. First, we ran an exploratory analysis showing the importance of quadratic effects of the variables (i.e., $y \sim x + x^2$) through a pair-wise comparison of the second-order Akaike information criterion (AICc; Akaike 1973) of two linear models: one with the linear terms ($y \sim x$) and the other with the quadratic term ($y \sim x + x^2$) (Burnham et al. 2011). When the AICc value of the model with the quadratic effect was lower than the model with the linear term, with a difference of at least two (ΔAICc ≤ 2), the quadratic effect was retained, otherwise, we retained the linear term (Burnham and Anderson 2002) (SM, Tables S1-S2). In the same way, we defined the importance of interaction terms between each variable and the watchsite (SM, Tables S1-S2).

Then, using the covariates retained in the exploratory analysis, we built four multiple linear regressions, (which could have different variable combinations), one for each species and one for each response variable (flight altitude, species flow). Because the response variables were not normally distributed, we log-transformed them, reaching normality (SM, Table S4). Correlation matrices showed that covariates were not highly correlated (r_S <|0.60|) (SM, Table S3). Considering the different units of measurements used in the analysis, the covariates were standardized by normalization (Quinn and Keough 2002). For all the models, we performed model selection using the Akaike Information Criterion (AIC) on all possible combinations of the covariates, with the function *dredge* of the package "MuMIn" (Bartoń 2018) for R 4.1.1 (R Core Team 2021). We selected models within the '95% confidence set' based on cumulative Akaike weights; these models were averaged and the importance of each variable in the set *sw* was calculated (Burnham and Anderson 2002). The goodness-of-fit of the model

obtained was assessed by correlating expected and observed values with Spearman's rank correlation test, while the variance explained was measured with the coefficient of determination R^2. Statistical analyses were computed with R software version 4.1.1 (R Core Team 2021) and the packages "MuMIn" (Bartoń 2018) and "car" (Fox and Weisberg 2011).

RESULTS

Wind pattern

During the study period, wind direction did not differ between watchsites (χ^2 = 0.348, df = 6, P = 1.00), with easterly and westerly winds prevailing (92.7%). The average wind speed was 7.9 m/s ± 4.04 (SD), with no significant differences between watchsites (Kruskal-Wallis test, χ^2 = 0.05, df = 2, P = 0.976). Calm days (wind speed < 2 m/s) were rare, accounting for only 7.4% of the total.

Species flow

Short-toed Snake Eagle

A total of 13,360 Short-toed Snake Eagles (Cazalla, N = 2,890; Punta Carnero, N = 10,470) were observed in 2022, and 13,497 (Cazalla, N = 2,058; Punta Carnero, N = 11,439) in 2023. The analysis showed that three models were considered in the set of best models, including all the weather variables (SM, Table S5). The strength of the eastward component of the wind had a negative effect at Cazalla, while positive at Punta Carnero. This means that at Cazalla more birds were seen with easterly winds than westerly; at Punta Carnero, contrarily, more eagles were seen with westerly winds (Table 1; Fig. 2). The temperature and the strength of the northward component of the

265 wind had very weak effects, negative and positive respectively. Moreover, watchsite
266 was important too, with a positive effect on Punta Carnero. The variance explained by
267 the model R^2 is equal to 30.2% and expected values are correlated with observed ones
268 (r_s = 0.535, P < 0.001). Averaged wind maps weighted by counts are reported in Fig. 3:
269 they show that Short-toed Snake Eagles crossed the Strait with weak average winds
270 when observed at Cazalla and a strong westerly component when observed at Punta
271 Carnero, with no relevant average meridional component.
272
273 *Booted Eagle*
274 A total of 8,209 Booted Eagles (Cazalla, N = 5,075; Punta Carnero, N = 3,134)
275 were observed in 2022, and 9,460 (Cazalla, N = 2,367; Punta Carnero, N = 7,093) in
276 2023. The analysis showed that only one model was considered as the best one,
277 including all the weather variables (Table 1). As regards horizontal winds, the strength
278 of the eastward component had a negative effect at Cazalla, while positive at Punta
279 Carnero. This means that, similarly to the Short-toed Snake Eagle, at Cazalla more birds
280 were seen with easterly winds, whereas at Punta Carnero with strong westerly winds
281 (Fig. 2). In relation to the northward component, the flow showed a non-linear effect,
282 with more Booted eagles passing with no wind; this effect is stronger at Punta Carnero
283 than at Cazalla. In addition, the temperature had a positive effect on the species flow.
284 The variance explained by the model R^2 is equal to 44.6% and expected values are
285 correlated with observed ones (r_s = 0.668, P < 0.001). In this case, the analysis is
286 confirmed by the averaged wind maps at about 500 m, weighted by counts, reported in
287 Fig. 3: Booted Eagles were seen crossing the Strait with weak average easterly winds
288 when observed at Cazalla and a stronger westerly component when observed at Punta

289 Carnero. A moderate northerly meridional component was present for passages at Punta
290 Carnero.
291
292 **Flight altitude**
293 We analysed the effect of the weather on 996 and 222 flight altitude records, for
294 the Short-toed Snake Eagle and Booted Eagle, respectively. In general, Short-toed
295 Snake Eagles were detected at higher altitude than Booted Eagles and higher at Punta
296 Carnero than at Trafico (Short-toed Snake Eagle: 150 m ± 139.12 (SD), range 1-670 m;
297 Booted Eagle: 100 m ± 90.32, range 18-465; Mann-Whitney *U* test, $P < 0.001$) and both
298 species arrived higher at Punta Carnero than at Trafico (Punta Carnero: 154 m ± 147.62;
299 Trafico: 107 m ± 76.82; Mann-Whitney *U* test, $P < 0.001$). This was true even
300 considering species separately (Short-toed Snake Eagle: Punta Carnero, 148 m ±
301 119.34, Trafico, 118 m ± 72.61, $P = 0.039$; Booted Eagle: Punta Carnero, 121 m ±
302 106.97, Trafico, 94 m ± 80.36; $P = 0.004$).
303
304 *Short-toed Snake Eagle*
305 Only one model was considered in the set of best ones, including all the weather
306 variables (Table 2). In particular, the cloud cover had a positive effect on the flight
307 altitude, especially at Punta Carnero; conversely, the temperature and the strength of the
308 thermal uplift had a weak negative effect (Fig. 4). Flight height at Punta Carnero tended
309 to be higher with easterly winds, while at Trafico birds reached higher altitudes with
310 winds blowing from the west (Fig. 4). The strength of the northward component had a
311 marked quadratic effect, with opposite patterns between watchpoints, even though this
312 species tended to fly higher with northerly winds than with strong southerly winds (Fig.

4). The variance explained by the model R^2 is equal to 30.8% and expected values are correlated with observed ones (r_s = 0.522, P < 0.001).

Booted Eagle

The analysis showed that three best models explained the flight altitude of the Booted Eagle (Table S6), and the average model included all the variables with the exclusion of the cloud cover (Table 3). In particular, the strength of the northward component of the wind showed a strong quadratic effect on the flight altitude, with a different pattern between watchsites, whereas the thermal uplift had a weak negative effect (Fig. 5). Indeed, at Punta Carnero the altitude was higher with strong winds, both northerly and southerly, whereas at Trafico Booted Eagles flew higher with strong northerly winds than with southerly winds (Fig. 5). The temperature and the strength of the eastward component of the wind had very little importance. The variance explained R^2 is equal to 26.3% and the expected values are correlated with the observed ones (r_s = 0.471, P < 0.001).

DISCUSSION

Among the factors affecting the sea-crossing performance of soaring migrants, wind conditions, starting altitude, and the bird's physiological state are crucial to success (Kerlinger 1989; Bildstein 2006; Santos et al. 2020; Panuccio et al. 2021). Strong crosswinds during a sea crossing force birds to extend their flapping flight and/or lengthen their flight path over water, whereas a higher starting altitude allows a soaring bird to glide for longer before resorting to flapping flight (Kerlinger 1989). In our study, Short-toed Snake Eagles reached the Spanish coast at higher altitudes than Booted

Eagles, likely because they attained much greater starting altitudes along the Moroccan coast, as expected for an obligate soaring species. However, we cannot rule out the possibility that these findings were influenced by size differences between the two eagle species: the Short-toed Snake Eagle typically weighs around 1.5–2.3 kg with a wingspan of 160–190 cm, whereas the Booted Eagle is smaller, weighing approximately 0.8–1.0 kg with a wingspan of 110–135 cm (Ferguson-Lees & Christie 2001).

Our results confirm the prediction regarding the influence of wind direction on migratory flow. Both eagle species were clearly affected by drift when flying in crosswinds. Their migratory flow was concentrated at Cazalla under weak wind conditions or light easterly winds, while more individuals reached Punta Carnero when winds had a westerly component. A substantial number of birds were observed even farther eastward as westerly winds increased in strength. Conversely, under stronger easterly winds, birds were seen west of Tarifa (see also Finlayson 2010). Wind maps indicate that migration at Punta Carnero typically occurs under a relevant westerly wind component, whereas the conditions at Cazalla are characterized by only weak easterly winds. Our findings align with those of De la Cruz et al. (2011b), who reported a very low passage of both Booted and Short-toed Snake Eagles at western observatories of the Strait under easterly winds. Therefore, our results suggest that under strong easterly winds, most individuals of both species either delay or abort the crossing to avoid being drifted on a longer route over water. On the Atlantic side the risk of being drifted towards the open sea is much higher than in the Mediterranean due to the geography of the coastline, heading more abruptly towards the north-west.

The strength of crosswinds negatively affected the flight altitude of the larger species, supporting our hypothesis: Short-toed Snake Eagles were detected at lower altitudes

361 when crosswinds were stronger, likely because they were forced to cover longer
362 distances over water, leading to altitude loss. However, under such conditions, we
363 cannot exclude the possibility that they deliberately flew closer to the sea surface where
364 winds were weaker, thereby reducing energy costs but increasing the risk of drowning.
365 Similarly, Black Kites, a facultative soaring species, have been shown to experience
366 drift when flying in crosswinds through the Strait of Gibraltar during autumn migration,
367 often reaching the Moroccan coast at 20–100 m a.s.l. (Santos et al. 2020). Black Kites
368 also lower their flight altitude over water and increase flapping effort when flying in
369 strong crosswinds (Santos et al. 2020). Conversely, the few Short-toed Snake Eagles
370 that reached the two watchpoints while compensating for drift were observed at higher
371 altitudes, likely because they relied more on flapping rather than gliding during the
372 crossing. This strategy was probably adopted by individuals in better physical condition,
373 who reduced their risk by maintaining a more direct route at a higher altitude, albeit at
374 the cost of increased energy expenditure. Since birds are expected to adopt a time-
375 minimization strategy during spring migration, they may continue flying even in
376 suboptimal weather conditions (Duerr et al. 2014).
377 Unexpectedly, both eagle species reached the European coast at lower altitudes under
378 atmospheric conditions ideal for the formation of strong thermals over land (e.g., high
379 temperatures and weak winds). A likely explanation is that, under such conditions, they
380 were able to gain higher starting altitudes (see also Panuccio et al. 2019) but extensively
381 used gliding flight during the crossing to minimize energy expenditure by reducing
382 flapping effort. Similarly, during rare occurrences of southerly winds at the Strait of
383 Gibraltar, both species reached the European coast at lower altitudes, particularly at
384 Trafico, and were detected in lower numbers at the two watchpoints. In contrast with

our expectations, Short-toed Snake Eagles were observed at higher altitudes when flying under high cloud cover, despite such conditions being unfavourable for strong thermal formation. Under these circumstances, they may have compensated by increasing flapping flight effort during the crossing. Finally, in agreement with our predictions, the few Booted Eagles crossing in strong headwinds preferred to fly at higher altitudes, minimizing the risk of drowning. Their ability to maintain altitude was likely facilitated by their ability to sustain flapping flight at a lower energetic cost. For the same reason, crosswind intensity did not significantly influence their flight altitude. As mentioned above, recent studies have revealed that thermal uplift can also occur over water, challenging the traditional assumption that soaring flight is largely unsupported over marine surfaces (Duriez et al. 2018; Nourani et al. 2021; Škrábal et al. 2023). Although our analyses did not include uplift potential over the sea, such findings suggest that thermals over water could occasionally aid even obliged soaring raptors at least during short sea-crossings. This phenomenon may partially explain some of the observed variability in flight altitude at the Strait of Gibraltar, particularly in individuals maintaining relatively high altitudes in suboptimal conditions.

In conclusion, this study demonstrates that the two eagle species adopt distinct strategies when facing adverse conditions during sea crossings, with the Short-toed Snake Eagle being more susceptible to altitude loss than the Booted Eagle. These findings enhance our understanding of the role of morphological and flight performance differences in shaping migratory behaviour under variable wind conditions. By providing ground-based observational evidence on non-tagged individuals, this study fills a gap in the existing literature concerning overwater flight strategies in soaring raptors. Future research integrating ground observations with high-resolution weather

models and GPS tracking will be essential to explore individual variability, physiological condition, and the influence of localized uplift on sea-crossing performance.

References

Agostini N, Panuccio M, Pasquaretta C (2015) Morphology, flight performance, and water crossing tendencies of Afro-Palearctic raptors during migration. Current Zoology, 61**:** 951-958.

Agostini N (2021) Short-toed snake eagle (*Circaetus gallicus*). In: Panuccio M, Agostini N, Mellone U (eds) Migration strategies of birds of prey in Western Palearctic. CRC Press, Boca Raton, FL (US), pp 68–75

Agostini N, Gustin M, Von Hardenberg J, Panuccio M (2016) Wind patterns affect migration flyways and flock size of a soaring bird over sea. Avian Biol Res 9:159–166. https://doi.org/10.3184/175815516X14627928448105

Akaike H (1973) Information theory as an extension of the maximum likelihood principle. In: Petrov BN, Csaki F (eds) Second International Symposium on Information Theory. Akademiai Kiado, Budapest, pp 267–281

Bartoń K (2018) Package MuMIn: Multi-Model Inference. www.cran.r-project.org, Wien

Bildstein KL (2006) Migrating raptors of the World: their ecology and conservation. Cornell University Press, Ithaca, NY

Burnham KP, Anderson DR (2002) Model selection and multimodel inference: a practical information-theoretic approach, 2nd ed. Springer, New York

Burnham KP, Anderson DR, Huyvaert KP (2011) AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. Behav Ecol Sociobiol 65:23–35. https://doi.org/10.1007/s00265-010-1029-6

456 De la Cruz A, Arroyo GM, Onrubia A, et al (2011a) Migración primaveral de aves
457 planeadoras en el Estrecho de Gibraltar. Diseño de un programa de seguimiento a
458 largo plazo. Migres 2:79–88
459 De la Cruz A, Onrubia A, Pérez B, et al (2011b) Seguimiento de la migración de las
460 aves en el estrecho de Gibraltar: resultados del Programa Migres 2009. Migres
461 2:65–78
462 Dodge S, Bohrer G, Weinzierl R, et al (2013) The environmental-data automated track
463 annotation (Env-DATA) system: linking animal tracks with environmental data.
464 Movement Ecol 1:3. https://doi.org/10.1186/2051-3933-1-3
465 Duerr AE, Miller TA, Lanzone M, et al (2015) Flight response of slope-soaring birds to
466 seasonal variation in thermal generation. Funct Ecol 29:779–790.
467 https://doi.org/10.1111/1365-2435.12381
468 Duriez O, Monti F, Fluhr J, Sforzi A, Sammuri G, Gremillet D (2018) Migrating
469 ospreys use thermal uplift over the open sea. Biol Lett 14, 20180687.
470 https://doi.org/10.1098/rsbl.2018.0687
471 Ferguson-Lees J, Christie DA (2001) *Raptors of the World*. Houghton Mifflin, Boston.
472 Finlayson C (2010) Birds of the Strait of Gibraltar. T & AD Poyser, London
473 Fox J, Monette G (1992) Generalized collinearity diagnostics. J Am Stat Assoc 87:178–
474 183. https://doi.org/10.2307/2290467
475 Fox J, Weisberg S (2011) Package car: an R Companion to Applied Regression.
476 www.cran.r-project.org, Wien
477 Hersbach H, Bell B, Berrisford P, et al (2023) ERA5 hourly data on pressure levels from
478 1940 to present. Copernicus Climate Change Service (C3S) Climate Data Store
479 (CDS). Accessed 27 Jan 2024

480 Hidalgo P, Gallego D (2019) A historical climatology of the easterly winds in the strait
481 of Gibraltar. Atmósfera 32:181–195. https://doi.org/10.20937/ATM.2019.32.03.02
482 Kerlinger P (1989) Flight strategies of migrating hawks. University of Chicago Press,
483 Chicago, US
484 Mellone U (2020) Sea crossing as a major determinant for the evolution of migratory
485 strategies in soaring birds. J Anim Ecol 89:1298–1301.
486 https://doi.org/10.1111/1365-2656.13241
487 Mellone U (2021) Booted eagle (*Hieraaetus pennatus*). In: Panuccio M, Agostini N,
488 Mellone U (eds) Migration strategies of birds of prey in Western Palearctic. CRC
489 Press, Boca Raton, FL (US), pp 101–107
490 Newton I (2023) The migration ecology of birds. Elsevier & Academic Press,
491 Amsterdam (NL)
492 Nourani E, Yamaguchi NM (2017) The effects of atmospheric currents on the migratory
493 behavior of soaring birds: A review. Ornithol Sci 16:5–15.
494 https://doi.org/10.2326/osj.16.5
495 Nourani E, Bohrer G, Becciu P, Bierregaard RO, Duriez O, Figuerola J, et al. (2021)
496 The interplay of wind and uplift facilitates over-water flight in facultative soaring
497 birds. Proc R Soc B 288, 20211603. https://doi.org/10.1098/rspb.2021.1603
498 Oppel S, Dobrev V, Arkumarev V, et al (2015) High juvenile mortality during migration
499 in a declining population of a long-distance migratory raptor. Ibis 157:545–557.
500 https://doi.org/10.1111/ibi.12258
501 Panuccio M, Gustin M, Lucia G, et al (2019) Flight altitude of migrating European
502 Honey Buzzards facing the open sea. Ornithol Sci 18:49–57.
503 https://doi.org/10.2326/osj.18.49

504 Panuccio M, Mellone U, Agostini N (2021) Migration strategies of birds of prey in
505 Western Palearctic. CRC Press, Boca Raton, FL
506 Quinn GP, Keough MJ (2002) Experimental design and data analysis for biologists.
507 Cambridge University Press, Cambridge
508 Santos CD, Silva JP, Muñoz A, et al (2020) The gateway to Africa: What determines sea
509 crossing performance of a migratory soaring bird at the Strait of Gibraltar? J Anim
510 Ecol 89:1317–1328. https://doi.org/10.1111/1365-2656.13201
511 Škrábal J, Krejčí Š, Raab R, Sebastián-González E, Literák I (2023) Soaring over open
512 waters: horizontal winds provide lift to soaring migrants in weak thermal
513 conditions. Mov Ecol 11, 76. https://doi.org/10.1186/s40462-023-00438-6
514 Wallace JM, Hobbs PV (2006) Atmospheric science: an introductory survey, 2nd ed.
515 Elsevier Academic Press, Amsterdam; Boston
516 Wikelski M, Kays R (2018) Movebank: archive, analysis and sharing of animal
517 movement data. Hosted by the Max Planck Institute for Ornithology.
518 www.movebank.org
519 Zu-Aretz S, Leshem Y (1983) The sea, a trap for gliding birds. Torgos 5:16–17
520 Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid common
521 statistical problems: Data exploration. Methods Ecol Evol 1:3–14.
522 https://doi.org/10.1111/j.2041-210X.2009.00001.x

523 **TABLES**

524 **Table 1.** The average model obtained by the linear regression investigating the
525 migration flow of both the Short-toed Snake Eagle and the Booted Eagle during the
526 spring migration at the Strait of Gibraltar (SE = standard error; LCI = 95% lower
527 confidence interval; UCI = 95% upper confidence interval; *sw* = relative importance).
528 When the watchsite is indicated, then the covariate is the interaction term with the
529 variable. For the watchsite the reference level is Cazalla.

Variable	**Watchsite**	**Estimate**	**SE**	**LCI**	**UCI**	***sw***
Short-toed snake-eagle						
Intercept		3.627	1.298	-	-	-
Watchsite	*P. Carnero*	0.644	0.547	-0.428	1.715	1.00
Temperature		-0.112	0.070	-0.249	0.025	0.63
Eastward wind	*P. Carnero*	0.225	0.072	0.083	0.366	1.00
	Cazalla	-0.152	0.049	-0.248	-0.057	
Northward wind		0.0001	0.141	-0.276	0.276	0.15
Booted eagle						
Intercept		0.479	0.991	-	-	-
Watchsite	*P. Carnero*	-0.089	0.408	-0.897	0.719	1.00
Temperature		0.141	0.059	0.024	0.258	1.00
Eastward wind	*P. Carnero*	-0.205	0.038	0.058	0.271	1.00
	Cazalla	0.165	0.054	-0.281	-0.129	
Northward wind	*P. Carnero*	-9.571	2.803	-15.127	-4.015	1.00
	Cazalla	1.827	2.474	-3.077	6.732	
Northward wind^2	*P. Carnero*	-6.814	2.606	-11.979	-1.648	
	Cazalla	-1.986	2.617	-7.173	3.200	

530

Table 2. The best model obtained by the linear regression investigating the effect of the weather on the flight altitude of the Short-toed Snake Eagles during the spring migration at the Strait of Gibraltar (SE = standard error; LCI = 95% lower confidence interval; UCI = 95% upper confidence interval). When the watchsite is indicated, then the covariate is the interaction term with the variable.

Variable	Watchsite	Estimate	SE	LCI	UCI
Intercept		2.094	0.020	-	-
Cloud cover	*P. Carnero*	0.056	0.012	0.033	0.079
	Trafico	0.022	0.022	-0.020	0.065
Temperature		-2.036	0.417	-2.854	-1.219
Temperature^2		0.330	0.341	-0.338	0.999
Thermal uplift		-0.051	0.012	-0.074	-0.028
Eastward wind	*P. Carnero*	-6.395	1.532	-9.402	-3.389
	Trafico	2.687	0.761	1.193	4.180
Eastward wind ^{2}	*P. Carnero*	0.994	1.110	-1.184	3.173
	Trafico	-0.074	0.587	-1.225	1.077
Northward wind	*P. Carnero*	-5.037	0.372	-5.768	-4.306
	Trafico	-7.169	1.261	-9.645	-4.694
Northward wind^2	*P. Carnero*	3.109	0.409	2.306	3.911
	Trafico	-2.939	0.898	-4.701	-1.177

537 **Table 3.** The average model obtained by the linear regression investigating the effect of
538 the weather on the flight altitude of the Booted Eagles during the spring migration at the
539 Strait of Gibraltar (SE = standard error; LCI = 95% lower confidence interval; UCI =
540 95% upper confidence interval; *sw* = relative importance). When the watchsite is
541 indicated, then the covariate is the interaction term with the variable.

Variable	**Watchsite**	**Estimate**	**SE**	**LCI**	**UCI**	***sw***
Intercept		1.878	0.018	-	-	-
Temperature		-0.038	0.044	-0.125	0.049	0.26
Thermal uplift		-0.973	0.368	-1.699	-0.248	1.00
Thermal uplift^2		0.564	0.306	-0.039	1.167	
Eastward wind		0.015	0.020	-0.025	0.055	0.23
Northward wind	*P. Carnero*	0.221	0.599	-0.960	1.401	1.00
	Trafico	-1.625	0.467	-2.546	-0.705	
Northward wind^2	*P. Carnero*	2.154	0.590	0.992	3.316	
	Trafico	-0.117	0.518	-1.137	0.904	

542

FIGURES

Figure 1 The watchsites used at the Strait of Gibraltar (Spain) to investigate daily counts (Cazalla and Punta Carnero) and the flight altitude (Trafico and Punta Carnero) of Short-toed Snake Eagles and Booted Eagles during the pre-breeding migration



549 **Figure 2** Response curves of the variables with higher relative importance (*sw* = 1.00)

550 selected by the linear regression investigating the migration flow of both the Short-toed

551 Snake Eagle (a) and the Booted Eagles (b-d) during the spring migration at the Strait of

552 Gibraltar. For the temperature (d) the effect includes both watchsites



553

Figure 3 Average wind vectors obtained by daily averaging ERA5 wind between 09:00 UTC and 17:00 UTC and weighting by the number of counts on observation days, for each species and observation site. The background colour reports the strength of the horizontal wind component (positive is westerly)



560 **Figure 4** Response curves obtained by the linear regression investigating the effect of
561 the weather on the flight altitude of the Short-toed Snake Eagles during the spring
562 migration at the Strait of Gibraltar



563

564 **Figure 5** Response curves of the variables with higher relative importance (*sw* = 1.00)
565 selected by the linear regression investigating the effect of the weather on the flight
566 altitude of the Booted Eagles during the spring migration at the Strait of Gibraltar



567