

Gevrey local solvability in locally integrable structures

Original

Gevrey local solvability in locally integrable structures / Malaspina, Francesco; Nicola, Fabio. - In: ANNALI DI MATEMATICA PURA ED APPLICATA. - ISSN 0373-3114. - STAMPA. - 193:5(2014), pp. 1491-1502. [10.1007/s10231-013-0340-z]

Availability:

This version is available at: 11583/2509875 since: 2015-12-28T11:52:14Z

Publisher:

SPRINGER-VERLAG

Published

DOI:10.1007/s10231-013-0340-z

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

GEVREY LOCAL SOLVABILITY IN LOCALLY INTEGRABLE STRUCTURES

FRANCESCO MALASPINA AND FABIO NICOLA

ABSTRACT. We consider a locally integrable real-analytic structure, and we investigate the local solvability in the category of Gevrey functions and ultradistributions of the complex d' naturally induced by the de Rham complex. We prove that the so-called condition $Y(q)$ on the signature of the Levi form, for local solvability of $d'u = f$, is still necessary even if we take f in the classes of Gevrey functions and look for solutions u in the corresponding spaces of ultradistributions.

1. INTRODUCTION AND STATEMENT OF THE RESULTS

Consider a real-analytic manifold M of dimension $m + n$. A real-analytic locally integrable structure on M , of rank n , is defined by a real analytic subbundle $\mathcal{V} \subset \mathbb{C}TM$ of rank n , satisfying the Frobenius condition and such that the subbundle $T' \subset \mathbb{C}T^*M$ orthogonal to \mathcal{V} is locally spanned by exact differentials. As usual we will denote by $T^0 = T' \cap T^*M$ the so-called *characteristic set*. For any open subset $\Omega \subset M$ and $s > 1$ the space $G^s(\Omega, \Lambda^{p,q})$ of (p, q) -forms with Gevrey coefficients of order s is then defined (see Section 3 below and Treves [28]) and the de Rham differential induces a map

$$d' : G^s(\Omega, \Lambda^{p,q}) \rightarrow G^s(\Omega, \Lambda^{p,q+1}).$$

Similarly, the de Rham differential induces a complex on the space of “ultra-currents” $\mathcal{D}'_s(\Omega, \Lambda^{p,q})$, i.e. forms with ultradistribution coefficients:

$$d' : \mathcal{D}'_s(\Omega, \Lambda^{p,q}) \rightarrow \mathcal{D}'_s(\Omega, \Lambda^{p,q+1}).$$

When $\mathcal{V} \cap \bar{\mathcal{V}} = 0$ the structure is called *CR* and d' is the so-called tangential Cauchy-Riemann operator.

We are interested in necessary conditions for the Gevrey local solvability problem for the complex d' to hold near a given point x_0 .

2000 *Mathematics Subject Classification.* 35S30, 47G30, 42C15.

Key words and phrases. Gevrey local solvability, locally integrable structures, Poincaré lemma, differential complexes, involutive structures.

Definition 1.1. *We say that the complex d' is locally solvable near x_0 and in degree q , $1 \leq q \leq n$, in the sense of ultradistribution of order s , if for every sufficiently small open neighborhood Ω of x_0 and every cocycle $f \in G^s(\Omega, \Lambda^{0,q})$ there exists an open neighbourhood $V \subset \Omega$ of x_0 and a ultradistribution section $u \in \mathcal{D}'_s(V, \Lambda^{0,q-1})$ solving $d'u = f$ in V .*

The analogous problem in the setting of smooth functions and Schwartz distributions has been extensively considered, see e.g. [1, 2, 3, 8, 9, 10, 11, 12, 20, 22, 23, 24, 26, 28], inspired by the results in [16, 17] for scalar operators of principal type; see also [18, 19] as general references for the problem of local solvability of scalar linear partial differential operators.

Several geometric invariants were there introduced, e.g. the signature of the Levi form recalled below, which represent obstructions to the solvability in the sense of distributions, that is, for some smooth $f \in C^\infty(U, \Lambda^{0,q})$ there is no distribution solution $u \in \mathcal{D}'(V, \Lambda^{0,q-1})$ to $d'u = f$ in V , for every neighbourhood $V \subset U$ of x_0 .

It is therefore natural to wonder whether, under the same condition as in the smooth category, d' is still non-solvable even if we choose f in the smaller class of Gevrey functions $G^s(U, \Lambda^{0,q}) \subset C^\infty(U, \Lambda^{0,q})$ and we look for solutions in the larger class of ultradistributions $\mathcal{D}'_s(V, \Lambda^{0,q}) \supset \mathcal{D}'(V, \Lambda^{0,q})$, as in Definition 1.1. In this note we present a result in this direction.

Let us note that general sufficient conditions for local solvability in the Gevrey category have been recently obtained in [5]; see also [4, 21].

We recall that at any point $(x_0, \omega_0) \in T^0$ it is well defined a sesquilinear form $\mathcal{B}_{(x_0, \omega_0)} : \mathcal{V}_{x_0} \times \mathcal{V}_{x_0} \rightarrow \mathbb{C}$, (\mathcal{V}_{x_0} is the fibre above x_0) by

$$\mathcal{B}_{(x_0, \omega_0)}(\mathbf{v}_1, \mathbf{v}_2) = \langle \omega_0, (2\iota)^{-1}[V_1, \overline{V_2}]|_{x_0} \rangle,$$

with $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{V}_{x_0}$, where V_1 and V_2 are smooth sections of \mathcal{V} such that $V_1|_{x_0} = \mathbf{v}_1$, $V_2|_{x_0} = \mathbf{v}_2$. The associated quadratic form $\mathcal{V}_{x_0} \ni \mathbf{v} \mapsto \mathcal{B}_{(x_0, \omega_0)}(\mathbf{v}, \mathbf{v})$, or $\mathcal{B}_{(x_0, \omega_0)}$ itself, is known as *Levi form*.

Here is our result.

Theorem 1.2. *Let $(x_0, \omega_0) \in T^0$, $\omega_0 \neq 0$. Suppose that $\mathcal{B}_{(x_0, \omega_0)}$ has exactly q positive eigenvalues, $1 \leq q \leq n$, and $n - q$ negative eigenvalues, and that its restriction to $\mathcal{V}_{x_0} \cap \overline{\mathcal{V}}_{x_0}$ is non-degenerate.*

Then, for every $s > 1$, d' is not locally solvable in the sense of ultradistributions of order s , near x_0 and in degree q .

This result therefore strengthens the analogous one in the category of smooth functions and Schwartz distributions, which was proved in [1] for *CR* manifolds and in [28, Theorem XVIII.3.1] for general locally integrable

structure; see also [15, 25] for partial results when the Levi form is degenerate. As general reference for related results about scalar operators on Gevrey spaces see [27].

2. PRELIMINARIES

2.1. Gevrey functions and ultradistributions. Let us briefly recall the definition of the classes of Gevrey functions and corresponding ultradistributions; see e.g. [27, Chapter 1] for details.

Let $s > 1$ be a real number and Ω be an open subset of \mathbb{R}^n ; let C be a positive constant. We denote by $G^s(\Omega, C)$ the space of smooth functions f in \mathbb{R}^n such that for every compact $K \subset \Omega$,

$$\|f\|_{K,C} := \sup_{\alpha} C^{-|\alpha|} (\alpha!)^{-s} \sup_{x \in K} |\partial^{\alpha} f(x)| < \infty.$$

This is a Fréchet space endowed with the above seminorms. We set $G^s(\Omega)$ for the usual Gevrey space of order s , i.e. $f \in G^s(\Omega)$ if f is smooth in Ω and for every compact $K \subset \Omega$ there exists $C > 0$ such that $\|f\|_{K,C} < \infty$. We will also consider the space $G_0^s(K, C)$ of functions in $G^s(\Omega, C)$ supported in the compact K ; it is a Banach space with the norm $\|u\|_{K,C}$. Finally we set

$$G_0^s(\Omega) = \bigcup_{K \subset \Omega, C > 0} G_0^s(K, C).$$

The space of $\mathcal{D}'_s(\Omega)$ of ultradistributions of order s in Ω is by definition the dual of $G_0^s(\Omega)$, i.e. an element $u \in \mathcal{D}'_s(\Omega)$ is a linear functional on $G_0^s(\Omega)$ such that for every compact $K \subset \Omega$ and every constant $C > 0$ there exists a constant $C' > 0$ such that

$$|\langle u, f \rangle| \leq C' \|f\|_{K,C},$$

namely $u \in (G_0^s(K, C))'$ for every K, C . Clearly, $\mathcal{D}'_s(\Omega)$ contains the usual space $\mathcal{D}'(\Omega)$ of Schwartz distributions.

We will need the following estimate for Gevrey seminorms of exponential functions.

Proposition 2.1. *Let ψ be a real-analytic function in a neighborhood Ω of 0 in \mathbb{R}^n ; then for every compact subset K of Ω and every $C > 0, s > s' > 1$, there exists a constant $C' > 0$ such that*

$$(2.1) \quad \|\exp(\iota\rho\psi)\|_{K,C} \leq C' \exp(a\rho + \rho^{1/s'})$$

for every $\rho > 0$, where $a = \sup\{-\text{Im}\psi(x) : x \in K\}$.

Proof. By the Faà di Bruno formula (see e.g. [14, page 16]) we have, for $|\alpha| \geq 1$,

$$\partial^\alpha e^{\iota\rho\psi(x)} = \sum_{j=1}^{|\alpha|} \frac{\exp(\iota\rho\psi)}{j!} \sum_{\substack{\gamma_1+\dots+\gamma_j=\alpha \\ |\gamma_k|\geq 1}} \frac{\alpha!}{\gamma_1!\dots\gamma_j!} |\partial^{\gamma_1}(\iota\rho\psi(x))| \dots |\partial^{\gamma_j}(\iota\rho\psi(x))|.$$

By assumption there exists a constant $C_1 > 0$ such that $|\partial^\gamma\psi(x)| \leq C_1^{|\gamma|}\gamma!$ for $x \in K$, $|\gamma| \geq 1$. Hence for every α ,

$$\sup_{x \in K} |\partial^\alpha e^{\iota\rho\psi(x)}| \leq e^{a\rho} \alpha! C_2^{|\alpha|} \sum_{j=0}^{|\alpha|} \frac{\rho^j}{j!}$$

with $C_2 = 2^{n+1}C_1$, where we used

$$\sum_{\substack{\gamma_1+\dots+\gamma_j=\alpha \\ |\gamma_k|\geq 1}} 1 \leq \prod_{k=1}^n \binom{\alpha_k + j - 1}{j - 1} \leq 2^{|\alpha|+n(j-1)} \leq 2^{(n+1)|\alpha|}.$$

Hence we have

$$\begin{aligned} \|\exp(\iota\rho\psi)\|_{K,C} &\leq e^{a\rho} (\alpha!)^{1-s} (C_2/C)^{|\alpha|} \sum_{j=0}^{|\alpha|} \frac{\rho^j}{j!} \\ &\leq e^{a\rho} (|\alpha|!)^{1-s} (C_2 C_3/C)^{|\alpha|} \sum_{j=0}^{|\alpha|} \frac{\rho^j}{j!}, \end{aligned}$$

because $|\alpha|! \leq n^{|\alpha|}\alpha!$. Now, we have $(|\alpha|!)^{1-s'} (j!)^{s'-1} \leq 1$ and by Stirling formula $(|\alpha|!)^{s'-s} (C_2 C_3/C)^{|\alpha|} \leq C'$, so that

$$\|\exp(\iota\rho\psi)\|_{K,C} \leq C' e^{a\rho} \sum_{j=0}^{|\alpha|} \frac{\rho^j}{(j!)^{s'}} \leq C' e^{a\rho} \sum_{j=0}^{|\alpha|} \left(\frac{\rho^{j/s'}}{j!}\right)^{s'} \leq C' e^{a\rho+s'\rho^{1/s'}}.$$

Since this holds for every $1 < s' < s$, we can replace the constant s' in front of $\rho^{1/s'}$ by 1, possibly for a new constant C' and for a slightly lower value of s' . Hence (2.1) is proved. \square

2.2. Locally integrable structures. Consider a real-analytic manifold M of dimension $N = m + n$. A real-analytic locally integrable structure on M , of rank n , is defined by a real analytic subbundle $\mathcal{V} \subset \mathbb{C}TM$ of rank n , satisfying the Frobenius condition and such that subbundle $T' \subset \mathbb{C}T^*M$ orthogonal to \mathcal{V} is locally spanned by exact differentials. As usual we will denote by $T^0 = T' \cap T^*M$ the so-called *characteristic set*. Let k be a positive

integer, we denote by $\Lambda^k \mathbb{C}T^*M$ the k -th exterior power of $\mathbb{C}T^*M$. Let us consider complex exterior algebra

$$\Lambda \mathbb{C}T^*M = \bigoplus_{k=0}^N \Lambda^k \mathbb{C}T^*M,$$

for any pair of positive integers p, q we denote by

$$T'^{p,q}$$

the homogeneous of degree $p+q$ in the ideal generated by the p -th exterior power of T' , $\Lambda^p T'$. We have the inclusion

$$T'^{p+1,q-1} \subset T'^{p,q}$$

which allows us to define

$$\Lambda^{p,q} = T'^{p,q} / T'^{p+1,q-1}.$$

If ϕ is a smooth section of T' over an open subset $\Omega \subset M$, its exterior derivative $d\phi$ is section of $T'^{1,1}$. In other words

$$dT' \subset T'^{1,1}.$$

It follows at once from this that, if σ is a smooth section of $T'^{p,q}$ over Ω , then $d\sigma$ is a section of $T'^{p,q+1}$ i.e.

$$dT'^{p,q} \subset T'^{p,q+1}.$$

Let $s > 1$, the space $G^s(\Omega, \Lambda^{p,q})$ of (p, q) -forms with Gevrey coefficients of order s is defined, as well as $G^s(\Omega, \mathbb{C}; \Lambda^{p,q})$, $G_0^s(K, \mathbb{C}; \Lambda^{p,q})$, etc, with notation analogous to the scalar case.

The de Rham differential induces then a map

$$d' : G^s(\Omega, \Lambda^{p,q}) \rightarrow G^s(\Omega, \Lambda^{p,q+1}).$$

(see Treves [28, Section I.6] for more details). Similarly, the de Rham differential induces a complex on the space of “ultra-currents” $\mathcal{D}'_s(\Omega, \Lambda^{p,q})$, i.e. forms with ultradistribution coefficients:

$$d' : \mathcal{D}'_s(\Omega, \Lambda^{p,q}) \rightarrow \mathcal{D}'_s(\Omega, \Lambda^{p,q+1}).$$

Namely, consider for simplicity the case when Ω is orientable (in fact, in the sequel we will work in a local chart). Stokes' theorem implies that

$$\int_{\Omega} d'u \wedge v = (-1)^{p+q-1} \int_{\Omega} u \wedge d'v$$

if $u \in G^s(\Omega, \Lambda^{p,q})$, $v \in G_0^s(\Omega, \Lambda^{m-p,n-q-1})$, and accordingly we can define

$$\langle d'u, v \rangle = (-1)^{p+q-1} \langle u, d'v \rangle$$

if $u \in \mathcal{D}'_s(\Omega, \Lambda^{p,q})$, $v \in G_0^s(\Omega, \Lambda^{m-p,n-q-1})$.

3. LOCAL SOLVABILITY ESTIMATES

We now show that local solvability implies an a priori-estimate. This is analogous to the estimates of Hörmander [16], Andreotti, Hill and Nacinovich [1], Treves [28, Lemma VIII.1.1], in the framework of Schwartz distributions.

Proposition 3.1. *Suppose that, for some $s > 1$, the complex d' is locally solvable near x_0 and in degree q , in the sense of ultradistributions of order s (see Definition 1.1). Then for every sufficiently small open neighborhood Ω of x_0 , every $C_1 > 0$, $0 < \epsilon < C_2$ there exist a compact $K \subset \Omega$, an open neighbourhood $\Omega' \subset\subset \Omega$ of x_0 and a constant $C' > 0$, such that*

$$(3.1) \quad \left| \int_{\Omega} f \wedge v \right| \leq C' \|f\|_{K, C_1} \|d'v\|_{\overline{\Omega'}, C_2},$$

for every cocycle $f \in G^s(\Omega, C_1; \Lambda^{0,q})$ and every $v \in G_0^s(\overline{\Omega'}, C_2 - \epsilon; \Lambda^{m, n-q})$.

It will follow from the proof that $\|d'v\|_{\overline{\Omega'}, C_2} < \infty$ if $v \in G_0^s(\overline{\Omega'}, C_2 - \epsilon; \Lambda^{m, n-q})$.

Proof. Let $V_{j+1} \subset V_j \subset\subset \Omega$, $j = 1, 2, \dots$, be a fundamental system of neighborhoods of x_0 . Fix $C_1 > 0$, $0 < \epsilon < C_2$ and consider the space

$$F_j = \{(f, u) \in G^s(\Omega, C_1; \Lambda^{0,q}) \times G_0^s(\overline{V_j}, C_2; \Lambda^{0, q-1})' : \\ d'f = 0 \text{ in } \Omega, \quad d'u = f \text{ in } G_0^s(\overline{V_j}, C_2 - \epsilon; \Lambda^{0,q})'\}.$$

The last condition means $\langle d'u, v \rangle = \langle f, v \rangle$ for every $v \in G_0^s(\overline{V_j}, C_2 - \epsilon; \Lambda^{m, n-q})$, which makes sense by transposition, because differentiation maps $G_0^s(\overline{V_j}, C_2 - \epsilon) \rightarrow G_0^s(\overline{V_j}, C_2)$ (see e.g. [27, Proposition 2.4.8]) and multiplication by analytic functions preserves the latter space.

Now, by direct inspection one sees that F_j is a closed subspace of $G^s(\Omega, C_1; \Lambda^{0, q-1}) \times G_0^s(\overline{V_j}, C_2; \Lambda^{0,q})'$, therefore Fréchet.

Let

$$\pi_j : F_j \rightarrow \{f \in G^s(\Omega, C_1; \Lambda^{0, q-1}) : d'f = 0\}$$

be the canonical projection $(f, u) \mapsto f$. The assumption of local solvability implies that

$$\{f \in G^s(\Omega, C_1; \Lambda^{0, q-1}) : d'f = 0\} = \cup_j \pi_j(F_j).$$

By the Baire theorem, there exists j_0 such that $\pi_{j_0}(F_{j_0})$ is of second category. By the open mapping theorem, we see that π_{j_0} is onto and open: there exists a compact $K \subset \Omega$ and a constant $C' > 0$ such that for every cocycle

$f \in G^s(\Omega, C_1; \Lambda^{0,q})$, there exists $u \in G_0^s(\overline{V}_{j_0}, C_2; \Lambda^{0,q-1})'$ satisfying $d'u = f$ in $G_0^s(\overline{V}_{j_0}, C_2 - \epsilon; \Lambda^{0,q})'$ and

$$|u|_{\overline{V}_{j_0}, C_2} := \sup_{\|v\|_{\overline{V}_{j_0}, C_2} = 1} |\langle u, v \rangle| \leq C' \|f\|_{K, C_1}.$$

Consider now the bilinear functional $(f, v) \mapsto \int_{\Omega} f \wedge v = \langle f, v \rangle$, for $f \in G^s(\Omega, C_1; \Lambda^{0,q})$ cocycle, and $v \in G_0^s(\overline{V}_{j_0}, C_2 - \epsilon; \Lambda^{m, n-q})$. Given such a f , we take u as before, and we get

$$|\langle f, v \rangle| = |\langle d'u, v \rangle| = |\langle u, d'v \rangle| \leq |u|_{\overline{V}_{j_0}, C_2} \|d'v\|_{\overline{V}_{j_0}, C_2} \leq C' \|f\|_{K, C_1} \|d'v\|_{\overline{V}_{j_0}, C_2}.$$

□

4. PROOF OF THEOREM 1.2

We work in a sufficiently small neighborhood Ω of the point x_0 (to be chosen later), where local solvability holds. We also take x_0 as the origin of the coordinates, i.e. $x_0 = 0$. Moreover we make use of the special coordinates, whose existence is proved in see section I.9 of [28]. Namely, let $n = \dim_{\mathbb{C}} \mathcal{V}_0$, $d = \dim_{\mathbb{R}} T_0^0$, $\nu = n - \dim_{\mathbb{C}}(\mathcal{V}_0 \cap \overline{V}_0)$. We have the following result.

Proposition 4.1. *Let $(0, \omega_0) \in T^0$, $\omega_0 \neq 0$, and suppose that the restriction of the Levi form $\mathcal{B}_{(0, \omega_0)}$ to $\mathcal{V}_0 \cap \overline{V}_0$ is non-degenerate. There exist real-analytic coordinates x_j, y_j, s_k and t_l , $j = 1, \dots, \nu$, $k = 1, \dots, d$, $l = 1, \dots, n - \nu$, and smooth real valued and real-analytic functions $\phi_k(x, y, s, t)$, $k = 1, \dots, d$, in a neighborhood \mathcal{O} of 0, satisfying*

$$(4.1) \quad \phi_k|_0 = 0 \quad \text{and} \quad d\phi_k|_0 = 0,$$

such that

$$\begin{cases} z_j := x_j + iy_j, & j = 1, \dots, \nu, \\ w_k := s_k + i\phi_k(x, y, s, t), & k = 1, \dots, d, \end{cases}$$

define a system of first integrals for \mathcal{V} , i.e. their differential span $T'|_{\mathcal{O}}$.

Moreover, with respect to the basis

$$\left\{ \frac{\partial}{\partial \bar{z}_j} \Big|_0, \frac{\partial}{\partial t_l} \Big|_0; j = 1, \dots, \nu, l = 1, \dots, n - \nu \right\}$$

of \mathcal{V}_0 the Levi form $\mathcal{B}_{(0, \omega_0)}$ reads

$$(4.2) \quad \sum_{j=1}^{p''} |\zeta_j|^2 - \sum_{j=p''+1}^{\nu} |\zeta_j|^2 + \sum_{l=1}^{p'} |\tau_l|^2 - \sum_{l=p'+1}^{n-\nu} |\tau_l|^2.$$

Remark 4.2. In particular

$$(4.3) \quad d'z_j = 0, \quad d'w_k = 0, \quad j = 1, \dots, \nu; \quad k = 1, \dots, d.$$

In these coordinates we have $T_0^0 = \text{span}_{\mathbb{R}}\{ds_k|_0; k = 1, \dots, d\}$, so that $\omega_0 = \sum_{k=1}^d \sigma_k ds_k|_0$, with $\sigma_k \in \mathbb{R}$. By (I.9.2) of [28] we have $\mathcal{B}_{(0, \omega_0)}(\mathbf{v}_1, \mathbf{v}_2) = \sum_{k=1}^d \sigma_k (V_1 \bar{V}_2 \phi_k)|_0$, with V_1 and V_2 smooth sections of \mathcal{V} extending \mathbf{v}_1 and \mathbf{v}_2 respectively. Upon setting $\Phi = \sum_{k=1}^d \sigma_k \phi_k$ we can suppose, in addition, that

$$(4.4) \quad \Phi = \sum_{j=1}^{p''} |z_j|^2 - \sum_{j=p''+1}^{\nu} |z_j|^2 + \frac{1}{2} \sum_{l=1}^{p'} t_l^2 - \frac{1}{2} \sum_{l=p'+1}^{n-\nu} t_l^2 + O(|s|(|z|+|s|+|t|) + |z|^3 + |t|^3);$$

see [28, Section I.9] and [28, (XVIII.3.2)] for details.

We can now prove Theorem 1.2. We may assume, without loss of generality, that $\sigma = (1, 0, \dots, 0)$. Consequently, from (4.4) (after the change of variables $t \mapsto t/\sqrt{2}$) we have

$$(4.5) \quad \phi_1(x, y, s, t) = |z'|^2 - |z''|^2 + |t'|^2 - |t''|^2 + O(|s|(|z|+|s|+|t|) + |z|^3 + |t|^3),$$

where we set

$$\begin{cases} z' = (z_1, \dots, z_{p''}), \\ z'' = (z_{p''+1}, \dots, z_{\nu}) \\ t' = (t_1, \dots, t_{p'}), \\ t'' = (t_{p'+1}, \dots, t_{n-\nu}). \end{cases}$$

Moreover, we choose a function $\chi(x, y, s, t)$ in $G_0^s(\mathbb{R}^{2\nu+d+(n-\nu)})$, $\chi = 0$ away from a neighborhood $V \subset\subset \Omega$ of 0 and $\chi = 1$ in a neighborhood $U \subset\subset V$ of 0, where V and U will be chosen later on. We set, for $\rho > 0, \lambda > 0$,

$$\begin{aligned} f_{\rho, \lambda} &= e^{\rho h_{1, \lambda}} d\bar{z}' \wedge dt' \\ v_{\rho, \lambda} &= \rho^{(m+n)/2} \chi e^{\rho h_{2, \lambda}} d\bar{z}'' \wedge dt'' \wedge dz \wedge dw, \end{aligned}$$

where, with $\lambda > 1$,

$$(4.6) \quad h_{1, \lambda} := -\iota s_1 + \phi_1 - 2|z'|^2 - 2|t'|^2 - \lambda \sum_{k=1}^d (s_k + \iota \phi_k)^2,$$

and

$$(4.7) \quad h_{2, \lambda} := \iota s_1 - \phi_1 - 2|z''|^2 - 2|t''|^2 - \lambda \sum_{k=1}^d (s_k + \iota \phi_k)^2.$$

Now, we have $\chi \in G_0^s(\bar{V}, C/2)$, for some $C > 0$. We then apply Proposition 3.1 with $f_{\rho,\lambda}$ and $v_{\rho,\lambda}$ in place of f and v respectively, and $C_1 = C_2 = C$, $\epsilon = C/2$, taking V small enough to be contained in the neighborhood Ω' which arises in the conclusion of Proposition 3.1. Observe that, in fact, $f_{\rho,\lambda} \in G^s(\Omega, C; \Lambda^{0,q})$ since this form is in fact real-analytic and $p'' + p' = q$ by hypothesis, whereas $v_{\rho,\lambda} \in G_0^s(\bar{V}, C/2; \Lambda^{m,n-q}) \subset G_0^s(\bar{\Omega}', C/2; \Lambda^{m,n-q})$ (recall, $m = \dim M - n$). We prove now that f_ρ is a cocycle (i.e. $d'f_{\rho,\lambda} = 0$), so that Proposition 3.1 can in fact be applied. However, we will show that (3.1) fails for every choice of C' when $\rho \rightarrow +\infty$, if λ is large enough, obtaining a contradiction.

Now, by (4.3)

$$d'f_{\rho,\lambda} = 0,$$

and

$$(4.8) \quad d'v_{\rho,\lambda} = \rho^{(m+n)/2} e^{\rho h_{2,\lambda}} d'\chi \wedge d\bar{z}'' \wedge dt'' \wedge dz \wedge dw.$$

In order to estimate the right hand side of (3.1) we observe that, by (4.5) and (4.6)

$$\operatorname{Re} h_{1,\lambda} = -|z'|^2 - |z''|^2 - |t|^2 - \lambda|s|^2 + \mathcal{R}(z, s, t) + O(|z|^3 + |t|^3) + \lambda O(|z|^4 + |t|^4),$$

where

$$(4.9) \quad |\mathcal{R}(z, s, t)| = O(|s|(|z| + |s| + |t|)) \leq \tilde{C} \left(\frac{\epsilon}{2} (|z| + |s| + |t|)^2 + \frac{1}{2\epsilon} |s|^2 \right),$$

for every $\epsilon > 0$. Hence, if ϵ and then $1/\lambda$ are small enough we see that, possibly after replacing Ω with a smaller neighborhood,

$$(4.10) \quad \sup_{\Omega} \operatorname{Re} h_{1,\lambda} \leq 0.$$

Similarly,

$$\begin{aligned} \operatorname{Re} h_{2,\lambda} &= -|z'|^2 - |z''|^2 - |t|^2 - \lambda|s|^2 \\ &\quad + \mathcal{R}'(z, s, t) + O(|z|^3 + |t|^3) + \lambda O(|z|^4 + |t|^4), \end{aligned}$$

with \mathcal{R}' satisfying the same estimate (4.9). Therefore if λ is sufficiently large, in Ω we have

$$\operatorname{Re} h_{2,\lambda} \leq -\frac{1}{2}(|z|^2 + |t|^2 + \lambda|s|^2) + \tilde{C}_1(|z|^3 + |t|^3) + \tilde{C}_2\lambda(|z|^4 + |t|^4).$$

Hence, possibly for a smaller V , since $U \subset\subset V$ is a neighborhood of 0, there exists a constant $c > 0$ such that

$$(4.11) \quad \sup_{\bar{V} \setminus U} h_{2,\lambda}(z, s, t) \leq -c.$$

As a consequence of Proposition 2.1 and (4.10), (4.11), for every compact subset $K \subset \Omega$ it turns out

$$(4.12) \quad \|f_{\rho,\lambda}\|_{K,C} \leq C' e^{\rho^{1/s'}},$$

$$(4.13) \quad \|d'v_{\rho,\lambda}\|_{K,C} \leq C'' \rho^{(m+n)/2} e^{-c\rho + \rho^{1/s'}},$$

for any $1 < s' < s$, where the constants C', C'' are independent of ρ . It follows that

$$(4.14) \quad \|f_{\rho,\lambda}\|_{K,C} \|d'v_{\rho,\lambda}\|_{K,C} \leq C' C'' \rho^{(m+n)/2} e^{-c\rho + 2\rho^{1/s'}} \longrightarrow 0 \text{ as } \rho \rightarrow +\infty,$$

because $s' > 1$. On the other hand, for the right-hand side of (3.1) it is easily seen that

$$\int f_{\rho,\lambda} \wedge v_{\rho,\lambda} \longrightarrow c' \neq 0,$$

as $\rho \rightarrow +\infty$ (see the end of the proof of [28, Theorem XVIII.3.1]), which together with (4.14) contradicts (3.1).

This completes the proof of Theorem 1.2.

Remark 4.3. The above machinery can be applied to prove other necessary conditions for Gevrey local solvability in the spirit of analogous results valid in the framework of smooth functions and Schwartz distributions.

As an example, consider the special case of local solvability when $q = n$, namely in top degree. In this case, the Cordaro-Hounie condition $(\mathcal{P})_{n-1}$ (see [7] and [9]) is known to be necessary for the local solvability in the smooth category, and it is conjectured to be sufficient as well. Consider the following analytic variant. Let $L_j, j = 1, \dots, n$, be real-analytic independent vector fields which generates \mathcal{V} at any point of Ω .

We say that the real-analytic condition $(\mathcal{P})_{n-1}$ is satisfied at x_0 if there exists an open neighbourhood $U \subset \Omega$ of x_0 such that, given any open set $V \subset U$ and given any real-analytic $h \in C^\infty(V)$ satisfying $L_j h = 0, j = 1, \dots, n$, then $\operatorname{Re} h$ does not assume a local minimum¹ over any nonempty compact subset of V .

One could then prove that *if the real-analytic condition $(\mathcal{P})_{n-1}$ is not satisfied at x_0 , then for every $s > 1$, d' is not locally solvable in the sense of ultradistributions of order s , near x_0 and in degree n .*

For the sake of brevity we omit the proof, which goes on along the same lines as that in [9, Theorem 1.2], using the local solvability estimates in Proposition 3.1, combined with Proposition 2.1.

¹A real-valued function f defined on a topological space X is said to assume a local minimum over a compact set $K \subset X$ if there exist $a \in \mathbb{R}$ and $K \subset V \subset X$ open such that $f = a$ on K and $f > a$ on $V \setminus K$.

REFERENCES

- [1] A. Andreotti, G. Fredricks, M. Nacinovich, *On the absence of the Poincaré lemma in tangential Cauchy-Riemann complexes*, Ann. Scuola Norm. Sup. Pisa, Sci. Fis. Mat., **8** (1981), 365–404.
- [2] A. Andreotti, C.D. Hill, *E.E. Levi convexity and the Hans Lewy problem, I and II*, Ann. Scuola Norm. Sup. Pisa, Sci. Fis. Mat., **26** (1972), 325–363, 747–806.
- [3] S. Berhanu, P.D. Cordaro, J. Hounie, *An introduction to involutive structures*, Cambridge University Press, 2008.
- [4] P.A.S. Caetano, *Classes de Gevrey em estruturas hipo-analíticas*, PhD thesis, University of Sao Paulo, Brazil, 2001.
- [5] P.A.S. Caetano, P.D. Cordaro, *Gevrey solvability and Gevrey regularity in differential complexes associated to locally integrable structures*, Trans. Amer. Math. Soc., **363** (2011), 185–201.
- [6] S. Chanillo, F. Trèves, *Local exactness in a class of differential complexes*, J. Amer. Math. Soc., **10**(2) (1997), 393–426
- [7] P.D. Cordaro, *Microlocal Analysis applied to locally integrable structures*, Lecture Notes for the Congress “Phase space analysis of Partial Differential Equations”, Centro di Ricerca Matematica “Ennio De Giorgi”, Pisa, April 2004.
- [8] P.D. Cordaro, J. Hounie, *On local solvability of undetermined systems of vector fields*, Amer. J. Math., **112** (1990), 243–270.
- [9] P.D. Cordaro, J. Hounie, *Local solvability for top degree forms in a class of systems of vector fields*, Amer. J. Math., **121** (1999), 487–495.
- [10] P.D. Cordaro, J. Hounie, *Local solvability for a class of differential complexes*, Acta Math., **187** (2001), 191–212.
- [11] P.D. Cordaro, F. Trèves, *Homology and cohomology in hypoanalytic structures of the hypersurfaces type*, J. Geometric Analysis, **1** (1991), 39–70.
- [12] P.D. Cordaro, F. Trèves, *Hyperfunctions on hypo-analytic manifolds*, Annals of Mathematics Studies, 136. Princeton University Press, Princeton, NJ, 1994.
- [13] S.-C. Chen, M.-C. Shaw, *Partial differential equations in several complex variables*, Studies in Advanced Mathematics, Vol. 19, Amer. Mat. Soc., International Press, Providence, RI, 2001.
- [14] T. V. Gramchev and P. R. Popivanov. *Partial differential equations*. Mathematical Research, 108, Wiley-VCH Verlag Berlin GmbH, Berlin, 2000.
- [15] C.D. Hill, M. Nacinovich, *On the failure of the Poincaré lemma for $\bar{\partial}_M$ II*, Math. Ann., **335** (2006), 193–219.
- [16] L. Hörmander, *Differential operators of principal type*, Math. Ann., **140** (1960), 124–146.
- [17] H. Lewy, *An example of a smooth linear partial differential equation without solution*, Ann. of Math., **66** (1957), 155–158.
- [18] N. Lerner, *Metrics on the Phase Space and Non-Selfadjoint Pseudodifferential Operators*, Birkhäuser, Basel, 2010.
- [19] M. Mascarello, L. Rodino, *Partial differential equations with multiple characteristics*, Wiley-VCH, Berlin 1997.
- [20] V. Michel, *Sur la régularité C^∞ du $\bar{\partial}$ au bord d’un domaine de \mathbb{C}^n dont la forme de Levi a exactement s valeurs propres négatives*, Math. Ann., **295** (1993), 135–161.

- [21] V. Michel, *Résolution locale du $\bar{\partial}$ avec régularité Gevrey au bord d'un domaine r -convexe*, Math. Z., **218** (1995), 305–317.
- [22] M. Nacinovich, *Poincaré lemma for tangential Cauchy-Riemann complexes*, Math. Ann., **268** (1984), 449–471.
- [23] M. Nacinovich, *On strict Levi q -convexity and q -concavity on domains with piecewise smooth boundaries*, Math. Ann., **281** (1988), 459–482.
- [24] F. Nicola, *On the absence of the one-sided Poincaré lemma in Cauchy-Riemann manifolds*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **4** (2005), no. 4, 587–600.
- [25] F. Nicola, *On Local solvability of certain differential complexes*, Proc. Amer. Math. Soc., **136**(1) (2008), 351–358.
- [26] M.M. Peloso, F. Ricci, *Tangential Cauchy-Riemann equations on quadratic CR manifolds*, Rend. Mat. Acc. Lincei, **13** (2002), 125–134.
- [27] L. Rodino, *Linear partial differential operators in Gevrey spaces*, 1993, World Scientific, Singapore.
- [28] F. Trèves, *Hypo-analytic structures: local theory*, Princeton University Press, Princeton, 1992.

DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY

DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY

E-mail address: francesco.malaspina@polito.it

E-mail address: fabio.nicola@polito.it