

On the generalized sum of the symmetric q-integers

Original

On the generalized sum of the symmetric q-integers / Sparavigna, Amelia Carolina. - ELETTRONICO. - (2018).
[10.5281/zenodo.1248959]

Availability:

This version is available at: 11583/2708001 since: 2018-05-22T11:12:23Z

Publisher:

Zenodo

Published

DOI:10.5281/zenodo.1248959

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

On the generalized sum of the symmetric q-integers

Amelia Carolina Sparavigna

Politecnico di Torino

Abstract Here we will show that the symmetric q-integers of the q-calculus have a generalized sum which is also the generalized sum that we can find in the κ -calculus proposed by G. Kaniadakis.

Keywords q-calculus, q-integers, Kaniadakis κ -entropy.

DOI: 10.5281/zenodo.1248959

Written in Turin, on May 17, 2018

Introduction In a previous work [1], we have discussed the group of the q-integers as defined by q-calculus. In the notation given in the book by Kac and Cheung [2], the q-integers are:

$$(1) \quad [n] = \frac{q^n - 1}{q - 1} = 1 + q + q^2 + \dots + q^{n-1} .$$

In [1], we defined the generalized sum of the group as:

$$(2) \quad [m] \oplus [n] = [m] + [n] + (q - 1)[m][n]$$

As a consequence, we have that the q-integers (1) with operation (2) form a multiplicative group. The generalized sum (2) is similar to the generalized sum that we find for the Tsallis entropies of independent systems [3].

In the q-calculus [2], it is also defined the symmetric q-integer in the following form (here we use a notation different from that given in the Ref.2):

$$(3) \quad [n]_s = \frac{q^n - q^{-n}}{q - q^{-1}}$$

Repeating the approach used in [1], we can determine the group of the symmetric q-integers.

Let us start from the q-integer $[m+n]_s$, which is according to (3):

$$[m+n]_s = \frac{q^{m+n} - q^{-(m+n)}}{q - q^{-1}}$$

and try to find it as a generalized sum of the q-integers $[m]_s$ and $[n]_s$.

By writing $q = \exp(\log q)$, the q-integer turns out into a hyperbolic sine:

$$(4) \quad [n]_s = \frac{q^n - q^{-n}}{q - q^{-1}} = \frac{e^{n \log q} - e^{-n \log q}}{q - q^{-1}} = 2 \frac{\sinh(n \log q)}{(q - q^{-1})}$$

Apart from a numerical factor, this is the form of the generalized numbers proposed by G. Kaniadakis in his κ -calculus [4-8].

From (4), we can write also:

$$\frac{1}{2}(q - q^{-1})[n]_s = \sinh(n \log q)$$

Therefore:

$$[m+n]_s = \frac{q^{m+n} - q^{-(m+n)}}{q - q^{-1}} = 2 \frac{\sinh((m+n) \log q)}{(q - q^{-1})}$$

Using the properties:

$$\sinh(x+y) = \sinh x \cosh y + \cosh x \sinh y \quad ; \quad \cosh x = \sqrt{1 + \sinh^2 x}$$

we obtain:

$$[m+n]_s = \frac{2}{(q - q^{-1})} [\sinh(m \log q) \cosh(n \log q) + \sinh(n \log q) \cosh(m \log q)]$$

$$[m+n]_s = [m]_s \cosh(n \log q) + [n]_s \cosh(m \log q)$$

$$[m+n]_s = [m]_s \sqrt{1 + \sinh^2(n \log q)} + [n]_s \sqrt{1 + \sinh^2(m \log q)}$$

Let us define: $k = (q - q^{-1})/2$ and then: $k[n]_s = \sinh(n \log q)$.

As a consequence we have the generalized sum of the symmetric q-integers as:

$$(5) \quad [m]_s \oplus [n]_s = [m]_s \sqrt{1 + k^2 [n]_s^2} + [n]_s \sqrt{1 + k^2 [m]_s^2}$$

Let us conclude stressing that (5) is also the generalized sum proposed by G. Kaniadakis in the framework of a calculus [5-8], the details of which are given in [8]. By means of (5), we can repeat the approach given in Ref.1 and study of the group of the symmetric q-integers.

References

1. Sparavigna, A. C. (2018). On the additive group of q-integers. Zenodo. DOI: 10.5281/zenodo.1245849
2. Kac, V., & Pokman Cheung (2002). Quantum Calculus, Springer, Berlin.
3. Tsallis, C. (1988). Possible Generalization of Boltzmann-Gibbs Statistics, Journal of Statistical Physics, 52: 479–487. DOI:10.1007/BF01016429
4. In a private discussion with the author, Giorgio Kaniadakis pointed out the form (4) of the q-integers as that of the generalized numbers given by Equation 9 of Ref. 8.
5. Kaniadakis, G. (2001). Non-linear kinetics underlying generalized statistics. Physica A, 296, 405–425.
6. Kaniadakis, G. (2002). Statistical mechanics in the context of special relativity. Phys. Rev. E, 66, 056125.
7. Kaniadakis, G. (2005). Statistical mechanics in the context of special relativity II. Phys. Rev. E, 72, 036108.
8. Kaniadakis, G. (2013). Theoretical Foundations and Mathematical Formalism of the Power-Law Tailed Statistical Distributions. Entropy, 15, 3983-4010.