

The quotient set of  $k$ -generalised Fibonacci numbers is dense in  $\mathbb{Q}_p$

*Original*

The quotient set of  $k$ -generalised Fibonacci numbers is dense in  $\mathbb{Q}_p$  / Sanna, Carlo. - In: BULLETIN OF THE AUSTRALIAN MATHEMATICAL SOCIETY. - ISSN 0004-9727. - STAMPA. - 96:1(2017), pp. 24-29. [10.1017/S0004972716001118]

*Availability:*

This version is available at: 11583/2722651 since: 2020-05-03T09:44:47Z

*Publisher:*

AUSTRALIAN MATHEMATICS PUBL ASSOC INC

*Published*

DOI:10.1017/S0004972716001118

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

# THE QUOTIENT SET OF $k$ -GENERALIZED FIBONACCI NUMBERS IS DENSE IN $\mathbb{Q}_p$

CARLO SANNA

ABSTRACT. The quotient set of  $A \subseteq \mathbb{N}$  is defined as  $R(A) := \{a/b : a, b \in A, b \neq 0\}$ . Using algebraic number theory in  $\mathbb{Q}(\sqrt{5})$ , Garcia and Luca proved that the quotient set of Fibonacci numbers is dense in the  $p$ -adic numbers  $\mathbb{Q}_p$ , for all prime numbers  $p$ . For any integer  $k \geq 2$ , let  $(F_n^{(k)})_{n \geq -(k-2)}$  be the sequence of  $k$ -generalized Fibonacci numbers, defined by the initial values  $0, 0, \dots, 0, 1$  ( $k$  terms) and such that each term afterwards is the sum of the  $k$  preceding terms. We use  $p$ -adic analysis to generalize Garcia and Luca's result, by proving that the quotient set of  $k$ -generalized Fibonacci numbers is dense in  $\mathbb{Q}_p$ , for any integer  $k \geq 2$  and any prime number  $p$ .

## 1. INTRODUCTION

Given a set of nonnegative integers  $A$ , the *quotient set* of  $A$  is defined as

$$R(A) := \{a/b : a, b \in A, b \neq 0\}.$$

The question of when  $R(A)$  is dense in  $\mathbb{R}^+$  is a classical topic and has been studied by many researchers (see, e.g., [1, 3, 4, 10, 12, 13, 17, 19]). On the other hand, the analog question of when  $R(A)$  is dense in the  $p$ -adic numbers  $\mathbb{Q}_p$ , for some prime number  $p$ , has been studied only recently [8, 9]. Let  $(F_n)_{n \geq 0}$  be the sequence of Fibonacci numbers, defined by  $F_0 = 0$ ,  $F_1 = 1$ , and  $F_n = F_{n-1} + F_{n-2}$ , for all integers  $n > 1$ . Using algebraic number theory in the field  $\mathbb{Q}(\sqrt{5})$ , Garcia and Luca [9] proved the following result.

**Theorem 1.1.** *For any prime  $p$ , the quotient set of Fibonacci numbers is dense in  $\mathbb{Q}_p$ .*

One of the many generalizations of the Fibonacci numbers is the sequence of  *$k$ -generalized Fibonacci numbers*  $(F_n^{(k)})_{n \geq -(k-2)}$ , also called *Fibonacci  $k$ -step sequence*, *Fibonacci  $k$ -sequence*, or  *$k$ -bonacci sequence*. For any integer  $k \geq 2$ , the sequence  $(F_n^{(k)})_{n \geq -(k-2)}$  is defined by

$$F_{-(k-2)}^{(k)} = \dots = F_0^{(k)} = 0, F_1^{(k)} = 1,$$

and

$$F_n^{(k)} = F_{n-1}^{(k)} + F_{n-2}^{(k)} + \dots + F_{n-k}^{(k)},$$

for all integers  $n > 1$ .

Usually, the study of the arithmetic properties of the  $k$ -generalized Fibonacci numbers is more difficult than that of Fibonacci numbers. Indeed, for  $k \geq 3$  the sequence of  $k$ -generalized Fibonacci numbers lacks several nice properties of the sequence of Fibonacci numbers, like: being a strong divisibility sequence [16, p. 9], having a Primitive Divisor Theorem [21], and having a simple formula for its  $p$ -adic valuation [14, 18].

We give the following generalization of Theorem 1.1.

**Theorem 1.2.** *For any integer  $k \geq 2$  and any prime number  $p$ , the quotient set of the  $k$ -generalized Fibonacci numbers is dense in  $\mathbb{Q}_p$ .*

It seems likely that Theorem 1.2 could be extended to other linear recurrences over the integers. However, in our proof we use some specific features of the  $k$ -generalized Fibonacci numbers sequence. Therefore, we leave the following open question to the interested readers:

---

2010 *Mathematics Subject Classification*. Primary: 11B39, Secondary: 11B37, 11B05.

*Key words and phrases*. Fibonacci numbers;  $k$ -generalized Fibonacci numbers;  $p$ -adic numbers; density.

Under which (reasonable) hypothesis is the quotient set of a linear recurrence over the integers dense in  $\mathbb{Q}_p$ , for some prime number  $p$ ?

## 2. PROOF OF THEOREM 1.2

From now on, fix an integer  $k \geq 2$  and a prime number  $p$ . In light of Theorem 1.1, we can suppose  $k \geq 3$ . Let

$$f_k(X) = X^k - X^{k-1} - \dots - X - 1$$

be the characteristic polynomial of the  $k$ -generalized Fibonacci numbers sequence.

It is known [20, Corollary 3.4] that  $f_k$  is separable. Let  $K$  be the splitting field of  $f_k$  over  $\mathbb{Q}_p$  and let  $\alpha_1, \dots, \alpha_k \in K$  be the  $k$  distinct roots of  $f_k$ . We have [5, Theorem 1]

$$(1) \quad F_n^{(k)} = \sum_{i=1}^k c_i \alpha_i^n,$$

for all integers  $n \geq 0$ , where

$$(2) \quad c_i := \frac{\alpha_i - 1}{(k+1)\alpha_i^2 - 2k\alpha_i},$$

for  $i = 1, \dots, k$ .

Now we shall interpolate a subsequence of  $(F_n^{(k)})_{n \geq 0}$  by a function analytic over  $\mathbb{Z}_p$ . This is a classical method in the study of linear recurrences, which goes back at least to the proof of Skolem–Mahler–Lech theorem [6, Theorem 2.1].

We refer the reader to [11, Ch. 4–6] for the  $p$ -adic analysis used hereafter. Let  $\mathcal{O}_K$  be the valuation ring of  $K$ ;  $e$  and  $f$  be the ramification index and the inertial degree of  $K$  over  $\mathbb{Q}_p$ , respectively; and  $\pi$  be a uniformizer of  $K$ .

Looking at the Newton's polygon of  $f_k$ , we get that  $|\alpha_i|_p = 1$  for all  $i = 1, \dots, k$ . Hence, in particular,  $\alpha_i \not\equiv 0 \pmod{\pi}$ . Thus, since  $\mathcal{O}_K/\pi\mathcal{O}_K$  is a finite field of  $p^f$  elements, we obtain that  $\alpha_i^{p^f-1} \equiv 1 \pmod{\pi}$ . Now pick any positive integer  $s$  such that  $p^s \geq e+1$ . Being  $|\pi|_p = p^{-1/e}$ , we have  $\pi^{p^s} \equiv 0 \pmod{p\pi}$ , and, in turn, it follows that  $\alpha_i^t \equiv 1 \pmod{p\pi}$ , where  $t := p^s(p^f - 1)$ . At this point,

$$(3) \quad |\alpha_i^t - 1|_p \leq |p\pi|_p = p^{-1-1/e} < p^{-1/(p-1)},$$

for  $i = 1, \dots, k$ .

Now let  $\log_p$  and  $\exp_p$  denote the  $p$ -adic logarithm and the  $p$ -adic exponential functions, respectively. Thanks to (3) we have that

$$\alpha_i^t = \exp_p(\log_p(\alpha_i^t)),$$

for  $i = 1, \dots, k$ , which together with (1) implies that  $F_{nt}^{(k)} = G(n)$  for all integer  $n \geq 0$ , where

$$G(z) := \sum_{i=1}^k c_i \exp_p(z \log_p(\alpha_i^t)),$$

is an analytic function over  $\mathbb{Z}_p$ .

Let  $r > 0$  be the radius of convergence of the Taylor series of  $G(z)$  at  $z = 0$ , and let  $\ell \geq 0$  be an integer. On the one hand, the radius of convergence of the Taylor series of  $G(p^\ell z)$  at  $z = 0$  is  $p^\ell r$ . On the other hand,

$$G(p^\ell z) = \sum_{i=1}^k c_i \exp_p(p^\ell z \log_p(\alpha_i^t)) = \sum_{i=1}^k c_i \exp_p(z \log_p(\alpha_i^{p^\ell t})).$$

Therefore, taking  $s$  sufficiently large, we can assume  $r > 1$ .

In particular, we have

$$(4) \quad G(z) = \sum_{j=0}^{\infty} \frac{G^{(j)}(0)}{j!} z^j,$$

for all  $z \in \mathbb{Z}_p$ .

Now we shall prove that  $G'(0) \neq 0$ . For the sake of contradiction, assume that

$$G'(0) = \sum_{i=1}^k c_i \log_p(\alpha_i^t) = 0.$$

Since  $f_k(0) = -1$  and  $t$  is even, we have  $\alpha_1^t \cdots \alpha_k^t = 1$ , so that

$$\log_p(\alpha_k^t) = -\log_p(\alpha_1^t) - \cdots - \log_p(\alpha_{k-1}^t),$$

and consequently

$$(5) \quad \sum_{i=1}^{k-1} (c_i - c_k) \log_p(\alpha_i^t) = 0.$$

We need the following lemma [7, Lemma 1], which is a special case of a general result of Mignotte [15] on Pisot numbers.

**Lemma 2.1.** *The roots  $\alpha_1, \dots, \alpha_{k-1}$  are multiplicatively independent, that is,  $\alpha_1^{e_1} \cdots \alpha_{k-1}^{e_{k-1}} = 1$  for some integers  $e_1, \dots, e_{k-1}$  if and only if  $e_1 = \cdots = e_{k-1} = 0$ .*

Thanks to Lemma 2.1, we know that  $\alpha_1^t, \dots, \alpha_{k-1}^t$  are multiplicatively independent. Hence,  $\log_p(\alpha_1^t), \dots, \log_p(\alpha_{k-1}^t)$  are linearly independent over  $\mathbb{Z}$ . Then by [2, Theorem 1] we get that  $\log_p(\alpha_1^t), \dots, \log_p(\alpha_{k-1}^t)$  are linearly independent over the algebraic numbers, hence (5) implies

$$(6) \quad c_1 = c_2 = \cdots = c_k.$$

At this point, from (2) and (6) it follows that  $\alpha_1, \dots, \alpha_k$  are all roots of the polynomial

$$c_1(k+1)X^2 - (2c_1k+1)X + 1,$$

but that is clearly impossible, since  $k \geq 3$ . Hence we have proved that  $G'(0) \neq 0$ .

Taking  $z = 1$  in (4), we find that  $\nu_p(G^{(j)}(0)/j!) \rightarrow +\infty$ , as  $j \rightarrow +\infty$ . In particular, there exists an integer  $\ell \geq 0$  such that  $\nu_p(G^{(j)}(0)/j!) \geq -\ell$ , for all integers  $j \geq 0$ . As a consequence of this, and since  $G(0) = F_0^{(k)} = 0$ , taking  $z = mp^h$  in (4) we get that

$$G(mp^h) = G'(0)mp^h + O(p^{2h-\ell}),$$

for all integers  $m, h \geq 0$ . Therefore, for  $h > h_0 := \ell + \nu_p(G'(0))$ , we have

$$\frac{G(mp^h)}{G(p^h)} - m = \frac{G'(0)mp^h + O(p^{2h-\ell})}{G'(0)p^h + O(p^{2h-\ell})} - m = \frac{O(p^{h-\ell})}{G'(0) + O(p^{h-\ell})} = O(p^{h-h_0}),$$

that is,

$$\lim_{h \rightarrow +\infty} \left| \frac{G(mp^h)}{G(p^h)} - m \right|_p = 0.$$

In conclusion, we have proved that

$$\lim_{v \rightarrow +\infty} \left| \frac{F^{(k)}(mp^v(p^f-1))}{F^{(k)}(p^v(p^f-1))} - m \right|_p = 0,$$

for all integers  $m \geq 0$ . In other words, the closure (respect to the  $p$ -adic topology) of the quotient set of  $k$ -generalized Fibonacci numbers contains the nonnegative integers  $\mathbb{N}$ .

The next easy lemma is enough to conclude.

**Lemma 2.2.** *Let  $A \subseteq \mathbb{N}$ . If the closure of  $R(A)$  contains  $\mathbb{N}$ , then  $R(A)$  is dense in  $\mathbb{Q}_p$ .*

*Proof.* Let  $C$  be the closure of  $R(A)$  as a subspace of  $\mathbb{Q}_p$ . Since  $\mathbb{N}$  is dense in  $\mathbb{Z}_p$ , we have  $\mathbb{Z}_p \subseteq C$ . Moreover, the inversion  $\iota : \mathbb{Z}_p^\times \rightarrow \mathbb{Q}_p : x \rightarrow x^{-1}$  is continuous and, obviously, sends nonzero elements of  $R(A)$  to  $R(A)$ , hence  $\iota(\mathbb{Z}_p) \subseteq C$ . Finally,  $\mathbb{Q}_p = \mathbb{Z}_p \cup \iota(\mathbb{Z}_p)$ , thus  $C = \mathbb{Q}_p$  and  $R(A)$  is dense in  $\mathbb{Q}_p$ .  $\square$

The proof of Theorem 1.2 is complete.

## REFERENCES

1. B. Brown, M. Dairyko, S. R. Garcia, B. Lutz, and M. Someck, *Four quotient set gems*, Amer. Math. Monthly **121** (2014), no. 7, 590–599.
2. A. Brumer, *On the units of algebraic number fields*, Mathematika **14** (1967), 121–124.
3. J. Bukor, T. Šalát, and J. T. Tóth, *Remarks on  $R$ -density of sets of numbers*, Tatra Mt. Math. Publ. **11** (1997), 159–165, Number theory (Liptovský Ján, 1995).
4. J. Bukor and J. T. Tóth, *On accumulation points of ratio sets of positive integers*, Amer. Math. Monthly **103** (1996), no. 6, 502–504.
5. G. P. B. Dresden and Z. Du, *A simplified Binet formula for  $k$ -generalized Fibonacci numbers*, J. Integer Seq. **17** (2014), no. 4, Article 14.4.7, 9.
6. G. Everest, A. van der Poorten, I. Shparlinski, and T. Ward, *Recurrence sequences*, Mathematical Surveys and Monographs, vol. 104, American Mathematical Society, Providence, RI, 2003.
7. C. Fuchs, C. Hutle, F. Luca, and L. Szalay, *Diophantine triples and  $k$ -generalized Fibonacci sequences*, Bull. Malays. Math. Sci. Soc. (2016), 1–17.
8. S. R. Garcia, Y. X. Hong, F. Luca, E. Pinsky, E. Schechter, and A. Starr,  *$p$ -adic quotient sets*, <https://arxiv.org/abs/1607.07951>.
9. S. R. Garcia and F. Luca, *Quotients of Fibonacci numbers*, Amer. Math. Monthly (to appear).
10. S. R. Garcia, V. Selhorst-Jones, D. E. Poore, and N. Simon, *Quotient sets and Diophantine equations*, Amer. Math. Monthly **118** (2011), no. 8, 704–711.
11. F. Q. Gouvêa,  *$p$ -adic numbers*, second ed., Universitext, Springer-Verlag, Berlin, 1997, An introduction.
12. S. Hedman and D. Rose, *Light subsets of  $\mathbb{N}$  with dense quotient sets*, Amer. Math. Monthly **116** (2009), no. 7, 635–641.
13. D. Hobby and D. M. Silberger, *Quotients of primes*, Amer. Math. Monthly **100** (1993), no. 1, 50–52.
14. T. Lengyel, *The order of the Fibonacci and Lucas numbers*, Fibonacci Quart. **33** (1995), no. 3, 234–239.
15. M. Mignotte, *Sur les conjugués des nombres de Pisot*, C. R. Acad. Sci. Paris Sér. I Math. **298** (1984), no. 2, 21.
16. P. Ribenboim, *My numbers, my friends*, Springer-Verlag, New York, 2000, Popular lectures on number theory.
17. T. Šalát, *On ratio sets of sets of natural numbers*, Acta Arith. **15** (1968/1969), 273–278.
18. C. Sanna, *The  $p$ -adic valuation of Lucas sequences*, Fibonacci Quart. **54** (2016), no. 2, 118–124.
19. O. Strauch and J. T. Tóth, *Asymptotic density of  $A \subset \mathbb{N}$  and density of the ratio set  $R(A)$* , Acta Arith. **87** (1998), no. 1, 67–78.
20. D. A. Wolfram, *Solving generalized Fibonacci recurrences*, Fibonacci Quart. **36** (1998), no. 2, 129–145.
21. M. Yabuta, *A simple proof of Carmichael’s theorem on primitive divisors*, Fibonacci Quart. **39** (2001), no. 5, 439–443.

UNIVERSITÀ DEGLI STUDI DI TORINO, DEPARTMENT OF MATHEMATICS, TURIN, ITALY

*E-mail address:* [carlo.sanna.dev@gmail.com](mailto:carlo.sanna.dev@gmail.com)

*URL:* <http://orcid.org/0000-0002-2111-7596>