

Critical Robertson–Walker universes

Original

Critical Robertson–Walker universes / Eshkobilov, Olimjon; Musso, Emilio; Nicolodi, Lorenzo. - In: DIFFERENTIAL GEOMETRY AND ITS APPLICATIONS. - ISSN 0926-2245. - ELETTRONICO. - 66:(2019), pp. 23-41. [10.1016/j.difgeo.2019.05.008]

Availability:

This version is available at: 11583/2734818 since: 2019-06-07T11:50:43Z

Publisher:

Elsevier

Published

DOI:10.1016/j.difgeo.2019.05.008

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Elsevier postprint/Author's Accepted Manuscript

© 2019. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:
<http://dx.doi.org/10.1016/j.difgeo.2019.05.008>

(Article begins on next page)

CRITICAL ROBERTSON–WALKER UNIVERSES

OLIMJON ESHKOBILOV, EMILIO MUSSO, AND LORENZO NICOLodi

ABSTRACT. The integral of the energy density function \mathfrak{m} of a closed Robertson–Walker (RW) spacetime with source a perfect fluid and cosmological constant Λ gives rise to an action functional on the space of scale functions of RW spacetime metrics. This paper studies closed RW spacetimes which are critical for this functional, subject to volume-preserving variations (\mathfrak{m} -critical RW spacetimes). A complete classification of \mathfrak{m} -critical RW spacetimes is given and explicit solutions in terms of Weierstrass elliptic functions and their degenerate forms are computed. The standard energy conditions (weak, dominant, and strong) as well as the cyclic property of \mathfrak{m} -critical RW spacetimes are discussed.

1. INTRODUCTION

Robertson–Walker (RW) spacetimes with source a perfect fluid are the most basic cosmological models in general relativity. Despite their old history [16, 23, 32, 36], RW cosmological models still provide a valuable testing ground for new ideas and theories. These include gravitational thermodynamics [9, 34, 39], relativistic diffusion [2], cyclic and conformal cyclic cosmology [3, 30, 31, 35], Lorentz conformal geometry [1, 4, 15], super-symmetry and string theory [7, 33].

In this work we will investigate a natural variational problem for closed RW spacetimes filled with a perfect fluid and satisfying the Einstein equation with a cosmological constant. More precisely, let $\mathcal{M}(\mathfrak{s}) = (\mathbf{I}_{\mathfrak{s}} \times \mathbb{S}^3, \ell_{\mathfrak{s}} = -dt^2 + \mathfrak{s}^2(t)g)$ be a closed RW spacetime, where \mathfrak{s} , the scale function, is a nonnegative smooth function on an open interval $\mathbf{I}_{\mathfrak{s}} \subset \mathbb{R}$ and g is the standard metric of the 3-sphere \mathbb{S}^3 . A closed RW cosmological model (universe) is a closed RW spacetime $\mathcal{M}(\mathfrak{s})$ whose RW metric $\ell_{\mathfrak{s}}$ satisfies the Einstein equation with cosmological constant Λ and source given by the stress-energy tensor of a perfect fluid, $T = \mathfrak{m}dt^2 + \mathfrak{p}\mathfrak{s}^2g$, where the energy density \mathfrak{m} and the pressure \mathfrak{p} are scalar functions of t only. Accordingly, the integral

$$(1.1) \quad \mathfrak{M}[\mathfrak{s}; \Lambda] = \int_{\mathcal{D}} \mathfrak{m} dV_{\ell_{\mathfrak{s}}}$$

defines an action functional on the space of scale functions of RW metrics. Here $dV_{\ell_{\mathfrak{s}}}$ stands for the volume element of the RW metric $\ell_{\mathfrak{s}}$ and $\mathcal{D} = [t_0, t_1] \times \mathbb{S}^3$ is a compact domain of $\mathbf{I}_{\mathfrak{s}} \times \mathbb{S}^3$. The critical points of (1.1) with respect to volume-preserving

Date: Version of August 14, 2018.

2010 Mathematics Subject Classification. 53C50, 53B30, 53Z05, 83C15, 33E05, 58E30.

Key words and phrases. Lorentz metrics, Robertson–Walker spacetimes, perfect fluid, conformal time, cyclic models in general relativity, weak, dominant, and strong energy conditions, elliptic functions and integrals, Weierstrass \wp -functions.

Authors partially supported by PRIN 2015-2018 “Varietà reali e complesse: geometria, topologia e analisi armonica”; by the GNSAGA of INDAM; and by the FFABR Grant 2017 of MIUR.

variations of the RW metric are called *m-critical RW universes* (or *critical models*, for short). After deriving the variational equation, which is a 2nd order ODE in the scale function \mathfrak{s} , and computing the equation of state fulfilled by the energy density \mathfrak{m} and the pressure \mathfrak{p} , we rewrite the variational equation in terms of the conformal time τ , defined by $d\tau = dt/\mathfrak{s}$. The advantage of this coordinate transformation is that the resulting *conformal scale function* can be described as a real form of a Weierstrass \wp -function. According to whether the discriminant of the cubic polynomial of such an associated Weierstrass \wp -function is different or equal to zero, a critical model is said to be of *general* or *exceptional* type. The critical models of exceptional type can be easily described in terms of trigonometric or hyperbolic functions (cf. Section 2). The main purpose of this paper is to study the critical models of general type. We will describe their structure and discuss their energy conditions and periodicity properties.

Special and elliptic functions have been already used in the literature to describe particular classes of RW universes [3, 6, 10, 11]. However, this was done by imposing the equation of state a priori, while our models are obtained from a rather natural variational principle. Weierstrass elliptic functions are also used in the study of the restricted evolution problem within the more general class of Lemaître–Tolman–Bondi spacetimes [5]. In addition, the theory of elliptic functions plays a relevant role in the study of null geodesics in Schwarzschild spacetime [18], higher-order variational problems for null curves in Lorentz space forms [14, 17, 20, 24, 25, 26, 27, 28], and conformally invariant variational problems for timelike curves in the Einstein static universe [12].

The paper is organized as follows. Section 2 collects some basic facts about closed RW cosmological models [19, 29, 37, 38] and derives the variational equation and the equation of state satisfied by the scale function of a critical model (cf. Proposition 1). As a byproduct, it follows that the critical scale functions depend on two internal parameters, denoted by \mathfrak{a} and \mathcal{H} . Proposition 2 writes the conformal scale function, the conformal energy density, and the conformal pressure,¹ in terms of real forms of Weierstrass \wp -functions. A critical model is said to be of *general* or *exceptional* type, according to whether the discriminant of the cubic polynomial of the associated Weierstrass \wp -function is different or equal to zero. The general critical models are further divided in two classes, namely Class I and Class II, depending on the positivity or negativity of the discriminant. The section terminates with the discussion and the explicit description of the exceptional critical models.

Section 3 studies the critical models of Class I. Theorem 3 explicitly describes the critical models of Class I, with $\mathcal{H} < 0$, and determines the bounds satisfied by the cosmological constant in order that the model fulfils the weak, dominant, or strong energy conditions. It is also shown that these solutions are cyclic and nondegenerate. Theorem 4 provides similar results for the critical models of Class I, with $\mathcal{H} > 0$. In the latter case, the scale function vanishes along the disjoint union of a countable family of totally umbilical, connected, spacelike hypersurfaces.

Section 4 is devoted to the analysis of critical models of Class II. We distinguish two possible types: negative or positive. Theorem 5 deals with the critical models of Class II and negative type. We analyze the bounds on the cosmological constant

¹i.e., the scale function, the energy density, and the pressure written in terms of the conformal time.

so that the weak, the dominant, or the strong energy conditions are satisfied. We also show that these models are cyclic and degenerate along countably many totally umbilical, connected, spacelike hypersurfaces. In Theorem 6, we consider critical models of Class II and positive type, with $\mathcal{H} > 0$. These models are not cyclic and their Lorentz quadratic forms are degenerate along a totally umbilical, connected, spacelike hypersurface. The conformal scale function tends to infinity in finite time. Finally, in Theorem 7 we prove similar results for critical models of Class II and positive type, but with $\mathcal{H} < 0$.

Summarizing, the only genuine cyclic \mathbf{m} -critical models are those corresponding to Class I, with $\mathcal{H} < 0$. In all other cases, the scale function either vanishes along a totally umbilical spacelike hypersurface, or tends to $+\infty$ in finite time.

As a basic reference for Weierstrass elliptic functions we use [22]. We acknowledge the use of the software MATHEMATICA for symbolic and numerical computations throughout the paper.

2. PRELIMINARIES

In this section, after recalling the basic definitions, we derive the variational equation for the scale function of an \mathbf{m} -critical RW universe and write the conformal scale function in terms of a real form of a Weierstrass \wp -function. We find the equation of state and classify the \mathbf{m} -critical RW universes into three classes.

2.1. Closed RW universes and their conformal models. Let g be the standard Riemannian metric of the unit 3-sphere $\mathbb{S}^3 \subset \mathbb{R}^4$. Consider a closed RW spacetime $\mathcal{M} := I_{\mathfrak{s}} \times_{\mathfrak{s}} \mathbb{S}^3$ with a *nonnegative scale function* $\mathfrak{s} : I_{\mathfrak{s}} \subset \mathbb{R} \rightarrow \mathbb{R}^+$, endowed with the (possibly degenerate) Lorentz metric $\ell_{\mathfrak{s}} = -dt^2 + \mathfrak{s}^2(t)g$, the *cosmological constant* Λ and a *perfect fluid stress-energy tensor*

$$T = \mathbf{m}dt^2 + \mathbf{p}\mathfrak{s}^2g,$$

where $\mathbf{m} : I_{\mathfrak{s}} \rightarrow \mathbb{R}^+ \cup \{+\infty\}$ is the *energy density function* and $\mathbf{p} : I_{\mathfrak{s}} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ is the *pressure function*. Let $\rho_{\mathfrak{s}}$ and $\sigma_{\mathfrak{s}}$ be, respectively, the Ricci tensor and the scalar curvature of $\ell_{\mathfrak{s}}$. Then the Einstein field equation²

$$\rho_{\mathfrak{s}} + \left(\Lambda - \frac{1}{2}\sigma_{\mathfrak{s}} \right) \ell_{\mathfrak{s}} = 8\pi T$$

is equivalent to

$$(2.1) \quad \begin{cases} \mathbf{m} = \frac{3}{8\pi} \left(\frac{\dot{\mathfrak{s}}^2}{\mathfrak{s}^2} + \frac{1}{\mathfrak{s}^2} - \frac{\Lambda}{3} \right), \\ \mathbf{p} = -\frac{1}{4\pi\mathfrak{s}} \left(\ddot{\mathfrak{s}} + \frac{1}{2\mathfrak{s}}(\dot{\mathfrak{s}}^2 + 1) - \frac{\Lambda}{2}\mathfrak{s} \right), \end{cases}$$

where dot denotes differentiation with respect to the cosmological time t . We implicitly assume that $I_{\mathfrak{s}}$ is the maximal domain of definition of the scale function, i.e., \mathfrak{s} cannot be extended to a nonnegative differentiable function whose interval of definition contains properly $I_{\mathfrak{s}}$.

Remark 1. The RW spacetime with $I_{\mathfrak{s}} = \mathbb{R}$, $\mathfrak{s} = 1$, $\mathbf{m} + \mathbf{p} = (4\pi)^{-1}$, and $\Lambda = 1 + 8\pi\mathbf{p}$ is the *Einstein static universe* (cf. [13]), which will be denoted by (\mathcal{E}, ℓ_1) .

Definition 1. There are three standard physical conditions that are usually imposed on a RW spacetime (cf. [19]):

²We use units in which the gravitational constant $G = 1$ and the speed of light $c = 1$.

- the *weak energy condition* : $-\mathfrak{m} \leq \mathfrak{p}$;
- the *dominant energy condition* : $-\mathfrak{m} \leq \mathfrak{p} \leq \mathfrak{m}$;
- the *strong energy condition* : $-\mathfrak{m}/3 \leq \mathfrak{p} \leq \mathfrak{m}$.

Assumption 1. If, on the one hand, the scale function \mathfrak{s} is allowed to vanish at some $t \in I_{\mathfrak{s}}$, we will assume that the *conformal time*

$$(2.2) \quad \tau_{\mathfrak{s}} : I_{\mathfrak{s}} \rightarrow \mathbb{R}, t \mapsto \tau_{\mathfrak{s}}(t) := \int_{t_0}^t \frac{du}{\mathfrak{s}(u)}$$

is a continuous, strictly increasing function.

Under the previous assumption, $J_{\mathfrak{s}} := \tau_{\mathfrak{s}}(I_{\mathfrak{s}})$ is an open interval and $\tau_{\mathfrak{s}}^{-1} : J_{\mathfrak{s}} \rightarrow \mathbb{R}$ is a strictly increasing differentiable function. Moreover, the map

$$(2.3) \quad \phi_{\mathfrak{s}} : J_{\mathfrak{s}} \times \mathbb{S}^3 \ni (\tau, p) \mapsto (\tau_{\mathfrak{s}}^{-1}(\tau), w) \in \mathcal{M}.$$

is a differentiable homeomorphism, such that

$$\phi_{\mathfrak{s}}^*(\ell_{\mathfrak{s}}) = S^2(-dt^2 + g),$$

where $S := \mathfrak{s} \circ \tau_{\mathfrak{s}}^{-1} : J_{\mathfrak{s}} \rightarrow \mathbb{R}$ is a nonnegative differentiable function, called the *conformal scale function*. Note that $J_{\mathfrak{s}}$ is the maximal domain of definition of S . Therefore, $\phi_{\mathfrak{s}}$ is a weakly conformal homeomorphism of the open domain $J_{\mathfrak{s}} \times \mathbb{S}^3$ of the Einstein static universe onto \mathcal{M} . The map $\phi_{\mathfrak{s}}$ fails to be of maximal rank at the points $(\tau, p) \in J_{\mathfrak{s}} \times \mathbb{S}^3$, such that $S(\tau) = 0$. The inverse map $\phi_{\mathfrak{s}}^{-1}$ is a homeomorphism which fails to be differentiable at the points $(t, p) \in I_{\mathfrak{s}} \times \mathbb{S}^3$, such that $\mathfrak{s}(t) = 0$.

Definition 2. Let $\ell_{\mathfrak{s}}$ denote the quadratic form $\phi_{\mathfrak{s}}^*(\ell_{\mathfrak{s}})$ and put $\widetilde{\mathcal{M}} = J_{\mathfrak{s}} \times \mathbb{S}^3$. We say that $(\widetilde{\mathcal{M}}, \ell_{\mathfrak{s}})$ is the *conformal model* of the RW spacetime with scale function \mathfrak{s} . If $J_{\mathfrak{s}} = \mathbb{R}$ and S is periodic, we say that the conformal model $(\widetilde{\mathcal{M}}, \ell_{\mathfrak{s}})$ is *cyclic*. The functions $\widetilde{\mathfrak{m}} := \mathfrak{m} \circ \tau_{\mathfrak{s}}^{-1} : J_{\mathfrak{s}} \rightarrow \mathbb{R} \cup \{+\infty\}$ and $\widetilde{\mathfrak{p}} := \mathfrak{p} \circ \tau_{\mathfrak{s}}^{-1} : J_{\mathfrak{s}} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ are called the *conformal energy density* and the *conformal pressure*, respectively.

In view of (2.1) and the fact that, by (2.2), $d\tau = dt/\mathfrak{s}$, the conformal energy density and the conformal pressure can be written as

$$(2.4) \quad \begin{cases} \widetilde{\mathfrak{m}} = \frac{3}{8\pi} \left(\frac{S'^2}{S^4} + \frac{1}{S^2} - \frac{\Lambda}{3} \right), \\ \widetilde{\mathfrak{p}} = -\frac{1}{4\pi S} \left(\frac{S''}{S^2} - \frac{S'^2}{2S^3} + \frac{1}{2S} - \frac{\Lambda S}{2} \right). \end{cases}$$

Here prime denotes differentiation with respect to the conformal time τ .

2.2. \mathfrak{m} -critical RW universes and their state equations. The total energy of the compact domain $\mathcal{D} = [t_0, t_1] \times \mathbb{S}^3 \subset \mathcal{M}$ is

$$\mathfrak{M} = \int_{\mathcal{D}} \mathfrak{m} dV_{\ell_{\mathfrak{s}}},$$

where $dV_{\ell_{\mathfrak{s}}}$ is the volume element of $\ell_{\mathfrak{s}}$.

Definition 3. A RW spacetime is said to be *\mathfrak{m} -critical* if its scale function is a critical point of the functional \mathfrak{M} with respect to volume-preserving variations of the RW quadratic form $\ell_{\mathfrak{s}}$.

Proposition 1. *A RW spacetime is \mathfrak{m} -critical if and only if its scale function \mathfrak{s} satisfies*

$$(2.5) \quad \mathfrak{s} \left(\dot{\mathfrak{s}}^2 + \frac{\mathfrak{a}}{3} \mathfrak{s}^2 - 1 \right) = \mathcal{H},$$

where \mathfrak{a} and \mathcal{H} are two arbitrary constants, called the internal parameters. In addition, the state equation of an \mathfrak{m} -critical spacetime with parameters \mathfrak{a} and \mathcal{H} is

$$(2.6) \quad \frac{8\pi}{3} \mathfrak{m} + 8\pi \mathfrak{p} - \frac{2}{3} (\Lambda + \mathfrak{a}) - \frac{\mathcal{H}}{2\sqrt{2}} (\Lambda + \mathfrak{a} - 8\pi \mathfrak{p})^{3/2} = 0.$$

Proof. The volume element of $\ell_{\mathfrak{s}}$ is $\mathfrak{s}^3 dt \wedge dv$, where dv is the volume element of the unit 3-sphere \mathbb{S}^3 . We then have

$$\mathfrak{M} = \frac{3\pi}{4} \int_{t_0}^{t_1} \mathfrak{s} \left(\dot{\mathfrak{s}}^2 - \frac{\Lambda}{3} \mathfrak{s}^2 + 1 \right) dt.$$

The constraint on the volume amounts to requiring that

$$(2.7) \quad \int_{t_0}^{t_1} \mathfrak{s}^3 dt = \text{const.}$$

Thus, a RW spacetime is \mathfrak{m} -critical if and only if its scale function \mathfrak{s} is critical for the action integral $\int_{t_0}^{t_1} L(\mathfrak{s}, \dot{\mathfrak{s}}) dt$ corresponding to the Lagrangian

$$L(\mathfrak{s}, \dot{\mathfrak{s}}) = \mathfrak{s} \left(\dot{\mathfrak{s}}^2 - \frac{\Lambda + \lambda}{3} \mathfrak{s}^2 + 1 \right),$$

where $\lambda \in \mathbb{R}$ is the Lagrange multiplier associated to the constraint (2.7) (see [8]). Equivalently, \mathfrak{s} is a solution of the Euler–Lagrange equation of the action,

$$(2.8) \quad \frac{d}{dt} \frac{\partial L}{\partial \dot{\mathfrak{s}}} - \frac{\partial L}{\partial \mathfrak{s}} = 2\mathfrak{s}\ddot{\mathfrak{s}} + \dot{\mathfrak{s}}^2 + \mathfrak{a}\mathfrak{s}^2 - 1 = 0,$$

where $\mathfrak{a} := \Lambda + \lambda$. Now, since L does not depend explicitly on time, taking the total time derivative of L and replacing $\frac{\partial L}{\partial \mathfrak{s}}$ by $\frac{d}{dt} \frac{\partial L}{\partial \dot{\mathfrak{s}}}$, in accordance with the Euler–Lagrange equation, yields

$$\frac{dL}{dt} = \frac{\partial L}{\partial \mathfrak{s}} \dot{\mathfrak{s}} + \frac{\partial L}{\partial \dot{\mathfrak{s}}} \ddot{\mathfrak{s}} = \frac{d}{dt} \left(\dot{\mathfrak{s}} \frac{\partial L}{\partial \dot{\mathfrak{s}}} \right).$$

This implies that

$$\dot{\mathfrak{s}} \frac{\partial L}{\partial \dot{\mathfrak{s}}} - L = \mathfrak{s} \left(\dot{\mathfrak{s}}^2 + \frac{\mathfrak{a}}{3} \mathfrak{s}^2 - 1 \right)$$

is a first integral of the motion. Therefore, \mathfrak{s} satisfies (2.8) if and only if there exists a constant \mathcal{H} such that

$$(2.9) \quad \mathfrak{s} \left(\dot{\mathfrak{s}}^2 + \frac{\mathfrak{a}}{3} \mathfrak{s}^2 - 1 \right) = \mathcal{H}.$$

In view of (2.8), the first equation of (2.1) becomes

$$(2.10) \quad \mathfrak{s}^2 = \frac{2}{\Lambda + \mathfrak{a} - 8\pi \mathfrak{p}}.$$

Thus $8\pi \mathfrak{p} \leq (\Lambda + \mathfrak{a})$. Using (2.1), (2.5) and (2.10) it is now an easy matter to check that the equation of state of an \mathfrak{m} -critical RW spacetime, with internal parameters \mathcal{H} and \mathfrak{a} , and cosmological constant Λ , is

$$(2.11) \quad \frac{8\pi}{3} \mathfrak{m} + 8\pi \mathfrak{p} - \frac{2}{3} (\Lambda + \mathfrak{a}) - \frac{\mathcal{H}}{2\sqrt{2}} (\Lambda + \mathfrak{a} - 8\pi \mathfrak{p})^{3/2} = 0.$$

This concludes the proof. \square

Remark 2. If $\mathcal{H} = 0$, then (2.5) can be easily integrated in terms of elementary functions. As a result we find

$$(2.12) \quad \begin{aligned} \mathfrak{s}(t) &= |\sqrt{3/\mathfrak{a}} \sin(\sqrt{\mathfrak{a}/3}t + c)|, \quad \mathfrak{a} > 0, \\ \mathfrak{s}(t) &= |\sqrt{3/\mathfrak{a}} \sinh(\sqrt{\mathfrak{a}/3}t + c)|, \quad \mathfrak{a} < 0, \\ \mathfrak{s}(t) &= |t + c|, \quad \mathfrak{a} = 0. \end{aligned}$$

From now on, we will assume that \mathcal{H} is different from zero and \mathfrak{s} is nonconstant.

2.3. The conformal scale function of a critical RW spacetime.

Definition 4. A *real form* of the Weierstrass elliptic function $\wp(z; g_2, g_3)$ with *real* invariants g_2 and g_3 is a nonconstant function $Q : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$, such that

$$(2.13) \quad Q'^2 = 4Q^3 - g_2Q - g_3.$$

Remark 3. The behavior of these functions depends on the discriminant $\Delta = 16(g_2^3 - 27g_3^2)$ of the cubic polynomial $P(x) = 4x^3 - g_2x - g_3$.

If $\Delta = 0$ (degenerate case), then

- if $g_3 = -8a^3 > 0$, then $Q(\tau) = -3a \tan(\sqrt{-3a}\tau + c)^2 - 2a$, $c \in \mathbb{R}$;
- if $g_3 = -8a^3 > 0$, then either
 - (1) $Q(\tau) = 3a \tanh(\sqrt{3a}\tau + c)^2 - 2a$, $c \in \mathbb{R}$, or
 - (2) $Q(\tau) = a(1 + 3\operatorname{csch}(\sqrt{3a}\tau + c)^2)$, $c \in \mathbb{R}$;
- if $g_3 = g_2 = 0$, then $Q(\tau) = (\tau + c)^2$, $c \in \mathbb{R}$.

If $\Delta \neq 0$, the Weierstrass \wp -function is doubly periodic, with primitive half-periods ω_1, ω_3 , such that $\omega_1 \in \mathbb{R}$, $\omega_1 > 0$ and $\operatorname{Im}(\omega_3/\omega_1) > 0$. If $\Delta > 0$, ω_3 is purely imaginary, while, if $\Delta < 0$, $\operatorname{Re}(\omega_3) \neq 0$. Its real forms are:

- if $\Delta > 0$, then either
 - (1) $Q(\tau) = \wp(\tau + 2m\omega_1 + 2n\omega_3 + c, g_2, g_3)$, $c \in \mathbb{R}$, $m, n \in \mathbb{Z}$, or
 - (2) $Q(\tau) = \wp(\tau + \omega_3 + 2m\omega_1 + 2n\omega_3 + c, g_2, g_3)$, $c \in \mathbb{R}$, $m, n \in \mathbb{Z}$;
- if $\Delta < 0$, then $Q(\tau) = \wp(\tau + 2m\omega_1 + 2n\omega_3 + c, g_2, g_3)$, $c \in \mathbb{R}$, $m, n \in \mathbb{Z}$.

The constants m, n and c are irrelevant and can be put equal to 0.

We can prove the following.

Proposition 2. *The conformal scale function of an \mathfrak{m} -critical RW universe with internal parameters \mathcal{H} and \mathfrak{a} , and cosmological constant Λ , is given by*

$$(2.14) \quad S(\tau) = \frac{3\mathcal{H}}{12Q(\tau) - 1},$$

where Q is a real form of the Weierstrass \wp -function with invariants

$$(2.15) \quad g_2 = \frac{1}{12}, \quad g_3 = \frac{1}{48} \left(\mathfrak{a}\mathcal{H}^2 - \frac{2}{9} \right).$$

The conformal energy density and the conformal pressure are given by

$$(2.16) \quad \begin{aligned} \tilde{\mathfrak{m}} &= \frac{1}{8\pi} \left(\frac{(1 - 12Q)^2(5 + 12Q)}{9\mathcal{H}^2} - \mathfrak{a} - \Lambda \right), \\ \tilde{\mathfrak{p}} &= \frac{1}{4\pi} \left(\frac{\mathfrak{a} + \Lambda}{2} - \frac{(12Q - 1)^2}{9\mathcal{H}^2} \right). \end{aligned}$$

Proof. Differentiating $S = \mathfrak{s} \circ \tau_{\mathfrak{s}}^{-1}$ and using (2.2) we have

$$(2.17) \quad \dot{\mathfrak{s}}|_{\tau_{\mathfrak{s}}^{-1}(\tau)} = S^{-1}(\tau)S'|_{\tau},$$

where \dot{f} is the derivative with respect to the cosmological time and h' the derivative with respect to the conformal time. Therefore, if $\mathcal{H} \neq 0$, the variational equation (2.5) is satisfied if and only if

$$(2.18) \quad S'^2 + \frac{\mathfrak{a}}{3}S^4 - S^2 - \mathcal{H}S = 0.$$

Let

$$(2.19) \quad Q := \frac{\mathcal{H}}{4S} + \frac{1}{12}.$$

It is an easy matter to check that (2.18) holds true if and only if

$$(2.20) \quad Q'^2 = 4Q^3 - \frac{1}{12}Q + \frac{1}{48}\left(\frac{2}{9} - \mathfrak{a}\mathcal{H}^2\right).$$

This proves the first part of the statement. Substituting (2.18) into the first equation of (2.4) and taking into account (2.14), we have

$$\begin{aligned} \tilde{\mathfrak{m}} &= \frac{3}{8\pi} \left[\frac{1}{S^4} \left(-\frac{\mathfrak{a}}{3}S^4 + S^2 + \mathcal{H}S \right) + \frac{1}{S^2} - \frac{\Lambda}{3} \right] \\ &= \frac{3}{8\pi} \left(\frac{2}{S^2} + \frac{\mathcal{H}}{S^3} - \frac{\mathfrak{a} + \Lambda}{3} \right) \\ &= \frac{1}{8\pi} \left(\frac{(1 - 12Q)^2(5 + 12Q)}{9\mathcal{H}^2} - \mathfrak{a} - \Lambda \right). \end{aligned}$$

Differentiating (2.18) yields

$$(2.21) \quad S'' = -\frac{2\mathfrak{a}}{3}S^3 + S + \frac{\mathcal{H}}{2}.$$

Substituting (2.18) and (2.21) into the second equation of (2.4), we obtain

$$\begin{aligned} \tilde{\mathfrak{p}} &= -\frac{1}{4\pi S} \left[\frac{1}{S^2} \left(-\frac{2\mathfrak{a}}{3}S^3 + S + \frac{\mathcal{H}}{2} \right) - \frac{1}{2S^3} \left(-\frac{\mathfrak{a}}{3}S^4 + S^2 + \mathcal{H}S \right) + \frac{1}{2S} - \frac{\Lambda}{2}S \right] \\ &= \frac{1}{4\pi} \left(\frac{\mathfrak{a} + \Lambda}{2} - \frac{1}{S^2} \right) = \frac{1}{4\pi} \left(\frac{\mathfrak{a} + \Lambda}{2} - \frac{(12Q - 1)^2}{9\mathcal{H}^2} \right). \end{aligned}$$

This concludes the proof. \square

Remark 4. The function $\tau_{\mathfrak{s}}^{-1}$ is given by the incomplete elliptic integral of the third kind

$$(2.22) \quad \tau_{\mathfrak{s}}^{-1}(\tau) = 3\mathcal{H} \int \frac{d\tau}{12Q(\tau) - 1}.$$

Actually, this integral can be computed in closed form using the σ and ζ Weierstrass functions (see [22], p. 173). The explicit expression of $\tau_{\mathfrak{s}}$ cannot be given in closed form. However, it can be evaluated using numerical solutions of the first order ODE $h' = 3\mathcal{H}^{-1}(12Q(h) - 1)$.

If g_2 and g_3 are as in (2.15), the discriminant $\Delta = 16(g_2^3 - 27g_3^2)$ takes the form

$$\Delta = \frac{1}{48}\mathfrak{a}\mathcal{H}^2(4 - 9\mathfrak{a}\mathcal{H}^2).$$

Definition 5. According to whether $\Delta \neq 0$ or $\Delta = 0$, an \mathbf{m} -critical RW universe is said to be *general* or *exceptional*. Moreover, a general \mathbf{m} -critical RW universe is said to be of Class I, if $\Delta > 0$, and of Class II, if $\Delta < 0$.

2.4. Exceptional \mathbf{m} -critical RW universes. The conformal scale functions of the exceptional \mathbf{m} -critical RW universes can be written in terms of trigonometric or hyperbolic functions.

- If $\mathcal{H}^2 \mathbf{a} = 4/9$, then $\mathcal{H} > 0$ and

$$S(\tau) = \frac{3\mathcal{H}}{3 \tan^2(\tau/2) + 1}.$$

The quadratic form ℓ_S vanishes if $\tau = (2h+1)\pi$, $h \in \mathbb{Z}$, and is of Lorentz type at all other points. The conformal energy density function $\tilde{\mathbf{m}}$ is non-negative if and only if $\Lambda \leq 1/3\mathcal{H}^2$. Assuming $\tilde{\mathbf{m}} \geq 0$, the weak energy condition is automatically fulfilled, while the dominant energy condition holds true if and only if $\Lambda \leq 1/18\mathcal{H}^2$. The strong energy condition is satisfied if and only if $-1/2\mathcal{H}^2 \leq \Lambda \leq 1/18\mathcal{H}^2$.

- If $\mathbf{a} = 0$ and $\mathcal{H} < 0$, then

$$S(\tau) = -\frac{\mathcal{H}}{2}(1 + \cosh \tau).$$

The quadratic form ℓ_S is nondegenerate of Lorentz type and $\tilde{\mathbf{m}}$ is nonnegative if and only if $\Lambda \leq 0$. Under this assumption the weak and the dominant energy conditions are automatically satisfied. The strong energy condition is never satisfied.

- If $\mathbf{a} = 0$ and $\mathcal{H} > 0$, then

$$S(\tau) = \mathcal{H} \sinh^2(\tau/2).$$

Then ℓ_S vanishes if $\tau = 0$ and is of Lorentz type if $\tau \neq 0$. The conformal energy density $\tilde{\mathbf{m}}$ is nonnegative if and only if $\Lambda \leq 0$. Under this hypothesis, the weak and the dominant energy conditions hold true, while the strong energy condition is satisfied if and only if $\Lambda = 0$.

3. CRITICAL RW UNIVERSES OF CLASS I

This section discusses \mathbf{m} -critical RW universes of Class I. Theorem 3 describes the main features of the critical models of Class I with negative \mathcal{H} , while Theorem 4 discusses the critical models of Class I with positive \mathcal{H} .

If $\Delta > 0$, the cubic polynomial $P(x)$ has three distinct real roots, say $e_1 > e_2 > e_3$, such that

$$e_1 > 0, \quad e_3 = -(e_1 + e_2) < 0, \quad 4(e_1^2 + e_1 e_2 + e_2^2) = 1/12.$$

We write e_1 and e_2 as functions of the *angular parameter* $\vartheta \in (-\pi/3, 0)$,

$$(3.1) \quad \begin{aligned} e_1(\vartheta) &= \frac{1}{4\sqrt{3}} \left(\frac{1}{\sqrt{3}} \cos \vartheta - \sin \vartheta \right), \\ e_2(\vartheta) &= \frac{1}{4\sqrt{3}} \left(\frac{1}{\sqrt{3}} \cos \vartheta + \sin \vartheta \right). \end{aligned}$$

The invariant g_3 and the parameter \mathfrak{a} can be written as functions of \mathcal{H} and ϑ , namely

$$(3.2) \quad g_3 = -\frac{\cos(3\vartheta)}{216}, \quad \mathfrak{a} = \frac{2}{9\mathcal{H}^2} (1 - \cos 3\vartheta).$$

Let \wp_ϑ be the Weierstrass \wp -function with invariants $g_2 = 1/12$ and $g_3(\vartheta)$. Let ω_ϑ and ω'_ϑ be its real and purely imaginary half-periods. Let

$$\tilde{\wp}_\vartheta(\tau) = \wp_\vartheta(\tau + \omega'_\vartheta), \quad \hat{\wp}_\vartheta(\tau) = \wp_\vartheta(\tau), \quad \forall \tau \in \mathbb{R},$$

be the real forms of \wp_ϑ .

3.1. Critical models of Class I with negative \mathcal{H} . With the notation introduced above, let f_1 , f_2 and f_3 be defined by

$$\begin{aligned} f_1(\vartheta, \mathcal{H}) &= \frac{1}{3\mathcal{H}^2} \left(1 - \cos \vartheta - \sqrt{3} \sin \vartheta \right)^2, \\ f_2(\vartheta, \mathcal{H}) &= \frac{2}{9\mathcal{H}^2} \sin^2 \left(\frac{\vartheta}{2} \right) \left(7 + \cos \vartheta - 2 \cos 2\vartheta - 5\sqrt{3} \sin \vartheta \right), \\ f_3(\vartheta, \mathcal{H}) &= -\frac{1}{6\mathcal{H}^2} (1 - 2 \cos \vartheta)(1 + 2 \cos \vartheta)^2. \end{aligned}$$

Note that

$$f_3(\vartheta, \mathcal{H}) \leq f_2(\vartheta, \mathcal{H}), \quad \forall (\vartheta, \mathcal{H}) \in (-\pi/3, \vartheta_I] \times (-\infty, 0),$$

where $\vartheta_I \approx -0.7706314502$ is the root in the interval $(-\pi/3, 0)$ of the equation

$$-10 + 22 \cos x + 11 \cos 2x + 4 \cos 3x + 10\sqrt{3} \sin x - 5\sqrt{3} \sin 2x = 0.$$

We can now state the following.

Theorem 3. *Let $\mathcal{M}(\mathfrak{s})$ be an \mathfrak{m} -critical RW spacetime of Class I, with cosmological constant Λ , angular parameter $\vartheta \in (-\pi/3, 0)$, and negative \mathcal{H} . Let $(\tilde{\mathcal{M}}_{\mathcal{H}, \vartheta}, \ell_{\mathcal{H}, \vartheta})$ denote its conformal model. Then,*

$$\Lambda \leq f_1(\vartheta, \mathcal{H}), \quad \tilde{\mathcal{M}}_{\mathcal{H}, \vartheta} = \mathbb{R} \times \mathbb{S}^3,$$

and $\ell_{\mathcal{H}, \vartheta}$ is the cyclic nondegenerate Lorentz metric

$$(3.3) \quad \ell_{\mathcal{H}, \vartheta} = \frac{9\mathcal{H}^2}{(12\tilde{\wp}_\vartheta - 1)^2} (-d\tau^2 + g).$$

The weak energy condition is automatically satisfied. The dominant energy condition is satisfied if and only if $\Lambda \leq f_2(\vartheta, \mathcal{H})$. The strong energy condition is satisfied if and only if $\vartheta \in (-\pi/3, \vartheta_I]$ and $f_3(\vartheta, \mathcal{H}) \leq \Lambda \leq f_2(\vartheta, \mathcal{H})$.

Proof. According to Proposition 1, we have $S = 3\mathcal{H}/(12Q - 1)$, where Q is one of the two possible real forms of \wp_ϑ . Since $S \geq 0$ and $\mathcal{H} < 0$, it follows that Q is bounded above by $1/12$. The roots

$$e_1(\vartheta) > e_2(\vartheta) > e_3(\vartheta) = -e_1(\vartheta) - e_2(\vartheta)$$

of the polynomial $4t^3 - t/12 - g_3$ satisfy

$$e_3(\vartheta) < e_2(\vartheta) < \frac{1}{12} < e_1(\vartheta).$$

Since

$$e_3(\vartheta) \leq \tilde{\wp}_\vartheta(\tau) \leq e_2(\vartheta), \quad e_1(\vartheta) \leq \hat{\wp}_\vartheta(\tau),$$

Q must coincide with $\tilde{\varphi}_\vartheta$. This implies that S is a strictly positive, real analytic periodic function with minimal period $2\omega_\vartheta$. Thus, the conformal model is $\mathbb{R} \times \mathbb{S}^3$ and ℓ_S coincides with (3.3).

It follows from (2.16) that the conformal energy density and the conformal pressure are given by

$$\begin{aligned}\tilde{\mathbf{m}} &= \frac{1}{8\pi} \left(\frac{(1 - 12\tilde{\varphi}_\vartheta)^2(5 + 12\tilde{\varphi}_\vartheta)}{9\mathcal{H}^2} - \mathbf{a} - \Lambda \right), \\ \tilde{\mathbf{p}} &= \frac{1}{4\pi} \left(\frac{\mathbf{a} + \Lambda}{2} - \frac{(12\tilde{\varphi}_\vartheta - 1)^2}{9\mathcal{H}^2} \right),\end{aligned}$$

where \mathbf{a} is as in (3.2). Thus, $\tilde{\mathbf{m}}$ and $\tilde{\mathbf{p}}$ are real analytic periodic functions of period $2\omega_\vartheta$. The function $\tilde{\mathbf{m}}$ achieves its maxima at the points $\tau_h^+ = 2h\omega_\vartheta$, $h \in \mathbb{Z}$, and its minima at $\tau_h^- = (2h+1)\omega_\vartheta$, $h \in \mathbb{Z}$. On the other hand, $\tilde{\mathbf{p}}$ achieves its maxima at $\tau_h^- = (2h+1)\omega_\vartheta$, $h \in \mathbb{Z}$, and its minima at $\tau_h^+ = 2h\omega_\vartheta$, $h \in \mathbb{Z}$.

Taking into account that

$$\tilde{\varphi}_\vartheta(\omega_\vartheta) = e_2(\vartheta), \quad \tilde{\varphi}_\vartheta(0) = -e_1(\vartheta) - e_2(\vartheta),$$

it follows that

$$\begin{aligned}(3.4) \quad \tilde{\mathbf{m}}(0) &= \frac{3 - 3\mathcal{H}^2\Lambda + 4\cos\vartheta + 2\cos 2\vartheta}{24\mathcal{H}^2\pi}, \\ \tilde{\mathbf{p}}(0) &= \frac{9\mathcal{H}^2\Lambda - 2(1 + 2\cos\vartheta)^2\cos\vartheta}{72\mathcal{H}^2\pi},\end{aligned}$$

and that

$$\begin{aligned}(3.5) \quad \tilde{\mathbf{m}}(\omega_\vartheta) &= \frac{1}{24\mathcal{H}^2\pi} \left((1 - \cos\vartheta - \sqrt{3}\sin\vartheta)^2 - 3\mathcal{H}^2\Lambda \right), \\ \tilde{\mathbf{p}}(\omega_\vartheta) &= \frac{1}{72\mathcal{H}^2\pi} (9\mathcal{H}^2\Lambda - 4 + 4\cos\vartheta + 2\cos 2\vartheta - 2\cos 3\vartheta \\ &\quad + 4\sqrt{3}(1 - \cos\vartheta)\sin\vartheta).\end{aligned}$$

From (3.5), it follows that

$$\min(\tilde{\mathbf{m}}) = \frac{1}{24\mathcal{H}^2\pi} \left((1 - \cos\vartheta - \sqrt{3}\sin\vartheta)^2 - 3\Lambda\mathcal{H}^2 \right).$$

This implies that $\tilde{\mathbf{m}} \geq 0$ if and only if $\Lambda \leq f_1(\vartheta, \mathcal{H})$, as claimed.

The function $\tilde{\mathbf{m}} + \tilde{\mathbf{p}}$ is real analytic and periodic, with period $2\omega_\vartheta$, and attains its minimum at the points τ_h^- . In particular, we compute

$$\min(\tilde{\mathbf{m}} + \tilde{\mathbf{p}}) = \frac{1}{18\mathcal{H}^2\pi} \sin^2\left(\frac{\vartheta}{2}\right) (5 + 5\cos\vartheta + 2\cos 2\vartheta - \sqrt{3}\sin\vartheta).$$

Now, the right-hand-side of the previous equation is strictly positive for every $\vartheta \in (-\pi/3, 0)$, which implies that the weak energy condition is automatically satisfied. Similarly, we have

$$\min(\tilde{\mathbf{m}} - \tilde{\mathbf{p}}) = -\frac{\Lambda}{4\pi} + \frac{\sin^2(\vartheta/2)}{18\mathcal{H}^2\pi} (7 + \cos\vartheta - 2\cos 2\vartheta - 5\sqrt{3}\sin\vartheta).$$

Thus $\min(\tilde{\mathbf{m}} - \tilde{\mathbf{p}}) \geq 0$ if and only if $\Lambda \leq f_2(\vartheta, \mathcal{H})$. Since $f_2(\vartheta, \mathcal{H}) < f_1(\vartheta, \mathcal{H})$, for each $\mathcal{H} < 0$ and for each $\vartheta \in (-\pi/3, 0)$, it follows that the dominant energy condition is satisfied if and only if $\Lambda \leq f_2(\vartheta, \mathcal{H})$.

The function $\tilde{\mathfrak{p}} + \tilde{\mathfrak{m}}/3$ is real analytic and periodic, with period $2\omega_\vartheta$, and attains its minimum at the points τ_h^+ . From (3.4), we have

$$\min(\tilde{\mathfrak{p}} + \tilde{\mathfrak{m}}/3) = \frac{\Lambda}{12\pi} + \frac{1}{72\mathcal{H}^2\pi}(1 - 2\cos\vartheta)(1 + 2\cos\vartheta)^2.$$

Thus, the strong energy condition is satisfied if and only if $\vartheta \in (-\pi/3, \vartheta_I]$ and $f_3(\vartheta, \mathcal{H}) \leq \Lambda \leq f_2(\vartheta, \mathcal{H})$. This concludes the proof. \square

3.2. Critical models of Class I with positive \mathcal{H} . With the notation introduced above, let \hat{f}_1 , \hat{f}_2 and \hat{f}_3 be given by

$$\begin{aligned}\hat{f}_1(\vartheta, \mathcal{H}) &= \frac{1}{3\mathcal{H}^2}(1 - \cos\vartheta + \sqrt{3}\sin\vartheta)^2, \\ \hat{f}_2(\vartheta, \mathcal{H}) &= \frac{2}{9\mathcal{H}^2}\sin^2\left(\frac{\vartheta}{2}\right)\left(7 + \cos\vartheta - 2\cos 2\vartheta + 5\sqrt{3}\sin\vartheta\right), \\ \hat{f}_3(\vartheta, \mathcal{H}) &= \frac{1}{6\mathcal{H}^2}\left(1 - 2\cos\vartheta - \cos 2\vartheta + 2\cos 3\vartheta + 2\sqrt{3}\sin\vartheta - \sqrt{3}\sin 2\vartheta\right).\end{aligned}$$

Note that

$$\hat{f}_3(\vartheta, \mathcal{H}) < 0 < \hat{f}_2(\vartheta, \mathcal{H}) < \hat{f}_1(\vartheta, \mathcal{H}), \quad \forall (\mathcal{H}, \vartheta) \in (0, +\infty) \times (-\pi/3, 0).$$

We can prove the following.

Theorem 4. *Let $\mathcal{M}(\mathfrak{s})$ be an \mathfrak{m} -critical RW space time of Class I, with cosmological constant Λ , angular parameter $\vartheta \in (-\pi/3, 0)$, and positive \mathcal{H} . Let $(\widetilde{\mathcal{M}}_{\mathcal{H},\vartheta}, \ell_{\mathcal{H},\vartheta})$ denote its conformal model. Then,*

$$\Lambda \leq \hat{f}_1(\vartheta, \mathcal{H}), \quad \widetilde{\mathcal{M}}_{\mathcal{H},\vartheta} = \mathbb{R} \times \mathbb{S}^3,$$

and $\ell_{\mathcal{H},\vartheta}$ is the cyclic quadratic form

$$(3.6) \quad \ell_{\mathcal{H},\vartheta} = \frac{9\mathcal{H}^2}{(12\hat{\varphi}_\vartheta - 1)^2}(-d\tau^2 + g).$$

The form $\ell_{\mathcal{H},\vartheta}$ vanishes along the totally umbilical spacelike hypersurfaces

$$\mathbb{S}_h^3 = \{(\tau, p) \in \widetilde{\mathcal{M}}_{\mathcal{H},\vartheta} \mid \tau = 2h\omega_\vartheta\},$$

$h \in \mathbb{Z}$, and is nondegenerate and of Lorentz type on the complement of $\bigcup_{h \in \mathbb{Z}} \mathbb{S}_h^3$. The weak energy condition is automatically satisfied. The dominant energy condition is satisfied if and only if $\Lambda \leq \hat{f}_2(\vartheta, \mathcal{H})$. The strong energy condition is satisfied if and only if $\hat{f}_3(\vartheta, \mathcal{H}) \leq \Lambda \leq \hat{f}_2(\vartheta, \mathcal{H})$.

Proof. Arguing as in the first part of the proof of Theorem 3, one sees that the function Q must coincide with $\hat{\varphi}_\vartheta$, and hence the conformal scale function

$$S(\tau) = \frac{3\mathcal{H}}{12\hat{\varphi}_\vartheta(\tau) - 1}.$$

Then, S is nonnegative, periodic, with minimal period $2\omega_\vartheta$, and has zeroes of second order at the points $2h\omega_\vartheta$, $h \in \mathbb{Z}$. This implies that $\ell_{\mathcal{H},\vartheta}$ is as in (3.6). Hence, $\ell_{\mathcal{H},\vartheta}$ is defined on $\widetilde{\mathcal{M}}_{\mathcal{H},\vartheta} = \mathbb{R} \times \mathbb{S}^3$, is nondegenerate and of Lorentz type on the strips

$(2h\omega_\vartheta, 2(h+1)\omega_\vartheta) \times \mathbb{S}^3$, $h \in \mathbb{Z}$, and vanishes when $\tau = 2h\omega_\vartheta$. Proceeding as in the proof of the previous theorem, one deduces that

$$(3.7) \quad \begin{aligned} \tilde{\mathfrak{m}} &= \frac{1}{8\pi} \left(\frac{(1 - 12\hat{\wp}_\vartheta)^2(5 + 12\hat{\wp}_\vartheta)}{9\mathcal{H}^2} - \mathfrak{a} - \Lambda \right), \\ \tilde{\mathfrak{p}} &= \frac{1}{4\pi} \left(\frac{\mathfrak{a} + \Lambda}{2} - \frac{(12\hat{\wp}_\vartheta - 1)^2}{9\mathcal{H}^2} \right), \end{aligned}$$

where \mathfrak{a} is as in (3.2). Thus, $\tilde{\mathfrak{m}}$ and $\tilde{\mathfrak{p}}$ are periodic function of period $2\omega_\vartheta$, real analytic on the intervals $(2h\omega_\vartheta, 2(h+1)\omega_\vartheta)$, with poles of finite order at $2h\omega_\vartheta$, $h \in \mathbb{Z}$.

The value of $\tilde{\mathfrak{m}}$ is minimum at $\tau_h = (2h+1)\omega_\vartheta$, $h \in \mathbb{Z}$. Since $\hat{\wp}_\vartheta(\omega_\vartheta) = e_1(\vartheta)$, we have

$$\min(\tilde{\mathfrak{m}}) = \frac{1}{24\pi\mathcal{H}^2} \left(-3\mathcal{H}^2\Lambda + \left(1 - \cos \vartheta + \sqrt{3} \sin \vartheta \right)^2 \right).$$

Thus $\tilde{\mathfrak{m}} \geq 0$ if and only if $\Lambda \leq \hat{f}_1(\vartheta, \mathcal{H})$. Similarly, $\tilde{\mathfrak{m}} + \tilde{\mathfrak{p}}$, $\tilde{\mathfrak{m}} - \tilde{\mathfrak{p}}$, and $\tilde{\mathfrak{m}}/3 + \tilde{\mathfrak{p}}$ are periodic with period $2\omega_\vartheta$, real analytic on $(2h\omega_\vartheta, 2(h+1)\omega_\vartheta)$, and have poles of finite order when $\tau = 2h\omega_\vartheta$. They achieve their minima at $\tau_h = (2h+1)\omega_\vartheta$, $h \in \mathbb{Z}$. This implies that

$$\min(\tilde{\mathfrak{m}} + \tilde{\mathfrak{p}}) = \frac{\sin^2(\vartheta/2)}{18\pi\mathcal{H}^2} \left(5 + 5 \cos \vartheta + 2 \cos 2\vartheta + \sqrt{3} \sin \vartheta \right).$$

Since the right-hand-side of the previous equation is positive, for each $\mathcal{H} > 0$ and for each $\vartheta \in (-\pi/3, 0)$, the weak energy condition is automatically satisfied. Similarly, we have

$$\min(\tilde{\mathfrak{m}} - \tilde{\mathfrak{p}}) = \frac{\sin^2(\vartheta/2)}{18\pi\mathcal{H}^2} \left(7 + \cos \vartheta + 2 \cos 2\vartheta + 5\sqrt{3} \sin \vartheta \right) - \frac{\Lambda}{4\pi}.$$

Therefore, $\tilde{\mathfrak{m}} - \tilde{\mathfrak{p}}$ is a nonnegative function if and only if $\Lambda \leq \hat{f}_2(\vartheta, \mathcal{H})$. On the other hand, $\hat{f}_2(\vartheta, \mathcal{H}) < \hat{f}_1(\vartheta, \mathcal{H})$, for every \mathcal{H} and for every $\vartheta \in (-\pi/3, 0)$. This implies that the dominant energy condition is satisfied if and only if $\Lambda \leq \hat{f}_2(\vartheta, \mathcal{H})$. Finally, the minimum of $\tilde{\mathfrak{m}}/3 + \tilde{\mathfrak{p}}$ is

$$\begin{aligned} \min(\tilde{\mathfrak{m}}/3 + \tilde{\mathfrak{p}}) &= \frac{1}{72\pi\mathcal{H}^2} \left(-1 + 2 \cos \vartheta + \cos 2\vartheta - 2 \cos 3\vartheta \right. \\ &\quad \left. - 2\sqrt{3} \sin \vartheta + \sqrt{3} \sin 2\vartheta \right) + \frac{\Lambda}{12\pi}. \end{aligned}$$

Hence, $\tilde{\mathfrak{m}}/3 + \tilde{\mathfrak{p}}$ is nonnegative if and only if $\hat{f}_3(\vartheta, \mathcal{H}) \leq \Lambda$. The strong energy condition is satisfied if and only if $\hat{f}_3(\vartheta, \mathcal{H}) \leq \Lambda \leq \hat{f}_2(\vartheta, \mathcal{H})$. \square

4. CRITICAL RW UNIVERSES OF CLASS II

This section discusses the \mathfrak{m} -critical RW universes of Class II. Two possible types are distinguished: negative and positive types. Theorem 5 provides a complete description of the critical models of negative type. Theorem 6 deals with the critical models of positive type with $\mathcal{H} > 0$, while Theorem 7 describes the main features of the critical models of positive type with $\mathcal{H} < 0$.

If $\Delta < 0$, the cubic polynomial $P(x)$ has two complex conjugate roots, $e_1 + ie_2$, $e_1 - ie_2$, $e_2 > 0$, and a real root, $-2e_1$, such that $144e_1^2 - 48e_2^2 = 1$. Consequently,

we can write

$$(4.1) \quad e_1 = \frac{\varepsilon}{12} \cosh \vartheta, \quad e_2 = \frac{1}{4\sqrt{3}} \sinh \vartheta, \quad \vartheta > 0, \quad \varepsilon = \pm 1.$$

If $\varepsilon = 1$, we say that the \mathfrak{m} -critical RW spacetime is of *positive type*, while, if $\varepsilon = -1$, we say that it is of *negative type*. We call $\vartheta > 0$ the *angular parameter*. The invariant g_3 and the parameter \mathfrak{a} can be written as

$$(4.2) \quad g_3 = -\frac{\varepsilon \cosh 3\vartheta}{216}, \quad \mathfrak{a} = \frac{2}{9\mathcal{H}^2} (1 - \varepsilon \cosh 3\vartheta).$$

Let $\wp_{\vartheta,\varepsilon}$ denote the Weierstrass \wp -function with invariants $g_2 = 1/12$ and g_3 . Let $\omega_{\vartheta,\varepsilon}$ be its real half-period. Then, $\wp_{\vartheta,\varepsilon}$ has the unique real form $\widehat{\wp}_{\vartheta,\varepsilon}(\tau) = \wp_{\vartheta,\varepsilon}(\tau)$, for every $\tau \in \mathbb{R}$. The function $\widehat{\wp}_{\vartheta,\varepsilon}$ is periodic, with minimal period $2\omega_{\vartheta,\varepsilon}$, is real analytic on the open intervals $(2h\omega_{\vartheta,\varepsilon}, 2(h+1)\omega_{\vartheta,\varepsilon})$, $h \in \mathbb{Z}$, and possesses double poles at the points $2h\omega_{\vartheta,\varepsilon}$, $h \in \mathbb{Z}$. Moreover, $\widehat{\wp}_{\vartheta,\varepsilon}(\tau) \geq -2e_1(\vartheta, \varepsilon)$, for all $\tau \in \mathbb{R}$, and $\widehat{\wp}_{\vartheta,\varepsilon}(\tau) = -2e_1(\vartheta, \varepsilon)$ if and only if $\tau = \tau_h = (2h+1)\omega_{\vartheta,\varepsilon}$, $h \in \mathbb{Z}$.

4.1. Critical models of Class II and negative type. With the notation above, let \widehat{h}_1 , \widehat{h}_2 and \widehat{h}_3 be defined by

$$\begin{aligned} \widehat{h}_1(\vartheta, \mathcal{H}) &= \frac{1}{3\mathcal{H}^2} (1 - 2 \cosh \vartheta)^2, \\ \widehat{h}_2(\vartheta, \mathcal{H}) &= \frac{1}{18\mathcal{H}^2} (3 - 2 \cosh \vartheta) (1 - 2 \cosh \vartheta)^2, \\ \widehat{h}_3(\vartheta, \mathcal{H}) &= -\frac{1}{6\mathcal{H}^2} (1 + 2 \cosh \vartheta) (1 - 2 \cosh \vartheta)^2. \end{aligned}$$

Note that $\widehat{h}_3(\vartheta, \mathcal{H}) < \widehat{h}_2(\vartheta, \mathcal{H}) < \widehat{h}_1(\vartheta, \mathcal{H})$, for every \mathcal{H} and for every $\vartheta > 0$.

We can prove the following.

Theorem 5. *Let $\mathcal{M}(\mathfrak{s})$ be an \mathfrak{m} -critical RW spacetime of Class II and negative type, with cosmological constant Λ , and angular parameter $\vartheta > 0$. (In this case, \mathcal{H} is necessarily positive.) Let $(\widetilde{\mathcal{M}}_{\mathcal{H},\vartheta}, \ell_{\mathcal{H},\vartheta})$ denote its conformal model. Then,*

$$\Lambda \leq \widehat{h}_1(\vartheta, \mathcal{H}), \quad \widetilde{\mathcal{M}}_{\mathcal{H},\vartheta} = \mathbb{R} \times \mathbb{S}^3,$$

and $\ell_{\mathcal{H},\vartheta}$ is the cyclic quadratic form

$$(4.3) \quad \ell_{\mathcal{H},\vartheta} = \frac{9\mathcal{H}^2}{(12\widehat{\wp}_{\vartheta,-1} - 1)^2} (-d\tau^2 + g).$$

The form $\ell_{\mathcal{H},\vartheta}$ vanishes along the totally umbilical spacelike hypersurfaces

$$\mathbb{S}_h^3 = \{(\tau, p) \in \widetilde{\mathcal{M}}_{\mathcal{H},\vartheta} \mid \tau = 2h\omega_{\vartheta,-1}\},$$

$h \in \mathbb{Z}$, and is nondegenerate and of Lorentz type on the complement of $\bigcup_{h \in \mathbb{Z}} \mathbb{S}_h^3$. The weak energy condition is automatically satisfied. The dominant energy condition is satisfied if and only if $\Lambda \leq \widehat{h}_2(\vartheta, \mathcal{H})$. The strong energy condition is satisfied if and only if $\widehat{h}_3(\vartheta, \mathcal{H}) \leq \Lambda \leq \widehat{h}_2(\vartheta, \mathcal{H})$.

Proof. The conformal scale function S is given by

$$S(\tau) = \frac{3\mathcal{H}}{12\widehat{\wp}_{\vartheta,-1}(\tau) - 1}.$$

Since $\widehat{\wp}_{\vartheta,-1} \geq -2e_1(\vartheta, \mathcal{H}) > 1/12$ and $S > 0$, then $\mathcal{H} > 0$. By construction, S is periodic, with minimal period $2\omega_{\vartheta,-1}$ and has zeroes of order 2 located at the points

$2h\omega_{\vartheta,-1}$, $h \in \mathbb{Z}$. This implies that $\ell_{\mathcal{H},\vartheta}$ is as in (3.6). Then, $\ell_{\mathcal{H},\vartheta}$ is smooth on $\widetilde{\mathcal{M}}_{\mathcal{H},\vartheta} = \mathbb{R} \times \mathbb{S}^3$, is nondegenerate and of Lorentz type on $(2h\omega_{\vartheta,-1}, 2(h+1)\omega_{\vartheta,-1}) \times \mathbb{S}^3$, $h \in \mathbb{Z}$, and vanishes at $2h\omega_{\vartheta,-1}$. The functions $\widetilde{\mathfrak{m}}$ and $\widetilde{\mathfrak{p}}$ are as in (3.7) and \mathfrak{a} is as in (4.2). Thus, $\widetilde{\mathfrak{m}}$ and $\widetilde{\mathfrak{p}}$ are periodic function of period $2\omega_{\vartheta,-1}$, real analytic on $(2h\omega_{\vartheta,-1}, 2(h+1)\omega_{\vartheta,-1})$, and with poles of finite order at $2h\omega_{\vartheta,-1}$. The function $\widetilde{\mathfrak{m}}$ achieves its minimum at $\tau_h = (2h+1)\omega_{\vartheta,-1}$, $h \in \mathbb{Z}$. Taking into account that

$$\widehat{\varphi}_{\vartheta,-1}(\omega_{\vartheta,-1}) = -2e_1(\vartheta) = \frac{1}{6} \cosh \vartheta$$

we have

$$\min(\widetilde{\mathfrak{m}}) = \frac{1}{24\pi\mathcal{H}^2}(-3\mathcal{H}^2\Lambda + (1 - 2\cosh \vartheta)^2).$$

Thus, $\widetilde{\mathfrak{m}}$ is nonnegative if and only if $\Lambda \leq \widehat{h}_1(\vartheta, \mathcal{H})$. The functions $\widetilde{\mathfrak{m}} + \widetilde{\mathfrak{p}}$, $\widetilde{\mathfrak{m}} - \widetilde{\mathfrak{p}}$, and $\widetilde{\mathfrak{m}}/3 + \widetilde{\mathfrak{p}}$ are periodic, with period $2\omega_{\vartheta,-1}$, and attain their minima at the points $\tau_h = (2h+1)\omega_{\vartheta,-1}$, $h \in \mathbb{Z}$. Then

$$\begin{aligned} \min(\widetilde{\mathfrak{m}} + \widetilde{\mathfrak{p}}) &= \frac{1}{72\pi\mathcal{H}^2} (1 - 2\cosh \vartheta)^2 (3 + 2\cosh \vartheta) > 0, \\ \min(\widetilde{\mathfrak{m}} - \widetilde{\mathfrak{p}}) &= -\frac{\Lambda}{4\pi} + \frac{1}{72\pi\mathcal{H}^2} (1 - 2\cosh \vartheta)^2 (3 - 2\cosh \vartheta), \\ \min(\widetilde{\mathfrak{m}}/3 + \widetilde{\mathfrak{p}}) &= \frac{\Lambda}{12\pi} + \frac{1}{72\pi\mathcal{H}^2} (1 - 2\cosh \vartheta)^2 (1 + 2\cosh \vartheta). \end{aligned}$$

These formulae imply that the weak energy condition is automatically satisfied, that the dominant energy condition is satisfied if and only if $\Lambda \leq \widehat{h}_2(\vartheta, \mathcal{H})$, and that the strong energy condition is satisfied if and only if $\widehat{h}_3(\vartheta, \mathcal{H}) \leq \Lambda \leq \widehat{h}_2(\vartheta, \mathcal{H})$. \square

4.2. Critical models of Class II and positive type. Retaining the notation introduced at the beginning of the section, we now consider the \mathfrak{m} -critical RW universes of Class II and positive type. In this case, the real root is

$$-2e_1(\vartheta) = -\frac{1}{6} \cosh \vartheta < \frac{1}{12}.$$

Thus, there exists a unique $\tau^* \in (0, \omega_{\vartheta,1})$, such that $\widehat{\varphi}_{\vartheta,1}(\tau^*) = 1/12$. The zeroes of the equation $\widehat{\varphi}_{\vartheta,1} = 1/12$ are

$$\tau_h^* = \tau^* + 2h\omega_{\vartheta,1}, \quad \widehat{\tau}_h^* = -\tau^* + 2h\omega_{\vartheta,1}, \quad h \in \mathbb{Z}.$$

If $\mathcal{H} > 0$, the conformal scale function S is positive on the intervals $I'_h = (\widehat{\tau}_h^*, \tau_h^*)$, $h \in \mathbb{Z}$, and is negative on the intervals $I''_h = (\tau_h^*, \widehat{\tau}_{h+1}^*)$, $h \in \mathbb{Z}$. On the other hand, if $\mathcal{H} < 0$, the conformal scale function S is positive on the intervals $I''_h = (\tau_h^*, \widehat{\tau}_{h+1}^*)$, $h \in \mathbb{Z}$, and is negative on the intervals $I'_h = (\widehat{\tau}_h^*, \tau_h^*)$, $h \in \mathbb{Z}$.

Taking into account that S is by definition a nonnegative function and that $\widehat{\varphi}_{\vartheta,1}$ is periodic with period $2\omega_{\vartheta,1}$, we may consider $I' := I'_0$ or $I'' := I''_0$ as the maximal intervals of definition for S , depending on whether \mathcal{H} is positive or negative.

For the case in which \mathcal{H} is positive, we have the following.

Theorem 6. *Let $\mathcal{M}(\mathfrak{s})$ be an \mathfrak{m} -critical RW space time of Class II and positive type, with cosmological constant Λ , angular parameter $\vartheta > 0$, and positive internal parameter \mathcal{H} . Let $(\widetilde{\mathcal{M}}_{\mathcal{H},\vartheta}, \ell_{\mathcal{H},\vartheta})$ denote its conformal model. Then,*

$$\Lambda \leq \frac{2\cosh 3\vartheta - 2}{9\mathcal{H}^2}, \quad \widetilde{\mathcal{M}}_{\mathcal{H},\vartheta} = I' \times \mathbb{S}^3,$$

and $\ell_{\mathcal{H},\vartheta}$ is the non-cyclic quadratic form

$$(4.4) \quad \ell_{\mathcal{H},\vartheta} = \frac{9\mathcal{H}^2}{(12\widehat{\varphi}_{\vartheta,1} - 1)^2}(-d\tau^2 + g).$$

The form $\ell_{\mathcal{H},\vartheta}$ vanishes along the totally umbilical spacelike hypersurface

$$\mathbb{S}_0^3 = \left\{ (\tau, p) \in \widetilde{\mathcal{M}}_{\mathcal{H},\vartheta} \mid \tau = 0 \right\}$$

and is nondegenerate of Lorentzian signature on the complement of \mathbb{S}_0^3 . The weak and the dominant energy conditions are automatically satisfied. The strong energy condition is satisfied if and only if $\Lambda = 2(\cosh 3\vartheta - 1)/9\mathcal{H}^2$.

Proof. Let $I' = I'_0$ be the maximal domain of definition of S . Since $\mathcal{H} > 0$, we have that $\widehat{\varphi}_{\vartheta,1}(\tau) > 1/12$, for each $\tau \in I'$. This implies that $\widetilde{\mathcal{M}}_{\mathcal{H},\vartheta} = I' \times \mathbb{S}^3$ and that the quadratic form $\ell_{\mathcal{H},\vartheta}$ is as in (4.4). The conformal factor $9\mathcal{H}^2/(12\widehat{\varphi}_{\vartheta,1} - 1)^2$ vanishes at $\tau = 0$, is strictly positive for each $\tau \in I'$, $\tau \neq 0$, and tends to $+\infty$ when $\tau \mapsto \pm\tau^*$. The energy density $\widetilde{\mathfrak{m}}$ and the pressure $\widetilde{\mathfrak{p}}$ are as in (3.7), with $\mathfrak{a} = \frac{2}{9\mathcal{H}^2}(1 - \cosh 3\vartheta)$. The density $\widetilde{\mathfrak{m}}$ has a pole at the origin and attains its minimum at $\pm\tau^*$. Taking into account that

$$\widehat{\varphi}_{\vartheta,1}(\pm\tau^*) = \frac{1}{12}, \quad \widehat{\varphi}'_{\vartheta,1}(\pm\tau^*) = \mp \frac{1}{6\sqrt{6}}\sqrt{\cosh 3\vartheta - 1}, \quad \widehat{\varphi}''_{\vartheta,1}(\pm\tau^*) = -\frac{1}{8},$$

we obtain

$$\min(\widetilde{\mathfrak{m}}) = \frac{1}{72\pi\mathcal{H}^2} (2 \cosh 3\vartheta - 9\mathcal{H}^2\Lambda - 2).$$

Therefore, $\widetilde{\mathfrak{m}} \geq 0$ if and only if

$$\Lambda \leq \frac{2 \cosh 3\vartheta - 2}{9\mathcal{H}^2}.$$

Similarly, $\widetilde{\mathfrak{m}} + \widetilde{\mathfrak{p}}$, $\widetilde{\mathfrak{m}} - \widetilde{\mathfrak{p}}$, and $\widetilde{\mathfrak{m}}/3 + \widetilde{\mathfrak{p}}$ attain their minima at $\pm\tau^*$. In particular, $\min(\widetilde{\mathfrak{p}}) = -\min(\widetilde{\mathfrak{m}})$. Consequently, we have

$$\min(\widetilde{\mathfrak{m}} + \widetilde{\mathfrak{p}}) = 0,$$

$$\min(\widetilde{\mathfrak{m}} - \widetilde{\mathfrak{p}}) = \frac{1}{36\pi\mathcal{H}^2} (2 \cosh 3\vartheta - 9\mathcal{H}^2\Lambda - 2),$$

$$\min(\widetilde{\mathfrak{m}}/3 + \widetilde{\mathfrak{p}}) = -\frac{1}{108\pi\mathcal{H}^2} (2 \cosh 3\vartheta - 9\mathcal{H}^2\Lambda - 2).$$

This implies that the weak and the dominant energy conditions are automatically satisfied, while the strong energy condition is enforced if and only if $\Lambda = 2(\cosh(3\vartheta) - 1)/9\mathcal{H}^2$. This proves the result. \square

As for the case in which \mathcal{H} is negative, we can prove the following.

Theorem 7. *Let $\mathcal{M}(\mathfrak{s})$ be an \mathfrak{m} -critical RW space time of Class II and positive type, with cosmological constant Λ , angular parameter $\vartheta > 0$, and negative internal parameter \mathcal{H} . Let $(\widetilde{\mathcal{M}}_{\mathcal{H},\vartheta}, \ell_{\mathcal{H},\vartheta})$ denote its conformal model. Then,*

$$\Lambda \leq \min \left(\frac{2 \cosh 3\vartheta - 2}{9\mathcal{H}^2}, \frac{(1 + 2 \cosh \vartheta)^2}{3\mathcal{H}^2} \right), \quad \widetilde{\mathcal{M}}_{\mathcal{H},\vartheta} = I'' \times \mathbb{S}^3,$$

and $\ell_{\mathcal{H},\vartheta}$ is the non-cyclic Lorentz metric

$$(4.5) \quad \ell_{\mathcal{H},\vartheta} = \frac{9\mathcal{H}^2}{(12\widehat{\varphi}_{\vartheta,1} - 1)^2}(-d\tau^2 + g).$$

The weak and the dominant energy conditions are satisfied if and only if $0 < \vartheta \leq \operatorname{arcosh}(3/2)$. The strong energy condition is satisfied if and only if $0 < \vartheta \leq \operatorname{arcosh}(3/2)$ and $\Lambda = 2(\cosh 3\vartheta - 1)/9\mathcal{H}^2$.

Proof. Let $I'' = I_0'' \subset (0, 2\omega_{\vartheta,1})$ be the maximal interval of definition of S . Since $\mathcal{H} < 0$, then $\widehat{\varphi}_{\vartheta,1}(\tau) < 1/12$, for every $\tau \in I''$. The conformal energy density $\widetilde{\mathfrak{m}}$ is real analytic on $(0, 2\omega_{\vartheta,1})$. If $\vartheta \in (0, \operatorname{arcosh}(3/2))$, then $\widetilde{\mathfrak{m}}$ has relative minima at $\tau = \tau^*, \widehat{\tau}^*$, and a relative maximum at $\omega_{\vartheta,1}$. We then have

$$(4.6) \quad \min_{I''}(\widetilde{\mathfrak{m}}) = \widetilde{\mathfrak{m}}(\tau^*) = \widetilde{\mathfrak{m}}(\widehat{\tau}^*) = \frac{2(\cosh 3\vartheta - 1) - 9\mathcal{H}^2\Lambda}{72\mathcal{H}^2\pi}.$$

Therefore, if $\vartheta \in (0, \operatorname{arcosh}(3/2))$, $\widetilde{\mathfrak{m}} \geq 0$ if and only if $\Lambda \leq 2(\cosh 3\vartheta - 1)/9\mathcal{H}^2$. If $\vartheta > \operatorname{arcosh}(3/2)$, then $\widetilde{\mathfrak{m}}$ has three relative minima on $(0, 2\omega_{\vartheta,1})$, located at $\tau^*, \widehat{\tau}^*$, and $\omega_{\vartheta,1}$. The value of $\widetilde{\mathfrak{m}}$ at $\tau^*, \widehat{\tau}^*$ is as in (4.6), while

$$\widetilde{\mathfrak{m}}(\omega_{\vartheta,1}) = \frac{(1 + 2\cosh \vartheta)^2 - 3\mathcal{H}^2\Lambda}{24\mathcal{H}^2\pi}.$$

Consequently, if $\vartheta \in (\operatorname{arcosh}(3/2), \operatorname{arcosh}(5/2))$, τ^* and $\widehat{\tau}^*$ are minimum points, and

$$\min_{I''}(\widetilde{\mathfrak{m}}) = \frac{2(\cosh 3\vartheta - 1) - 9\mathcal{H}^2\Lambda}{72\mathcal{H}^2\pi}.$$

If $\vartheta = \operatorname{arcosh}(5/2)$, the values of $\widetilde{\mathfrak{m}}$ on $\tau^*, \widehat{\tau}^*$, and $\omega_{\vartheta,1}$ do coincide. If $\vartheta > \operatorname{arcosh}(5/2)$, the minimum of $\widetilde{\mathfrak{m}}$ is attained at $\omega_{\vartheta,1}$, and

$$\min_{I''}(\widetilde{\mathfrak{m}}) = \frac{(1 + 2\cosh \vartheta)^2 - 3\mathcal{H}^2\Lambda}{24\mathcal{H}^2\pi}.$$

Then, if $\vartheta \in (\operatorname{arcosh}(3/2), \operatorname{arcosh}(5/2))$, $\widetilde{\mathfrak{m}} \geq 0$ if and only if $\Lambda \leq 2(\cosh 3\vartheta - 1)/9\mathcal{H}^2$; and if $\vartheta > \operatorname{arcosh}(5/2)$, $\widetilde{\mathfrak{m}} \geq 0$ if and only if $\Lambda \leq (1 + 2\cosh \vartheta)^2/3\mathcal{H}^2$. Since $2(\cosh 3\vartheta - 1) \leq 3(1 + 2\cosh \vartheta)^2$ on $[0, \operatorname{arcosh}(5/2)]$ and $3(1 + 2\cosh \vartheta)^2 \leq 2(\cosh 3\vartheta - 1)$ on $[\operatorname{arcosh}(5/2), +\infty)$, it follows that $\widetilde{\mathfrak{m}}$ is nonnegative on I'' if and only if $\Lambda \leq \min\left(\frac{2\cosh 3\vartheta - 1}{9\mathcal{H}^2}, \frac{(1 + 2\cosh \vartheta)^2}{3\mathcal{H}^2}\right)$.

The function $\widetilde{\mathfrak{m}} + \widetilde{\mathfrak{p}}$ has relative minima at $\tau^*, \widehat{\tau}^*$, and $\omega_{\vartheta,1}$ in the interval $(0, 2\omega_{\vartheta,1})$. In particular, $\widetilde{\mathfrak{m}} + \widetilde{\mathfrak{p}}$ vanishes at τ^* and $\widehat{\tau}^*$, while its value at $\omega_{\vartheta,1}$ is

$$\frac{(3 - 2\cosh \vartheta)(1 + 2\cosh \vartheta)^2}{72\mathcal{H}^2\pi}.$$

Thus $\widetilde{\mathfrak{m}} + \widetilde{\mathfrak{p}}$ is nonnegative if and only if $\vartheta \in (0, \operatorname{arcosh}(3/2)]$. This means that the weak energy condition is satisfied if and only if $0 < \vartheta \leq \operatorname{arcosh}(3/2)$.

For $\vartheta \in (0, \operatorname{arcosh}(3/2)]$, the function $\widetilde{\mathfrak{m}} - \widetilde{\mathfrak{p}}$ has two minima at $\tau^*, \widehat{\tau}^*$, and

$$\widetilde{\mathfrak{m}}(\tau^*) - \widetilde{\mathfrak{p}}(\tau^*) = \widetilde{\mathfrak{m}}(\widehat{\tau}^*) - \widetilde{\mathfrak{p}}(\widehat{\tau}^*) = -\frac{9\mathcal{H}^2\Lambda - 2(\cosh 3\vartheta - 1)}{36\mathcal{H}^2\pi}.$$

This implies that the dominant energy condition is automatically satisfied.

For $\vartheta \in (0, \operatorname{arcosh}(3/2)]$, the minimum of $\widetilde{\mathfrak{m}}/3 + \widetilde{\mathfrak{p}}$ on the interval $(\tau^*, \widehat{\tau}^*)$ is attained at the half period $\omega_{\vartheta,1}$, and

$$\frac{\widetilde{\mathfrak{m}}(\omega_{\vartheta,1})}{3} + \widetilde{\mathfrak{p}}(\omega_{\vartheta,1}) = \frac{9\mathcal{H}^2\Lambda - 2(\cosh 3\vartheta - 1)}{108\mathcal{H}^2\pi}.$$

Accordingly, the strong energy condition is satisfied if and only if $\Lambda = 2(\cosh 3\vartheta - 1)/9\mathcal{H}^2$, as claimed. \square

REFERENCES

- [1] D. ALEKSEEVSKY, Lorentzian manifolds with transitive conformal group, *Note Mat.* **37** (2017), suppl. 1, 35–47; arXiv:1611.03447 [math.DG].
- [2] J. ANGST, Asymptotic behavior of a relativistic diffusion in Robertson–Walker space-times, *Ann. Inst. Poincaré Probab. Statist.* **52** (2016), no. 1, 376–411; arXiv:1405.0142[math.PR].
- [3] L. BANCHI AND F. CARAVELLI, Geometric phases and cyclic isotropic cosmologies, *Classical Quantum Gravity* **33** (2016), no. 10, 105003, 23 pp.; arXiv:1411.7553[gr-qc].
- [4] T. BARBOT, V. CHARETTE, T. DRUMM, W. M. GOLDMAN, AND K. MELNICK, A premier on the (2+1)-Einstein universe, in *Recent developments in pseudo-Riemannian geometry*, 179–229, ESI Lect. Math. Phys., Eur. Math. Soc., Zürich, 2008; arXiv:0706.3055[math.DG].
- [5] I. BOCHICCHIO, S. CAPOZZIELLO, AND E. LASERRA, The Weierstrass criterion and the Lemaître–Tolman–Bondi models with cosmological constant Λ , *Int. J. Geom. Methods Mod. Phys.* **8** (2011), no. 7, 1653–1666; arXiv:111.0819 [gr-qc].
- [6] K. BAMBA, K. YESMAKHANOVA, K. YERZHANOV, AND R. MYRZAKULOV, Reconstruction of the equation of state for cyclic universes in homogeneous and isotropic cosmology, *Cent. Eur. J. Phys.* **11** (2013), no. 4, 397–411; arXiv:1203.3401 [gr-qc].
- [7] K. BAMBA, A. N. MAKARENKO, A. N. MYAGKY, AND S. D. ODINTSOV, Bounce universe from string-inspired Gauss-Bonnet gravity, *J. Cosmol. Astropart. Phys.* 2015, no. 4, 001, 27 pp.; arXiv:1411.3852 [hep-th].
- [8] M. BADIALE, E. SERRA, *Semilinear Elliptic Equations for Beginners. Existence Results via the Variational Approach*, Universitext, Springer, London, 2011.
- [9] R.-G. CAI AND S. P. KIM, First law of thermodynamics and Friedmann equations of Friedmann–Robertson–Walker universe, *J. High Energy Phys.* 2005, no. 2, 050, 13 pp.
- [10] J. D’AMBROISE, Applications of elliptic and theta functions to Friedmann–Robertson–Lemaître–Walker cosmology with cosmological constant, in: *A Window into Zeta and Modular Physics*, K. Kirsten and F. L. Williams eds., MSRI Publications, **57**, Cambridge University Press (2010), 279–293; arXiv:0908.2481 [gr-qc].
- [11] J. D’AMBROISE AND F. L. WILLIAMS, Parametric solutions of certain nonlinear differential equations in cosmology, *J. Nonlinear Math. Phys.*, **18** (2011), no. 2, 269–278; arXiv:1202.4422 [gr-qc].
- [12] A. DZHALILOV, E. MUSSO, AND L. NICOLodi, Conformal geometry of timelike curves in the (1+2)-Einstein universe, *Nonlinear Anal.*, **143** (2016), 224–255.
- [13] A. EINSTEIN, Kosmologische Betrachtungen zur allgemeinen Relativitätstheorie, *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*, 142–152, Berlin, 1917.
- [14] A. FERRÁNDEZ, A. GIMÉNEZ, AND P. LUCAS, Geometrical particle models on 3D null curves, *Phys. Lett. B* **543** (2002), no. 3-4, 311–317; hep-th/0205284.
- [15] C. FRANCES, Sur les variétés lorentziennes dont le group conforme est essentiel, *Math. Ann.* **332** (2005), no. 1, 103–119.
- [16] A. FRIEDMANN, Über die Möglichkeit einer Welt mit konstanter negativer Krümmung des Raumes, *Zeitschrift für Physik A*, **21** (1924) (1), 326–332.
- [17] J. D. GRANT AND E. MUSSO, Coisotropic variational problems, *J. Geom. Phys.* **50** (2004), 303–338.
- [18] G. W. GIBBONS AND M. VYSKA, The application of Weierstrass elliptic functions to Schwarzschild null geodesics, *Classical Quantum Gravity* **29** (2012), no. 6, 065016, 19 pp.; arXiv:1110.6508 [gr-qc].
- [19] S. W. HAWKING AND G. F. R. ELLIS, *The Large Scale Structure of Space-time*, Cambridge University Press, Cambridge, UK, 1973.
- [20] Y. A. KUZNETSOV AND M. S. PLYUSHCHAY, (2+1)-dimensional models of relativistic particles with curvature and torsion, *J. Math. Phys.* **35** (1994), no. 6, 2772–2778.
- [21] L. D. LANDAU AND E. M. LIFSHITZ, *Course of theoretical physics. Vol. 1. Mechanics*, third edition, Pergamon Press, Oxford-New York-Toronto, Ont., 1976.
- [22] D. F. LAW DEN, *Elliptic functions and applications*, Series in Applied Mathematical Science, **80**, Springer-Verlag, New York, 1989.
- [23] G. LEMAITRE, Expansion of the universe, A homogeneous universe of constant mass and increasing radius accounting for the radial velocity of extra-galactic nebulae, *Monthly Notices of the Royal Astronomical Society*, **91** (1931), 483–490.

- [24] E. MUSSO AND L. NICOLodi, Closed trajectories of a particle model on null curves in anti-de Sitter 3-space, *Classical Quantum Gravity* **24** (2007), no. 22, 5401–5411.
- [25] E. MUSSO AND L. NICOLodi, Reduction for constrained variational problems on 3-dimensional null curves, *SIAM J. Control Optim.* **47** (2008), no. 3, 1399–1414.
- [26] V. V. NESTERENKO, A. FEOLI, AND G. SCARPETTA, Complete integrability for Lagrangians dependent on acceleration in a spacetime of constant curvature, *Classical Quantum Gravity* **13** (1996), 1201–1211.
- [27] A. NERSESSIAN, R. MANVELYAN, AND H. J. W. MÜLLER-KIRSTEN, Particle with torsion on 3d null-curves, *Nuclear Phys. B* **88** (2000), 381–384; hep-th/9912061.
- [28] A. NERSESSIAN AND E. RAMOS, Massive spinning particles and the geometry of null curves, *Phys. Lett. B* **445** (1998), 123–128.
- [29] B. O’NEILL, *Semi-Riemannian Geometry with Applications to Relativity*, Pure and Applied Mathematics, 103, Academic Press, Inc., New York, 1983.
- [30] R. PENROSE, *Cycles of time: An extraordinary new view of the universe*, Alfred A. Knopf, Inc., New York, 2010.
- [31] R. PENROSE, On the gravitization of quantum mechanics 2: Conformal cyclic cosmology, *Found. Phys.* **44** (2014), 873–890.
- [32] H. P. ROBERTSON, Kinematics and world-structure, *Astrophys. J.* **82** (1935), 284–301.
- [33] H. C. ROSU AND P. OJEDA-MAY, Supersymmetry of FRW barotropic cosmologies, *International Journal of Theoretical Physics*, **45**, n.6, 873–890 (2006); arXiv:0510004 [gr-qc].
- [34] D. W. TIAN AND I. BOOTH, Apparent horizon and gravitational thermodynamics of the Universe: solutions to the temperature and entropy confusion, and extension to modified gravity, *Physical Review D*, **92**, n.2, (2015), arXiv:1411.6547 [gr-qc]
- [35] P. TOD, Penrose’s Weyl curvature hypothesis and conformally-cyclic cosmology, *Journal of Physics: Conference Series* **229** (2010), 1–5.
- [36] A. G. WALKER, On Milne’s theory of world-structure, *Proc. London Math. Soc.* **42** (1937), 90–127.
- [37] R. M. WALD, *General Relativity*, University of Chicago Press, Chicago, IL, 1984.
- [38] S. WEINBERG, *Gravitation and Cosmology*, John Wiley & Sons, New York, 1972.
- [39] T. ZHU, J.-R. REN, AND M.-F. LI, Corrected Entropy of Friedmann–Robertson–Walker Universe in Tunneling Method, *J. Cosmol. Astropart. Phys.* 2009; arXiv:0905.1838 [hep-th]

(O. ESHKOBILOV) DIPARTIMENTO DI MATEMATICA “GIUSEPPE PEANO”, UNIVERSITÀ DI TORINO, VIA CARLO ALBERTO 10, I-10123 TORINO, ITALY

E-mail address: olimjon.eshkobilov@edu.unito.it, olimjon.eshkobilov@polito.it

(E. MUSSO) DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI 24, I-10129 TORINO, ITALY

E-mail address: emilio.musso@polito.it

(L. NICOLodi) DIPARTIMENTO DI SCIENZE MATEMATICHE, FISICHE E INFORMATICHE, UNIVERSITÀ DI PARMA, PARCO AREA DELLE SCIENZE 53/A, I-43124 PARMA, ITALY

E-mail address: lorenzo.nicolodi@unipr.it