

Lipoplexes for miRNA delivery in myocardial regeneration

*Original*

Lipoplexes for miRNA delivery in myocardial regeneration / Nicoletti, Letizia; Tarricone, Giulia; Paoletti, Camilla; Divieto, Carla; Andreana, Ilaria; Stella, Barbara; Arpicco, Silvia; Mattu, Clara; Chiono, Valeria. - ELETTRONICO. - (2019), pp. 136-136. (Intervento presentato al convegno Nano-Day IV tenutosi a Milan nel December, 11-14).

*Availability:*

This version is available at: 11583/2776917 since: 2020-01-03T15:13:11Z

*Publisher:*

Cinsa and University of Parma

*Published*

DOI:

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

## Lipoplexes for miRNA delivery in myocardial regeneration

Letizia Nicoletti<sup>1</sup>, Giulia Tarricone<sup>1</sup>, Camilla Paoletti<sup>1</sup>, Carla Divieto<sup>2</sup>, Ilaria Andreana<sup>3</sup>,  
Barbara Stella<sup>3</sup>, Silvia Arpicco<sup>3</sup>, Clara Mattu<sup>1</sup>, Valeria Chiono<sup>1\*</sup>

<sup>1</sup>*Department of Mechanical and Aerospace Engineering, Politecnico di Torino, Corso Duca degli Abruzzi 24, Turin, Italy*

<sup>2</sup>*Division of Metrology for Quality of Life, Istituto Nazionale di Ricerca Metrologica, Strada delle Cacce 91, Turin, Italy*

<sup>3</sup>*Department of Drug Science and Technology, University of Turin, via P. Giuria 9, 10125, Turin, Italy*

\**valeria.chiono@polito.it*

Myocardial infarction is one of the major causes of mortality in industrialized countries. Several strategies have been studied to regenerate myocardial infarction, reaching a limited success [1].

Direct reprogramming of cardiac fibroblasts populating post-infarct scar into cardiomyocytes could represent an alternative promising approach [1-3]. In 2012, mouse fibroblasts transfected with a combination of four microRNAs (miRNAs), termed “miRcombo”, were transdifferentiated into cardiomyocytes [1-3].

In this work, nanotechnology-based approaches were explored to efficiently deliver miRNAs to human cardiac fibroblasts in the perspective of their direct reprogramming. Lipoplexes containing negmiR or miR-1 (miRVana™ miRNA Mimic, Life Technologies) were prepared at different N:P ratios (3.0; 1.75; 0.70; 0.35). Lipoplexes showed an average size between 405 nm and 560 nm and an average zeta potential between +21 mV and -29 mV. Encapsulation efficiency and morphology (cryoTEM) were also analyzed.

In vitro tests with human cardiac fibroblasts showed a significantly higher miR-1 expression in the case of transfection with lipoplexes compared to a commercial agent, as evaluated by digital droplet PCR.

As a conclusion, in this work new lipoplexes were developed showing efficient delivery of miR-1 to human cardiac fibroblasts, for future use in direct reprogramming.

BIORECAR project is acknowledged: it has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme grant agreement No 772168

[1] C. Paoletti et al. *Cells* 2018;7: 114.

[2] T. M. Jayawardena et al. *Circ. Res.* 2012; 110: 1465.

[3] T. M. Jayawardena et al. *Circ. Res.* 2015; 116: 418.