

Measuring Progress towards a Circular Economy: A Monitoring Framework for Economy-wide Material Loop Closing in the EU28

Original

Measuring Progress towards a Circular Economy: A Monitoring Framework for Economy-wide Material Loop Closing in the EU28 / Mayer, A.; Haas, W.; Wiedenhofer, D.; Krausmann, F.; Nuss, P.; Blengini, G. A.. - In: JOURNAL OF INDUSTRIAL ECOLOGY. - ISSN 1088-1980. - ELETTRONICO. - 23:1(2019), pp. 62-76. [10.1111/jiec.12809]

Availability:

This version is available at: 11583/2813064 since: 2020-04-17T14:30:38Z

Publisher:

Blackwell Publishing

Published

DOI:10.1111/jiec.12809

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Measuring Progress towards a Circular Economy

A Monitoring Framework for Economy-wide Material Loop Closing in the EU28

Andreas Mayer ¹, Willi Haas ¹, Dominik Wiedenhofer ¹, Fridolin Krausmann ¹, Philip Nuss ² and Gian Andrea Blengini ³


¹Institute of Social Ecology (SEC), Department of Economics and Social Sciences, University of Natural Resources and Life Sciences (BOKU), Vienna, Austria

²German Environment Agency (UBA), Dessau-Roßlau, Germany

³European Commission Directorate-General Joint Research Centre Sustainable Resources Directorate, Ispra, Italy

Keywords:

circular economy
environmental pressures
industrial ecology
recycling
secondary materials
waste management

 Supporting information is linked to this article on the JIE website



For information about the data openness badges, see bit.ly/JIE_data_trans

Summary

The concept of a circular economy (CE) is gaining increasing attention from policy makers, industry, and academia. There is a rapidly evolving debate on definitions, limitations, the contribution to a wider sustainability agenda, and a need for indicators to assess the effectiveness of circular economy measures at larger scales. Herein, we present a framework for a comprehensive and economy-wide biophysical assessment of a CE, utilizing and systematically linking official statistics on resource extraction and use and waste flows in a mass-balanced approach. This framework builds on the widely applied framework of economy-wide material flow accounting and expands it by integrating waste flows, recycling, and downcycled materials. We propose a comprehensive set of indicators that measure the scale and circularity of total material and waste flows and their socioeconomic and ecological loop closing. We applied this framework in the context of monitoring efforts for a CE in the European Union (EU28) for the year 2014. We found that 7.4 gigatons (Gt) of materials were processed in the EU and only 0.71 Gt of them were secondary materials. The derived input socioeconomic cycling rate of materials was therefore 9.6%. Further, of the 4.8 Gt of interim output flows, 14.8% were recycled or downcycled. Based on these findings and our first efforts in assessing sensitivity of the framework, a number of improvements are deemed necessary: improved reporting of wastes, explicit modeling of societal in-use stocks, introduction of criteria for ecological cycling, and disaggregated mass-based indicators to evaluate environmental impacts of different materials and circularity initiatives. This article met the requirements for a gold – gold JIE data openness badge described at <http://jie.click/badges>.

Introduction

The idea of a more circular economy involves that the value and utility of products are extended and that production and consumption wastes are utilized as secondary resources, promis-

ing solutions and co-benefits to a range of economic and environmental issues (Kirchherr et al. 2017; Winans et al. 2017). Thus, the concept of a circular economy (CE) has gained increasing attention from policy makers, industry, and academia

Conflict of interest statement: The authors declare no conflict of interest.

[Several small corrections were made to table 1 after initial online publication.]

Address correspondence to: Gian Andrea Blengini, European Commission Directorate-General Joint Research Centre Sustainable Resources Directorate, Unit D3 – Land Resources, Via Enrico Fermi 2749, Ispra, VA 21027, Italy. Email: gianandrea.blengini@ec.europa.eu

© 2018 The Authors. *Journal of Industrial Ecology*, published by Wiley Periodicals, Inc., on behalf of Yale University. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.
DOI: 10.1111/jiec.12809

Editor managing review: Guillaume Majeau-Bettez

(Geisendorf and Pietrulla 2017; Bocken et al. 2017a). Nevertheless, a broadly accepted and precise definition of a CE is still lacking; rather, the concept is applied in different ways by stakeholders, depending on their specific interests (Murray et al. 2017; Bocken et al. 2017b; Lieder and Rashid 2016; MacArthur 2013). For European countries, for example, Bocken and colleagues (2017a) have identified a focus of CE applications on business opportunities along with resource efficiency implications, while in China CE was developed around issues of pollution and in the context of China's rapid growth. Also, positive employment effects are frequently mentioned as an important co-benefit of the CE (Stahel 2016; Wijkman and Skanberg 2015; MacArthur 2013). Additionally, the environmental implications, benefits, and trade-offs of a more circular economy are also widely debated (Pauliuk 2018; Geissdoerfer et al. 2017; Geisendorf and Pietrulla 2017).

The European Union embraced the concept of a CE as its key strategy towards a more sustainable use of natural resources. In 2015 the European Commission adopted a strengthened CE package that aims at maintaining the value of products, materials, and resources in the economy for as long as possible, and at minimizing the generation of waste as an essential contribution to the European Union's (EU28) efforts to develop a sustainable, low-carbon, resource-efficient, and competitive economy (COM 2017). There, the transition from a "take-make-consume and dispose" economy towards a "recycle-and-reuse" economy was put into the center. In 2018, and as part of continuous efforts to transform the EU28 economy towards more sustainability, the European Commission adopted a range of CE-related policy measures on, for example, plastics and improved legislation on waste or critical raw materials (European Commission 2018). Additionally, EU28-wide monitoring efforts have been established.

While the basic idea of the CE is intuitive and convincing and the notion is widely used in policy documents, the assessment of progress towards a CE is an issue of ongoing debate (EASAC 2016; Haupt et al. 2017; Kovanda 2014; Hashimoto et al. 2009; Yuan et al. 2006). The majority of CE research has focused on individual products or specific substances within regions or nations, or at the global scale (Huysman et al. 2017). For some substances, mainly metals, the knowledge base on anthropogenic cycles has indeed improved in recent years (BIO by Deloitte 2015; Cullen and Allwood 2013; Graedel et al. 2013; Reck et al. 2008; Wang et al. 2007). A growing body of research has also focused on the company or industry level (Pauliuk 2018; BSI 2017; Lieder and Rashid 2016). However, comprehensive macro-scale assessments of national-level circularity, including derived policy indicators, are very rare (Nuss et al. 2017; Haas et al. 2015; Hashimoto et al. 2004).

Typically, two types of loop closing are distinguished in CE strategies (Haas et al. 2015; MacArthur 2013; UNEP 2012; Braungart et al. 2007; McDonough and Braungart 2003): socioeconomic loop closing by recycling waste materials as secondary material inputs and ecological loop closing by using renewable biomass. If these strategies lead to a reduction in the demand for primary resources, a more circular economy

can lower pressures related to resource extraction, and by reducing the amount of wastes and emissions returned to the environment, it can mitigate pressures on the output side of societies' metabolism (Geissdoerfer et al. 2017). To achieve a more sustainable economy, it is insufficient to only increase recycling and focus on (partial) improvements in the degree of circularity (Cullen 2017; Zink and Geyer 2017), but it is essential to also achieve absolute reductions in resource extraction and consumption, that is, to downsize the socioeconomic metabolism (Akenji et al. 2016). This has implications for the assessment of CE strategies and monitoring tools needed to measure both the degree of loop closing and the overall in- and outflows of societies' metabolism (Krausmann et al. 2017a).

Thus, appropriate monitoring tools need to be able to capture different critical issues related to CE strategies. First, not all recycling activities are necessarily reducing overall resource demand, but they can result in problem shifting. Under certain circumstances, recycling indirectly may require more material and/or energy than the direct use of primary materials (Cullen 2017; Geyer et al. 2016; Behera et al. 2014). Second, a CE is often promoted as an environmentally friendly strategy to facilitate business opportunities and green economic growth, that is, a decoupling of resource use and environmental impacts from economic growth (Ekins 2002; Geng et al. 2013). Consequently, it needs to be critically assessed whether these promises can be realized (Cullen 2017; Zink and Geyer 2017). Third, in-use stocks of manufactured capital are growing in most countries and require an increasing share of overall resource use, which substantially limits possibilities for loop closing (Krausmann et al. 2017b). Only if in-use stocks are steady does a substantial closing of loops become possible because then end-of-life (EoL) flows from demolitions and discards can equal materials used for the maintenance and replacement of in-use stocks (O'Neill 2015; Wiedenhofer et al. 2015). So far, only few studies have attempted to assess the CE at a more comprehensive level taking overall material flows into account (Nuss et al. 2017; Haas et al. 2015; Haupt et al. 2017; Kovanda 2014).

This article aims at contributing to the establishment of monitoring tools of material flows in a CE at the macro level, with the premise that a more circular economy should contribute to the reduction of environmental pressures instigated by resource use. Our proposal is to go beyond the level of individual products, substances, or industrial symbiosis but monitor progress towards a CE from an economy-wide perspective at the national or higher scale. Only at this scale is it possible to also capture system-wide effects such as displacement or rebound effects (Geyer et al. 2016) and to assess whether absolute reductions in resource use and waste flows were achieved. We aim at contributing to the debate of potentials and limitations of a more circular economy by developing a mass-based monitoring framework for the European Union (European Commission 2018; Eurostat 2018). We based our framework upon previous attempts at system-wide assessment of a CE at the global and European level by some of the authors (Haas et al. 2015). However, Haas and colleagues (2015) developed their assessment solely from economy-wide material




Figure 1 Framework and throughput indicators for an economy-wide CE assessment. This framework applies from individual materials (e.g., DE of corn or iron) to aggregated material categories (e.g., PM of biomass, fossil energy carriers) to the total material level (e.g., total DE). Colors indicate data sources used: orange = official data from economy-wide material flow accounts (Eurostat 2017c), blue = official waste and emissions statistics (Eurostat 2017b), green = mass-balanced modeling. Please note that a shift from green to blue color indicates a combination of statistical data and modelling. See table S1 and figure S1 in the supporting information on the Journal's website for an overview, definitions, and results for all flows and variables shown in figure 1. CE = circular economy.

flow (ew-MFA) data on material consumption and their uses. The circularity investigation by Nuss and colleagues (2017) focused on metals and nonmetallic minerals and nonenergy and nonfood biomass. Herein, we expand these efforts by, first, developing a comprehensive approach including all biomass, metals, nonmetallic minerals, and fossil energy carriers and, second, by reconciling officially available datasets published by the Statistical Office of European Communities (EUROSTAT) on material use, waste, and recycling (Eurostat 2017b, 2017c). Innovatively, we linked these datasets in a fully consistent and mass-balanced way in order to achieve a systematic monitoring of resource use, waste, and recycling through the socioeconomic system. From this framework, we derived a set of indicators that measure the scale of input and output flows as well as socioeconomic and ecological loop closing.

The remainder of this article is organized as follows. The next section outlines the general accounting framework, methods and indicators for monitoring the CE, and a first sensitivity assessment. In the following sections, we then present the results in the form of Sankey diagrams of flows of materials through the EU28 economy in 2014 and discuss our findings. In the final section, we discuss the relevance and limitations of the proposed framework and sketch out the way ahead.

Developing a Monitoring Framework for Material Loop Closing in the Circular Economy: Method, Data Sources, and Robustness

We developed a CE monitoring framework at the macro level, built on previous research that proposed an expansion of

ew-MFA by including flows of secondary materials to allow for monitoring socioeconomic loop closing in national economies (Haas et al. 2015). Secondary materials refer to materials recovered through all forms of recycling, reuse, and remanufacturing but also downcycling (e.g., backfilling) or cascading use. We built upon a systems and material perspective of the economy, and as a substantial advancement, we based the assessment as far as possible on statistical data from the official environmental reporting system of the EU (Eurostat 2017c, 2017b) and systematically mass-balanced material inputs with waste flows reported in the different statistical sources. While recovered materials were reported in waste statistics and could be directly quantified, this was not possible for other CE strategies like extending product lifetimes, reusing and remanufacturing, or sharing. In our framework, these strategies would result in an increase of the service lifetime of in-use stocks and potentially a stabilization of in-use stock growth, as indicated by the net additions to stocks (NAS; figure 1). Thus, even though these strategies were difficult to measure directly, their effects on the size of inflows, additions to stock, and outflows can be substantial and are observable via this CE monitoring framework (figure 1).

Quantifying and Tracing Material, Energy, and Waste Flows through the Socioeconomic System

The accounting framework shown in figure 1 traces materials by main material groups from their extraction to major uses within the socioeconomic system and towards discard and either material recovery or deposition to nature as wastes and emissions. In figure 1, the main physical stages of the flow of materials through the entire system are marked by

throughput indicators, represented as boxes. These include the source of material inputs (e.g., domestic extraction, imports), major material transformation processing stages within the system (e.g., processed materials; energetic use; material use, in-use stocks of materials; waste treatment; EoL waste) and the destination of outflows (e.g., exports, domestic processed output to the environment). Flows of materials are displayed as arrows between these boxes; the colors of flows indicate the type of data source (e.g., orange for ew-MFA, blue for waste statistics).

Direct material input (DMI) into the socioeconomic system comprises materials that were extracted from the domestic environment (domestic extraction [DE]) and imports of raw materials and manufactured products. Exports to other economies were deducted to calculate domestic material consumption (DMC). Processed materials (PMs) were defined as the sum total of DMC and secondary material (SM) inputs.

PMs were allocated to either energetic or material use (see table S2 in supporting information S1 available on the Journal's website for detailed allocation tables). Energetic use (eUse) not only comprises materials used to provide technical energy (fuel wood and biofuels) but also feed and food, the primary energy sources for livestock and humans (Haberl 2001; Krausmann et al. 2016). The majority of all fossil energy carriers were allocated to eUse with the exception of, for example, petrochemical feedstock. All other materials were assigned to material use (mUse). mUse comprises all metal ores and metals and nonmetallic minerals, as well as those fractions of fossil and biomass materials not used for energy provision. The biomass fractions assigned to material use include industrial roundwood, crops for uses other than food, and feed including seeds and most harvested crop residues.

mUse was split into extractive waste, materials used for stock building (i.e., gross additions to in-use stocks of materials [GAS]), and throughput materials (Krausmann et al. 2017b; Haas et al. 2015; Lauk et al. 2012). Extractive waste refers to waste material that occurs during early stages of the processing of domestically extracted ores (European Parliament, Council of the European Union 2006) and directly goes from PM to interim output (IntOut). Stock building materials comprise all materials that accumulate in buildings, infrastructures, or durable goods with a lifetime of more than one year (e.g., concrete, asphalt, or steel). The share of stock-building materials in mUse was estimated based on information from industry and production statistics, results from material flow studies and assumptions (see table S2 in supporting information S1 on the Web). Throughput materials comprise materials that do not accumulate in in-use stocks, and can be split into two types of materials: first, materials used deliberately in a dissipative way such as salt or fertilizer minerals, and losses that occur during material processing (wastage, not reported in waste statistics); and second, short-lived products such as packaging or newspaper, manufacturing wastes, and food waste (reported in waste statistics).

All materials that are neither added to stocks nor recycled are converted into gaseous, solid, or liquid outputs within the year of extraction. Together with demolition and discard from

in-use stocks that have reached the end of their service lifetime, these outflows were denoted as interim outputs (IntOut) in figure 1. IntOuts were split into emissions, comprising all gaseous emissions (e.g., carbon dioxide [CO₂], sulfur dioxide [SO₂], methane [CH₄]) including water vapor and into EoL waste, including all solid (and liquid) outputs. Information on outflows was either sourced from Eurostat waste statistics or modeled and mass-balanced with input flows (see below and supporting information S1 and supplementary data S2 on the Web for details). Emissions cannot be recycled and go straight into domestic processed output (DPO). A fraction of total EoL waste, reported as RCV_B – (recovery other than energy recovery—backfilling) and RCV_O (recovery other than energy recovery—except backfilling) in Eurostat (2017b) waste statistics, is reentering socioeconomic processes as secondary materials. Note that imported or exported secondary materials (e.g., scrap, waste paper) were not explicitly accounted for as secondary materials but included in trade flows; they are therefore also not reflected in circularity indicators. The remaining EoL waste (after subtracting SM) is returned to the environment as DPO waste and either landfilled, incinerated, or deliberately applied (e.g., manure, fertilizer). DPO emissions and DPO waste together are DPO.

Closing the Mass Balance between Material Inputs and Outputs of Wastes and Emissions

To close the material balance between input and output flows we combined data from statistical reporting (i.e., from ew-MFA and waste statistics) with modeling. This was done separately for eUse and for the mUse components in two balancing calculations. The following equations summarize the mass balancing for eUse (equation 1) and mUse (equation 2).

$$\text{DPO emissions} = \text{eUse} - \text{solid and liquid wastes} \quad (1)$$

$$\begin{aligned} \text{Demolition and discard} = & \text{EoL waste from mUse} \\ & - \text{throughput materials in waste} \quad (2) \end{aligned}$$

eUse Balancing

We assumed that all materials used to provide energy were converted into DPO emissions (including water vapor) and solid waste within the year of extraction. We used data for solid waste from combustion reported in waste statistics and estimated the amount of solid waste from human and animal metabolism (excrements) by applying appropriate coefficients reflecting the nondigestible fraction of food and feed intake. DPO emissions were then calculated as the difference between eUse and the outflow of solid waste. Note that so-called balancing items (oxygen uptake from air during combustion and water consumed by humans and livestock) were excluded. This means that all outflows from eUse include only the materials contained in actual inputs as comprised in PM (e.g., CO₂ or SO₂ in terms of C or S content; excrements at the average water content of food and feed intake). Closing the mass balance

for eUse in this way implies that all inaccuracies in statistical data and assumptions that result in inconsistencies between input and output flows accrued in DPO emissions (DPOe). For the combustion of fossil energy carriers we cross-checked the calculated DPO emissions with data from emission statistics. We found that our estimate is roughly 6% above emission statistics, which is a reasonably good fit, considering that emission statistics do not provide a comprehensive coverage of all emissions (see table S5 in supporting information S1 on the Web).

mUse Balancing

Due to a lack of knowledge of actual in-use stocks, we used the following approach to close the material balance: In a first step, a consistent split of total EoL waste from mUse into waste flows resulting from discard and demolition and throughput materials was required. Total EoL waste from mUse was derived from waste statistics. While waste statistics report information on construction and demolition waste, this waste flow was not fully consistent with EoL waste from discard and demolition, which also contains waste flows from discarded long-living products such as furniture, cars, or electric appliances. In a second step, we calculated the amount of discard and demolition as the difference between EoL waste from mUse reported in waste statistics and the fraction of throughput materials (i.e., materials with a life span < 1 year) in mUse (e.g., waste from packaging, paper, food waste, etc.). In a third step, NAS were calculated as the difference between additions to stocks and discard and demolition. Closing the mass balance in this way implies that all inaccuracies in statistical data and assumptions that result in inconsistencies between input and output flows for mUse accrue in demolition and discard flows as residual flow category, and consequently in the value for NAS. Since no information on NAS in the EU28 was available, a cross-check with independent data was not possible. Krausmann et al. (2017b) have estimated NAS in industrial countries in 2014 to amount to 6.7 tons per capita per year (t/cap/yr). For the EU28 we arrived at 5.2 t/cap/yr. Given that the two estimates refer to different country groups and apply different methods, we conclude that our estimate is in a plausible range.

All flows and indicators were calculated for the four main material groups distinguished in ew-MFA: nonmetallic minerals, metal ores and metals, fossil energy carriers, and biomass. The calculation at the level of material groups was challenging because waste statistics of Eurostat [*Regulation (EC) No 2150/2002*; see European Commission and Eurostat (2013)] follow a classification that refers to economic sectors and activity (NACE classification), different collection systems, and/or hazard potential. Waste materials reported in one category typically comprise multiple material categories in ew-MFA, which required an allocation of output to input flows. Waste flows reported in waste statistics needed adjustments to the system boundaries used in ew-MFA to ensure that input and output flows can be mass balanced (see table S4 in supporting information S1 on the Web for detailed allocation tables).

Deriving Mass-Based Indicators for a Circular Economy

Indicators highly depend on the underlying definition of a CE. Here, we follow a definition that was first developed by Braungart and colleagues (2007) and summarized by the Global Energy Outlook (UNEP 2012) and defines a CE as an economy in which material flows are made up either of biological materials, which after discard are integrated into ecological cycles, or of materials designed to circulate within the socioeconomic system (MacArthur 2013). Based on this general definition, we developed a set of six indicator pairs, which allows us to measure progress towards a CE in terms of closing material cycles (table 1). The indicators presented here are based on ew-MFA principles and build on proposals from previous research (Nuss et al. 2017; Haas et al. 2015; Hashimoto et al. 2004; Kovanda 2014). We distinguished between scale indicators, which provide measures for the overall size of the socioeconomic metabolism (O'Neill 2015), and circularity rates, which measure socioeconomic and ecological cycling relative to input and output flows. Providing independent measures for flows on both the input and output sides is necessary because of the delaying effect that in-use stocks of materials have on output flows.

We used three pairs of indicators to measure the scale of material and waste flows: DMC measures all materials directly used in a national production system and is regarded as proxy for the aggregated pressure the economy exerts on the environment (Krausmann et al. 2017a). DPO measures the total amount of outflow of wastes and emissions from a national economy. In order to be able to capture displacement effects related to imports and exports, we include a consumption-based indicator in the framework. For this purpose, Eurostat (2017a) provides the indicator raw material consumption (RMC), which is similar to the material footprint (Wiedmann et al. 2015) and measures global material use associated with domestic final consumption (Krausmann et al. 2017a). The final pair of scale indicators takes the flow of secondary materials into account, which is not presented in conventional ew-MFA indicators: On the input side, the indicator PM measures the sum total of DMC plus the input of secondary materials, and on the output side, IntOut measures wastes and emissions before materials for recycling and down-cycling are diverted. Even in industrial countries, stocks are growing and interim outflows in a given year are much smaller than the amount of PM in that year, which further inhibits loop closing at present, producing a delaying effect for potential recycling of these materials after their lifetime has ended in the future.

As indicators for the degree of loop closing that has been achieved, we propose three pairs of circularity rates, which measure material flows relative to interim flows PM and IntOut. The socioeconomic cycling rates measure the contribution of secondary materials to PM (input socioeconomic cycling rate [ISC_r]) and the share of IntOut that is diverted to be used as secondary materials (output socioeconomic cycling rate [OSC_r]). For biomass, derived circularity indicators are more intricate. Due to the absence of a clear definition and recognized criteria for sustainably produced biomass, as well as a lack of

Table 1 Mass-based circular economy indicators for the EU28 in 2014 where scale indicators measure the absolute size of input and outputs flows in tons and circularity rates measure socioeconomic and ecological cycling relative to input and output flows in percentage¹

	<i>Dimension</i>	<i>Input-side indicator</i>	<i>Output-side indicator</i>
Scale indicators (t)	In- and output flows	Domestic material consumption DMC	Domestic processed outputs DPO
	Consumption based perspective	Raw material consumption RMC	n.a.
	Interim flows	Processed materials PM = DMC + secondary materials	Interim outputs IntOut = EoL waste + DPO emissions
Circularity rates (%)	Socioeconomic cycling SC	Input socioeconomic cycling rate ISCr = Share of secondary materials in PM	Output socioeconomic cycling rate OSCr = Share of secondary materials in IntOut
	Ecological cycling potential EC	Input ecological cycling rate potential IECrp = Share of DMC of primary biomass in PM	Output ecological cycling rate potential OECrp = Share of DPO biomass in IntOut
	Non-circularity NC	Input non-circularity rate INCr = Share of eUse of fossil energy carriers in PM	Output non-circularity rate ONCr = Share of eUse of fossil energy carriers in IntOut

Note: n.a. = not applicable.

related data, we use the share of primary biomass (i.e., biomass DMC) in PM for the input ecological cycling rate potential (IECrp) and the share of DPO from biomass in IntOut for the output ecological cycling rate potential (OECrp). Because ecological cycling is a crucial part of CE strategies, data and adequate indicators have to be developed so that socioeconomic and ecological cycling rates indicate the overall circularity of an economy. Finally, the noncircularity indicators measure the share of eUse of fossil energy carriers in PM and IntOut, thus quantifying the share of material flows that do not qualify neither for socioeconomic and ecological loop closing. Due to unreliable information on dissipation rates of fertilizers or salt for deicing roads, for example, we did not allocate these materials to noncircularity flows. The difference between 100% and the sum total of the three circularity rates serve as a measure for the unexploited potential for socioeconomic cycling.

The indicators proposed herein were developed to detect and monitor economy-wide improvements and trade-offs for economy-wide circularity. Since specific materials or material categories are interconnected (Bleischwitz et al. 2018), enhancing circularity for one material has an impact upon the use of other materials and energy. In the case of plastic, construction minerals, or metals, recycling might mean high energy investments in recycling operations, infrastructure, and transport. Consequently, increasing recovery rates, for example, for plastic waste, might reduce overall circularity through higher energy use. Thus, we used the shares of socioeconomic, ecological, and noncircular flows in PM and IntOut as circularity indicators that are able to monitor system-wide implications of circularity initiatives.

Validation and Sensitivity: How Robust Are the Derived Circular Economy Indicators?

Generally, for ew-MFA a systematic assessment of uncertainty is in its infancy (Patrício et al. 2015; Laner et al. 2014). It has been shown that material flow accounts using different sources and estimation procedures typically differ in their results for global DE by approximately 5% to 20% (Schandl et al. 2017; Krausmann et al. 2017a; Fischer-Kowalski et al. 2011). For waste statistics, harmonization of data reporting across countries is a major challenge, and data quality differs across EU28 countries (Tisserant et al. 2017; Nicolli et al. 2012).

We employed a one-at-a-time sensitivity assessment (Saltelli et al. 2000) to quantify the significance of variations in all major data sources and allocation steps used in the CE assessment for four scale indicators (PM, IntOut, DPO, DPOe) and two circularity indicators (ISCr and OSCr) from our framework and applied a $\pm 20\%$ sensitivity test on the underlying statistical data (a–c) and allocation schemes (d–e). We conducted such tests for (a) the ew-MFA data, (b) waste statistics, (c) secondary materials, (d) the allocation between energetic and material use, and (e) the matching of waste flows to ew-MFA categories (figure 2).

We found that all indicators were most sensitive to changes in the underlying statistical data, with material flow data causing the largest sensitivities (sensitivities a–c, see figure 2). DPOe showed the highest and same directional sensitivity with $\pm 26\%$ ($\pm 20\%$ variation in ew-MFA data), followed by lower sensitivities for PM, DPO, and IntOut. A $\pm 20\%$ variation in waste statistics caused sensitivities less than 10% for all scale indicators, and $\pm 20\%$ variations in the amount of secondary




Figure 2 Results for the one-at-a-time (OOAT) $\pm 20\%$ sensitivity tests of the CE assessment for four scale and two circularity indicators. Due to the modeling framework architecture, we performed sensitivity e for industrial minerals and construction minerals, respectively. Results are presented as the relative change of each indicator due to the $\pm 20\%$ sensitivity test for the identified main sources of uncertainty. For detailed results, see table S3 in supporting information S1 on the Web. ew-MFA = EUROSTAT ew-MFA dataset (Eurostat 2017c). env_wasrt = EUROSTAT waste treatment statistics (Eurostat 2017b), mUse = material use, SM = secondary materials.

materials showed lower sensitivities of less than 4%. ISCr and OSCr responded nonlinearly to changes in input data, again in declining magnitude from sensitivities a through d (except sensitivity d for OSCr).

All indicators were much less sensitive to changes in the allocation schemes (sensitivities d–e). The output-related scale indicators responded with $\pm 1.4\%$ to $\pm 2.6\%$ to a $\pm 20\%$ change in the assumptions used for the allocation of PM to eUse and mUse, with the OSCr showing a sensitivity less than 1.5%. All other allocation sensitivities were below $\pm 1.1\%$ for the selected indicators.

Overall, the results of the one-at-a-time sensitivity test indicated that the CE indicators and main results of the assessment are quite robust against variations in the allocation of waste flows to main material categories for the assessed material flows and indicators. The strongest relative changes in the assessed flows and indicators relate to the quality of input data.

Results

Material Flows through the EU28 Economy

Applying the CE monitoring framework to the EU28 in 2014 shows that 5.8 Gt (1 gigaton = 10^9 tons) of raw materials, which

were used in the EU, originated from domestic extraction, 0.9 Gt from net imports, and 0.7 Gt from secondary materials (figure 3). Of the 7.4 Gt of PM, 3.1 Gt were used to provide energy and 4.3 Gt for material use. Throughput materials including extractive waste accounted for 0.7 Gt, while the majority of 3.5 Gt were used to expand and maintain material in-use stocks of buildings, infrastructure, and other long-lived material products. Overall, 2.6 Gt have been net additions to in-use stocks. Thus, nearly one third of all processed materials in 2014 increased the material stocks in the EU28 and will remain in use for years to decades, thereby shaping future waste flows and potentials for closing material loops, but also requiring energy use for their operation. A total of 2.2 Gt of EoL wastes resulted from demolition and discard, throughput materials, and solid wastes from energetic use, of which one third were recovered and reused as secondary materials. The remaining two thirds were landfilled or incinerated and thus eventually released to the environment. DPO emissions to air accounted for 2.5 Gt, of which 1.1 Gt originated from food, feed, and biomass energy, and 1.4 Gt from fossil energy carriers.

Figure 4a–d show results for the four main material categories (see supporting information S1 and supplementary data S2 on the Web for additional results). The largest material flow in the EU28 were nonmetallic minerals, which amounted to 3.5 Gt




Figure 3 Material flows through the EU28 economy in 2014. In this Sankey diagram, the width of the arrows is proportional to the size of material flows (dark blue); the numbers show the size of the material flows in Gt/yr and the bars their composition (share of four main material groups in %). Note that numbers may not always sum up to total due to rounding. EU28 = European Union; Gt/yr = gigatons per year.




Figure 4 Material flows through the EU28 economy in 2014 for four main material groups in Gt/yr: (a) Nonmetallic minerals, including both construction and industrial minerals; (b) fossil energy carriers; (c) biomass; (d) metal ores and metals (note that the scale for metal ores is factor 5 lower). Balancing flows of oxygen and water are not included. Note that numbers may not always sum up to total due to rounding. DE = domestic extraction, DPO = domestic processed output; EU28 = European Union; Gt/yr = gigatons per year.

or 50% of all processed materials and are shown in figure 4a. The majority of nonmetallic minerals were used to maintain and expand societal in-use stocks, resulting in 2.4 Gt of net additions to material stocks. Of the total EoL waste from nonmetallic minerals, one third (0.35 Gt) were recovered, equivalent to a material-specific output cycling rate of 33.8%, and an input cycling rate of 9.5%, caused by the large net additions to stocks.

Only a small fraction (0.05 Gt) of processed fossil energy carriers were used for nonenergetic purposes (figure 4b). Of the 0.15 Gt of solid and liquid residues from fossil energy materials/carriers use, 42% were recovered, mainly bitumen and plastics, although the material-specific output circularity is only 4.1%, due to the majority of fossil energy carriers used for energy provision. The 1.7 Gt of eUse of fossil energy carriers did not qualify for socioeconomic cycling and lead to 1.5 Gt of emissions to air (including water vapor but excluding oxygen from air), which are a major driver for anthropogenic climate change (IPCC 2013).

For biomass (figure 4c), we found that 90% of the 2 Gt PM were sourced from extraction within the EU28 in 2014, of which roughly one fifth were used for material purposes; most importantly, wood, with 0.3 Gt. 78% of all biomass (1.6 Gt), was used as energy source to feed livestock and humans and to provide technical energy, leading to 1.1 Gt of emissions. A total of 0.2 Gt (or 26% of EoL waste of biomass) were recovered, leaving 0.6 Gt DPO waste flows to be returned to the environment through landfills or incineration.

Metal ores and metals are materials of high economic and thermodynamic value. Additionally, they also have high strategic importance, given that 90% (0.21 Gt) of all processed metals were imported from other world regions (figure 4d). Domestic extraction of metals contributed only 0.02 Gt to the PM of 0.22 Gt but generated 0.17 Gt of extractive waste. The majority of

processed metal ores and metals, of which nearly one third were secondary materials, have been integrated into material stocks.

Circularity Indicators

In 2014, 65% (4.8 Gt) of PM were converted into IntOut. The remaining 35% were added to in-use stocks of buildings, infrastructure, and durable goods, which have grown in 2014 by 2.6 Gt (NAS) (figure 5 and table 2). Even though nearly one third of the total EoL waste was recovered and used as secondary resources, the OSCr remained low, at only 14.8%. The input socioeconomic cycling rate, measuring the recycled and downcycled materials that were reprocessed as secondary material inputs into the domestic economy was even lower, at 9.6%. Ecological cycling, although indicated only as theoretical potential, was comparatively high; the IECrp, indicating the maximum share of PM that qualifies for ecological cycling, was 24.6%; the OECrp was even higher, at 35.3%. The high significance of fossil energy carriers in the EU28 primary energy supply, flows that cannot be recycled or reused, led to an input noncircularity rate (INCr) of 21.2%, and an output noncircularity rate (ONCr) of 32.8%.

While our assessment provided a snapshot for a single year, Eurostat material flow analysis and waste statistics allow deriving some rough indications on the direction of the development. For the years 2010 and 2012, consistent information on DMC and the flow of recycled and downcycled materials was available, which was used to calculate the ISCr (table 3). We found a slight decrease in DMC and a more or less stable flow of secondary materials, which resulted in a slight growth in ISCr between 2010 and 2014. Slight progress towards a CE was rather due to reductions in material input, while socioeconomic cycling showed little improvement. Providing indicators for ecological loop closing was more intricate, but the slightly




Figure 5 Input- and output-side CE indicators. Left bar depicts processed materials; right bar depicts interim outputs. Percentage denotes the share in relation to processed materials (left bar) and interim outputs (right bar).

Table 2 Mass-based circular economy indicators for the EU28 in 2014

Input-side indicator			Output-side indicator	
Scale indicators (Gt/yr)	Domestic material consumption (DMC)	6.7	Domestic processed outputs (DPO)	4.1
	Raw material consumption (RMC)	7.2	NA	
	Processed materials (PM)	7.4	Interim outputs (IntOut)	4.8
Circularity rates (%)	Input socioeconomic cycling rate (ISCr; share of PM)	9.6	Output socioeconomic cycling rate (OSCr; share of IntOut)	14.8
	Input ecological cycling rate potential (IECrp)	24.6	Output ecological cycling rate potential (OECrp)	35.3
	Input non-circularity rate (INCr)	21.2	Output noncircularity rate (ONCr)	32.8

Note: Indicators differentiate between input- and output-side and in the scale of flows and related circularity rates. Scale indicators presented in Gt/yr, circularity rates in percentage.

Gt/yr = gigatons per year; NA = not applicable.

increasing share of biomass in DMC indicates a growing ecological cycling potential.

Discussion

Assessment of Circularity

We investigated the status quo of the CE in the EU28 in 2014 and developed a set of headline indicators on the scale and circularity of material and waste flows. Of 7.4 Gt of PMs, 6.7 Gt were primary resources and led to 4.8 Gt of IntOuts. All circularity indicators showed higher rates when compared to outflows, and these differences were due to the growth of in-use stocks in the EU28 economy. Nearly one tenth of all PMs (9.6%), and 14.8% of all IntOuts were secondary resources. Acknowledging the caveats in calculating and interpreting ecological cycling rates, we found potentials of 24.6% and 35.3% for input and output ecological cycling rate potentials, respectively. The low socioeconomic loops we found in our study are also lower than in Haas and colleagues (2015), who estimated an ISCr of 13% for the EU27 in 2005 (see table S6 in supporting information S1 on the Web for a more detailed comparison). This study did not use data from actual waste statistics but was based on a simple leaching model for outflows from in-use stocks and assumed that all abandoned stocks after EoL turn into demolition and

discard waste, thus ignoring the issue of hibernating stocks. This resulted in an overestimation of waste flows and, by applying recycling rates rather than statistical data, to an overestimation of circularity rates.

Overall, input and output socioeconomic cycling was surprisingly low, considering the fact that the EU28 has strict waste regulations, elaborate waste collection and recovery systems, and high material category-specific recovery rates that range from 25% for biomass to 70% for metals. While it is important to further improve the recycling and downcycling of EoL waste, our results emphasize that achieving a CE goes far beyond increasing reuse and recycling. Major obstacles for substantially increasing socioeconomic cycling are the ongoing expansion of in-use stocks and the high share of noncircular fossil energy carriers in PM (INCr). Increasing the lifetime and a more intensive use of material stocks, as envisaged via increasing value and utility of products, are important measures towards a CE and need to be developed in this direction. Additionally, improved recycling technology and changes in product design to enhance recyclability are important challenges (Ciacci et al. 2015; Reck and Graedel 2012; Gaustad et al. 2010). Clearly, also a reduction of fossil fuel consumption is urgently required to mitigate climate change, which would also increase socioeconomic circularity (Ghisellini et al. 2016). For renewable biomass resources, CE strategies should focus on less wasteful, more efficient, and cascading uses as well as a production system that fosters and sustains ecological cycles, rather than simply increasing biomass inputs to substitute other materials (Smith et al. 2014; Haberl and Geissler 2000).

Table 3 Comparison of DMC, secondary materials (SM), processed materials (PM), input socioeconomic cycling rate (ISCr), and share of biomass in DMC for 2010, 2012, and 2014

	2010	2012	2014
DMC ^a (Gt)	7.0	6.8	6.7
Secondary materials ^b (Gt)	0.72	0.73	0.71
PM (Gt)	7.7	7.5	7.4
ISCr = Secondary materials/PM (%)	9.3%	9.7%	9.6%
DMC biomass/DMC (%)	23.2%	24.4%	27.2%

^aData source for 2010 and 2012: Eurostat 2017c.

^bData source for 2010 and 2012: Eurostat 2017b.

Gt = gigatons.

Further Advancing Data and Indicators for Circular Economy Assessments

We introduced an economy-wide and mass-based framework to monitor progress towards a CE and shown that data from ew-MFA and waste statistics can be used to calculate a set of headline indicators. Our research also revealed a number of limitations and weaknesses, which need to be addressed to better exploit the potential of this approach and to provide better and more reliable assessments of a CE in the future.

Improving Statistical Reporting

Consistently integrating data from different statistical sources reveals the need to improve statistical reporting and to explicitly address uncertainties in reported quantities. Similar to other statistical data sources, it would be beneficial to develop an official concordance between ew-MFA and waste statistics. In particular, the quality of waste statistics calls for improvements, since it is unclear how complete the coverage of waste flows in the different countries is and reported data seem to be of very different quality. Discrepancies between the amount of recovered and actually recycled materials replacing primary material are not well understood and reported and lead to an overestimation of socioeconomic cycling rates. Schiller and colleagues (2017) conclude that for Germany, only 48% of outflows of concrete and bricks are suitable for high-quality recycling, reducing the recycling share in fresh concrete to a maximum of 32%. A Swiss study of polyethylene terephthalate (PET) bottles, tinplate, aluminum, paper and cardboard, and glass reveals that actual recycling rates are substantially lower than reported in official statistics (Haupt et al. 2017), and the authors expect similar differences in other countries. Thus, the EU28's continuous efforts towards a CE calls for improvements and harmonization in statistical reporting.

Improving knowledge about material stocks would enable filling gaps in waste statistics and cross-validating mass balances, explicitly considering so-called hibernating stocks, and gaining insights about the future dynamics of EoL waste potentials. Currently, waste statistics (incompletely) report those end-of-life materials which are collected and enter waste treatment facilities. On-site waste flows often downcycled are not covered, as are so-called hibernating stocks (Hashimoto et al. 2009), which results in an overestimation of NAS and an underestimation of outflows from stocks, and adds considerable uncertainty about indicators for socioeconomic loop closing. Lack of knowledge about the aging dynamics of in-use stocks means that future EoL flows cannot be prospectively modeled, nor can spatial patterns of waste flows be derived. Such information would be useful to inform waste management planning and circularity potentials. Closing the knowledge gap about in-use stocks requires records of in-use vintages of material stocks and their service lifetimes, which can be achieved via different modeling approaches, depending on data availability (Miatto et al. 2017; Augiseau and Barles 2017; Fishman et al. 2014; Pauliuk et al. 2012).

Broadening the Scope of Circular Economy Monitoring

The proposed framework provides a set of mass-based indicators for monitoring the progress towards a more circular economy in terms of material loop closing. The strength of this approach is the system-wide perspective, which provides a comprehensive picture of material loop closing in a CE across all materials used by society. The proposed indicators allow us to detect problem shifting and can be used to determine target values to downscale the industrial metabolism (Bringezu 2015), which, in addition to targeted measures for specifically problematic flows, must be the ultimate aim of a circular economy. The aggregate indicators, however, obscure flows of in-

dividual materials or substances, which may be of high interest for CE policies due to criticality or specific environmental pressures but have only a low share in terms of mass. Plastics, for example, are subsumed under fossil energy carriers or critical metals (European Commission 2010) under metal ores and metals. Tracing changes in these flows requires additional data, which can be obtained from more detailed material flow analysis (Nuss and Blengini 2018; Brooks et al. 2018) and should be integrated consistently into the proposed framework. The mass-based indicators deliver a robust data basis that is extendable with more in-depth assessments of individual materials, but also with social, economic, and environmental indicators. For example, Van der Voet and colleagues (2009) proposed an environmentally weighted material consumption indicator (EMC) to measure the total environmental impact of material flows, and Pauliuk (2018) proposed a dashboard of CE core indicators for organizations that partly builds upon MFA methods. Additionally, a more explicit measurement of energy flows (Cleveland et al. 2000) or differences in energy quality (Ayres and Warr 2009) can be modeled based on this framework. Also, a better understanding of the GHG implications of ecological cycling is required.

Criteria for Ecological Loop Closing

Currently, it is assumed that all used biomass contributes to ecological loop closing; this is the case only if biomass production and discharge maintains the regenerative capacities of eco-systems. A proper assessment of ecological loop closing is, however, not possible due to a lack of data and appropriate indicators. Criteria to be considered refer to soil degradation, overexploitation of water resources, efficient management of plant nutrients and livestock manure (Zanten et al. 2018; Dawson and Hilton 2011), the regional scale of ecological loop closing (Therond et al. 2017), and net carbon losses related to cultivation (Erb et al. 2017; Fargione et al. 2008). This issue is of particular importance because climate change mitigation and bioeconomy strategies foster the use of biomass.

Global Integration

Our assessment focuses on material flows and circularity within the EU28; however, as trade with waste and secondary resources is rapidly increasing, it is important to also capture the global cross-country dimension of circularity in national assessments (Beukering 2013; Berglund and Söderholm 2003; Frazzoli et al. 2010). While consumption-based indicators such as the RMC capture the global dimension of domestic final consumption (Wiedmann et al. 2015), exports and imports of waste and secondary materials must be better documented and integrated in the assessment framework and derived indicators.

Conclusions and the Way Forward

The concept of a CE is a powerful bridging concept to foster the fundamental links among resource use, waste, and emissions, and to contribute to integrating environmental

(output-related) and economic (input-related) policies. Environmental pressures resulting from the scale and structure of the industrial metabolism require concerted action on both ends. Improved collaboration between these currently isolated policy domains could realize co-benefits among environment, employment, and security of supply. Beyond the policy arena, well-attuned concerted actions of policy makers with industries ranging from production to waste management is a further necessary strand for pushing the CE. At the same time, all these initiatives require monitoring frameworks, which provide indicators to assess links between a CE and sustainability goals on all scales. The combination of a systematic and mass-balanced approach with regularly published statistical data calls for improvements of data quality, standardizing waste statistics, and consolidating them with ew-MFA data. Such improvements allow better understanding of not only the level of circularity, but the quality of the circularity as well, which would, in turn, allow for a better understanding of the true contribution to sustainability goals.

Acknowledgments

The authors thank colleagues from the Directorate-General Internal Market, Industry, Entrepreneurship and SMEs (DG GROW) and Directorate-General Eurostat (DG ESTAT) for their valuable inputs. The views expressed in the article are personal and do not necessarily reflect an official position of the European Commission. Finally, we thank four anonymous reviewers who provided competent comments that helped to improve the manuscript.

Funding Information

The authors gratefully acknowledge funding by the Austrian Science Fund (FWF; project MISO P27590), and the European Commission – Directorate-General Joint Research Centre (DG JRC) under Expert Contract Number CT-EX2016D288092-101-1 for Willi Haas.

Notes

1. Gigatons and tons refer to *metric tons* throughout the manuscript and supporting information files.

References

- Akenji, L., M. Bengtsson, R. Bleischwitz, A. Tukker, and H. Schandl. 2016. Ossified materialism: Introduction to the special volume on absolute reductions in materials throughput and emissions. *Journal of Cleaner Production* 132(Supplement C): Absolute Reductions in Material Throughput, Energy Use and Emissions: 1–12.
- Augiseau, V. and S. Barles. 2017. Studying construction materials flows and stock: A review. *Resources, Conservation and Recycling* 123: 153–164.
- Ayres, R.U. and B. Warr. 2009. *The economic growth engine: How energy and work drive material prosperity*. Cheltenham, UK; Northampton, MA: Edward Elgar.
- Behera, M., S.K. Bhattacharyya, A.K. Minocha, R. Deoliya, and S. Maiti. 2014. Recycled aggregate from C&D waste & its use in concrete—A breakthrough towards sustainability in construction sector: A review. *Construction and Building Materials* 68: 501–516.
- Berglund, C. and P. Söderholm. 2003. An econometric analysis of global waste paper recovery and utilization. *Environmental and Resource Economics* 26(3): 429–456.
- Beukering, P.J. van. 2013. *Recycling, international trade and the environment: An empirical analysis*. Dordrecht: Springer Science & Business Media, March 9.
- BIO by Deloitte. 2015. *Study on data for a raw material system analysis: Roadmap and test of the fully operational MSA for raw materials*. Prepared for the European Commission. DG GROW. https://ec.europa.eu/assets/jrc/msa/images/msa_final_report.pdf. Accessed August 14, 2018.
- Bleischwitz, Raimund, Holger Hoff, Catalina Spatar, Ester van derVoet, and Stacy VanDeveer, eds. 2018. *Routledge handbook of the resource nexus*. Routledge. www.beck-shop.de/Bleischwitz-Hoff-Spataru-Routledge-Handbook-of-Resource-Nexus/productview.aspx?product=18971823. Accessed January 5, 2017.
- Bocken, N.M.P., E.A. Olivetti, J.M. Cullen, J. Potting, and R. Lifset. 2017a. Taking the circularity to the next level: A special issue on the circular economy. *Journal of Industrial Ecology* 21: 476–482.
- Bocken, N.M.P., P. Ritala, and P. Huotari. 2017b. The circular economy: Exploring the introduction of the concept among S&P 500 firms. *Journal of Industrial Ecology* 21(3): 487–490.
- Braungart, M., W. McDonough, and A. Bollinger. 2007. Cradle-to-cradle design: Creating healthy emissions—a strategy for eco-effective product and system design. *Journal of Cleaner Production* 15(13–14): 1337–1348.
- Bringezu, S. 2015. Possible target corridor for sustainable use of global material resources. *Resources* 4(1): 25–54.
- Brooks, A.L., S. Wang, and J.R. Jambeck. 2018. The Chinese import ban and its impact on global plastic waste trade. *Science Advances* 4(6): eaat0131.
- BSI. 2017. *BS 8001:2017. Framework for implementing the principles of the circular economy in organizations—Guide*. London: The British Standards Institution. www.bsigroup.com/en-GB/standards/benefits-of-using-standards/becoming-more-sustainable-with-standards/BS8001-Circular-Economy/. Accessed April 9, 2018.
- Ciacchi, L., B.K. Reck, N.T. Nassar, and T.E. Graedel. 2015. Lost by design. *Environmental Science & Technology* 49(16): 9443–9451.
- Cleveland, C.J., R.K. Kaufman, and D.I. Stern. 2000. Aggregation and the role of energy in the economy. *Ecological Economics* 32(2): 301–317.
- COM. 2017. *Report from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions on the implementation of the Circular Economy Action Plan*. Brussels: European Commission. http://ec.europa.eu/environment/circular-economy/implementation_report.pdf. Accessed April 21, 2017.
- Cullen, J.M. 2017. Circular economy: Theoretical benchmark or perpetual motion machine? *Journal of Industrial Ecology* 21(3): 483–486.
- Cullen, J.M. and J.M. Allwood. 2013. Mapping the global flow of aluminum: From liquid aluminum to end-use goods. *Environmental Science & Technology* 47(7): 3057–3064.

- Dawson, C.J. and J. Hilton. 2011. Fertiliser availability in a resource-limited world: Production and recycling of nitrogen and phosphorus. *Food Policy* 36: S14–S22.
- EASAC, ed. 2016. *Indicators for a circular economy*. EASAC policy report 30. Halle (Saale), Germany: EASAC Secretariat, Deutsche Akademie der Naturforscher Leopoldina.
- Ekins, P. 2002. *Economic growth and environmental sustainability: The prospects for green growth*. London and New York: Routledge, January 4.
- Erb, K.-H., T. Kastner, C. Plutzer, A.L.S. Bais, N. Carvalhais, T. Fetzel, S. Gingrich., et al. 2017. Unexpectedly large impact of forest management and grazing on global vegetation biomass. *Nature*. www.nature.com/doi/10.1038/nature25138. Accessed December 25, 2017.
- European Commission. 2010. Critical raw materials for the EU: Report of the Ad-hoc Working Group on defining critical raw materials. <https://ec.europa.eu/growth/tools-databases/eip-raw-materials/en/community/document/critical-raw-materials-eu-report-ad-hoc-working-group-defining-critical-raw>. Accessed August 14, 2018.
- European Commission. 2018. Circular economy—Implementation of the Circular Economy Action Plan. http://ec.europa.eu/environment/circular-economy/index_en.htm. Accessed January 17, 2018.
- European Commission and Eurostat. 2013. *Manual on waste statistics: A handbook for data collection on waste generation and treatment*. Luxembourg: Publications Office. <http://dx.publications.europa.eu/10.2785/4198>. Accessed March 15, 2017.
- European Parliament, Council of the European Union. 2006. EUR-Lex - 32006L0021 - EN - EUR-Lex. <https://eur-lex.europa.eu/eli/dir/2006/21/oj>. Accessed June 24, 2018.
- Eurostat. 2017a. Material flow accounts in raw material equivalents - modelling estimates [env_ac_rme]. *Eurostat: Statistics explained—Abbreviations*. <http://ec.europa.eu/eurostat/statistics-explained/index.php/Category:Abbreviations>. Accessed May 29, 2017.
- Eurostat. 2017b. Waste generation and treatment (env_wastrt). October 17. http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=env_wastrt&lang=en. Accessed August 14, 2018.
- Eurostat. 2017c. Material flow accounts (env_ac_mfa). October 17. http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=env_ac_mfa&lang=en. Accessed August 14, 2018.
- Eurostat. 2018. Eurostat—Statistics explained. Circular economy. <http://ec.europa.eu/eurostat/web/circular-economy/material-flow-diagram>. Accessed August 14, 2018.
- Fargione, J., J. Hill, D. Tilman, S. Polasky, and P. Hawthorne. 2008. Land clearing and the biofuel carbon debt. *Science* 319(5867): 1235–1238.
- Fischer-Kowalski, M., F. Krausmann, S. Giljum, S. Lutter, A. Mayer, S. Bringezu, Y. Moriguchi, H. Schütz, H. Schandl, and H. Weisz. 2011. Methodology and indicators of economy-wide material flow accounting. *Journal of Industrial Ecology* 15(6): 855–876.
- Fishman, T., H. Schandl, H. Tanikawa, P. Walker, and F. Krausmann. 2014. Accounting for the material stock of nations. *Journal of Industrial Ecology* 18(3): 407–420.
- Frazzoli, C., O.E. Orisakwe, R. Dragone, and A. Mantovani. 2010. Diagnostic health risk assessment of electronic waste on the general population in developing countries' scenarios. *Environmental Impact Assessment Review* 30(6): 388–399.
- Gaustad, G., E. Olivetti, and R. Kirchain. 2010. Design for recycling. *Journal of Industrial Ecology* 14(2): 286–308.
- Geisendorf, S. and F. Pietrulla. 2017. The circular economy and circular economic concepts: A literature analysis and redefinition. *Thunderbird International Business Review*. <https://doi.org/10.1002/tie.21924>. Accessed August 8, 2017.
- Geissdoerfer, M., P. Savaget, N.M.P. Bocken, and E.J. Hultink. 2017. The circular economy—A new sustainability paradigm? *Journal of Cleaner Production* 143: 757–768.
- Geng, Y., J. Sarkis, S. Ulgiati, and P. Zhang. 2013. Measuring China's circular economy. *Science* 339(6127): 1526–1527.
- Geyer, R., B. Kuczenski, T. Zink, and A. Henderson. 2016. Common misconceptions about recycling. *Journal of Industrial Ecology* 20(5): 1010–1017.
- Ghisellini, P., C. Cialani, and S. Ulgiati. 2016. A review on circular economy: The expected transition to a balanced interplay of environmental and economic systems. *Journal of Cleaner Production* 114. Towards Post Fossil Carbon Societies: Regenerative and Preventative Eco-Industrial Development: 11–32.
- Graedel, T.E., E.M. Harper, N.T. Nassar, and B.K. Reck. 2013. On the materials basis of modern society. *Proceedings of the National Academy of Sciences*: 201312752. <https://doi.org/10.1073/pnas.1312752110>.
- Haas, W., F. Krausmann, D. Wiedenhofer, and M. Heinz. 2015. How circular is the global economy? An assessment of material flows, waste production, and recycling in the European Union and the world in 2005. *Journal of Industrial Ecology* 19: 765–777.
- Haberl, H. 2001. The energetic metabolism of societies part I: Accounting concepts. *Journal of Industrial Ecology* 5(1): 11–33.
- Haberl, H. and S. Geissler. 2000. Cascade utilization of biomass: Strategies for a more efficient use of a scarce resource. *Ecological Engineering* 16: 111–121.
- Hashimoto, S., Y. Moriguchi, A. Saito, and T. Ono. 2004. Six indicators of material cycles for describing society's metabolism: Application to wood resources in Japan. *Resources, Conservation and Recycling* 40(3): 201–223.
- Hashimoto, S., H. Tanikawa, and Y. Moriguchi. 2009. Framework for estimating potential wastes and secondary resources accumulated within an economy—A case study of construction minerals in Japan. *Waste Management* 29(11): 2859–2866.
- Haupt, M., C. Vadenbo, and S. Hellweg. 2017. Do we have the right performance indicators for the circular economy? Insight into the Swiss waste management system. *Journal of Industrial Ecology*. 21(3): 615–627.
- Huysman, S., J. De Schaepmeester, K. Ragaert, J. Dewulf, and S. De Meester. 2017. Performance indicators for a circular economy: A case study on post-industrial plastic waste. *Resources, Conservation and Recycling* 120: 46–54.
- IPCC, 2013. Climate Change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change. Edited by T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P.M. Midgley. Cambridge, UK and New York, NY: Cambridge University Press, 1535 pp, <https://doi.org/10.1017/CBO9781107415324>.
- Kirchherr, J., D. Reike, and M. Hekkert. 2017. Conceptualizing the circular economy: An analysis of 114 definitions. *Resources, Conservation and Recycling* 127: 221–232.
- Kovanda, J. 2014. Incorporation of recycling flows into economy-wide material flow accounting and analysis: A case study for the Czech Republic. *Resources, Conservation and Recycling* 92(Supplement C): 78–84.

- Krausmann, F., A. Schaffartzik, A. Mayer, N. Eisenmenger, S. Gingrich, H. Haberl, and M. Fischer-Kowalski. 2016. Long-term trends in global material and energy use. In *Social Ecology: Society-Nature Relations across Time and Space*, edited by H. Haberl et al., Human-environment interactions, Vol. 5: 199–211. Switzerland: Springer.
- Krausmann, F., H. Schandl, N. Eisenmenger, S. Giljum, and T. Jackson. 2017a. Material flow accounting: Measuring global material use for sustainable development. *Annual Review of Environment and Resources* 42(1): 647–675.
- Krausmann, F., D. Wiedenhofer, C. Lauk, W. Haas, H. Tanikawa, T. Fishman, A. Miatto, H. Schandl, and H. Haberl. 2017b. Global socioeconomic material stocks rise 23-fold over the 20th century and require half of annual resource use. *Proceedings of the National Academy of Sciences*: 201613773. <http://www.pnas.org/content/114/8/1880>.
- Laner, D., H. Rechberger, and T. Astrup. 2014. Systematic evaluation of uncertainty in material flow analysis. *Journal of Industrial Ecology* 18(6): 859–870.
- Lauk, C., H. Haberl, K.-H. Erb, S. Gingrich, and F. Krausmann. 2012. Global socioeconomic carbon stocks in long-lived products 1900–2008. *Environmental Research Letters* 7(3): 034023.
- Lieder, M. and A. Rashid. 2016. Towards circular economy implementation: A comprehensive review in context of manufacturing industry. *Journal of Cleaner Production* 115: 36–51. <https://www.sciencedirect.com/science/article/pii/S0959652615018661?via%3Dihub>
- MacArthur, E. 2013. *Towards the circular economy: Economic and business rationale for an accelerated transition*. Cowes, UK: Ellen MacArthur Foundation.
- McDonough, W. and M. Braungart. 2003. *Cradle to cradle: Remaking the way we make things*. 1st ed. New York: Macmillan USA, June 2.
- Miatto, A., H. Schandl, and H. Tanikawa. 2017. How important are realistic building lifespan assumptions for material stock and demolition waste accounts? *Resources, Conservation and Recycling* 122: 143–154.
- Murray, A., K. Skene, and K. Haynes. 2017. The circular economy: An interdisciplinary exploration of the concept and application in a global context. *Journal of Business Ethics* 140(3): 369–380.
- Nicolli, F., M. Mazzanti, and V. Iafolla. 2012. Waste dynamics, country heterogeneity and European environmental policy effectiveness. *Journal of Environmental Policy & Planning* 14(4): 371–393.
- Nuss, P., G.A. Blengini, W. Haas, A. Mayer, V. Nita, and D. Pennington. 2017. *Development of a Sankey diagram of material flows in the EU economy based on Eurostat data*, EUR 28811 EN. JRC technical reports. Luxembourg: Publications Office of the European Union, November 7. <https://publications.europa.eu/en/publication-detail/-/publication/40dffbce-c439-11e7-9b01-01aa75ed71a1>. Accessed November 8, 2017.
- Nuss, P. and G.A. Blengini. 2018. Towards better monitoring of technology critical elements in Europe: Coupling of natural and anthropogenic cycles. *Science of the Total Environment* 613–614: 569–578.
- O'Neill, D. 2015. What should be held steady in a steady-state economy? Interpreting Daly's definition at the national level. *Journal of Industrial Ecology* 19(4): 552–563.
- Patrício, J., Y. Kalmykova, L. Rosado, and V. Lisovskaja. 2015. Uncertainty in material flow analysis indicators at different spatial levels. *Journal of Industrial Ecology* 19(5): 837–852.
- Pauliuk, S. 2018. Critical appraisal of the circular economy standard BS 8001: 2017 and a dashboard of quantitative system indicators for its implementation in organizations. *Resources, Conservation and Recycling* 129: 81–92.
- Pauliuk, S., T. Wang, and D.B. Müller. 2012. Moving toward the circular economy: The role of stocks in the Chinese steel cycle. *Environmental Science & Technology* 46(1): 148–154.
- Reck, B.K. and T.E. Graedel. 2012. Challenges in metal recycling. *Science* 337(6095): 690–695.
- Reck, B.K., D.B. Müller, K. Rostkowski, and T.E. Graedel. 2008. Anthropogenic nickel cycle: Insights into use, trade, and recycling. *Environmental Science & Technology* 42(9): 3394–3400.
- Saltelli, A., K. Chan, and E.M. Scott. 2000. *Sensitivity analysis*. Vol. 1. New York: Wiley.
- Schandl, H., M. Fischer-Kowalski, J. West, S. Giljum, M. Dittrich, N. Eisenmenger, A. Geschke, et al. 2017. Global material flows and resource productivity: Forty years of evidence: Global material flows and resource productivity. *Journal of Industrial Ecology*. <https://doi.org/10.1111/jiec.12626>. Accessed December 14, 2017.
- Schiller, G., K. Gruhler, and R. Ortler. 2017. Quantification of anthropogenic metabolism using spatially differentiated continuous MFA. *Change and Adaptation in Socio-Ecological Systems* 3(1): 119–132.
- Smith, P., H. Clark, H. Dong, E.A. Elsidig, H. Haberl, R. Harper, J. House, et al. 2014. Chapter 11: Agriculture, forestry and other land use (AFOLU). In *Climate Change 2014: Mitigation of Climate Change. IPCC Working Group III Contribution to AR5*. Cambridge, UK; New York: Cambridge University Press, November. www.ipcc.ch/pdf/assessment-report/ar5/wg3/ipcc_wg3_ar5_chapter11.pdf. Accessed April 10, 2017.
- Stahel, W.R. 2016. The circular economy. *Nature News* 531(7595): 435.
- Therond, O., M. Duru, J. Roger-Estrade, and G. Richard. 2017. A new analytical framework of farming system and agriculture model diversities: A review. *Agronomy for Sustainable Development* 37(3): 1–24.
- Tisserant, A., S. Pauliuk, S. Merciai, J. Schmidt, J. Fry, R. Wood, and A. Tukker. 2017. Solid waste and the circular economy: A global analysis of waste treatment and waste footprints: Global analysis of solid waste and waste footprint. *Journal of Industrial Ecology* 21(3): 628–640.
- UNEP. 2012. *GEO 5: global environment outlook: Environment for the future we want*. Nairobi: United Nations Environment Programme.
- Voet, E., van der, S. de Bruyn, and A. Tukker. 2009. *Environmental impact of the use of natural resources and products*. CML Report 184. Leiden University. www.leidenuniv.nl/cml/ssp/publications/eurostat_indicators_final_report_version_141009.pdf. Accessed August 14, 2018.
- Wang, T., D.B. Müller, and T.E. Graedel. 2007. Forging the anthropogenic iron cycle. *Environmental Science & Technology* 41(14): 5120–5129.
- Wiedenhofer, D., J.K. Steinberger, N. Eisenmenger, and W. Haas. 2015. Maintenance and expansion: Modeling material stocks and flows for residential buildings and transportation networks in the EU25: Stocks and flows in the EU25. *Journal of Industrial Ecology* 19(4): 538–551.
- Wiedmann, T.O., H. Schandl, M. Lenzen, D. Moran, S. Suh, J. West, and K. Kanemoto. 2015. The material footprint of nations. *Proceedings of the National Academy of Sciences* 112(20): 6271–6276.

- Wijkman, A. and K. Skanberg. 2015. The circular economy and benefits for society: Swedish case study shows jobs and climate as clear winners. *An Interim Report by the Club of Rome with Support from the MAVA Foundation and the Swedish Association of Recycling Industries*. <https://groenomstilling.erhvervsstyrelsen.dk/sites/default/files/media/final-version-swedish-study-13-04-15-till-tryck-ny.pdf>. Accessed August 14, 2018.
- Winans, K., A. Kendall, and H. Deng. 2017. The history and current applications of the circular economy concept. *Renewable and Sustainable Energy Reviews* 68: 825–833.
- Yuan, Z., J. Bi, and Y. Moriguchi. 2006. The circular economy: A new development strategy in China. *Journal of Industrial Ecology* 10(1–2): 4–8.
- Zanten, H.H.E.V., M. Herrero, O.V. Hal, E. Röss, A. Muller, T. Garnett, P.J. Gerber, C. Schader, and I.J.M.D. Boer. 2018. Defining a land boundary for sustainable livestock consumption. *Global Change Biology*. <https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.14321>. Accessed June 25, 2018.
- Zink, T. and R. Geyer. 2017. Circular economy rebound. *Journal of Industrial Ecology*. <https://doi.org/10.1111/jiec.12545>. Accessed March 3, 2017.

Supporting Information

Supporting information is linked to this article on the *JIE* website:

Supporting Information S1: This supporting information provides details of all assumptions, primary and secondary data sources, and various calculation factors referred to in the article.

Supplementary Data S2: This supporting information provides details of all data referred to in the article, along with a complete system description. Color coding in this SI file is as follows: green = data as used in figures; yellow = final circular economy system description; blue = calculation steps; red = raw data files as downloaded.