

On the third coefficient of TYZ expansion for radial scalar flat metrics

Original

On the third coefficient of TYZ expansion for radial scalar flat metrics / Loi, A.; Salis, F.; Zuddas, F.. - In: JOURNAL OF GEOMETRY AND PHYSICS. - ISSN 0393-0440. - STAMPA. - 133:(2018), pp. 210-218.
[10.1016/j.geomphys.2018.07.013]

Availability:

This version is available at: 11583/2862036 since: 2021-03-10T09:45:07Z

Publisher:

Elsevier B.V.

Published

DOI:10.1016/j.geomphys.2018.07.013

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

ON THE THIRD COEFFICIENT OF TYZ EXPANSION FOR RADIAL SCALAR FLAT METRICS

ANDREA LOI, FILIPPO SALIS, FABIO ZUDDAS

ABSTRACT. We classify radial scalar flat metrics with constant third coefficient of its TYZ expansion. As a byproduct of our analysis we provide a characterization of Simanca's scalar flat metric.

CONTENTS

1.	Introduction	1
2.	Radial metrics with constant a_1 e a_2	3
3.	Proof of Theorem 1.1 and Corollary 1.2	8
	References	12

1. INTRODUCTION

A Kähler metric g on a complex manifold M is said to be *projectively induced* if there exists a Kähler (isometric and holomorphic) immersion of (M, g) into the complex projective space $(\mathbb{C}P^N, g_{FS})$, $N \leq +\infty$, endowed with the Fubini–Study metric g_{FS} , namely the metric whose associated Kähler form is given in homogeneous coordinates by $\omega_{FS} = \frac{i}{2} \partial \bar{\partial} \log(|Z_0|^2 + \dots + |Z_N|^2)$. Since requirement that a Kähler metric is projectively induced is a somehow strong assumption, one could try to approximate an integral Kähler form on a complex manifold with suitable normalized projectively induced Kähler forms through the following construction.

Assume that there exists a Hermitian line bundle (L, h) over M whose Ricci curvature $\text{Ric}(h)$ equals ω (this is always possible in the compact case). For every positive integer m one considers the holomorphic line bundle $L^m = L^{\otimes m}$ endowed with the Hermitian metric h_m induced on L^m by h , such that $\text{Ric}(h_m) = m\omega$.

2010 *Mathematics Subject Classification.* 53C55; 58C25; 58F06.

Key words and phrases. Kähler manifolds; TYZ asymptotic expansion; projectively induced metrics.

The first author was supported by Prin 2015 – Real and Complex Manifolds; Geometry, Topology and Harmonic Analysis – Italy by INdAM. GNSAGA - Gruppo Nazionale per le Strutture Algebriche, Geometriche e le loro Applicazioni and by Fondazione di Sardegna and Regione Autonoma della Sardegna.

Let \mathcal{H}_m be the separable complex Hilbert space consisting of global holomorphic sections of L^m such that

$$\langle s, s \rangle_m = \int_M h_m(s, s) \frac{\omega^n}{n!} < \infty.$$

If $s_j, j = 0, \dots, N_m, N_m + 1 = \dim \mathcal{H}_m \leq \infty$ is an orthonormal basis of \mathcal{H}_m , the smooth function

$$\epsilon_{mg}(x) = \sum_{j=0}^{N_m} h_m(s_j(x), s_j(x))$$

is globally defined on M and, as suggest by the notation, it does not depend on the orthonormal basis or the Hermitian metric chosen (see e.g. [5]). This function, also known in literature as *distortion function*, constitutes a tool to evaluate how a Kähler metric differs from being projectively induced. Indeed, if there exists a sufficiently large integer m such that, for every point of M , we can find at least one element of an orthonormal basis $\{s_j\}_{j=0, \dots, N_m}$ of \mathcal{H}_m which does not vanishes at this point (the free based point condition in the compact case), *coherent state map*, namely the holomorphic map

$$\begin{aligned} \varphi_m : M &\rightarrow \mathbb{C}P^{N_m} \\ x &\mapsto [s_0(x), \dots, s_{N_m}(x)], \end{aligned}$$

is well-defined and, moreover (see, e.g. [1] for a proof), it satisfies

$$\varphi_m^* \omega_{FS} = m\omega + \frac{i}{2} \partial \bar{\partial} \log \epsilon_{mg}.$$

Although not all Kähler metrics are projectively induced, Tian ([24]) and Ruan ([22]) solved a conjectured by Yau by proving that any polarized metric on a compact complex manifold is the C^∞ -limit of normalized projectively induced metrics $\frac{\varphi_m^* \omega_{FS}}{m}$. Then, Catlin ([8]) and Zelditch ([27]) independently generalized the Tian-Ruan theorem by proving the existence of a complete asymptotic expansion for the distortion function related to any polarized metric defined on compact complex manifold. This asymptotic expansion

$$\epsilon_{mg}(x) \sim \sum_{j=0}^{\infty} a_j(x) m^{n-j}, \tag{1}$$

is called *Tian-Yau-Zelditch expansion* and it means that there exists a positive constant $C_{l,r}$ depending on two positive integer constants l and r such that

$$\left\| \epsilon_{mg}(x) - \sum_{j=0}^l a_j(x) m^{n-j} \right\|_{C^r} \leq \frac{C_{l,r}}{m^{l+1}},$$

where $a_0(x) = 1$ and $a_j(x), j = 1, \dots$ are smooth functions on M . In particular, Z. Lu [20] computed the first three TYZ coefficients. The expression of the first two

coefficients are (for the rather involved expression of third coefficient a_3 see (13) below):

$$\begin{cases} a_1(x) = \frac{1}{2}\rho \\ a_2(x) = \frac{1}{3}\Delta\rho + \frac{1}{24}(|R|^2 - 4|\text{Ric}|^2 + 3\rho^2) \end{cases} \quad (2)$$

where ρ , R , Ric denote respectively the scalar curvature, the curvature tensor and the Ricci tensor of (M, g) ,

Due to Donaldson’s work (cf. [9], [10] and [1]) in the compact case and respectively to the theory of quantization in the noncompact case (see e.g. [3], [6] and [7]), it is natural to study metrics with the Tian-Yau-Zelditch coefficients being prescribed. For instance, the vanishing of this coefficients for large enough indexes turns out to be related to some important problems in the theory of pseudoconvex manifolds (cf. [21], [2]). Furthermore, in the noncompact case, one can find in [19] a characterization of the flat metric as a Taub-NUT metric with $a_3 = 0$, while Z. Feng and Z. Tu [13] solve a conjecture formulated in [26] by showing that the complex hyperbolic space is the only Cartan-Hartogs domain where the coefficient a_2 is constant. In [18] A. Loi and M. Zedda prove that a locally hermitian symmetric space with vanishing a_1 and a_2 is flat.

The present paper deals with *radial* Kähler metrics, namely those Kähler metrics admitting a Kähler potential which depends only on the sum $|z|^2 = |z_1|^2 + \dots + |z_n|^2$ of the moduli of a local coordinates’ system z_i .

Our main results are the following

Theorem 1.1. *The third Tian-Yau-Zelditch coefficient a_3 of a radial Kähler metric with constant scalar curvature (cscK metric) is constant if and only if the second Tian-Yau-Zelditch coefficient a_2 is constant.*

Corollary 1.2. *The flat metric and the Simanca metric are the only radial projectively induced metrics with $a_1 = a_3 = 0$.*

The proof of Theorem 1.1 is based on Feng’s work [12] which classifies the radial metrics whose first and second coefficients of TYZ expansion are constant functions. Since the proof of Feng’s classification theorem given in [12] is quite technical and not so easy to read we have provided in the next section a more readable (to the authors’ opinion) proof. In the last section we prove Theorem 1.1 and Corollary 1.2.

2. RADIAL METRICS WITH CONSTANT a_1 E a_2

Theorem 2.1 (Z. Feng [12]). *The only radial Kähler potentials defined on a complex domain of dimension n which have constant first and second TYZ coefficient are:*

1. *the Euclidean metric,*
2. *constant multiple of the hyperbolic metric defined on $\mathcal{B}_{\frac{1}{\lambda}}^n$,*
3. *constant multiple of the Fubini-Study metric.*

Moreover, if $n = 2$

4. $|z|^2 + \lambda \log |z|^2$ on $\mathbb{C}^2 \setminus \{0\}$,
5. $\mu|z|^2 - \lambda \log |z|^2$ on $\mathbb{C}^2 \setminus \overline{\mathcal{B}}_{\frac{1}{\mu}}^2$,
6. $-\mu(\log(1 - \xi|z|^{2\zeta}) + \frac{1-\zeta}{2} \log |z|^2)$ on $\mathcal{B}_{(\frac{1}{\xi})^{\frac{1}{\zeta}}}^2 \setminus \overline{\mathcal{B}}_{((\frac{1}{\xi})(\frac{1-\zeta}{1+\zeta}))^{\frac{1}{\zeta}}}^2$,
7. $-\mu(\log(1 - \xi|z|^{2(\lambda+1)}) - \frac{\lambda}{2} \log |z|^2)$ on $\mathcal{B}_{(\frac{1}{\xi})^{\frac{1}{\lambda+1}}}^2 \setminus \{0\}$,
8. $\mu(\log(1 + \xi|z|^{-2(\lambda+1)}) + \frac{2+\lambda}{2} \log |z|^2)$ on $\mathbb{C}^2 \setminus \mathcal{B}_{((\frac{1}{\xi})(\frac{\lambda}{2+\lambda}))^{\frac{1}{\lambda+1}}}^2$,
9. $\mu(\log(1 + |z|^{-2\zeta}) + \frac{1+\zeta}{2} \log |z|^2)$ on $\mathbb{C}^2 \setminus \{0\}$,
- 10a. $-\mu(\log(-\log |z|^2 + \kappa) + \frac{\log |z|^2}{2})$ on $\mathcal{B}_{e^\kappa}^2 \setminus \overline{\mathcal{B}}_{e^{\kappa-2}}^2$,
- 11a. $-\mu(\log |\cos(\lambda \log |z|^2 + \kappa)| + \frac{\log |z|^2}{2})$ on $\mathcal{B}_{r_1(h, \kappa, \lambda)}^2 \setminus \overline{\mathcal{B}}_{r_3(h, \kappa, \lambda)}^2$,

or if $n = 1$

- 10b. $-\mu \log |-\log |z|^2 + \kappa|$ on $\mathbb{C} \setminus (\partial \mathcal{B}_{e^\kappa} \cup \{0\})$,
- 11b. $-\mu \log |\cos(\lambda \log |z|^2 + \kappa)|$, on $\mathcal{B}_{r_1(h, \kappa, \lambda)} \setminus \overline{\mathcal{B}}_{r_2(h, \kappa, \lambda)}$.

Where of \mathcal{B}_r^n denotes the ball of radius r in \mathbb{C}^n , $\partial \mathcal{B}_r^n$ denotes the boundary of \mathcal{B}_r^n , $\overline{\mathcal{B}}_r^n = \mathcal{B}_r^n \cup \partial \mathcal{B}_r^n$, $\mu, \lambda, \xi, \zeta \in \mathbb{R}^+$, $0 < \zeta < 1$, $\kappa \in \mathbb{R}$, $h \in \mathbb{Z}$, $r_1(h, \kappa) = e^{\frac{1}{\lambda}(\frac{2h+1}{2}\pi - \kappa)}$, $r_2(h, \kappa, \lambda) = e^{\frac{1}{\lambda}(\frac{2h-1}{2}\pi - \kappa)}$ and $r_3(h, \kappa, \lambda) = e^{\frac{1}{\lambda}(h\pi + \arctan(\frac{1}{2\lambda}) - \kappa)}$.

Proof. Let $\Phi(\log r)$ be a Kähler potential defined on a radial complex domain of complex dimension n , where $r = |z_1|^2 + \dots + |z_n|^2$.

Firstly, we classify Kähler metrics related to such kind of potentials which have constant scalar curvature (namely a_1 is constant). By definition, the metric tensor reads as

$$g_{i\bar{j}} = \frac{\partial^2 \Phi(\log r)}{\partial z_i \partial \bar{z}_j} = \frac{\Phi'' - \Phi'}{r^2} \bar{z}_i z_j + \frac{\Phi'}{r} \delta_{ij}, \quad (3)$$

where δ_{ij} is the Kronecker delta and Φ' represents the first derivative of Φ with respect to

$$t = \log r.$$

Since Riemannian metrics are positive definite, Φ'' needs to be a positive function. Hence we can consider the following substitutions

$$\begin{cases} y = \Phi'(t), \\ \psi(y) = \Phi''(t). \end{cases}$$

From eq. (3), we easily get

$$\det(g_{i\bar{j}}) = \frac{(\Phi')^{n-1}\Phi''}{r^n}.$$

By considering that

$$\frac{d}{dt}((n-1)\log\Phi' + \log\Phi'') = (n-1)\frac{\Phi''}{\Phi'} + \frac{\Phi'''}{\Phi''} = (n-1)\frac{\psi}{y} + \psi',$$

we introduce the further substitution

$$\sigma(y) = \frac{(y^{n-1}\psi)'}{y^{n-1}} = (n-1)\frac{\psi(y)}{y} + \psi'(y),$$

and we compute the Ricci tensor's components:

$$\text{Ric}_{i\bar{j}} = -\frac{\partial^2 \log \det(g_{i\bar{j}})}{\partial z_i \partial \bar{z}_j} = \frac{-\sigma'\psi + \sigma - n}{r^2} \bar{z}_i z_j + \frac{n - \sigma}{r} \delta_{ij}, \quad (4)$$

to be in the position to represent the scalar curvature as a function of y :

$$\rho = g^{i\bar{j}} \text{Ric}_{i\bar{j}} = \frac{n(n-1)}{y} - \frac{(n-1)\sigma + \sigma'y}{y}.$$

Thus we reach our initial objective by solving the ODE obtained by imposing ρ to be constant, namely

$$\frac{n(n-1)}{y} - \frac{(y^{n-1}\psi)''}{y^{n-1}} = -An(n+1),$$

whose solutions are

$$\psi(y) = Ay^2 + y + \frac{B}{y^{n-2}} + \frac{C}{y^{n-1}}, \quad (5)$$

where $A, B, C \in \mathbb{R}$.

Now, we determinate which conditions have to be satisfied so that the previous metrics also verify the PDE

$$a_2 = K, \quad (6)$$

where K is a real constant.

We are going to represent the first term in the previous PDE (see eq. 2) as a function of y , $\psi(y)$ and its derivatives, in order to convert it to an ODE.

Riemann tensor's components $R_{i\bar{j}k\bar{l}}$ are by definition equal to $\frac{\partial^2 g_{i\bar{l}}}{\partial z_k \partial \bar{z}_j} - g^{p\bar{q}} \frac{\partial g_{i\bar{p}}}{\partial z_k} \frac{\partial g_{q\bar{l}}}{\partial \bar{z}_j}$. Therefore, the following derivatives (cf. eq. 3)

$$\begin{aligned} \frac{\partial g_{i\bar{l}}}{\partial z_k} &= \frac{\Phi'' - \Phi'}{r^2} (\delta_{kl} \bar{z}_i + \delta_{il} \bar{z}_k) + \frac{\Phi''' - 3\Phi'' + 2\Phi'}{r^3} \bar{z}_i z_l \bar{z}_k \\ \frac{\partial^2 g_{i\bar{l}}}{\partial z_k \partial \bar{z}_j} &= \frac{\Phi'' - \Phi'}{r^2} (\delta_{kl} \delta_{ij} + \delta_{il} \delta_{kj}) + \frac{\Phi'''' - 6\Phi''' + 11\Phi'' - 6\Phi'}{r^4} \bar{z}_i z_j z_l \bar{z}_k \\ &\quad + \frac{\Phi''' - 3\Phi'' + 2\Phi'}{r^3} (\delta_{kj} \bar{z}_i z_l + \delta_{kl} z_j \bar{z}_i + \delta_{il} z_j \bar{z}_k + \delta_{ij} z_l \bar{z}_k) \end{aligned} \quad (7)$$

allow us to state that the unique (up to consider tensor's symmetries) nonvanishing components in $(z_1, 0, \dots, 0)$ are equal to

$$\begin{aligned} R_{1\bar{1}\bar{1}\bar{1}} &= \frac{\psi'' \psi^2}{r^2}, \\ R_{1\bar{1}\bar{i}\bar{i}} &= \frac{\psi' y - \psi}{y r^2} \psi, \\ R_{i\bar{i}\bar{i}\bar{i}} &= 2R_{i\bar{i}j\bar{j}} = 2 \frac{\psi - y}{r^2}, \end{aligned} \quad (8)$$

where $i \neq j$ and $i, j \neq 1$. By taking into account the invariance of $|R|^2$ under the action of the unitary group, we can use Riemann tensor's components of formula (8) and the metric tensor's components evaluated in $(z_1, 0, \dots, 0)$ (cf. eq. 3) to get the general formula

$$\begin{aligned} |R|^2 &= (g^{1\bar{1}})^4 (R_{1\bar{1}\bar{1}\bar{1}})^2 + 4(n-1)(g^{1\bar{1}})^2 (g^{i\bar{i}})^2 (R_{1\bar{1}\bar{i}\bar{i}})^2 \\ &\quad + (g^{i\bar{i}})^4 \left(4 \frac{(n-2)(n-1)}{2} (R_{i\bar{i}j\bar{j}})^2 + (n-1) (R_{i\bar{i}\bar{i}\bar{i}})^2 \right) \\ &= (\psi'')^2 + 4(n-1) \left(\frac{\psi' y - \psi}{y^2} \right)^2 + 2n(n-1) \left(\frac{\psi - y}{y^2} \right)^2. \end{aligned} \quad (9)$$

Similarly, we compute

$$\begin{aligned} |Ric|^2 &= (g^{1\bar{1}})^2 (Ric_{1\bar{1}})^2 + (n-1)(g^{i\bar{i}})^2 (Ric_{i\bar{i}})^2 \\ &= (\sigma')^2 + (n-1) \left(\frac{\sigma - n}{y} \right)^2. \end{aligned} \quad (10)$$

Therefore the equation (6) is equivalent to

$$\begin{aligned} 4(n-1) \left(\frac{\psi' y - \psi}{y^2} \right)^2 + 2n(n-1) \left(\frac{\psi - y}{y^2} \right)^2 \\ - 4 \left((\sigma')^2 + (n-1) \left(\frac{\sigma - n}{y} \right)^2 \right) + (\psi'')^2 \\ = 24K - 3A^2 n^2 (n+1)^2. \end{aligned}$$

By imposing in the previous ODE, ψ to be equal to the solution (5), we get

$$\begin{aligned} (n-1) \left(2BCn^2 y^{-2n-1} (n+1) + C^2 n y^{-2(n+1)} (n+1)(n+2) \right. \\ \left. + B^2 n y^{-2n} (n+1)(n-2) + A^2 n (n+1)(3n+2) \right) - 24K = 0. \end{aligned}$$

Hence, the previous equation is satisfied if and only if

$$\begin{cases} C = B = 0, K = A^2 n (n+1)(n-1)(3n+2)/24 & \text{if } n \geq 2 \\ C = 0, K = 2A^2 & \text{if } n = 2 \\ K = 0 & \text{if } n = 1 \end{cases}$$

By definition of ψ and y and by renaming the constants for more convenience, one gets that $a_2 = K$ if and only if

$$\Phi'' = \begin{cases} a(\Phi')^2 + \Phi' & \text{if } n \geq 2, \\ a(\Phi')^2 + \Phi' + c & \text{if } n = 2, \\ a(\Phi')^2 + b\Phi' + c & \text{if } n = 1. \end{cases} \quad (11)$$

In order to solve this equation let us first suppose $a \neq 0$. Therefore

$$\Phi''(t) = a \left(\left(\Phi' + \frac{b}{2a} \right)^2 + \frac{c}{a} - \frac{b^2}{4a^2} \right). \quad (12)$$

We distinguish now three different cases.

Case 1: $\frac{c}{a} - \frac{b^2}{4a^2} = -D^2$, where $D \in \mathbb{R}^+$. By solving this ODE, we get

$$\left| \frac{\Phi'(t) + \frac{b}{2a} - D}{\Phi'(t) + \frac{b}{2a} + D} \right| = \xi e^{2aDt},$$

where $\xi \in \mathbb{R}^+$. Thus we have to distinguish two further different possibilities.

1a) $\Phi' + \frac{b}{2a} - D < 0 < \Phi' + \frac{b}{2a} + D$.

Hence

$$\Phi'(t) = -\frac{\xi e^{2aDt}}{\xi e^{2aDt} + 1} \left(D + \frac{b}{2a} \right) + \frac{D - \frac{b}{2a}}{\xi e^{2aDt} + 1},$$

namely

$$\Phi(\log r) = -\frac{1}{a} \left(\log(1 + \xi r^{2aD}) + \left(\frac{b}{2} - Da \right) \log r \right).$$

Since $\Phi'' > 0$, then $a < 0$. If $n \geq 2$, also Φ' has to be positive by definition of Kähler potential. This condition is trivially satisfied if $n > 2$, while if $n = 2$ it is equivalent to $(2aD + 1)\xi r^{2aD} > 2aD - 1$. Moreover, if $n = 1$, we recall that $\partial\bar{\partial} \log r = 0$.

1b) $\Phi' + \frac{b}{2a} - D > 0$ or $\Phi' + \frac{b}{2a} + D < 0$. Therefore

$$\Phi'(t) = -\frac{\xi e^{2aDt}}{\xi e^{2aDt} - 1} \left(D + \frac{b}{2a} \right) - \frac{D - \frac{b}{2a}}{\xi e^{2aDt} - 1},$$

namely

$$\Phi(\log r) = -\frac{1}{a} \left(\log |1 - \xi r^{2aD}| + \left(\frac{b}{2} - Da \right) \log r \right).$$

Since $\Phi'' > 0$, then $a > 0$. If $n \geq 2$, also Φ' has to be positive. This condition is satisfied if and only if $\xi r^{2aD} < 1$ and $\xi r^{2aD} > \frac{1-2aD}{2aD+1}$ if $n = 2$.

Case 2: $\frac{c}{a} - \frac{b^2}{4a^2} = 0$. This case may occur only if $n = 1$ or $n = 2$. By solving the initial ODE (12), we get

$$\Phi(\log r) = -\frac{1}{a} \log |\log r + \kappa| - \frac{b}{2a} \log r,$$

where $\kappa \in \mathbb{R}$. Since $\Phi'' > 0$, then $a > 0$. If $n = 2$, also Φ' has to be positive, hence $-2 - \kappa < \log r < -\kappa$ (in this case $b = 1$).

Case 3: $\frac{c}{a} - \frac{b^2}{4a^2} = D^2$. Also this case may occur only if $n = 1$ or $n = 2$. By solving the initial ODE (12), we get

$$\Phi(\log r) = -\frac{1}{a} \log |\cos(aD \log r + \kappa)| - \frac{b}{2a} \log r,$$

where $\kappa \in \mathbb{R}$. Since $\Phi'' > 0$, then $a > 0$. If $n = 2$, also Φ' has to be positive, hence $\frac{1}{aD}(\arctan(\frac{1}{2aD}) + h\pi - \kappa) < \log r < \frac{1}{aD}(\frac{\pi}{2} + h\pi - \kappa)$, where $h \in \mathbb{Z}$.

To conclude we have to consider that a may also be equal to 0, hence we have to solve

$$\Phi''(t) = b\Phi' + c.$$

If $b \neq 0$, we get

$$\Phi(\log r) = \frac{1}{b^2} \xi r^b - \frac{c}{b} \log r.$$

If $n \geq 2$, Φ' has to be positive, hence $\xi r^b > \frac{c}{b}$. This condition is always satisfied if $n > 2$, since $c = 0$. Moreover Φ'' is always positive.

Instead, if $b = 0$ (this case may occur only if $n = 1$), we get

$$\Phi(\log r) = \frac{c}{2} (\log r)^2 + \kappa \log r,$$

which is the Euclidean metric (already considered before). \square

3. PROOF OF THEOREM 1.1 AND COROLLARY 1.2

In order to prove Theorem 1.1 we recall the expression of the term a_3 in Tian-Yau-Zelditch expansion:

$$\begin{aligned} a_3(x) &= \frac{1}{8} \Delta \Delta \rho + \frac{1}{24} \operatorname{div} \operatorname{div}(R, \operatorname{Ric}) - \frac{1}{6} \operatorname{div} \operatorname{div}(\rho \operatorname{Ric}) \\ &\quad + \frac{1}{48} \Delta(|R|^2 - 4|\operatorname{Ric}|^2 + 8\rho^2) + \frac{1}{48} \rho(\rho^2 - 4|\operatorname{Ric}|^2 + |R|^2) \\ &\quad + \frac{1}{24} (\sigma_3(\operatorname{Ric}) - \operatorname{Ric}(R, R) - R(\operatorname{Ric}, \operatorname{Ric})), \end{aligned} \quad (13)$$

where we are using the following notations (in local coordinates z_1, \dots, z_n):

$$\begin{aligned}
|D' \rho|^2 &= g^{j\bar{i}} \frac{\partial \rho}{\partial z_i} \frac{\partial \rho}{\partial \bar{z}_j}, \\
|D' \text{Ric}|^2 &= g^{\alpha\bar{i}} g^{j\bar{\beta}} g^{\gamma\bar{k}} \text{Ric}_{i\bar{j},k} \overline{\text{Ric}}_{\alpha\bar{\beta},\gamma}, \\
|D' R|^2 &= g^{\alpha\bar{i}} g^{j\bar{\beta}} g^{\gamma\bar{k}} g^{l\bar{\delta}} g^{\epsilon\bar{p}} R_{i\bar{j}k\bar{l},p} \overline{R}_{\alpha\bar{\beta}\gamma\bar{\delta},\epsilon}, \\
\text{divdiv}(\rho \text{Ric}) &= 2|D' \rho|^2 + g^{\beta\bar{i}} g^{j\bar{\alpha}} \text{Ric}_{i\bar{j}} \frac{\partial^2 \rho}{\partial z_\alpha \partial \bar{z}_\beta} + \rho \Delta \rho, \\
\text{divdiv}(R, \text{Ric}) &= -g^{\beta\bar{i}} g^{j\bar{\alpha}} \text{Ric}_{i\bar{j}} \frac{\partial^2 \rho}{\partial z_\alpha \partial \bar{z}_\beta} - 2|D' \text{Ric}|^2 \\
&\quad + g^{\alpha\bar{i}} g^{j\bar{\beta}} g^{\gamma\bar{k}} g^{l\bar{\delta}} R_{i\bar{j},k\bar{l}} R_{\beta\bar{\alpha}\delta\bar{\gamma}} - R(\text{Ric}, \text{Ric}) - \sigma_3(\text{Ric}), \\
R(\text{Ric}, \text{Ric}) &= g^{\alpha\bar{i}} g^{j\bar{\beta}} g^{\gamma\bar{k}} g^{l\bar{\delta}} R_{i\bar{j}k\bar{l}} \text{Ric}_{\beta\bar{\alpha}} \text{Ric}_{\delta\bar{\gamma}}, \\
\text{Ric}(R, R) &= g^{\alpha\bar{i}} g^{j\bar{\beta}} g^{\gamma\bar{k}} g^{\delta\bar{p}} g^{q\bar{\epsilon}} \text{Ric}_{i\bar{j}} R_{\beta\bar{\gamma}p\bar{q}} R_{k\bar{\alpha}\epsilon\bar{\delta}}, \\
\sigma_3(\text{Ric}) &= g^{\delta\bar{i}} g^{j\bar{\alpha}} g^{\beta\bar{\gamma}} \text{Ric}_{i\bar{j}} \text{Ric}_{\alpha\bar{\beta}} \text{Ric}_{\gamma\bar{\delta}},
\end{aligned}$$

where the $g^{j\bar{i}}$'s denote the entries of the inverse matrix of the metric (i.e. $g_{k\bar{i}} g^{j\bar{i}} = \delta_{kj}$) and “,p” represents the covariant derivative in the direction $\frac{\partial}{\partial z_p}$ and we are using the summation convention for repeated indices. The reader is also referred to [14] and [15] for a recursive formula for the coefficients a_j 's and an alternative computation of a_j for $j \leq 3$ using Calabi's diastasis function (see also [25] for a graph-theoretic interpretation of this recursive formula). Moreover note that, given any Kähler manifold (M, g) it makes sense to call the a_j 's Tian-Yau-Zelditch coefficients, regardless of the existence of Tian-Yau-Zelditch expansion.

Proof of Theorem 1.1. The same arguments about unitary invariance of terms of a_2 shown in the proof of Theor. 2.1 hold true also in this case for the term a_3 . Therefore, we can easily compute by using eq. (8) and (3) evaluated at $(z_1, 0, \dots, 0)$ the following terms:

$$\begin{aligned}
\text{Ric}(R, R) &= (g^{1\bar{1}})^2 \text{Ric}_{1\bar{1}} ((g^{1\bar{1}})^3 (R_{1\bar{1}1\bar{1}})^2 + 2(n-1)g^{1\bar{1}}(g^{i\bar{i}})^2 (R_{1\bar{1}i\bar{i}})^2) \\
&\quad + (g^{i\bar{i}})^5 \text{Ric}_{i\bar{i}} ((n-1)(R_{i\bar{i}i\bar{i}})^2 + 2(n-1)(n-2)(R_{i\bar{i}j\bar{j}})^2) \\
&\quad + 2(n-1)(g^{1\bar{1}})^2 (g^{i\bar{i}})^3 \text{Ric}_{i\bar{i}} (R_{1\bar{1}i\bar{i}})^2 \\
&= 2(n-1) \frac{n-\sigma}{y^5} \left(n(\psi-y)^2 + (\psi'y - \psi)^2 \right) \\
&\quad - \sigma' \left((\psi'')^2 + 2(n-1) \frac{(\psi'y - \psi)^2}{y^4} \right);
\end{aligned} \tag{14}$$

$$\begin{aligned}
R(\text{Ric}, \text{Ric}) &= (g^{1\bar{1}})^4 (\text{Ric}_{1\bar{1}})^2 R_{1\bar{1}1\bar{1}} + 2(n-1)(g^{1\bar{1}})^2 (g^{i\bar{i}})^2 \text{Ric}_{1\bar{1}} \text{Ric}_{i\bar{i}} R_{1\bar{1}i\bar{i}} \\
&\quad + (g^{i\bar{i}})^4 (\text{Ric}_{i\bar{i}})^2 ((n-1)R_{i\bar{i}i\bar{i}} + (n-1)(n-2)R_{i\bar{i}j\bar{j}}) \\
&= (\sigma')^2 \psi'' - 2(n-1) \frac{(\psi'y - \psi)(n-\sigma)\sigma'}{y^3} + n(n-1) \frac{(\psi-y)(n-\sigma)^2}{y^4}.
\end{aligned} \tag{15}$$

By using formulas (9) and (10), we define

$$\chi(y) = |R|^2 - 4|\text{Ric}|^2,$$

to get

$$\begin{aligned}\Delta(|R|^2 - 4|\text{Ric}|^2) &= g^{i\bar{j}} \frac{\partial^2 \chi}{\partial z_j \partial \bar{z}_i} \\ &= g^{i\bar{j}} \left(\frac{\chi' \psi}{r} \delta_{ji} + \frac{\chi'' \psi + \chi' \psi' - \chi'}{r^2} \psi \bar{z}_j z_i \right) = (\chi' \psi)' + \frac{n-1}{y} \chi' \psi.\end{aligned}\quad (16)$$

Since Christoffel's symbols Γ_{ki}^p are equal to $g^{p\bar{q}} \frac{\partial g_{i\bar{q}}}{\partial z_k}$, we easily deduce from (7) that they are all equal to zero when evaluated at $(z_1, 0, \dots, 0)$ except for:

$$\begin{aligned}\Gamma_{11}^1 &= \left(\frac{\Phi'' - \Phi'}{r^2} \right) \left(\frac{r}{\Phi''} \right) \bar{z}_1 = \frac{\psi' - 1}{r} \bar{z}_1, \\ \Gamma_{1i}^i &= \left(\frac{\Phi'' - \Phi'}{r^2} \right) \left(\frac{r}{\Phi'} \right) \bar{z}_1 = \frac{\psi - y}{yr} \bar{z}_1.\end{aligned}$$

We also compute that the unique first derivatives of Ricci tensor's components different from zero in $(z_1, 0, \dots, 0)$ (cf. (4)) are:

$$\begin{aligned}\frac{\partial}{\partial \bar{z}_1} Ric_{1\bar{1}} &= \frac{\sigma' \psi - (\sigma' \psi)' \psi}{r^2} z_1, \\ \frac{\partial}{\partial \bar{z}_1} Ric_{i\bar{i}} &= \frac{\partial}{\partial \bar{z}_i} Ric_{i\bar{1}} = \frac{\sigma - \sigma' \psi - n}{r^2} z_1,\end{aligned}$$

and we use them to evaluate Ricci tensor's first covariant derivatives at $(z_1, 0, \dots, 0)$, which are defined as $Ric_{i\bar{j},k} = \frac{\partial}{\partial z_k} Ric_{i\bar{j}} - Ric_{p\bar{j}} \Gamma_{ki}^p$, and we get

$$\begin{aligned}|D' \text{Ric}|^2 &= (g^{1\bar{1}})^3 |Ric_{1\bar{1},1}|^2 + 2(n-1)g^{1\bar{1}}(g^{i\bar{i}})^2 |Ric_{i\bar{i},1}|^2 \\ &= (\sigma'')^2 \psi + 2(n-1) \frac{\psi}{y^2} (\sigma' + \frac{n-\sigma}{y})^2.\end{aligned}\quad (17)$$

Because some Riemann tensor's components in $(z_1, 0, \dots, 0)$ are equal to zero (cf. eq. (8)), we need to compute just some Ricci tensor's second covariant derivatives, of which we recall the definition: $Ric_{i\bar{j},k\bar{l}} = \partial_{\bar{l}} \partial_k Ric_{i\bar{j}} + \Gamma_{ki}^q \Gamma_{\bar{l}\bar{j}}^{\bar{p}} Ric_{q\bar{p}} - \Gamma_{ki}^p \partial_{\bar{l}} Ric_{p\bar{j}} - \partial_{\bar{l}} \Gamma_{ki}^p Ric_{p\bar{j}} - \Gamma_{\bar{l}\bar{j}}^{\bar{p}} \partial_k Ric_{i\bar{p}}$. In particular, we preliminary evaluate at $(z_1, 0, \dots, 0)$ the following derivatives $(\partial_{\bar{l}} \Gamma_{ki}^p = g^{p\bar{q}} \frac{\partial^2 g_{i\bar{q}}}{\partial \bar{z}_l \partial z_k} - g^{p\bar{\gamma}} g^{\alpha\bar{q}} \frac{\partial g_{\alpha\bar{\gamma}}}{\partial \bar{z}_l} \frac{\partial g_{i\bar{q}}}{\partial z_k})$

$$\begin{aligned}\partial_{\bar{1}} \Gamma_{11}^1 &= \frac{\psi'' \psi}{r}, \\ \partial_{\bar{i}} \Gamma_{i1}^1 &= \frac{\psi' y - \psi}{yr}, \\ \partial_{\bar{1}} \Gamma_{i1}^i &= \frac{\psi' y - \psi}{y^2 r} \psi, \\ \partial_{\bar{i}} \Gamma_{ii}^i &= 2 \partial_{\bar{j}} \Gamma_{ji}^i = 2 \frac{\psi - y}{yr}\end{aligned}$$

and

$$\begin{aligned}\frac{\partial^2}{\partial z_1 \partial \bar{z}_1} Ric_{1\bar{1}} &= \frac{-((\sigma' \psi)' \psi)' \psi + 2(\sigma' \psi)' \psi - \sigma' \psi}{r^2}, \\ \frac{\partial^2}{\partial z_1 \partial \bar{z}_1} Ric_{i\bar{i}} &= \frac{\partial^2}{\partial z_i \partial \bar{z}_1} Ric_{1\bar{i}} = \frac{-(\sigma' \psi)' \psi + 2\sigma' \psi - \sigma + n}{r^2}, \\ \frac{\partial^2}{\partial z_i \partial \bar{z}_i} Ric_{i\bar{i}} &= 2 \frac{\partial^2}{\partial z_j \partial \bar{z}_j} Ric_{i\bar{i}} = 2 \frac{\partial^2}{\partial z_j \partial \bar{z}_i} Ric_{i\bar{j}} = 2 \frac{-\sigma' \psi + \sigma - n}{r^2},\end{aligned}$$

to be in the position to express

$$\begin{aligned}
 & (g^{1\bar{1}})^4 R_{1\bar{1}1\bar{1}} Ric_{1\bar{1},1\bar{1}} + 2(n-1)(g^{1\bar{1}})^2 (g^{i\bar{i}})^2 R_{1\bar{1}i\bar{i}} (Ric_{1\bar{1},i\bar{i}} + Ric_{i\bar{i},1\bar{1}}) \\
 & \quad + (g^{i\bar{i}})^4 ((n-1)R_{i\bar{i}i\bar{i}} Ric_{i\bar{i},i\bar{i}} + 2(n-1)(n-2)R_{i\bar{i}j\bar{j}} Ric_{i\bar{i},j\bar{j}}) \\
 & = \frac{\psi''}{\psi} \left(-((\sigma'\psi)'\psi)' - (\psi')^2 \sigma' + 2\psi'(\sigma'\psi)' + \sigma'\psi''\psi \right) \\
 & \quad + 2(n-1) \frac{\psi' y - \psi}{y^3} \left(-2(\sigma'\psi)' + 2 \frac{\psi}{y^2} (n - \sigma) + 4 \frac{\psi}{y} \sigma' \right) \\
 & \quad + 2(n-1) \frac{\psi' y - \psi}{y^3} \left(\psi' - \frac{\psi}{y} \right) \left(\sigma' + \frac{\sigma - n}{y} \right) \\
 & \quad + 2n \frac{\psi}{y^5} (n-1)(\psi - y)(\sigma - n - \sigma' y)
 \end{aligned} \tag{18}$$

as a function of y , ψ and its derivatives.

Therefore, thanks to the formulas (9), (10), (15), (14), (16), (17) and (18), using (5), we convert the PDE

$$a_3 = K, \tag{19}$$

where K is a real constant, to the identity:

$$\begin{aligned}
 & n(n-1)(n+1) \left(A^3 n y^{3n+3} (n+1)(n-2) + AB^2 n y^{n+3} (2-n)(n+3) - 2Bn y^{n+2} (AC(n+1)(n+4) + B(n^2-4)) \right. \\
 & \quad - C y^{n+1} (n+1)(n+2) (AC(n+6) + 4Bn) - 2C^2 y^n (n+1)(n+2)(n+3) + 2B^3 n y^3 (2-n)(2n+1) \\
 & \quad \left. - 6B^2 C n^2 y^2 (2n+1) - 6BC^2 y (n+1)(2n^2 + 3n + 2) - 2C^3 (n+1)(n+2)(2n+3) \right) = -48K y^{3n+3}.
 \end{aligned}$$

Therefore, if $n = 1$ and $K = 0$ the previous equation holds true independently from A , B and C . If $n \neq 1$, C needs to be equal to 0. By putting $C = 0$ the equality is satisfied if $K = 0$ and $n = 2$ or $n > 2$, $B = 0$ and $K = -\frac{A^3}{48} n^2 (n+1)^2 (n-1)(n-2)$.

To sum up, we have just proved that $a_3 = K$ if and only if

$$\psi(y) = \begin{cases} Ay^2 + y & \text{if } n \geq 2, \\ Ay^2 + y + B & \text{if } n = 2, \\ Ay^2 + By + C & \text{if } n = 1. \end{cases}$$

By recalling that $y = \Phi'$ and eq. (11), we get that a_2 is constant if and only if $a_3 = K$ is constant, which concludes the proof of Theorem 1.1. \square

In order to prove Corollary 1.2 we need the following result, proved by the authors in [17] (Lemma 2.2 and Corollary 2.3).

Lemma 3.1. *Let $n \geq 2$ and $p = (s, 0, \dots, 0)$, with $s \in \mathbb{R}$, $s \neq 0$, be a point of the complex domain $U \subset \mathbb{C}^n \setminus \{0\}$ on which is defined a radial metric g with radial Kähler potential $\Phi : U \rightarrow \mathbb{R}$ and corresponding diastasis $D_p : U \rightarrow \mathbb{R}$. Let $f : \tilde{U} \rightarrow \mathbb{R}$ defined by $f(r) = \Phi(z)$, $z = (z_1, \dots, z_n)$ where $\tilde{U} = \{r = |z|^2 = |z_1|^2 + \dots + |z_n|^2 \mid z \in U\}$ and, for $h \in \mathbb{N}$, let $g_h : \tilde{U} \rightarrow \mathbb{R}$ given by:*

$$g_h(r) = \frac{d^h e^{f(r)}}{dr^h} e^{-f(r)}. \tag{20}$$

If there exists $r \in \tilde{U}$ and $h \in \mathbb{N}$ such that the function given by (20) is negative, namely $g_h(r) < 0$, then the metric g is not projectively induced.

Proof of Corollary 1.2. Let us first recall that the *Simanca metric* is the Kähler metric g_S on $\mathbb{C}^2 \setminus \{0\}$ defined by the Kähler potential

$$|z|^2 + \log |z|^2 \quad (21)$$

This metric, used by Simanca in [23] to construct constant scalar curvature metrics on blowups, was studied by the authors in [17] (Theorem 1.3), where it was proved that it is projectively induced.

Now, Theorem 1.1 implies that the Simanca metric, the flat metric on \mathbb{C}^n and the metric on $\{z \in \mathbb{C}^2 \mid |z|^2 > 1\}$ given by the potential

$$|z|^2 - \log |z|^2 \quad (22)$$

are, up to homothety, the only radial Kähler metrics g with $a_1 = a_3 = 0$. Indeed, more precisely, by Theorem 1.1, if $a_1 = a_3 = 0$ then g must be one of the metrics of the list in the statement of Theorem 2.1. In particular, since $a_1 = 0$ we are in the case $\Phi'' = b\Phi' + c$ (being Φ a Kähler potential of g) considered in the proof at the end of Section 2: after an easy integration of this ODE, as seen above, one gets the Euclidean metric and the metrics given by 4. and 5. of the list in the statement of Theorem 2.1, which, respectively after the change of variables $w = \frac{z}{\sqrt{\lambda}}$ and $w = \sqrt{\frac{\mu}{\lambda}}z$, proves our claim.

Since it is well-known and easily verified that the Euclidean metric is projectively induced (see e.g. [4]), in order to conclude the proof of Theorem 1.2 we need to show that the Kähler metrics whose Kähler forms are given by $\omega = \lambda \frac{i}{2} \partial \bar{\partial} (|z|^2 - \log |z|^2)$ are not projectively induced, for any $\lambda > 0$.

Set $f(r) = \lambda(r - \log r)$, $r = |z|^2$. Thus $e^{f(r)} = e^{\lambda r} r^{-\lambda}$ and $g_3(r)$ in the previous Lemma reads as:

$$g_3(r) = \frac{de^f}{dr^3} e^{-f} = \lambda r^{-3} [\lambda^2 r^3 - 3\lambda^2 r^2 + 3\lambda(\lambda + 1)r - (\lambda + 1)(\lambda + 2)]. \quad (23)$$

Since the metrics are defined on $\{z \in \mathbb{C}^2 \mid |z|^2 > 1\}$, we can let r tend to 1 in (23) and get

$$\lim_{r \rightarrow 1^+} g_3(r) = \lambda[\lambda^2 - 3\lambda^2 + 3\lambda^2 + 3\lambda - \lambda^2 - 3\lambda - 2] = -2\lambda < 0$$

which, by Lemma 3.1 proves our claim and concludes the proof of Corollary 1.2. \square

REFERENCES

- [1] C. Arezzo, A. Loi, *Moment maps, scalar curvature and quantization of Kähler manifolds*, Comm. Math. Phys. 243 (2004), 543-559.

- [2] C. Arezzo, A. Loi, F. Zuddas, *Szegő kernel, regular quantizations and spherical CR-structures*, Math. Z. (2013) 275, 1207-1216.
- [3] F. A. Berezin, *Quantization*, Izv. Akad. Nauk SSSR Ser. Mat. 38 (1974), 1116–1175 (Russian).
- [4] E. Calabi, *Isometric Imbedding of Complex Manifolds*, Ann. of Math. 58 (1953), 1–23.
- [5] M. Cahen, S. Gutt and J. H. Rawnsley. *Quantization of Kähler manifolds I: Geometric interpretation of Berezin's quantization*, J. Geom. Phys. 7(1990), 43-62.
- [6] M. Cahen, S. Gutt, J. H. Rawnsley, *Quantization of Kähler manifolds III*, Lett. Math. Phys. 30 (1994), 291-305.
- [7] M. Cahen, S. Gutt, J. H. Rawnsley, *Quantization of Kähler manifolds IV*, Lett. Math. Phys. 34 (1995), 159-168.
- [8] D. Catlin. *The Bergman kernel and a theorem of Tian*, Analysis and geometry in several complex variables - Proceedings of the 40th Taniguchi Symposium, 1-23, 1999.
- [9] S. Donaldson, *Scalar Curvature and Projective Embeddings, I*, J. Diff. Geom. 59 (2001), 479-522.
- [10] S. Donaldson, *Scalar Curvature and Projective Embeddings, II*, Q. J. Math. 56 (2005), 345–356.
- [11] M. Engliš, G. Zhang, *Ramadanov conjecture and line bundles over compact Hermitian symmetric spaces*, Math. Z., vol. 264, no. 4 (2010), 901-912.
- [12] Z. Feng. *On the first two coefficients of the Bergman function expansion for radial metrics*, J. Geom. Phys. 119 (2017), 254–271.
- [13] Z. Feng, Z. Tu, *On canonical metrics on Cartan-Hartogs domains*, Math. Zeit. 278 (2014), Issue 1-2, 301–320.
- [14] A. Loi, *The Tian–Yau–Zelditch asymptotic expansion for real analytic Kähler metrics*, Int. J. of Geom. Methods Mod. Phys. 1 (2004), 253-263.
- [15] A. Loi, *A Laplace integral, the T-Y-Z expansion and Berezin's transform on a Kaehler manifold*, Int. J. of Geom. Methods Mod. Phys. 2 (2005), 359-371.
- [16] A. Loi, R. Mossa, *Berezin quantization of homogeneous bounded domains*, Geom. Ded. 161 (2012), 1, 119-128.
- [17] A. Loi, F. Salis, F. Zuddas, *Two conjectures on Ricci-flat Kähler metrics*.
- [18] A. Loi, M. Zedda, *On the coefficients of TYZ expansion of locally Hermitian symmetric spaces*, Manuscr. Math., 148 (2015), no. 3, 303-315.
- [19] A. Loi, M. Zedda, F. Zuddas, *Some remarks on the Kähler geometry of the Taub-NUT metrics*, Ann. Global Anal. Geom. 41 (2012), no. 4, 515-533.
- [20] Z. Lu, *On the lower terms of the asymptotic expansion of Tian–Yau–Zelditch*, Amer. J. Math. 122 (2000), 235-273.
- [21] Z. Lu and G. Tian, *The log term of Szegő Kernel*, Duke Math. J. Volume 125, No 2 (2004), 351-387.
- [22] W. D. Ruan. *Canonical coordinates and Bergmann metrics*, Comm. Anal. Geom. 28(1977), 403–415.
- [23] S.R. Simanca, *Kähler metrics of constant scalar curvature on bundles over $\mathbb{C}P^{n-1}$* , Math. Ann. 291 (1991) 239-246.
- [24] G. Tian. *On a set of polarized Kähler metrics on algebraic manifolds*, J. Differential Geom. 32(1990), 99-130.
- [25] H. Xu, *A closed formula for the asymptotic expansion of the Bergman kernel*, Comm. Math. Phys. 314 (2012), no. 3, 555-585.

- [26] M. Zedda, *Canonical metrics on Cartan-Hartogs domains*, Int. J. Geom. Methods Mod. Phys. 9 (2012), no. 1.
- [27] S. Zelditch. *Szegő kernels and a theorem of Tian*, Int. Mat. Res. Notice 6(1998), 317-331.

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI CAGLIARI, VIA OSPEDALE 72,
09124 CAGLIARI (ITALY)

E-mail address: `loi@unica.it`, `filippo.salis@gmail.com`, `fabio.zuddas@unica.it`