

MICRORNA-RELEASING LIPOPLEXES FOR DIRECT CARDIAC REPROGRAMMING IN
PATHOLOGICAL CARDIAC TISSUE

Original

MICRORNA-RELEASING LIPOPLEXES FOR DIRECT CARDIAC REPROGRAMMING IN PATHOLOGICAL CARDIAC TISSUE / Nicoletti, Letizia; Paoletti, Camilla; Tarricone, Giulia; Divieto, Carla; Andreana, Ilaria; Stella, Barbara; Arpicco, Silvia; Mattu, Clara; Chiono, Valeria. - ELETTRONICO. - (2020). (Intervento presentato al convegno Nature Conferences - RNA at the Bench and Bedside II).

Availability:

This version is available at: 11583/2872073 since: 2021-02-19T19:19:43Z

Publisher:

Nature Conferences

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

MICRORNA-RELEASING LIPOPLEXES FOR DIRECT CARDIAC REPROGRAMMING IN PATHOLOGICAL CARDIAC TISSUE

Letizia Nicoletti^{1,2,3}, Camilla Paoletti^{1,2,3}, Giulia Tarricone^{1,2,3}, Carla Divieto⁴, Ilaria Andreana⁵, Barbara Stella⁵, Silvia Arpicco⁵, Clara Mattu^{1,2,3}, Valeria Chiono^{1,2,3}

¹Department of Mechanical and Aerospace Engineering, Politecnico di Torino, Turin, Italy;

²Interuniversity Center for the promotion of the 3Rs principles in teaching and research, Turin, Italy; ³POLITO Biomedlab, Politecnico di Torino, Turin, Italy; ⁴Division of Advanced Materials and Life Sciences, Istituto Nazionale di Ricerca Metrologica, Turin, Italy;

⁵Department of Drug Science and Technology, University of Turin, Turin, Italy

Cardiovascular diseases are the major cause of death globally. They include ischemic heart disease as a consequence of myocardial infarction. During an infarct, billions of cardiomyocytes irreversibly die, replaced by non-contractile fibrotic tissue. Among the approaches under investigation for cardiac regeneration, the release of specific microRNAs (miRNAs) to the infarcted area has been reported to promote cardiomyocyte proliferation or the direct reprogramming of cardiac fibroblasts (CFs) into induced cardiomyocytes (iCMs). Our group has recently demonstrated that a combination of four miRNAs called miRcombo (miR-1, 133, 208, 499), previously identified by others in mouse model, can trigger trans-differentiation of human CFs into iCMs. CFs were transfected with miRcombo using a commercially available agent and then cultured up to 30 days. After 2 weeks culture time, 11% of the cells expressed cardiac troponin T (cTnT), while after 4 weeks culture time, 38% of the cells showed spontaneous calcium transients. Subsequently, new lipoplexes based on a mixture of cationic and helper lipids were formulated for efficient encapsulation and delivery of microRNAs to human CFs, aimed at enhancing direct reprogramming efficiency.

Lipoplexes loaded with one microRNA (negmiR or miR-1) were initially prepared showing 99% encapsulation efficiency and an average hydrodynamic diameter increasing from 372 nm to 876 nm and a Z-potential decreasing from + 40 mV to -26 mV with decreasing N:P ratios (3.0-0.35). Based on physicochemical characteristics and stability experiments at different temperatures (4°C and 37°C), lipoplexes with a N:P ratio of 3 were selected for further *in vitro* tests with human CFs, showing biocompatibility and significantly higher miR-1 expression, compared to a commercial agent. Encapsulation of miRcombo within the new lipoplexes is in progress with the aim to validate the nanocarriers as efficient vectors for direct cardiac reprogramming.

This project is supported from European Research Council (ERC) under European Union's Horizon 2020 research and innovation program (grant agreement No 7721).