

Perturbations of the orbital elements due to the magnetic-like part of the field of a plane gravitational wave

Original

Perturbations of the orbital elements due to the magnetic-like part of the field of a plane gravitational wave / Iorio, L.; Ruggiero, M. L. - In: INTERNATIONAL JOURNAL OF MODERN PHYSICS D. - ISSN 0218-2718. - 30:12(2021), p. 2150088. [10.1142/S0218271821500887]

Availability:

This version is available at: 11583/2927350 since: 2021-09-26T12:08:24Z

Publisher:

World Scientific

Published

DOI:10.1142/S0218271821500887

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

World Scientific preprint/submitted version

(Article begins on next page)

International Journal of Modern Physics D
 © World Scientific Publishing Company

Perturbations of the orbital elements due to the magnetic-like part of the field of a plane gravitational wave

LORENZO IORIO*

*Ministero dell'Istruzione, dell'Università e della Ricerca (M.I.U.R.)
 lorenzo.iorio@libero.it*

MATTEO LUCA RUGGIERO

*Politecnico di Torino, Torino (Italy) and INFN, Laboratori Nazionali di Legnaro, Legnaro
 (Italy)
 matteo.ruggiero@polito.it*

Received Day Month Year

Revised Day Month Year

We focus on the secular changes of the orbital elements of a planet in the solar system, determined by the magnetic-like part of a gravitational wave field. Using Fermi coordinates we show that the total force acting on a test particle is made of two contributions: a gravito-electric one and a gravito-magnetic one. While the electric-like force has been thoroughly discussed in the past, the effect of the gravito-magnetic force, which depends on the velocity of the test particle, has not been considered yet. We obtain approximated results to some orders in the orbital eccentricity and show that these effects are much smaller than the corresponding gravito-electric ones.

Keywords: general relativity and gravitation; celestial mechanics; gravitational waves

PACS numbers:

1. Introduction

Einstein's theory of gravity, General Relativity (GR), naturally predicts the existence of gravitational waves (see Ref. ? for a review). Indeed, already in 1905, Poincaré[?] discussing Lorentz transformations, suggested that gravitational interactions could not be instantaneous, as in Newtonian theory, but should propagate at the speed of light. Actually, it is possible to show (see e.g. Ref. ?) that the basic properties of the gravitational waves (GWs) can be obtained by combining Newtonian gravity with the retardation effects due to the finite size of the speed of light. Einstein^{?,?} made the first calculations defining the characteristics of the quadrupole formula for the emission of GWs. According to this formula, in order to have substantial emission of GWs, huge masses moving at relativistic velocities

*Permanent address for correspondence: Viale Unità di Italia 68, 70125, Bari (BA).

are needed. These sources are far away from the Earth and the weakness of the gravitational interaction entails that what can be actually measured by our devices are very small perturbations or ripples in the fabric of space-time. Indirect evidence for gravitational waves was first obtained by observing the orbital decay of the pulsar B1913+16.^{?,?} It took 100 years from Einstein's first calculation to have, in 2016, the first direct detection of GWs h ,[?] which was the beginning of *gravitational waves astronomy*. More in general, the study and the detection of GWs could give information not only on GR, but also on the reliability of extended theories of gravity.^{?,?}

As far as we know (see e.g. Ref. ?), the spectrum of gravitational waves spreads over some 18 orders of magnitude, thus encompassing astrophysical and cosmological events on very different scales. In particular, while high frequency waves ($10-10^4$ Hz) are in the target of terrestrial detectors, there is no hope to detect on the Earth low-frequency ($10^{-5}-1$ Hz) and very low frequency waves ($10^{-9}-10^{-7}$ Hz): in fact the ground motions and the coupling to fluctuations in the local mass distribution limit the sensitivity up to 1 Hz (see Ref. ?). While GWs in the low frequency regime are ideal target of the LISA space mission,^{?,?} as for the very low frequency regime possible sources could be unresolved coalescing massive binary black holes or binaries that are either too massive to radiate in the LISA band, or else are inspiralling towards the LISA band.[?]

Bertotti[?] first suggested to use gravitationally bound systems to detect secular effects due to a GW background: a Newtonian binary system consisting of point masses orbiting each other on Keplerian ellipses is acted upon by a gravitational wave, which causes a perturbation of the orbital elements. There have been several subsequent approaches (see e.g. Refs. ?, ?, ?, ?) that are thoroughly discussed by Iorio.[?] Mashhoon[?] first developed a description of the tidal effects associated to the passage of a gravitational wave using Fermi coordinates, then in Ref. ? he studied the interaction of a weak gravitational wave and a Newtonian binary system.

Fermi coordinates^{?,?,?} are defined around the world-line of an observer and are the coordinates he would naturally use to make space and time measurements. It is possible to show[?] that using Fermi coordinates the effects of a plane gravitational wave can be described by gravito-electromagnetic fields:^{?,?} in other words, the wave field is equivalent to the action of a gravito-electric and a gravito-magnetic field, that are transverse to the propagation direction and orthogonal to each other. In particular, while current detectors, such LIGO and VIRGO, or future ones, such as LISA, reveal the interaction of test masses with the gravito-electric component of the wave, there are also gravito-magnetic interactions that could be used to detect the effect of the gravitational wave on moving masses and spinning particles.^{?,?,?} We point out that the effect of the angular momentum of gravitational waves on the rotation of inertial frames is studied in Refs. ?, ?, while the theoretical possibility of measuring by a Sagnac detector the gravito-magnetic effects caused by a gravitational wave is investigated by Frauendiener.[?]

In this paper we study the effect of the gravito-magnetic part of a gravitational

wave, in the very low frequency regime, on the long-term variations of all the six standard Keplerian orbital elements of a solar system planet. This work extends to the magnetic-like components of the wave the study of the interaction with a Newtonian binary system, according to an approach similar to Ref. ?, where it is also pointed out that very low frequency waves are important to get information on astrophysics of galaxies and black holes, early Universe cosmology and on the extension of the standard model.

The paper is organised as follows: in Section ?? we briefly review gravito-electromagnetic approach to the description of the GWs field, in the Fermi frame; the perturbations of the orbital elements are worked out in Section ??, while conclusions are eventually drawn in Section ??.

2. Gravito-electromagnetic fields in the Fermi frame

In this Section we are going to briefly review the construction of the Fermi frame to describe the dynamics of test particles using the gravito-electromagnetic analogy. In order to define the Fermi frame, we use the approach described in Refs. ?, ?.

We consider a background spacetime describing the gravitational field where we use a set of coordinates^a x^μ . Using these coordinates, the world-line $x^\mu(\tau)$ of a reference observer, as function of the proper time τ , is determined by the following equation

$$\frac{Dx^\mu}{d\tau} = \ddot{x}^\mu + \Gamma^\mu_{\nu\sigma} \dot{x}^\nu \dot{x}^\sigma = a^\mu, \quad (1)$$

where D is the covariant derivative along the world-line, a dot means derivative with respect to τ and a^μ is the four-acceleration. In the tangent space along the world-line $x^\mu(\tau)$ it is possible to define the orthonormal tetrad of the observer $e^\mu_{(\alpha)}(\tau)$, such that $e^\mu_{(0)}(\tau)$ is the unit vector tangent to his world-line and $e^\mu_{(i)}(\tau)$ (for $i=1,2,3$) are the spatial vectors orthogonal to each other and, also, orthogonal to $e^\mu_{(0)}(\tau)$. The equation of motion of the tetrad is $\frac{De^\mu_{(\alpha)}}{d\tau} = -\Omega^{\mu\nu} e_{\nu(\alpha)}$ where $\Omega^{\mu\nu} = a^\mu \dot{x}^\nu - a^\nu \dot{x}^\mu + \dot{x}_\alpha \Omega_\beta \epsilon^{\alpha\beta\mu\nu}$. In the latter equation Ω^α is the four-rotation of the tetrad. In particular, we notice that for a geodesic ($a^\mu = 0$) and non rotating ($\Omega^\alpha = 0$) tetrad we have $\Omega^{\mu\nu} = 0$: consequently, in this case the tetrad is parallel transported. If $\Omega = 0$ and $a^\mu \neq 0$, the tetrad is Fermi-Walker transported. We remember that Fermi-Walker transport enables to define, in a natural way, a non rotating moving frame for an accelerated observer.[?] That being said, Fermi coordinates are defined as follows: the observer along the congruence measures time intervals according to the proper time, so the time coordinate is defined by $T = \tau$; the spatial coordinates X, Y, Z are defined by space-like geodesics, with unit tangent vectors n^μ , whose components, with respect to the orthonormal tetrad are $n^{(i)} = n_{(i)} = n_\mu e^\mu_{(i)}(\tau)$,

^aGreek indices refer to spacetime coordinates, and assume the values 0, 1, 2, 3, while Latin indices refer to spatial coordinates and assume the values 1, 2, 3, usually corresponding to the coordinates x, y, z .

4 *Lorenzo Iorio, Matteo Luca Ruggiero*

and $n^{(0)} = 0$. The reference observer's frame equipped with Fermi coordinates is the *Fermi frame*. Fermi coordinates in the neighbourhood of the world-line of an observer in accelerated motion with rotating tetrads are studied in Refs. ?, ?, ?.

Fermi coordinates prove useful to develop a gravito-electromagnetic analogy. Indeed, as discussed by Mashhoon,[?] the spacetime element in Fermi coordinates in the vicinity of the observer's world-line can be recast in terms of the gravito-electromagnetic potentials (Φ, \mathbf{A})

$$ds^2 = - \left(1 - 2 \frac{\Phi}{c^2} \right) c^2 dT^2 - \frac{4}{c} (\mathbf{A} \cdot d\mathbf{X}) dt + \delta_{ij} dX^i dX^j, \quad (2)$$

and this peculiarity allows us to apply the corresponding formalism^b. Indeed, as shown by,[?] it is possible to distinguish between the *inertial* and the *curvature* contributions to the gravito-electromagnetic potentials: the former depend on the inertial features of the Fermi frame (i.e. the observer's acceleration and tetrad rotation), while the latter on the Riemann curvature tensor. Here, we are concerned with the curvature effects only, since we refer our results to an inertial frame: for instance, if we consider planetary motion in the solar system, its origin is at the position of the Sun. Consequently, the *gravito-electric* (GE) potential $\Phi = \Phi(T, \mathbf{X})$ is

$$\Phi(T, \mathbf{X}) = -\frac{1}{2} R_{0i0j}(T) X^i X^j \quad (3)$$

and the components of the *gravito-magnetic* potential (GM) $\mathbf{A} = \mathbf{A}(T, \mathbf{X})$ turn out to be

$$A_i(T, \mathbf{X}) = \frac{1}{3} R_{0jik}(T) X^j X^k. \quad (4)$$

In close analogy to electromagnetism, the gravito-electric and gravito-magnetic fields \mathbf{E} and \mathbf{B} are defined in terms of the potentials by

$$\mathbf{E} = -\nabla\Phi - \frac{1}{c} \frac{\partial}{\partial T} \left(\frac{1}{2} \mathbf{A} \right), \quad \mathbf{B} = \nabla \times \mathbf{A}. \quad (5)$$

Using the definitions (??) and (??) we obtain (up to linear order in $|X^i|$) the following components

$$E_i^C(T, \mathbf{X}) = c^2 R_{0i0j}(T) X^j, \quad (6)$$

and

$$B_i^C(T, \mathbf{X}) = -\frac{c^2}{2} \epsilon_{ijk} R^{jk}_{\quad 0l}(T) X^l. \quad (7)$$

The electromagnetic analogy is also useful to describe the motion of free test particles: namely, the geodesic equation of the space-time metric (??) can be written in the form of a Lorentz-like force equation[?]

$$m \frac{d^2 \mathbf{X}}{dT^2} = q_E \mathbf{E} + q_B \frac{\mathbf{V}}{c} \times \mathbf{B}, \quad (8)$$

^bHere and henceforth \mathbf{X} is the position vector in the Fermi frame.

up to linear order in the particle velocity $\mathbf{V} = \frac{d\mathbf{X}}{dT}$ (which is indeed the *relative velocity* with respect to the reference mass at the origin of the frame). In the Lorentz-like force equation, $q_E = -m$ is the GE charge, and $q_B = -2m$ is the GM one (the minus sign takes into account that the gravitational force is always attractive). As a consequence, the Lorentz-like force equation becomes

$$m \frac{d^2\mathbf{X}}{dT^2} = -m\mathbf{E} - 2m \frac{\mathbf{V}}{c} \times \mathbf{B}. \quad (9)$$

In the following Section we are going to work out the effect of the gravito-magnetic field of the wave on a moving test particle. For the sake of simplicity, we consider plane monochromatic circularly polarised GWs propagating with frequency ν_g along the direction Z . Moreover, in what follows we suppose that the extension of the reference frame is much smaller than the wavelength, so that we may neglect the spatial variation of the wave field: consequently the components of the Riemann tensor are evaluated at the origin of our frame. Accordingly, as shown in,[?] the corresponding gravito-magnetic field is

$$B_Z = 0, \quad (10)$$

$$B_X = -\frac{A\nu_g^2}{2} (-X \cos \chi + Y \sin \chi), \quad (11)$$

$$B_Y = -\frac{A\nu_g^2}{2} (X \sin \chi + Y \cos \chi), \quad (12)$$

with $\chi \doteq \nu_g T$.

Hence, the gravito-magnetic force acting upon a point-like mass m is obtained by

$$m \frac{d^2\mathbf{X}}{dT^2} = -2m \frac{\mathbf{V}}{c} \times \mathbf{B}. \quad (13)$$

Note that, being B dimensionally an acceleration, the scaling parameter A is dimensionless.

3. Perturbations due to the gravito-magnetic field of the wave

To study the effect of the magnetic-like part of the gravitational field of the wave on a planet of the solar system we proceed as follows. We consider a Fermi coordinate system, whose origin is located at the Sun position; accordingly, we treat the planet as a test particle moving around the Sun.^c Let us assume that the wave's frequency

^cMore in general, studying a binary system, the origin of the Fermi frame should be located at the center of mass of the system, and what we would consider the relative position of the two masses (see e.g. Ref. ?). The approximation considered is reasonable for most of the bodies of the solar system.

6 *Lorenzo Iorio, Matteo Luca Ruggiero*

ν_g is much lower than the orbital one n_b , so that the averaged rates of change of the orbital elements of the test particle can be analytically calculated with the Gauss perturbing equations^{?, ?, ?, ?, ?, ?} by keeping $\chi = \text{const}$ in the integration over the true anomaly f from 0 to 2Π of the latter. The radial, transverse and normal components of the acceleration of Equation (??) turn out to be

$$A_r = -\frac{A a^2 \sqrt{1-e^2} n_b \nu_g^2 \sin I}{c} [\cos I \cos(\chi + 2\Omega) \sin u + \cos u \sin(\chi + 2\Omega)], \quad (14)$$

$$A_t = \frac{A e a^2 \sqrt{1-e^2} n_b \nu_g^2 \sin I \sin f}{c (1+e \cos f)} [\cos I \cos(\chi + 2\Omega) \sin u + \cos u \sin(\chi + 2\Omega)], \quad (15)$$

$$\begin{aligned} A_n = & \frac{A a^2 \sqrt{1-e^2} n_b \nu_g^2}{16 c (1+e \cos f)} (2e \cos(f - \chi - 2\Omega) - e \cos(f - 2I - \chi - 2\Omega) - \\ & - 2 \cos(2I - \chi - 2\Omega) + 2 [2 - \cos 2I + 2 (3 + \cos 2I) \cos 2u] \cos(\chi + 2\Omega) + \\ & + 4 \cos I (\sin I - 4 \sin 2u) \sin(\chi + 2\Omega) + \\ & + e \{-\cos(f + 2I - \chi - 2\Omega) + 2 [(3 + \cos 2I) \cos(f - \chi + 2\omega - 2\Omega) + \\ & + \cos(f + \chi + 2\Omega) + 3 \cos(f + \chi + 2\varpi) - 8 \cos I \sin(f + 2\omega) \sin(\chi + 2\Omega) - \\ & - 2 \cos 2I \sin \omega \sin(f + \chi + \omega + 2\Omega)]\}), \quad (17) \end{aligned}$$

where a is the semimajor axis, e is the eccentricity, I is the inclination to the reference $\{x, y\}$ plane, Ω is the longitude of the ascending node, ω is the argument of pericentre, $u \doteq \omega + f$ is the argument of latitude, and $\varpi \doteq \Omega + \omega$ is the longitude of pericentre. An exact calculation in e turns out to be exceedingly cumbersome; thus, we will only show approximate results to some orders in e .

To the order of $\mathcal{O}(e^{-1})$, nonzero secular rates occur for the argument of pericentre and the mean anomaly at epoch η . They are

$$\dot{\omega} = \frac{a A \nu_g^2 \sin I (\cos I \cos(\chi + 2\Omega) \sin \omega + \cos \omega \sin(\chi + 2\Omega))}{2 c e}, \quad (18)$$

$$\dot{\eta} = -\dot{\omega}. \quad (19)$$

To the order zero in e , only the eccentricity experiences a long-term rate of

change

$$\dot{e} = \frac{a A \nu_g^2 \sin I (-\cos I \cos \omega \cos(\chi + 2\Omega) + \sin \omega \sin(\chi + 2\Omega))}{2c}. \quad (20)$$

To the order of $\mathcal{O}(e)$, the inclination I , the node Ω , the pericentre ω and the mean anomaly at epoch η undergo secular precessions given by

$$\begin{aligned} \dot{I} = & \frac{e a A \nu_g^2}{16c} [-2 \cos(I - \chi - \omega - 2\Omega) + \cos(2I - \chi - \omega - 2\Omega) + 2 \cos(I - \chi + \omega - 2\Omega) + \\ & + \cos(2I - \chi + \omega - 2\Omega) + 2 \cos \omega (-6 \cos(\chi + 2\Omega) + \cos(2I + \chi + 2\Omega)) + \\ & + 4 \sin \omega \sin(I + \chi + 2\Omega)], \end{aligned} \quad (21)$$

$$\begin{aligned} \dot{\Omega} = & \frac{e a A \nu_g^2 \csc I}{8c} \{-\sin(I - \chi - \omega - 2\Omega) - \sin(2I - \chi - \omega - 2\Omega) - \sin(I - \chi + \omega - 2\Omega) + \\ & + \sin(2I - \chi + \omega - 2\Omega) + \sin(I + \chi - \omega + 2\Omega) - \sin(2I + \chi - \omega + 2\Omega) + \\ & + \sin(I + \chi + \omega + 2\Omega) + \sin(2I + \chi + \omega + 2\Omega)\}, \end{aligned} \quad (22)$$

$$\dot{\omega} = \frac{e a A \nu_g^2}{4c} [-2 \cos(\chi + 2\Omega) \cot I \sin \omega + (-3 + \cos 2I) \cos \omega \csc I \sin(\chi + 2\Omega)], \quad (23)$$

$$\dot{\eta} = -\frac{7 e a A \nu_g^2 \sin I}{4c} (\cos I \cos(\chi + 2\Omega) \sin \omega + \cos \omega \sin(\chi + 2\Omega)) \quad (24)$$

If, instead, it is $\nu_g \gg n_b$, it turns out that there are no net orbital shifts averaged over a wave's full cycle.

In the previous results, we made no assumptions on the orbital inclination I ; we notice that, for $I = 0$, i.e. the incidence of the gravitational wave is normal to the orbital plane, we see that there is no secular variation of the eccentricity, and the same holds true for the mean anomaly at epoch.

4. Conclusions

In this paper we studied the averaged rates of change of the orbital elements of a test particle orbiting a central body, such as a planet orbiting the Sun, determined by the magnetic-like force due to the passage of a monochromatic plane gravitational wave. In doing so, we assumed that the wavelength is larger than the orbit size and, also, that the frequency is much smaller than the orbital frequency of the test particle.

To fix ideas, in the solar system we remember that Mercury has the highest orbital frequency, 1.3×10^{-7} Hz. Hence, in this case our results apply to the very low frequency regime of gravitational waves. However, our results could be applied to other astrophysical situations, such as planets in extrasolar systems, where higher orbital frequencies are possible.

Since exact calculations are complicated, in order to give an idea of the effects, we worked out approximated results to some orders in the orbital eccentricity e . Our results showed that all orbital elements, except the semi-major axis, undergo secular variations; in particular, we proved that to zero order in the eccentricity e , only the eccentricity itself experiences a long-term rate of change. This variation is a purely gravito-magnetic effect: in fact, as discussed in Ref. ?, if we consider the gravito-electric perturbation only, the variation of the eccentricity is proportional to the eccentricity itself, so that circular orbits do not change their shape. More in general, orbits with given eccentricities change their shape and orientation due to the passage of the wave: all variations determined by the gravito-magnetic effects are proportional to $\epsilon = \frac{aA\nu_g^2}{c}$, which is a very small quantity, depending on the wave amplitude A , the orbital frequency ν_g and semimajor axis a , and the speed of light c . However, even if these gravito-magnetic effects are in principle present, it is reasonably expected that they are negligible with respect to the gravito-electric ones. This can be explained by inspecting Eq. (??): in fact, while both the gravito-electric and magnetic fields have the same order of magnitude, the gravito-magnetic force is reduced, with respect to the gravito-electric one, by a factor $\frac{|\mathbf{V}|}{c}$, where the magnitude of the orbital speed $|\mathbf{V}|$ is much smaller than the speed of light.

In conclusion, the aim of this paper was to focused on the gravito-magnetic effects provoked by the passage of a plane gravitational wave; as discussed by one of us,[?] there are also gravito-electric effects which, according to the above discussion, have a larger impact. However, the separation of the two effects is purely academic, since a point mass experiences a total force in the form (??). As a consequence, to make predictions on the impact of a gravitational wave on orbital elements we need to consider the whole perturbing acceleration determined by the passage of the wave: this will be the subject of a future work. These predictions could be used to make a comparison with planetary data available from the solar system which can, in principle, give constraints on all unknowns wave parameters. This formalism could be applied not only in the solar system, but to arbitrary gravitationally bound two-body systems, such as extrasolar systems, and with suitable modifications, it can be used to test models of gravity alternative to General Relativity, thus adding new perspectives to the field of gravitational waves astronomy.

References

1. E. E. Flanagan and S. A. Hughes, *New Journal of Physics* **7** (2005) 204.
2. H. Poincaré, *Comptes rendus hebdomadaires des séances de l'Académie des sciences* **T140** (1905) 1504.

3. B. F. Schutz, *American Journal of Physics* **52** (1984) 412.
4. A. Einstein, *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)* **1916** (1916) 688.
5. A. Einstein, *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)* **1918** (1918) 154.
6. R. Hulse and J. Taylor, *Astrophys. J. Lett.* **195** (1975) L51.
7. J. M. Weisberg and J. H. Taylor, *ASP Conf. Ser.* **328** (2005) 25, [arXiv:astro-ph/0407149](#).
8. B. P. Abbott, R. Abbott, T. Abbott, M. Abernathy, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. Adhikari *et al.*, *Physical review letters* **116** (2016) 061102.
9. S. Capozziello and M. De Laurentis, *Phys. Rept.* **509** (2011) 167, [arXiv:1108.6266 \[gr-qc\]](#).
10. S. Capozziello and V. Faraoni, *Beyond Einstein gravity: A Survey of gravitational theories for cosmology and astrophysics* (Springer Science & Business Media, 2010).
11. T. A. Prince, The promise of low-frequency gravitational wave astronomy (2009).
12. M. Tinto and S. V. Dhurandhar, *Living Reviews in Relativity* **8** (Jul 2005).
13. P. Amaro-Seoane, S. Aoudia, S. Babak, P. Binetruy, E. Berti, A. Bohe, C. Caprini, M. Colpi, N. J. Cornish, K. Danzmann *et al.*, *Classical and Quantum Gravity* **29** (2012) 124016.
14. B. Bertotti, *Astrophysical Letters* **14** (January 1973) 51.
15. V. N. Rudenko, *Soviet Astronomy* **19** (October 1975) 270.
16. M. S. Turner, *Astrophysical Journal* **233** (October 1979) 685.
17. A. V. Ivashchenko, *Soviet Astronomy* **31** (February 1987) 76.
18. L. Hui, S. T. McWilliams and I.-S. Yang, *Phys. Rev. D* **87** (2013) 084009, [arXiv:1212.2623 \[gr-qc\]](#).
19. L. Iorio, *ScienceOpen Res. SOR-ASTRO* (2014) [arXiv:1104.4853 \[gr-qc\]](#).
20. B. Mashhoon, *The Astrophysical Journal* **216** (1977) 591.
21. B. Mashhoon, *The Astrophysical Journal* **223** (1978) 285.
22. E. Fermi, *Rend. Lincci, 1922, 31(1), pp. 21-23, 51-52, 101-103 (in Italian)* **31** (December 1922) 21.
23. F. Manasse and C. W. Misner, *Journal of mathematical physics* **4** (1963) 735.
24. W.-T. Ni and M. Zimmermann, *Physical Review D* **17** (1978) 1473.
25. M. L. Ruggiero and A. Ortolan, *Journal of Physics Communications* **4** (May 2020) 055013.
26. M. L. Ruggiero and A. Tartaglia, *Nuovo Cim.* **B117** (2002) 743, [arXiv:gr-qc/0207065 \[gr-qc\]](#).
27. B. Mashhoon (2003) [arXiv:gr-qc/0311030 \[gr-qc\]](#).
28. D. Bini, A. Geralico and A. Ortolan, *Phys. Rev. D* **95** (May 2017) 104044.
29. M. L. Ruggiero and A. Ortolan, *Phys. Rev. D* **102** (2020) 101501, [arXiv:2011.01663 \[gr-qc\]](#).
30. J. c. v. Bičák, J. Katz, T. c. v. Ledvinka and D. Lynden-Bell, *Phys. Rev. D* **85** (Jun 2012) 124003.
31. W. Barker, *Classical and Quantum Gravity* **34** (Jul 2017) 167001.
32. J. Frauendiener, *Classical and Quantum Gravity* **37** (Feb 2020) 05LT01.
33. C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation* (San Francisco: WH Freeman and Co., 1973).
34. K.-P. Marzlin, *Phys. Rev. D* **50** (Jul 1994) 888.
35. W.-T. Ni and M. Zimmermann, *Physical Review D - Particles and Fields* **17** (March 1978) 1473.
36. W.-Q. Li and W.-T. Ni, *Journal of Mathematical Physics* **20** (July 1979) 1473.
37. C. Chicone, B. Mashhoon and D. G. Retzliff, Gravitational ionization: periodic orbits of binary systems perturbed by gravitational radiation, in *Annales de l'IHP Physique*

10 *Lorenzo Iorio, Matteo Luca Ruggiero*

théorique, (1) (1996), pp. 87–125.

38. M. H. Soffel, *Relativity in Astrometry, Celestial Mechanics and Geodesy* (Springer, Heidelberg, 1989).
39. V. A. Brumberg, *Essential Relativistic Celestial Mechanics* (Adam Hilger, Bristol, 1991).
40. B. Bertotti, P. Farinella and D. Vokrouhlický, *Physics of the Solar System* (Kluwer, Dordrecht, October 2003).
41. A. E. Roy, *Orbital Motion. Fourth Edition* (Institute of Physics Publishing, Bristol, 2005).
42. S. Kopeikin, M. Efroimsky and G. Kaplan, *Relativistic Celestial Mechanics of the Solar System* (Weinheim: Wiley-VCH, September 2011).
43. E. Poisson and C. M. Will, *Gravity* (Cambridge: Cambridge Univ. Press, May 2014).
44. M. H. Soffel and W.-B. Han, *Applied General Relativity* Astronomy and Astrophysics Library, Astronomy and Astrophysics Library (Springer Nature Switzerland, Cham, 2019).