

Three-dimensional exotic Newtonian supergravity theory with cosmological constant

Original

Three-dimensional exotic Newtonian supergravity theory with cosmological constant / Concha, P.; Ravera, L.; Rodriguez, E.. - In: THE EUROPEAN PHYSICAL JOURNAL. C, PARTICLES AND FIELDS. - ISSN 1434-6044. - ELETTRONICO. - 81:646(2021), pp. 1-15. [10.1140/epjc/s10052-021-09456-3]

Availability:

This version is available at: 11583/2949098 since: 2022-01-12T12:11:47Z

Publisher:

Springer Science and Business Media Deutschland GmbH

Published

DOI:10.1140/epjc/s10052-021-09456-3

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Three-dimensional exotic Newtonian supergravity theory with cosmological constant

Patrick Concha^{1,a}, Lucrezia Ravera^{2,3,b}, Evelyn Rodríguez^{4,c}

¹ Departamento de Matemática y Física Aplicadas, Universidad Católica de la Santísima Concepción, Alonso de Ribera 2850, Concepción, Chile

² DISAT, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Turin, Italy

³ INFN, Sezione di Torino, Via P. Giuria 1, 10125 Turin, Italy

⁴ Departamento de Física, Universidad del Bío-Bío, Avenida Collao 1202, Casilla 5-C, Concepción, Chile

Received: 11 May 2021 / Accepted: 12 July 2021 / Published online: 22 July 2021
 © The Author(s) 2021

Abstract We present a supersymmetric extension of the exotic Newtonian Chern–Simons gravity theory in three spacetime dimensions. The underlying new non-relativistic superalgebra is obtained by expanding the $\mathcal{N} = 2$ AdS superalgebra and can be written as two copies of the enhanced Nappi–Witten algebra, one of which is augmented by supersymmetry. We show that the exotic Newtonian superalgebra allows to introduce a cosmological constant to the extended Newtonian supergravity. Interestingly, the obtained supergravity action contains the extended Newton–Hooke supergravity as a sub-case.

1 Introduction

There has been a growing interest in exploring Newtonian (super)gravity theories due to their use in strongly coupled condensed matter systems [1–11] and non-relativistic effective field theories [12–16]. The construction of Newtonian gravity, describing the physical gravitational force at non-relativistic level, requires to consider the so-called Newton–Cartan geometry [17, 18]. Such geometrical framework is necessary to covariantize the Poisson equation of Newtonian gravity. Nevertheless, a principle action for Newtonian gravity was recently presented in [19] which has required to extend the Bargmann algebra [20–27] by including three additional generators. Subsequently, a three-dimensional Chern–Simons (CS) action has been constructed in [28] which is invariant under a central extension of the symmetry group that leaves the recently constructed Newtonian gravity action invariant. The novel symmetry has been denoted as extended Newtonian algebra and can be recovered by means of a contraction of a bi-metric model being the sum of Einstein gravity in the Lorentzian and Euclidean signatures. Interestingly, unlike the Newtonian gravity of [19], the matter coupling of the extended Newtonian gravity theory admits backgrounds with non-trivial curvature whenever matter is present, similarly to the matter-coupled extended Bargmann gravity [27].

The introduction of a cosmological constant in gravity theories is known to have important applications, such as, e.g., the ones recently presented in [29]. In non-relativistic gravity theories the inclusion of a cosmological constant in non-relativistic gravity theories is done considering the Newton–Hooke symmetry [30–36]. However, an extension of the extended Newton–Hooke algebra [37, 38] is needed to include a cosmological constant to the extended Newtonian gravity theory [39]. The novel symmetry is denoted as

Contents

1 Introduction	1
2 Exotic Newtonian gravity in three spacetime dimensions	2
3 Exotic Newtonian superalgebra and semigroup expansion method	4
3.1 Exotic Newtonian superalgebra by expanding the $\mathcal{N} = 2$ AdS superalgebra	4
4 Exotic Newtonian supergravity action and flat limit	7
5 Conclusions	10
A Exotic Newtonian superalgebra by expanding an enhanced Nappi–Witten superalgebra	11
B Supersymmetry gauge transformations	12
References	13

^a e-mail: patrick.concha@ucsc.cl (corresponding author)

^b e-mail: lucrezia.ravera@polito.it

^c e-mail: ekrodriguez@ubiobio.cl

exotic Newtonian algebra and can be seen as an enhanced Bargmann–Newton–Hooke algebra [40]. Both extended and exotic Newtonian gravity theories can be recovered as the non-relativistic limit of the coadjoint Poincaré $\oplus \mathfrak{u}(1)^2$ and coadjoint AdS $\oplus \mathfrak{u}(1)^2$ gravity theories [40].

Supersymmetric extensions of three-dimensional non-relativistic gravity models have been recently approached in [27, 41, 42] and subsequently studied in [28, 43–47]. In particular, a CS action based on the supersymmetric extension of the extended Newtonian algebra has been presented in [28]. Although a cosmological constant has been accommodated in a non-relativistic supergravity theory through the extended Newton–Hooke superalgebra [44], the possible supersymmetric extensions of the exotic Newtonian gravity remain unexplored. Unlike bosonic non-relativistic gravity, the construction of an action based on a non-relativistic superalgebra is non-trivial and requires the introduction of additional bosonic generators. Furthermore, the non-relativistic limit is often ambiguous when supercharges are present. One way to circumvent this difficulty is through the expansion method based on Maurer–Cartan forms [48] and semigroups [49], which have proved to be useful to obtain known and new non-relativistic supergravity theories from relativistic ones [43, 44, 46].

In this work, we present a supersymmetric extension of the three-dimensional exotic Newtonian CS gravity introduced in [39] by applying the semigroup expansion (S -expansion) [49] to the $\mathcal{N} = 2$ AdS supergravity theory. The motivation to consider the CS formalism in three spacetime dimensions is twofold. On one hand, three-dimensional CS gravity can be seen as a toy model to approach higher-dimensional theories. On the other hand, the construction of a non-relativistic (super)gravity action is much simpler and affordable through the CS formalism which allows us to write a gauge-invariant action. The novel Newtonian supergravity theory is invariant under an exotic Newtonian superalgebra which can be written as two copies of the enhanced Nappi–Witten algebra [40, 50], one of which is augmented by supersymmetry. Interestingly, the extended Newton–Hooke supergravity action appears as a sub-case of the exotic Newtonian supergravity theory. Moreover, we show that the extended Bargmann and extended Newtonian supergravities are recovered in the vanishing cosmological constant limit (i.e. the flat limit).

The paper is organized as follows: In Sect. 2 we review the exotic Newtonian algebra and the corresponding three-dimensional CS action. Sections 3 and 4 contain our main results. In Sect. 3, we present an exotic Newtonian superalgebra by considering an S -expansion of the $\mathcal{N} = 2$ super AdS algebra. In Sect. 4, we construct the exotic Newtonian CS supergravity action and analyze its vanishing cosmological constant limit. Section 5 is devoted to discussion and future outlook.

2 Exotic Newtonian gravity in three spacetime dimensions

In this section, we briefly review the three-dimensional CS exotic Newtonian gravity theory presented in [39]. The exotic Newtonian algebra, also denoted as extended post-Newtonian algebra [51], is spanned by the set of generators $\{J, G_a, H, P_a, M, S, B_a, T_a\}$ along with two central charges Y and Z . The non-vanishing commutation relations of the exotic Newtonian algebra are given by

$$\begin{aligned} [J, G_a] &= \epsilon_{ab} G_b, & [G_a, G_b] &= -\epsilon_{ab} S, \\ [H, G_a] &= \epsilon_{ab} P_b, & [J, P_a] &= \epsilon_{ab} P_b, & [G_a, P_b] &= -\epsilon_{ab} M, \\ [H, B_a] &= \epsilon_{ab} T_b, & [J, B_a] &= \epsilon_{ab} B_b, & [G_a, B_b] &= -\epsilon_{ab} Z, \\ [S, G_a] &= \epsilon_{ab} B_b, & [J, T_a] &= \epsilon_{ab} T_b, & [G_a, T_b] &= -\epsilon_{ab} Y, \\ [S, P_a] &= \epsilon_{ab} T_b, & [M, G_a] &= \epsilon_{ab} T_b, & [P_a, B_b] &= -\epsilon_{ab} Y, \\ [H, P_a] &= \frac{1}{\ell^2} \epsilon_{ab} G_b, & [H, T_a] &= \frac{1}{\ell^2} \epsilon_{ab} B_b, & [P_a, P_b] &= -\frac{1}{\ell^2} \epsilon_{ab} S, \\ [M, P_a] &= \frac{1}{\ell^2} \epsilon_{ab} B_b, & [P_a, T_b] &= -\frac{1}{\ell^2} \epsilon_{ab} Z. \end{aligned} \quad (2.1)$$

Let us note that ℓ is a length parameter related to the cosmological constant through $\Lambda \propto \pm \frac{1}{\ell^2}$. In particular, in the vanishing cosmological constant limit $\ell \rightarrow \infty$, the algebra corresponds to the extended Newtonian algebra introduced in [28]. One can notice that the extended Newton–Hooke algebra [37, 38] appears setting B_a , T_a , Y and Z to zero.

The exotic Newtonian algebra admits the extended Newton–Hooke non-vanishing components of the invariant tensor [37, 38]

$$\begin{aligned} \langle JS \rangle &= -\alpha_0, \\ \langle G_a G_b \rangle &= \alpha_0 \delta_{ab}, \\ \langle JM \rangle &= \langle HS \rangle = -\alpha_1, \\ \langle G_a P_b \rangle &= \alpha_1 \delta_{ab}, \\ \langle HM \rangle &= -\frac{\alpha_0}{\ell^2}, \\ \langle P_a P_b \rangle &= \frac{\alpha_0}{\ell^2} \delta_{ab}, \end{aligned} \quad (2.2)$$

along with [39]

$$\begin{aligned} \langle SS \rangle &= \langle JZ \rangle = -\beta_0, \\ \langle G_a B_b \rangle &= \beta_0 \delta_{ab}, \\ \langle MS \rangle &= \langle HZ \rangle = \langle JY \rangle = -\beta_1, \end{aligned}$$

$$\begin{aligned}\langle P_a B_b \rangle &= \langle G_a T_b \rangle = \beta_1 \delta_{ab}, \\ \langle MM \rangle &= \langle HY \rangle = -\frac{\beta_0}{\ell^2}, \\ \langle P_a T_b \rangle &= \frac{\beta_0}{\ell^2} \delta_{ab}.\end{aligned}\quad (2.3)$$

Here, α_0 , α_1 , β_0 and β_1 are arbitrary independent constants. In particular, the components of the invariant tensor proportional to α_0 and β_0 are related to the “exotic” sector of the theory [52]. On the other hand, the flat limit $\ell \rightarrow \infty$ reproduces the invariant tensor of the extended Bargmann algebra [27,38] and the most general extended Newtonian one [39]. Let us note that, at the level of the exotic Newtonian algebra, the invariant tensor (2.2) alone is degenerate. Nevertheless, one obtain a non-degenerate bilinear form considering both families of invariant tensor given by (2.2) and (2.3).

The CS action based on the exotic Newtonian algebra can be obtained considering the following gauge connection one-form:

$$\begin{aligned}A &= \tau H + e^a P_a + \omega J + \omega^a G_a + m M + s S + t^a T_a \\ &\quad + b^a B_a + y Y + z Z,\end{aligned}\quad (2.4)$$

and the non-vanishing components of the invariant tensor (2.2) and (2.3) into the general expression of a three-dimensional CS action,

$$I_{CS} = \frac{k}{4\pi} \int \left\langle AdA + \frac{2}{3} A^3 \right\rangle. \quad (2.5)$$

Here k is the CS level of the theory which, for gravitational theories, is related to the gravitational constant G through $k = 1/(4G)$. The three-dimensional exotic Newtonian CS gravity action reads, up to boundary terms, as follows [39]:

$$I_{\text{Exotic-N}} = \frac{k}{4\pi} \int \mathcal{L}_{\text{Extended-NH}} + \mathcal{L}_{\text{Enhanced-BNH}}, \quad (2.6)$$

where

$$\begin{aligned}\mathcal{L}_{\text{Extended-NH}} &= \alpha_0 \left[\omega_a R^a(\omega^b) - 2s R(\omega) + \frac{1}{\ell^2} e_a R^a(e^b) \right. \\ &\quad \left. - \frac{2}{\ell^2} m R(\tau) \right] + \alpha_1 \left[e_a R^a(\omega^b) \right. \\ &\quad \left. + \omega_a R^a(e^b) - 2m R(\omega) - 2s R(\tau) \right],\end{aligned}\quad (2.7)$$

and

$$\begin{aligned}\mathcal{L}_{\text{Enhanced-BNH}} &= \beta_0 \left[b_a R^a(\omega^b) + \omega_a R^a(b^b) - 2z R(\omega) \right. \\ &\quad \left. - s ds - \frac{2}{\ell^2} y R(\tau) - \frac{1}{\ell^2} m dm \right. \\ &\quad \left. + \frac{1}{\ell^2} t_a R^a(e^b) + \frac{1}{\ell^2} e_a R^a(t^b) \right] \\ &\quad + \beta_1 \left[e_a R^a(b^b) + b_a R^a(e^b) \right. \\ &\quad \left. + t_a R^a(\omega^b) + \omega_a R^a(t^b) - 2y R(\omega) \right. \\ &\quad \left. - 2z R(\tau) - 2mds \right].\end{aligned}\quad (2.8)$$

Here we have used the curvatures

$$\begin{aligned}R(\omega) &= d\omega, \\ R(\tau) &= d\tau, \\ R^a(\omega^b) &= d\omega^a + \epsilon^{ac} \omega \omega_c + \frac{1}{\ell^2} \epsilon^{ac} \tau e_c, \\ R^a(e^b) &= de^a + \epsilon^{ac} \omega e_c + \epsilon^{ac} \tau \omega_c, \\ R^a(t^b) &= dt^a + \epsilon^{ac} \omega t_c + \epsilon^{ac} \tau b_c + \epsilon^{ac} s e_c + \epsilon^{ac} m \omega_c, \\ R^a(b^b) &= db^a + \epsilon^{ac} \omega b_c + \epsilon^{ac} s \omega_c + \frac{1}{\ell^2} \epsilon^{ac} \tau t_c \\ &\quad + \frac{1}{\ell^2} \epsilon^{ac} m e_c.\end{aligned}\quad (2.9)$$

Moreover, one can also introduce the curvatures

$$\begin{aligned}R(m) &= dm + \epsilon^{ac} \omega_a e_c, \\ R(s) &= ds + \frac{1}{2} \epsilon^{ac} \omega_a \omega_c + \frac{1}{2\ell^2} \epsilon^{ac} e_a e_c, \\ R(y) &= dy + \epsilon^{ac} \omega_a t_c + \epsilon^{ac} e_a b_c, \\ R(z) &= dz + \epsilon^{ac} \omega_a b_c + \frac{1}{\ell^2} \epsilon^{ac} e_a t_c.\end{aligned}\quad (2.10)$$

While the first Lagrangian (2.7) corresponds to the extended Newton-Hooke gravity Lagrangian [37,38], the second one (2.8) coincides with the enhanced Bargmann-Newton-Hooke gravity Lagrangian [40] plus an exotic sector proportional to β_0 . The extended Newton-Hooke gravity can be seen as the non-relativistic limit of the $\text{AdS} \oplus \mathfrak{u}(1)^2$ gravity. On the other hand, the enhanced Bargmann-Newton-Hooke gravity theory appears as a non-relativistic limit of the coadjoint $\text{AdS} \oplus \mathfrak{u}(1)^2$ gravity [40]. Both Lagrangians define the exotic Newtonian gravity theory which in the vanishing cosmological constant limit $\ell \rightarrow \infty$ reproduces the extended Newtonian gravity theory [28,39].

3 Exotic Newtonian superalgebra and semigroup expansion method

In this section, we show that the exotic Newtonian superalgebra can be obtained by considering the S -expansion [49] of the relativistic $\mathcal{N} = 2$ AdS superalgebra. Our approach provides not only the (anti-)commutation relations of the exotic Newtonian superalgebra but also the non-vanishing components of the invariant tensor, which are essential for the construction of a CS action. Before presenting the explicit derivation of the exotic Newtonian superalgebra, we first review the S -expansion procedure and its main features.

The expansion method was first introduced in [53] and subsequently developed in [48, 49] to obtain a new (bigger) Lie algebra \mathfrak{G} from an original one \mathfrak{g} . In particular, the expansion based on semigroups [49, 54–59] allows to define a new Lie algebra, $\mathfrak{G} = S \times \mathfrak{g}$, by combining the structure constants of a Lie algebra \mathfrak{g} with the elements of a semigroup S . The S -expanded Lie (super)algebra satisfies

$$[T_{(A,\alpha)}, T_{(B,\beta)}] = K_{\alpha\beta}{}^\gamma C_{AB}{}^C T_{(C,\gamma)}, \quad (3.1)$$

where $C_{AB}{}^C$ are the structure constants of the original Lie (super)algebra \mathfrak{g} , $T_{(A,\alpha)} = \lambda_\alpha T_A$ denote the generators of the expanded (super)algebra \mathfrak{G} , and $K_{\alpha\beta}{}^\gamma$ is the so-called 2-selector, which satisfies

$$K_{\alpha\beta}{}^\gamma = \begin{cases} 1 & \text{when } \lambda_\alpha \lambda_\beta = \lambda_\gamma, \\ 0 & \text{otherwise.} \end{cases} \quad (3.2)$$

A smaller (super)algebra can be extracted from the expanded one, called as 0_S -reduced (super)algebra, by considering a semigroup with zero element $0_S \in S$ and imposing the condition $0_S T_A = 0$. An alternative method to extract a smaller (super)algebra from the expanded one \mathfrak{G} requires to consider a subset decomposition of the semigroup $S = \bigcup_{p \in I} S_p$ which satisfies the same structure than a subspace decomposition of the original (super)algebra $\mathfrak{g} = \bigoplus_{p \in I} V_p$, where I denotes a set of indices. Then, the subalgebra

$$\mathfrak{G}_R = \bigoplus_{p \in I} S_p \times V_p, \quad (3.3)$$

is called the resonant subalgebra of the expanded (super) algebra \mathfrak{G} .

It is important to mention that, unlike the power series expansion carried out on the Maurer–Cartan forms [48], the S -expansion method is defined entirely on the Lie (super)algebra \mathfrak{g} without considering the group manifold. Furthermore, for a particular choice of the semigroup S , the S -expansion procedure can reproduce the expanded (super)algebras obtained through the Maurer–Cartan forms power series expansion. Moreover, the S -expansion method provides us with the non-vanishing components of the invariant tensor of the S -expanded (super)algebra in terms of the invariant tensor of the original (super)algebra. It is important

to mention that both expansion mechanisms have been useful not only to construct new relativistic (super)gravity theories [60–73] but also at the non-relativistic level [43, 47, 50, 51, 74–80].

3.1 Exotic Newtonian superalgebra by expanding the $\mathcal{N} = 2$ AdS superalgebra

Here we explore a supersymmetric extension of the exotic Newtonian algebra by considering the S -expansion of the $\mathcal{N} = 2$ AdS superalgebra given by an $\mathfrak{so}(2)$ extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra. To this end, we consider the procedure used in [46, 51] to obtain a proper exotic Newtonian superalgebra which admits an invariant supertrace.

Let us first consider an $\mathfrak{so}(2)$ extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra which is spanned by the set of generators $\{\tilde{J}_A, \tilde{P}_A, \tilde{T}, \tilde{U}, \tilde{Q}_\alpha^i\}$ and satisfies the following (anti-)commutation relations [81]:

$$\begin{aligned} [\tilde{J}_A, \tilde{J}_B] &= \epsilon_{ABC} \tilde{J}^C, \\ [\tilde{J}_A, \tilde{P}_B] &= \epsilon_{ABC} \tilde{P}^C, \\ [\tilde{P}_A, \tilde{P}_B] &= \frac{1}{\ell^2} \epsilon_{ABC} \tilde{J}^C, \\ [\tilde{J}_A, \tilde{Q}_\alpha^i] &= -\frac{1}{2} (\gamma_A)_\alpha{}^\beta \tilde{Q}_\beta^i, \\ [\tilde{P}_A, \tilde{Q}_\alpha^i] &= -\frac{1}{2\ell} (\gamma_A)_\alpha{}^\beta \tilde{Q}_\beta^i, \\ [\tilde{T}, \tilde{Q}_\alpha^i] &= \frac{1}{2} \epsilon^{ij} \tilde{Q}_\beta^j, \\ \{\tilde{Q}_\alpha^i, \tilde{Q}_\beta^j\} &= -\frac{\delta_{ij}}{\ell} (\gamma^A C)_{\alpha\beta} \tilde{J}_A - \delta_{ij} (\gamma^A C)_{\alpha\beta} \tilde{P}_A \\ &\quad - C_{\alpha\beta} \epsilon^{ij} \left(\tilde{U} + \frac{1}{\ell} \tilde{T} \right), \end{aligned} \quad (3.4)$$

where $\alpha, \beta = 1, 2$, $i, j = 1, 2$ denotes the number of supercharges and $A, B, C = 0, 1, 2$ are the Lorentz indices which are raised and lowered with the Minkowski metric $\eta_{AB} = (-1, 1, 1)$. Here, C is the charge conjugation matrix,

$$C_{\alpha\beta} = C^{\alpha\beta} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad (3.5)$$

which satisfies $C^T = -C$ and $C\gamma^A = (C\gamma^A)^T$ with γ^A being the Dirac matrices in three spacetime dimensions. Let us note that the presence of the generators $\{\tilde{T}, \tilde{U}\}$ is required not only to construct an exotic Newtonian superalgebra with non-degenerate invariant tensor, but also to establish a proper flat limit $\ell \rightarrow \infty$. The non-degeneracy of the invariant tensor is crucial to get a well-defined CS (super)gravity action and is related to the Physical requirement that the CS action involves a kinematical term for each gauge field. In particular, the non-vanishing components of the invariant tensor for the $\mathfrak{so}(2)$

extension of the $\mathfrak{osp}(2, 2) \otimes \mathfrak{sp}(2)$ superalgebra read

$$\begin{aligned} \langle \tilde{J}_A \tilde{J}_B \rangle &= \mu_0 \eta_{AB}, \\ \langle \tilde{J}_A \tilde{P}_B \rangle &= \mu_1 \eta_{AB}, \\ \langle \tilde{P}_A \tilde{P}_B \rangle &= \frac{\mu_0}{\ell^2} \eta_{AB}, \\ \langle \tilde{T} \tilde{T} \rangle &= \mu_0, \\ \langle \tilde{T} \tilde{U} \rangle &= \mu_1, \\ \langle \tilde{U} \tilde{U} \rangle &= -\frac{\mu_1}{\ell^2}, \\ \langle \tilde{Q}_\alpha^i \tilde{Q}_\beta^j \rangle &= 2 \left(\mu_1 + \frac{\mu_0}{\ell} \right) C_{\alpha\beta} \delta^{ij}, \end{aligned} \quad (3.6)$$

where μ_0 and μ_1 are arbitrary constants. In the flat limit $\ell \rightarrow \infty$, the invariant tensor corresponds to the $\mathcal{N} = 2$ super-Poincaré ones [81].

Before considering the S -expansion of the $\mathcal{N} = 2$ AdS superalgebra (3.4), it is convenient to decompose the indices A, B as

$$A \rightarrow (0, a) \quad (3.7)$$

with $a = 1, 2$. Then let $V_0 = \{\tilde{J}_0, \tilde{P}_0, \tilde{T}, \tilde{U}, \tilde{Q}_\alpha^+\}$ and $V_1 = \{\tilde{J}_a, \tilde{P}_a, \tilde{Q}_\alpha^-\}$ be a subset decomposition of the $\mathcal{N} = 2$ AdS superalgebra where

$$\tilde{Q}_\alpha^\pm = \frac{1}{\sqrt{2}} \left(\tilde{Q}_\alpha^1 \pm \epsilon_{\alpha\beta} \tilde{Q}_\beta^2 \right). \quad (3.8)$$

One can see that the subspace decomposition of the $\mathcal{N} = 2$ AdS superalgebra satisfies

$$[V_0, V_0] \subset V_0, \quad [V_0, V_1] \subset V_1, \quad [V_1, V_1] \subset V_0. \quad (3.9)$$

Let us consider now $S_E^{(4)} = \{\lambda_0, \lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5\}$ as the relevant abelian semigroup whose elements satisfy the following multiplication law:

$$\lambda_\alpha \lambda_\beta = \begin{cases} \lambda_{\alpha+\beta} & \text{if } \alpha + \beta < 5, \\ \lambda_5 & \text{if } \alpha + \beta \geq 5, \end{cases} \quad (3.10)$$

with $\lambda_5 = 0_S$ being the zero element of the semigroup which satisfies $0_S \lambda_i = 0_S$. Then, let $S_E^{(4)} = S_0 \cup S_1$ be a resonant semigroup decomposition where

$$S_0 = \{\lambda_0, \lambda_2, \lambda_4, \lambda_5\}, \quad (3.11)$$

$$S_1 = \{\lambda_1, \lambda_3, \lambda_5\}. \quad (3.12)$$

One can see that such semigroup decomposition satisfies the same algebraic structure than the subspace decomposition,

$$S_0 \cdot S_0 \subset S_0, \quad S_0 \cdot S_1 \subset S_1, \quad S_1 \cdot S_1 \subset S_0. \quad (3.13)$$

Then one finds a new expanded superalgebra after extracting a resonant subalgebra from the $S_E^{(4)}$ -expansion of the $\mathcal{N} = 2$ AdS superalgebra,

$$\mathfrak{G}_R = S_0 \times V_0 \oplus S_1 \times V_1, \quad (3.14)$$

and considering a 0_S -reduction. The novel superalgebra is spanned by the exotic Newtonian bosonic generators

$$\{J, G_a, S, B_a, Z, H, P_a, M, T_a, Y\} \quad (3.15)$$

along with

$$\{Y_1, Y_2, Y_3, U_1, U_2, U_3, Q_\alpha^+, Q_\alpha^-, R_\alpha, W_\alpha^-, W_\alpha^+\}. \quad (3.16)$$

The expanded generators (3.15) and (3.16) are related to the $\mathcal{N} = 2$ super AdS ones through

λ_5		
λ_4	$Z, Y, Y_3, U_3, W_\alpha^+$	
λ_3		B_a, T_a, W_α^-
λ_2	S, M, Y_2, U_2, R_α	
λ_1		G_a, P_a, Q_α^-
λ_0	$J, H, Y_1, U_1, Q_\alpha^+$	
	$\tilde{J}_0, \tilde{P}_0, \tilde{T}, \tilde{U}, \tilde{Q}_\alpha^+$	$\tilde{J}_a, \tilde{P}_a, \tilde{Q}_\alpha^-$

(3.17)

Using the (anti-)commutation relations of the $\mathcal{N} = 2$ AdS superalgebra (3.4) and the multiplication law of the semigroup (3.10), one can show that the expanded generators satisfy the exotic Newtonian algebra (2.1) along with the following commutation relations:

$$\begin{aligned} [J, Q_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta Q_\beta^\pm, & [J, R_\alpha] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta R_\beta, \\ [J, W_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta W_\beta^\pm, \\ [S, Q_\alpha^+] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta R_\beta, & [S, Q_\alpha^-] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta W_\beta^-, \\ [S, R_\alpha] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta W_\beta^+, \\ [H, Q_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta Q_\beta^\pm, & [H, R_\alpha] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta R_\beta, \\ [H, W_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta W_\beta^\pm, \\ [M, Q_\alpha^+] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta R_\beta, & [M, Q_\alpha^-] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta W_\beta^-, \\ [M, R_\alpha] &= -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta W_\beta^+, \\ [G_a, Q_\alpha^+] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta Q_\beta^-, & [G_a, Q_\alpha^-] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta R_\beta, \\ [G_a, R_\alpha] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta W_\beta^-, \\ [G_a, W_\alpha^-] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta W_\beta^+, & [B_a, Q_\alpha^\pm] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta W_\beta^\mp, \end{aligned}$$

$$\begin{aligned}
[P_a, Q_\alpha^+] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta Q_\beta^-, \\
[P_a, Q_\alpha^-] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta R_\beta, \quad [P_a, R_\alpha] = -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta W_\beta^-, \\
[P_a, W_\alpha^-] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta W_\beta^+, \\
[T_a, Q_\alpha^\pm] &= -\frac{1}{2\ell} (\gamma_a)_\alpha^\beta W_\beta^\mp, \quad [Y_1, Q_\alpha^\pm] = \pm \frac{1}{2} (\gamma_0)_{\alpha\beta} Q_\beta^\pm, \\
[Y_1, R_\alpha] &= \frac{1}{2} (\gamma_0)_{\alpha\beta} R_\beta, \\
[Y_1, W_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_{\alpha\beta} W_\beta^\pm, \quad [Y_2, Q_\alpha^+] = \frac{1}{2} (\gamma_0)_{\alpha\beta} R_\beta, \\
[Y_2, Q_\alpha^-] &= -\frac{1}{2} (\gamma_0)_{\alpha\beta} W_\beta^-, \\
[Y_2, R_\alpha] &= \frac{1}{2} (\gamma_0)_{\alpha\beta} W_\beta^+, \quad [Y_3, Q_\alpha^+] = \frac{1}{2} (\gamma_0)_{\alpha\beta} W_\beta^+, \\
[Z, Q_\alpha^+] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta W_\beta^+, \quad [Y, Q_\alpha^+] = -\frac{1}{2\ell} (\gamma_0)_\alpha^\beta W_\beta^+, \\
\end{aligned} \tag{3.18}$$

and the following anti-commutation relations:

$$\begin{aligned}
\{Q_\alpha^+, Q_\beta^+\} &= -\frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} J - (\gamma^0 C)_{\alpha\beta} H \\
&\quad - \frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} Y_1 - (\gamma^0 C)_{\alpha\beta} U_1, \\
\{Q_\alpha^+, Q_\beta^-\} &= -\frac{1}{\ell} (\gamma^a C)_{\alpha\beta} G_a - (\gamma^a C)_{\alpha\beta} P_a, \\
\{Q_\alpha^-, Q_\beta^-\} &= -\frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} S - (\gamma^0 C)_{\alpha\beta} M \\
&\quad + \frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} Y_2 + (\gamma^0 C)_{\alpha\beta} U_2, \\
\{Q_\alpha^+, R_\beta\} &= -\frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} S - (\gamma^0 C)_{\alpha\beta} M \\
&\quad - \frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} Y_2 - (\gamma^0 C)_{\alpha\beta} U_2, \\
\{Q_\alpha^-, R_\beta\} &= -\frac{1}{\ell} (\gamma^a C)_{\alpha\beta} B_a - (\gamma^a C)_{\alpha\beta} T_a, \\
\{Q_\alpha^+, W_\beta^-\} &= -\frac{1}{\ell} (\gamma^a C)_{\alpha\beta} B_a - (\gamma^a C)_{\alpha\beta} T_a, \\
\{Q_\alpha^-, W_\beta^-\} &= -\frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} Z - (\gamma^0 C)_{\alpha\beta} Y \\
&\quad + \frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} Y_3 + (\gamma^0 C)_{\alpha\beta} U_3, \\
\{Q_\alpha^+, W_\beta^+\} &= -\frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} Z - (\gamma^0 C)_{\alpha\beta} Y \\
&\quad - \frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} Y_3 - (\gamma^0 C)_{\alpha\beta} U_3, \\
\{R_\alpha, R_\beta\} &= -\frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} Z - (\gamma^0 C)_{\alpha\beta} Y \\
&\quad - \frac{1}{\ell} (\gamma^0 C)_{\alpha\beta} Y_3 - (\gamma^0 C)_{\alpha\beta} U_3. \tag{3.19}
\end{aligned}$$

The superalgebra given by (2.1), (3.18) and (3.19) corresponds to a supersymmetric extension of the exotic Newtonian algebra [39]. Let us note that the expansion proce-

dures implies the presence of additional bosonic generators $\{Y_1, Y_2, Y_3, U_1, U_2, U_3\}$ which, as we shall see, ensure having not only a non-degenerate invariant tensor but also a well-defined non-vanishing cosmological constant limit $\ell \rightarrow \infty$. Indeed, in the flat limit, one can see that the superalgebra reduces to the extended Newtonian superalgebra introduced in [28] in presence of the additional generators $\{Y_1, Y_2, Y_3\}$ and central charges $\{U_1, U_2, U_3\}$. The extra bosonic generators $\{Y_1, Y_2, Y_3\}$ are expansions of the relativistic R -symmetry generator \mathcal{T} and act non-trivially on the fermionic charges.

Interestingly, the exotic Newtonian superalgebra (2.1), (3.18) and (3.19) can be written as two copies of the so-called enhanced Nappi–Witten algebra [40, 47], one of which is augmented by supersymmetry. Indeed, let us consider the following redefinition of the generators:

$$\begin{aligned}
G_a &= L_a - \tilde{L}_a, \quad P_a = \frac{1}{\ell} (L_a + \tilde{L}_a), \quad Q_\alpha^+ = \sqrt{\frac{2}{\ell}} \mathcal{Q}_\alpha^+, \\
B_a &= N_a - \tilde{N}_a, \quad T_a = \frac{1}{\ell} (N_a + \tilde{N}_a), \quad Q_\alpha^- = \sqrt{\frac{2}{\ell}} \mathcal{Q}_\alpha^-, \\
J &= L + \tilde{L}, \quad H = \frac{1}{\ell} (L - \tilde{L}), \quad R_\alpha = \sqrt{\frac{2}{\ell}} \mathcal{R}_\alpha, \\
S &= N + \tilde{N}, \quad M = \frac{1}{\ell} (N - \tilde{N}), \\
W_\alpha^+ &= \sqrt{\frac{2}{\ell}} \mathcal{W}_\alpha^+, \quad Z = B + \tilde{B}, \quad Y = \frac{1}{\ell} (B - \tilde{B}), \\
W_\alpha^- &= \sqrt{\frac{2}{\ell}} \mathcal{W}_\alpha^-, \\
Y_1 &= X_1 - \tilde{X}_1, \quad Y_2 = X_2 - \tilde{X}_2, \\
Y_3 &= X_3 - \tilde{X}_3, \\
U_1 &= \frac{1}{\ell} \tilde{X}_1, \quad U_2 = \frac{1}{\ell} \tilde{X}_2, \\
U_3 &= \frac{1}{\ell} \tilde{X}_3, \tag{3.20}
\end{aligned}$$

where the set of generators

$$\{L, N, B, L_a, N_a, X_1, X_2, X_3, \mathcal{Q}_\alpha^+, \mathcal{Q}_\alpha^-, \mathcal{R}_\alpha, \mathcal{W}_\alpha^-, \mathcal{W}_\alpha^+\} \tag{3.21}$$

satisfies a supersymmetric extension of the enhanced Nappi–Witten algebra. In particular, the generators of the enhanced Nappi–Witten superalgebra obey the following commutation relations:

$$\begin{aligned}
[L, L_a] &= \epsilon_{ab} L_b, \quad [L_a, L_b] = -\epsilon_{ab} N, \\
[N, L_a] &= \epsilon_{ab} N_b, \\
[L, N_a] &= \epsilon_{ab} N_b, \quad [L_a, N_b] = -\epsilon_{ab} B, \\
[L, \mathcal{Q}_\alpha^\pm] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathcal{Q}_\beta^\pm,
\end{aligned}$$

$$\begin{aligned}
 [L, \mathcal{R}_\alpha] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathcal{R}_\beta, \quad [L, \mathcal{W}_\alpha^\pm] = -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathcal{W}_\beta^\pm, \\
 [N, \mathcal{Q}_\alpha^+] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathcal{R}_\beta, \\
 [N, \mathcal{Q}_\alpha^-] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathcal{W}_\beta^-, \quad [N, \mathcal{R}_\alpha] = -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathcal{W}_\beta^+, \\
 [B, \mathcal{Q}_\alpha^+] &= -\frac{1}{2} (\gamma_0)_\alpha^\beta \mathcal{W}_\beta^+, \\
 [X_1, \mathcal{Q}_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_{\alpha\beta} \mathcal{Q}_\beta^\pm, \quad [X_1, \mathcal{R}_\alpha] = \frac{1}{2} (\gamma_0)_{\alpha\beta} \mathcal{R}_\beta, \\
 [X_1, \mathcal{W}_\alpha^\pm] &= \pm \frac{1}{2} (\gamma_0)_{\alpha\beta} \mathcal{W}_\beta^\pm, \\
 [X_2, \mathcal{Q}_\alpha^+] &= \frac{1}{2} (\gamma_0)_{\alpha\beta} \mathcal{R}_\beta, \quad [X_2, \mathcal{Q}_\alpha^-] = -\frac{1}{2} (\gamma_0)_{\alpha\beta} \mathcal{W}_\beta^-, \\
 [X_2, \mathcal{R}_\alpha] &= \frac{1}{2} (\gamma_0)_{\alpha\beta} \mathcal{W}_\beta^+, \\
 [X_3, \mathcal{Q}_\alpha^+] &= \frac{1}{2} (\gamma_0)_{\alpha\beta} \mathcal{W}_\beta^+, \quad [L_a, \mathcal{Q}_\alpha^+] = -\frac{1}{2} (\gamma_a)_\alpha^\beta \mathcal{Q}_\beta^-, \\
 [L_a, \mathcal{Q}_\alpha^-] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta \mathcal{R}_\beta, \\
 [L_a, \mathcal{R}_\alpha] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta \mathcal{W}_\beta^-, \quad [L_a, \mathcal{W}_\alpha^-] = -\frac{1}{2} (\gamma_a)_\alpha^\beta \mathcal{W}_\beta^+, \\
 [N_a, \mathcal{Q}_\alpha^+] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta \mathcal{W}_\beta^-, \\
 [N_a, \mathcal{Q}_\alpha^-] &= -\frac{1}{2} (\gamma_a)_\alpha^\beta \mathcal{W}_\beta^+, \quad (3.22)
 \end{aligned}$$

along with the following anti-commutation relations:

$$\begin{aligned}
 \{\mathcal{Q}_\alpha^+, \mathcal{Q}_\beta^+\} &= -(\gamma_0 C)_{\alpha\beta} L - (\gamma_0 C)_{\alpha\beta} X_1, \\
 \{\mathcal{Q}_\alpha^-, \mathcal{Q}_\beta^-\} &= -(\gamma_0 C)_{\alpha\beta} N + (\gamma_0 C)_{\alpha\beta} X_2, \\
 \{\mathcal{Q}_\alpha^+, \mathcal{R}_\beta\} &= -(\gamma_0 C)_{\alpha\beta} N - (\gamma_0 C)_{\alpha\beta} X_2, \\
 \{\mathcal{Q}_\alpha^-, \mathcal{W}_\beta^-\} &= -(\gamma_0 C)_{\alpha\beta} B + (\gamma_0 C)_{\alpha\beta} X_3, \\
 \{\mathcal{Q}_\alpha^+, \mathcal{W}_\beta^+\} &= -(\gamma_0 C)_{\alpha\beta} B - (\gamma_0 C)_{\alpha\beta} X_3, \\
 \{\mathcal{R}_\alpha, \mathcal{R}_\beta\} &= -(\gamma_0 C)_{\alpha\beta} B - (\gamma_0 C)_{\alpha\beta} X_3, \\
 \{\mathcal{Q}_\alpha^+, \mathcal{Q}_\beta^-\} &= -(\gamma^a C)_{\alpha\beta} L_a, \\
 \{\mathcal{Q}_\alpha^-, \mathcal{R}_\beta\} &= -(\gamma^a C)_{\alpha\beta} N_a, \\
 \{\mathcal{Q}_\alpha^+, \mathcal{W}_\beta^-\} &= -(\gamma^a C)_{\alpha\beta} N_a. \quad (3.23)
 \end{aligned}$$

While the Nappi–Witten superalgebra [46,47] requires three fermionic generators $\{\mathcal{Q}_\alpha^+, \mathcal{Q}_\alpha^-, \mathcal{R}_\alpha\}$, the supersymmetric extension of the enhanced Nappi–Witten algebra requires the introduction of five fermionic generators $\{\mathcal{Q}_\alpha^+, \mathcal{Q}_\alpha^-, \mathcal{R}_\alpha, \mathcal{W}_\alpha^+, \mathcal{W}_\alpha^-\}$. The presence of additional Majorana fermionic charges is due to the extra bosonic content and such fermionic generators are necessary to satisfy the Jacobi identity.¹ On the other hand, the set of generators

¹ Let us mention, here, that the presence of extra fermionic generators (namely besides the gravitino) in supergravity has been considered also in higher-dimensional (relativistic) cases, see e.g. [69,82–85].

$$\{\tilde{L}, \tilde{N}, \tilde{B}, \tilde{L}_a, \tilde{N}_a, \tilde{X}_1, \tilde{X}_2, \tilde{X}_3\} \quad (3.24)$$

satisfies the bosonic enhanced Nappi–Witten algebra [40,47], that is

$$\begin{aligned}
 [\tilde{L}, \tilde{L}_a] &= \epsilon_{ab} \tilde{L}_b, \quad [\tilde{L}_a, \tilde{L}_b] = -\epsilon_{ab} \tilde{N}, \\
 [\tilde{N}, \tilde{L}_a] &= \epsilon_{ab} \tilde{N}_b, \\
 [\tilde{L}, \tilde{N}_a] &= \epsilon_{ab} \tilde{N}_b, \quad [\tilde{L}_a, \tilde{N}_b] = -\epsilon_{ab} \tilde{B}, \quad (3.25)
 \end{aligned}$$

supplemented with the additional bosonic generators $\{\tilde{X}_1, \tilde{X}_2, \tilde{X}_3\}$. Although the extra bosonic generators $\{\tilde{X}_1, \tilde{X}_2, \tilde{X}_3\}$ do not appear explicitly into the commutation relations, they are essential to define a non-degenerate invariant tensor. It is important to mention that the exotic Newtonian superalgebra can alternatively be obtained by expanding the enhanced Nappi–Witten superalgebra (3.22)–(3.23), following the same procedure used in [47] (more details about the expansion of the enhanced Nappi–Witten superalgebra and its invariant tensor are given in “Appendix A”).

4 Exotic Newtonian supergravity action and flat limit

In this section, we shall explore the explicit construction of a CS action based on the exotic Newtonian superalgebra given by (2.1), (3.18) and (3.19). As we shall see, the exotic Newtonian superalgebra allows us to accommodate a cosmological constant to the Newtonian supergravity.

The non-vanishing components of the invariant tensor for the exotic Newtonian superalgebra can be obtained from the $\mathcal{N} = 2$ super AdS ones considering Theorem VII of Ref. [49]. In particular, the exotic Newtonian superalgebra admits the non-vanishing components of the invariant tensor for the extended Newton–Hooke superalgebra [47],

$$\begin{aligned}
 \langle JS \rangle &= -\alpha_0, & \langle G_a G_b \rangle &= \alpha_0 \delta_{ab}, \\
 \langle JM \rangle &= \langle HS \rangle = -\alpha_1, & \langle G_a P_b \rangle &= \alpha_1 \delta_{ab}, \\
 \langle HM \rangle &= -\frac{\alpha_0}{\ell^2}, & \langle P_a P_b \rangle &= \frac{\alpha_0}{\ell^2} \delta_{ab}, \\
 \langle Y_1 Y_2 \rangle &= \alpha_0 & \langle \mathcal{Q}_\alpha^- \mathcal{Q}_\beta^- \rangle &= \langle \mathcal{Q}_\alpha^+ \mathcal{R}_\beta \rangle \\
 & & &= 2 \left(\alpha_1 + \frac{\alpha_0}{\ell} \right) C_{\alpha\beta}, \\
 \langle U_1 U_2 \rangle &= -\frac{\alpha_1}{\ell^2}, & \langle Y_1 U_2 \rangle &= \langle Y_2 U_1 \rangle = \alpha_1, \quad (4.1)
 \end{aligned}$$

along with

$$\begin{aligned}
 \langle SS \rangle &= \langle JZ \rangle = -\beta_0, \\
 \langle G_a B_b \rangle &= \beta_0 \delta_{ab}, \\
 \langle MS \rangle &= \langle HZ \rangle = \langle JY \rangle = -\beta_1, \\
 \langle P_a B_b \rangle &= \langle G_a T_b \rangle = \beta_1 \delta_{ab},
 \end{aligned}$$

$$\begin{aligned}
\langle MM \rangle &= \langle HY \rangle = -\frac{\beta_0}{\ell^2}, \quad \langle P_a T_b \rangle = \frac{\beta_0}{\ell^2} \delta_{ab}, \\
\langle Y_1 Y_3 \rangle &= \langle Y_2 Y_2 \rangle = \beta_0, \\
\langle Q_\alpha^+ W_\beta^+ \rangle &= \langle Q_\alpha^- W_\beta^- \rangle = \langle R_\alpha R_\beta \rangle = 2 \left(\beta_1 + \frac{\beta_0}{\ell} \right) C_{\alpha\beta}, \\
\langle U_1 U_3 \rangle &= \langle U_2 U_2 \rangle = -\frac{\beta_1}{\ell^2}, \\
\langle Y_1 U_3 \rangle &= \langle Y_3 U_1 \rangle = \langle Y_2 U_2 \rangle = \beta_1,
\end{aligned} \quad (4.2)$$

where the extended Newtonian parameters are related to the relativistic $\mathcal{N} = 2$ super AdS ones through the semigroup elements as

$$\begin{aligned}
\alpha_0 &= \lambda_2 \mu_0, \quad \alpha_1 = \lambda_2 \mu_1, \\
\beta_0 &= \lambda_4 \mu_0, \quad \beta_1 = \lambda_4 \mu_1.
\end{aligned} \quad (4.3)$$

One can notice that the invariant tensor is given by two families proportional to α and β , respectively. In particular, the components proportional to α_0 and β_0 are related to an exotic sector of a non-relativistic supergravity theory. In the vanishing cosmological constant limit $\ell \rightarrow \infty$, we recover the invariant tensor for the extended Bargmann [27] and the extended Newtonian superalgebras [28]. Although one could consider additional components coming from the expansion with the λ_0 element, we shall omit such components since they are related to trivial CS terms. On the other hand, although the invariant tensor (4.1) is non-degenerate for the extended Newton–Hooke superalgebra, the non-degeneracy of the invariant supertrace for the exotic Newtonian superalgebra requires also to consider the components given by (4.2). As we shall see, the non-degeneracy of the bilinear invariant supertrace implies that the CS action involves a kinematical term for each gauge field.

The gauge connection one-form for the exotic Newtonian superalgebra is given by

$$\begin{aligned}
A &= \tau H + e^a P_a + \omega J + \omega^a G_a + m M + s S + t^a T_a \\
&\quad + b^a B_a + y Y + z Z + y_1 Y_1 + y_2 Y_2 + y_3 Y_3 \\
&\quad + u_1 U_1 + u_2 U_2 + u_3 U_3 + \bar{\psi}^+ Q^+ + \bar{\psi}^- Q^- + \bar{\rho} R \\
&\quad + \bar{\phi}^+ W^+ + \bar{\phi}^- W^-.
\end{aligned} \quad (4.4)$$

The corresponding curvature two-form $F = dA + \frac{1}{2} [A, A]$ reads

$$\begin{aligned}
F &= R(\tau) H + R^a(e^b) P_a + R(\omega) J + R^a(\omega^b) G_a \\
&\quad + R(m) M + R(s) S + R^a(t^b) T_a \\
&\quad + F^a(b^b) B_a + F(y) Y + F(z) Z + F(y_1) Y_1 \\
&\quad + F(y_2) Y_2 + F(y_3) Y_3 + F(u_1) U_1
\end{aligned}$$

$$\begin{aligned}
&+ F(u_2) U_2 + F(u_3) U_3 + \nabla \bar{\psi}^+ Q^+ + \nabla \bar{\psi}^- Q^- \\
&+ \nabla \bar{\rho} R + \nabla \bar{\phi}^+ W^+ + \nabla \bar{\phi}^- W^-.
\end{aligned} \quad (4.5)$$

Here, the bosonic curvature two-forms are given by

$$\begin{aligned}
F(\omega) &= R(\omega) + \frac{1}{2\ell} \bar{\psi}^+ \gamma^0 \psi^+, \\
F(y_1) &= dy_1 + \frac{1}{2\ell} \bar{\psi}^+ \gamma^0 \psi^+, \\
F^a(\omega^b) &= R^a(\omega^b) + \frac{1}{\ell} \bar{\psi}^+ \gamma^a \psi^-, \\
F(y_2) &= dy_2 - \frac{1}{2\ell} \bar{\psi}^- \gamma^0 \psi^- + \frac{1}{\ell} \bar{\psi}^+ \gamma^0 \rho, \\
F(\tau) &= R(\tau) + \frac{1}{2} \bar{\psi}^+ \gamma^0 \psi^+, \\
F(y_3) &= dy_3 - \frac{1}{\ell} \bar{\psi}^- \gamma^0 \phi^- + \frac{1}{\ell} \bar{\psi}^+ \gamma^0 \phi^+ + \frac{1}{2\ell} \bar{\rho} \gamma^0 \rho, \\
F^a(e^b) &= R^a(e^b) + \bar{\psi}^+ \gamma^a \psi^-, \\
F(u_1) &= du_1 + \frac{1}{2} \bar{\psi}^+ \gamma^0 \psi^+, \\
F(s) &= R(s) + \frac{1}{2\ell} \bar{\psi}^- \gamma^0 \psi^- + \frac{1}{\ell} \bar{\psi}^+ \gamma^0 \rho, \\
F(u_2) &= du_2 - \frac{1}{2} \bar{\psi}^- \gamma^0 \psi^- + \frac{1}{2} \bar{\psi}^+ \gamma^0 \rho, \\
F^a(b^b) &= R^a(b^b) + \frac{1}{\ell} \bar{\psi}^- \gamma^a \rho + \frac{1}{\ell} \bar{\psi}^+ \gamma^a \phi^-, \\
F(u_3) &= du_3 - \bar{\psi}^- \gamma^0 \phi^- + \bar{\psi}^+ \gamma^0 \phi^+ + \frac{1}{2} \bar{\rho} \gamma^0 \rho, \\
F(m) &= R(m) + \frac{1}{2} \bar{\psi}^- \gamma^0 \psi^- + \bar{\psi}^+ \gamma^0 \rho, \\
F^a(t^b) &= R^a(t^b) + \bar{\psi}^- \gamma^a \rho + \bar{\psi}^+ \gamma^a \phi^-, \\
F(z) &= R(z) + \frac{1}{\ell} \bar{\psi}^- \gamma^0 \phi^- + \frac{1}{\ell} \bar{\psi}^+ \gamma^0 \phi^+ + \frac{1}{2\ell} \bar{\rho} \gamma^0 \rho, \\
F(y) &= R(y) + \bar{\psi}^- \gamma^0 \phi^- + \bar{\psi}^+ \gamma^0 \phi^+ + \frac{1}{2} \bar{\rho} \gamma^0 \rho,
\end{aligned} \quad (4.6)$$

where $R(\omega)$, $R^a(\omega^b)$, $R(\tau)$, $R^a(e^b)$, $R(s)$, $R^a(b^b)$, $R(m)$, $R^a(t^b)$, $R(z)$ and $R(y)$ are the respective bosonic curvature two-forms (2.9)–(2.10) of the exotic Newtonian algebra. On the other hand, the fermionic curvature two-forms reads

$$\begin{aligned}
\nabla \psi^+ &= d\psi^+ + \frac{1}{2} \omega \gamma_0 \psi^+ + \frac{1}{2\ell} \tau \gamma_0 \psi^+ - \frac{1}{2} y_1 \gamma_0 \psi^+ \\
\nabla \psi^- &= d\psi^- + \frac{1}{2} \omega \gamma_0 \psi^- + \frac{1}{2\ell} \tau \gamma_0 \psi^- + \frac{1}{2} \omega^a \gamma_a \psi^- \\
&\quad + \frac{1}{2\ell} e^a \gamma_a \psi^+ + \frac{1}{2} y_1 \gamma_0 \psi^- \\
\nabla \rho &= d\rho + \frac{1}{2} \omega \gamma_0 \rho + \frac{1}{2} s \gamma_0 \psi^+ + \frac{1}{2} \omega^a \gamma_a \psi^-
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2\ell} \tau \gamma_0 \rho + \frac{1}{2\ell} m \gamma_0 \psi^+ + \frac{1}{2\ell} e^a \gamma_a \psi^- \\
& - \frac{1}{2} y_1 \gamma_0 \rho - \frac{1}{2} y_2 \gamma_0 \psi^+, \\
\nabla \phi^+ & = d\phi^+ + \frac{1}{2} \omega \gamma_0 \psi^+ + \frac{1}{2} s \gamma_0 \rho + \frac{1}{2} \omega^a \gamma_a \phi^- \\
& + \frac{1}{2} b^a \gamma_a \psi^- + \frac{1}{2\ell} \tau \gamma_0 \phi^+ + \frac{1}{2\ell} m \gamma_0 \rho \\
& + \frac{1}{2\ell} e^a \gamma_a \phi^- + \frac{1}{2\ell} t^a \gamma_a \psi^- - \frac{1}{2} y_1 \gamma_0 \phi^+ \\
& - \frac{1}{2} y_2 \gamma_0 \rho - \frac{1}{2} y_3 \gamma_0 \psi^+, \\
\nabla \phi^- & = d\phi^- + \frac{1}{2} \omega \gamma_0 \phi^- + \frac{1}{2} s \gamma_0 \psi^- \\
& + \frac{1}{2} \omega^a \gamma_a \rho + \frac{1}{2} b^a \gamma_a \psi^+ + \frac{1}{2\ell} \tau \gamma_0 \phi^- + \frac{1}{2\ell} m \gamma_0 \psi^- \\
& + \frac{1}{2\ell} e^a \gamma_a \rho + \frac{1}{2\ell} t^a \gamma_a \psi^+ + \frac{1}{2} y_1 \gamma_0 \phi^- \\
& + \frac{1}{2} y_2 \gamma_0 \psi^- + \frac{1}{2} z \gamma_0 \psi^+ + \frac{1}{2\ell} y \gamma_0 \psi^+. \quad (4.7)
\end{aligned}$$

The gauge-invariant CS supergravity action based on the super exotic Newtonian algebra given by (2.1), (3.18) and (3.19) can be obtained by combining the gauge connection one-form (4.4) and the non-vanishing components of the invariant tensor (4.1) and (4.2) into the general expression for the CS action (2.5). In particular, the exotic Newtonian (EN) CS supergravity action can be written in terms of the super Extended Newton–Hooke (ENH) and super Enhanced Bargmann–Newton–Hooke (EBNH) supergravity Lagrangians, that is

$$I_{\text{Super-EN}} = \frac{k}{4\pi} \int \mathcal{L}_{\text{Super ENH}} + \mathcal{L}_{\text{Super EBNH}}, \quad (4.8)$$

where

$$\begin{aligned}
\mathcal{L}_{\text{Super-ENH}} & = \mathcal{L}_{\text{Extended-NH}} + \alpha_0 \left[2y_1 dy_2 - \frac{2}{\ell} \bar{\psi}^- \nabla \psi^- \right. \\
& \quad \left. - \frac{2}{\ell} \bar{\psi}^+ \nabla \rho - \frac{2}{\ell} \bar{\rho} \nabla \psi^+ \right] \\
& + \alpha_1 \left[2y_1 du_2 + 2y_2 du_1 + \frac{2}{\ell^2} u_1 du_2 \right. \\
& \quad \left. - 2\bar{\psi}^- \nabla \psi^- - 2\bar{\psi}^+ \nabla \rho - 2\bar{\rho} \nabla \psi^+ \right], \quad (4.9)
\end{aligned}$$

and

$$\begin{aligned}
\mathcal{L}_{\text{Super-EBNH}} & = \mathcal{L}_{\text{Enhanced-BNH}} + \beta_0 \left[2y_1 dy_3 + y_2 dy_2 \right. \\
& \quad \left. - \frac{2}{\ell} \bar{\psi}^+ \nabla \phi^+ \right. \\
& \quad \left. - \frac{2}{\ell} \bar{\phi}^+ \nabla \psi^+ - \frac{2}{\ell} \bar{\psi}^- \nabla \phi^- \right. \\
& \quad \left. - \frac{2}{\ell} \bar{\phi}^- \nabla \psi^- - \frac{2}{\ell} \bar{\rho} \nabla \rho \right]
\end{aligned}$$

$$\begin{aligned}
& + \beta_1 [2y_1 du_3 + 2y_3 du_1 + 2y_2 du_2 \\
& + \frac{2}{\ell^2} u_1 du_3 + \frac{1}{\ell^2} u_2 du_2 - 2\bar{\psi}^+ \nabla \phi^+ \\
& - 2\bar{\phi}^+ \nabla \psi^+ - 2\bar{\psi}^- \nabla \phi^- - 2\bar{\phi}^- \nabla \psi^- \\
& - 2\bar{\rho} \nabla \rho]. \quad (4.10)
\end{aligned}$$

Here, $\mathcal{L}_{\text{Extended-NH}}$ and $\mathcal{L}_{\text{Enhanced-BNH}}$ correspond to the Extended Newton–Hooke and Enhanced Bargmann–Newton–Hooke gravity Lagrangians which are given by (2.7) and (2.8), respectively. The exotic Newtonian supergravity CS action (4.8) generalizes the extended Newtonian supergravity theory [28] by introducing a cosmological constant and considering six additional bosonic gauge fields y_1, y_2, y_3, u_1, u_2 and u_3 . Interestingly, a supersymmetric extension of the extended Newtonian gravity [28, 39] appears in the vanishing cosmological constant limit $\ell \rightarrow \infty$. In such limit, the curvature two-forms reduce to the extended Newtonian ones. Furthermore, the fermionic gauge fields do not contribute anymore to the exotic sectors, along α_0 and β_0 , in the flat limit. Such exotic terms can be seen as the non-relativistic version of the exotic supergravity Lagrangian [52]. In particular, the exotic Newtonian supergravity CS action (4.8) can alternatively be recovered as an $S_E^{(4)}$ -expansion of the relativistic $\mathcal{N} = 2$ AdS supergravity CS action [81, 86, 87],

$$\begin{aligned}
I_{\text{AdS}}^{\mathcal{N}=2} & = \int \mu_0 \left[\omega_A d\omega^A + \frac{1}{3} \epsilon_{ABC} \omega^A \omega^B \omega^C + \frac{1}{\ell^2} e_A T^A \right. \\
& \quad \left. + \mathfrak{t} d\mathfrak{t} - \frac{2}{\ell} \tilde{\psi}^i \nabla \tilde{\psi}^i \right] \\
& + \mu_1 \left[2e_A R^A + \frac{1}{3\ell^2} \epsilon_{ABC} e^A e^B e^C + \mathfrak{t} du \right. \\
& \quad \left. - \frac{1}{\ell^2} u d\mathfrak{u} - 2\tilde{\psi}^i \nabla \tilde{\psi}^i \right], \quad (4.11)
\end{aligned}$$

where

$$\begin{aligned}
R^A & = d\omega^A + \frac{1}{2} \epsilon^{ABC} \omega_B \omega_C, \\
T^A & = de^A + \frac{1}{2} \epsilon^{ABC} \omega_B e_C, \\
\nabla \tilde{\psi}^i & = d\tilde{\psi}^i + \frac{1}{2} \omega^A \gamma_A \tilde{\psi}^i + \frac{1}{2\ell} e^A \gamma_A \tilde{\psi}^i + \mathfrak{t} \epsilon^{ij} \tilde{\psi}^j. \quad (4.12)
\end{aligned}$$

Indeed, the exotic Newtonian CS supergravity action appears after considering the $S_E^{(4)}$ -expansion of the $\mathcal{N} = 2$ super AdS gauge fields as

$$\begin{aligned}
\omega &= \lambda_0 \omega_0, \quad s = \lambda_2 \omega_0, \quad z = \lambda_4 \omega_0, \\
\omega_a &= \lambda_1 \omega_a, \quad b_a = \lambda_3 \omega_a, \\
\tau &= \lambda_0 e_0, \quad m = \lambda_2 e_0, \quad y = \lambda_4 e_0, \\
e_a &= \lambda_1 e_a, \quad t_a = \lambda_3 e_a, \\
\psi^+ &= \lambda_0 \tilde{\psi}^+, \quad \rho = \lambda_2 \tilde{\psi}^+, \quad \phi^+ = \lambda_4 \tilde{\psi}^+, \\
\psi^- &= \lambda_1 \tilde{\psi}^-, \quad \phi^- = \lambda_3 \tilde{\psi}^-, \\
y_1 &= \lambda_0 \mathfrak{t}, \quad y_2 = \lambda_2 \mathfrak{t}, \quad y_3 = \lambda_4 \mathfrak{t}, \\
u_1 &= \lambda_0 \mathfrak{u}, \quad u_2 = \lambda_2 \mathfrak{u}, \quad u_3 = \lambda_4 \mathfrak{u},
\end{aligned} \tag{4.13}$$

along with the expanded parameters (4.3) and defining

$$\tilde{\psi}_\alpha^\pm = \frac{1}{\sqrt{2}} \left(\tilde{\psi}_\alpha^1 \pm \epsilon_{\alpha\beta} \tilde{\psi}_\beta^2 \right). \tag{4.14}$$

On the other hand, the non-degeneracy of the invariant tensor (4.1) and (4.2) implies that the field equations derived from the CS action (4.8) are given by the vanishing of the curvature two-forms (4.6)–(4.7). Such curvatures transform covariantly under the supersymmetry transformation laws (the explicit supersymmetry transformation laws of the exotic Newtonian superalgebra are given in “Appendix B”).

5 Conclusions

In this work we have presented an exotic Newtonian CS supergravity theory in presence of a cosmological constant. The underlying non-relativistic superalgebra, which we have called as exotic Newtonian superalgebra, is obtained as an S -expansion of the $\mathcal{N} = 2$ AdS superalgebra considering a particular semigroup. Interestingly, we showed that the exotic Newtonian superalgebra can be written as two copies of the so-called enhanced Nappi–Witten algebra [40,50], one of which is supersymmetric. The new Newtonian supergravity action contains the extended Newton–Hooke CS supergravity term [44] as a sub-case. Furthermore, we showed that the most general extended Newtonian supergravity action appears in the flat limit $\ell \rightarrow \infty$ which contains the extended Bargmann supergravity [27] as a particular sub-case. It would be interesting to explore the matter coupling of the exotic Newtonian supergravity theory presented here. Although the exotic Newtonian supergravity theory differs from the extended Newtonian one at the action level, one could expect that the matter couplings of both theories behave similarly.

It would be worth considering further extensions of the Newtonian supergravity theory obtained here and in [28] by considering a Maxwellian generalization. One could expect that such generalizations are given by the respective supersymmetric extensions of the enlarged and Maxwellian extended Newtonian gravity presented in [50]. On the other hand, following [44], one could explore a Schrödinger extension [42,88] of the exotic Newtonian superalgebra. Moving towards Newtonian supergravity the-

ories and their Schrödinger extension can be useful, for instance, to approach supersymmetric field theories on non-relativistic curved backgrounds via localization [89–91].

Our results and those obtained in [46,51] could be extended to other relativistic (super)algebras. Indeed, it seems that the $S_E^{(4)}$ semigroup allows to obtain the respective Newtonian version of a relativistic (super)algebra. In particular, the procedure used here could be useful in presence of supersymmetry, where the study of the non-relativistic limit is highly non-trivial. It is interesting to notice that the exotic Newtonian superalgebra can alternatively be recovered by expanding the enhanced Nappi–Witten superalgebra (see “Appendix A”). Although both methods are based on the semigroup expansion method [49], they present subtle differences which could lead to diverse extensions of our results. Indeed, to obtain diverse Newtonian (super)algebras from an enhanced Nappi–Witten (super)algebra, we need to consider diverse semigroups as in [50]. On the other hand, the derivation of various Newtonian (super)algebras by expanding a relativistic (super)algebra requires to consider different original algebras without modifying the semigroup.

Acknowledgements This work was funded by the National Agency for Research and Development ANID (ex-CONICYT) - PAI grant No. 77190078 (P.C.). This work was supported by the Research project Code DIREG_09/2020 (P.C.) of the Universidad Católica de la Santísima Concepción, Chile. P.C. would like to thank to the Dirección de Investigación and Vice-rectoría de Investigación of the Universidad Católica de la Santísima Concepción, Chile, for their constant support. L.R. would like to thank the Department of Applied Science and Technology of the Polytechnic University of Turin, and in particular Laura Andrianopoli and Francesco Raffa, for financial support.

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors’ comment: The results obtained and presented in the manuscript did not require any data.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP³.

A Exotic Newtonian superalgebra by expanding an enhanced Nappi–Witten superalgebra

As was shown in [50], diverse generalizations of the extended Newtonian algebra appear as S -expansions of an enhanced Nappi–Witten algebra. Here, we show that a supersymmetric extension of the enhanced Nappi–Witten algebra allows to obtain the exotic Newtonian superalgebra (2.1), (3.18) and (3.19) considering a particular semigroup. Let

$$\mathfrak{g} = \{\hat{L}, \hat{N}, \hat{B}, \hat{L}_a, \hat{N}_a, \hat{X}_1, \hat{X}_2, \hat{X}_3, \hat{Q}_\alpha^+, \hat{Q}_\alpha^-, \hat{R}_\alpha, \hat{W}_\alpha^-, \hat{W}_\alpha^+\} \quad (\text{A.1})$$

be a supersymmetric extension of the enhanced Nappi–Witten algebra whose generators satisfy (3.22) and (3.23). The presence of the additional generators $\{\hat{X}_1, \hat{X}_2, \hat{X}_3\}$ allows us not only to make contact with the exotic Newtonian superalgebra obtained here but also to establish a non-degenerate invariant tensor. In particular, the enhanced Nappi–Witten superalgebra (3.22)–(3.23) admits the following non-vanishing components of the invariant tensor:

λ_2	$L, N, B, L_a, N_a, X_1, X_2, X_3$	$Q_\alpha^+, Q_\alpha^-, R_\alpha, W_\alpha^+, W_\alpha^-$
λ_1	$\tilde{L}, \tilde{N}, \tilde{B}, \tilde{L}_a, \tilde{N}_a, \tilde{X}_1, \tilde{X}_2, \tilde{X}_3$	$\tilde{Q}_\alpha^+, \tilde{Q}_\alpha^-, \tilde{R}_\alpha, \tilde{W}_\alpha^+, \tilde{W}_\alpha^-$
λ_0	$\hat{L}, \hat{N}, \hat{B}, \hat{L}_a, \hat{N}_a, \hat{X}_1, \hat{X}_2, \hat{X}_3$	$\hat{Q}_\alpha^+, \hat{Q}_\alpha^-, \hat{R}_\alpha, \hat{W}_\alpha^+, \hat{W}_\alpha^-$

(A.5)

$$\begin{aligned} \langle \hat{L} \hat{N} \rangle &= -\gamma_1, \quad \langle \hat{N} \hat{N} \rangle = \langle \hat{N} \hat{B} \rangle = -\gamma_2, \\ \langle \hat{L}_a \hat{L}_b \rangle &= \gamma_1 \delta_{ab}, \quad \langle \hat{L}_a \hat{N}_b \rangle = \gamma_2 \delta_{ab}, \\ \langle \hat{X}_1 \hat{X}_2 \rangle &= \varrho_1, \quad \langle \hat{X}_2 \hat{X}_2 \rangle = \langle \hat{X}_1 \hat{X}_3 \rangle = \varrho_2, \\ \langle \hat{Q}_\alpha^- \hat{Q}_\beta^- \rangle &= \langle \hat{Q}_\alpha^+ \hat{R}_\beta \rangle = 2\gamma_1 C_{\alpha\beta}, \\ \langle \hat{Q}_\alpha^+ \hat{W}_\beta^+ \rangle &= \langle \hat{Q}_\alpha^- \hat{W}_\beta^- \rangle = \langle \hat{R}_\alpha \hat{R}_\beta \rangle = 2\gamma_2 C_{\alpha\beta}, \end{aligned} \quad (\text{A.2})$$

where $\gamma_1, \varrho_1, \varrho_2$ and γ_2 are arbitrary independent constants. Let us note that the components proportional to γ_1 and ρ_1 correspond to those of the super Nappi–Witten ones [47]. Nevertheless, the complete set of components (A.2) is required to ensure the non-degeneracy of the bilinear invariant trace.

Before considering the S -expansion of the enhanced Nappi–Witten superalgebra, it is convenient to consider a subspace decomposition $\mathfrak{g} = V_0 \oplus V_1$ where $V_0 = \{\hat{L}, \hat{N}, \hat{B}, \hat{L}_a, \hat{N}_a, \hat{X}_1, \hat{X}_2, \hat{X}_3\}$ and $V_1 = \{\hat{Q}_\alpha^+, \hat{Q}_\alpha^-, \hat{R}_\alpha, \hat{W}_\alpha^-, \hat{W}_\alpha^+\}$. Such subspace decomposition satisfies (3.9). On the other hand, let $S_L^{(1)} = \{\lambda_0, \lambda_1, \lambda_2\}$ be the relevant semi-

group whose elements satisfy the following multiplication law [73]:

$$\begin{array}{c|ccc} \lambda_2 & \lambda_2 & \lambda_2 & \lambda_2 \\ \lambda_1 & \lambda_2 & \lambda_1 & \lambda_2 \\ \lambda_0 & \lambda_0 & \lambda_2 & \lambda_2 \\ \hline & \lambda_0 & \lambda_1 & \lambda_2 \end{array} \quad (\text{A.3})$$

where $\lambda_2 = 0_S$ is the zero element of the semigroup. Then, let us consider a semigroup decomposition $S_L^{(1)} = S_0 \cup S_1$ with $S_0 = \{\lambda_0, \lambda_1, \lambda_2\}$ and $S_1 = \{\lambda_1, \lambda_2\}$, which is said to be resonant since it satisfies the same algebraic structure as the subspace decomposition (3.9). After considering a resonant subalgebra of the $S_L^{(1)}$ -expansion of the enhanced Nappi–Witten superalgebra and performing a 0_S -reduction, one finds an expanded superalgebra which is spanned by the set of generators

$$\{L, N, B, L_a, N_a, X_1, X_2, X_3, \tilde{L}, \tilde{N}, \tilde{B}, \tilde{L}_a, \tilde{N}_a, \tilde{X}_1, \tilde{X}_2, \tilde{X}_3, Q_\alpha^+, Q_\alpha^-, R_\alpha, W_\alpha^-, W_\alpha^+\}. \quad (\text{A.4})$$

The expanded generators are related to the original ones through the semigroup elements as

Two enhanced Nappi–Witten algebras, one of which is augmented by supersymmetry, are obtained using the multiplication law of the semigroup (A.3) and the (anti-)commutation relations of the original superalgebra (3.22)–(3.23). In particular, the set of generators

$$\{L, N, B, L_a, N_a, X_1, X_2, X_3, Q_\alpha^+, Q_\alpha^-, R_\alpha, W_\alpha^-, W_\alpha^+\} \quad (\text{A.6})$$

satisfies the enhanced Nappi–Witten superalgebra (3.22)–(3.23). On the other hand, the set of generators

$$\{\tilde{L}, \tilde{N}, \tilde{B}, \tilde{L}_a, \tilde{N}_a, \tilde{X}_1, \tilde{X}_2, \tilde{X}_3\} \quad (\text{A.7})$$

satisfies the bosonic enhanced Nappi–Witten algebra (3.25). Then, the exotic Newtonian superalgebra (2.1), (3.18) and (3.19) appears by considering the redefinition of the generators as in (3.20). Let us note that the non-vanishing components of the invariant tensor for the two copies of the enhanced Nappi–Witten algebra, one of which is supersymmetric, can be obtained following the definitions of [49]. Indeed, the expanded superalgebra admits the following non-vanishing components of the invariant tensor:

$$\begin{aligned}
\langle L_a L_b \rangle &= \mu_1 \delta_{ab}, \quad \langle \tilde{L}_a \tilde{L}_b \rangle = \nu_1 \delta_{ab}, \\
\langle LN \rangle &= -\mu_1, \quad \langle \tilde{L} \tilde{N} \rangle = -\nu_1, \\
\langle X_1 X_2 \rangle &= \rho_1, \quad \langle \tilde{X}_1 \tilde{X}_2 \rangle = \tilde{\rho}_1, \\
\langle L_a N_b \rangle &= \mu_2 \delta_{ab}, \quad \langle \tilde{L}_a \tilde{N}_b \rangle = \nu_2 \delta_{ab}, \\
\langle NN \rangle &= \langle NB \rangle = -\mu_2, \quad \langle \tilde{N} \tilde{N} \rangle = \langle \tilde{N} \tilde{B} \rangle = -\nu_2, \\
\langle X_2 X_2 \rangle &= \langle X_1 X_3 \rangle = \rho_2, \quad \langle \tilde{X}_2 \tilde{X}_2 \rangle = \langle \tilde{X}_1 \tilde{X}_3 \rangle = \tilde{\rho}_2, \\
\langle Q_\alpha^- Q_\beta^- \rangle &= \langle Q_\alpha^+ R_\beta \rangle = 2\mu_1 C_{\alpha\beta}, \quad \langle Q_\alpha^+ W_\beta^+ \rangle = \langle Q_\alpha^- W_\beta^- \rangle \\
&= \langle R_\alpha R_\beta \rangle = 2\mu_2 C_{\alpha\beta}, \quad (A.8)
\end{aligned}$$

where the constants are related to the enhanced super Nappi–Witten ones through the semigroup elements as

$$\begin{aligned}
\mu_1 &= \lambda_1 \gamma_1, \quad \mu_2 = \lambda_1 \gamma_2, \quad \rho_1 = \lambda_1 \varrho_1, \quad \rho_2 = \lambda_1 \varrho_2, \\
\nu_1 &= \lambda_0 \gamma_1, \quad \nu_2 = \lambda_0 \gamma_2, \quad \tilde{\rho}_1 = \lambda_0 \varrho_1, \quad \tilde{\rho}_2 = \lambda_0 \varrho_2. \quad (A.9)
\end{aligned}$$

Interestingly, the non-vanishing components of the invariant tensor for the exotic Newtonian superalgebra (4.1) and (4.2) can be alternatively recovered from the invariant tensor (A.8), considering the redefinition of the generators as in (3.20) and setting

$$\begin{aligned}
\alpha_0 &= \mu_1 + \nu_1, \quad \alpha_1 = \frac{1}{\ell} (\mu_1 - \nu_1), \\
\rho_1 &= \alpha_0 + \alpha_1, \quad \tilde{\rho}_1 = -\alpha_1, \\
\beta_0 &= \mu_2 + \nu_2, \quad \beta_1 = \frac{1}{\ell} (\mu_2 - \nu_2), \\
\rho_2 &= \beta_0 + \beta_1, \quad \tilde{\rho}_2 = -\beta_1. \quad (A.10)
\end{aligned}$$

B Supersymmetry gauge transformations

The supersymmetry transformation laws under which the curvature two-forms (4.6) and (4.7) of the exotic Newtonian superalgebra transform in a covariant way read

$$\begin{aligned}
\delta\omega &= -\frac{1}{\ell} \bar{\epsilon}^+ \gamma^0 \psi^+, \\
\delta\omega^a &= -\frac{1}{\ell} \bar{\epsilon}^+ \gamma^a \psi^- - \frac{1}{\ell} \bar{\epsilon}^- \gamma^a \psi^+, \\
\delta\tau &= -\bar{\epsilon}^+ \gamma^0 \psi^+, \\
\delta e^a &= -\bar{\epsilon}^+ \gamma^a \psi^- - \bar{\epsilon}^- \gamma^a \psi^+, \\
\delta s &= -\frac{1}{\ell} \bar{\epsilon}^- \gamma^0 \psi^- - \frac{1}{\ell} \bar{\epsilon}^+ \gamma^0 \rho - \frac{1}{\ell} \bar{\eta} \gamma^0 \psi^+, \\
\delta m &= -\bar{\epsilon}^- \gamma^0 \psi^- - \bar{\epsilon}^+ \gamma^0 \rho - \bar{\rho} \gamma^0 \psi^+, \\
\delta b^a &= -\frac{1}{\ell} \bar{\epsilon}^- \gamma^a \rho - \frac{1}{\ell} \bar{\eta} \gamma^a \psi^- - \frac{1}{\ell} \bar{\epsilon}^+ \gamma^a \phi^- - \frac{1}{\ell} \bar{\zeta}^- \gamma^a \psi^+, \\
\delta t^a &= -\bar{\epsilon}^- \gamma^a \rho - \bar{\eta} \gamma^a \psi^- - \bar{\epsilon}^+ \gamma^a \phi^- - \bar{\zeta}^- \gamma^a \psi^+, \\
\delta z &= -\frac{1}{\ell} \bar{\epsilon}^- \gamma^0 \phi^- - \frac{1}{\ell} \bar{\zeta}^- \gamma^0 \psi^- - \frac{1}{\ell} \bar{\epsilon}^+ \gamma^0 \phi^+
\end{aligned}$$

$$\begin{aligned}
& -\frac{1}{\ell} \bar{\zeta}^+ \gamma^0 \psi^+ - \frac{1}{\ell} \bar{\eta} \gamma^0 \rho, \\
\delta y &= -\bar{\epsilon}^- \gamma^0 \phi^- - \bar{\zeta}^- \gamma^0 \psi^- - \bar{\epsilon}^+ \gamma^0 \phi^+ - \bar{\zeta}^+ \gamma^0 \psi^+ \\
& - \bar{\eta} \gamma^0 \rho, \\
\delta y_1 &= -\frac{1}{\ell} \bar{\epsilon}^+ \gamma^0 \psi^+, \\
\delta y_2 &= +\frac{1}{\ell} \bar{\epsilon}^- \gamma^0 \psi^- - \frac{1}{\ell} \bar{\epsilon}^+ \gamma^0 \rho - \frac{1}{\ell} \bar{\eta} \gamma^0 \psi^+, \\
\delta y_3 &= +\frac{1}{\ell} \bar{\epsilon}^- \gamma^0 \phi^- + \frac{1}{\ell} \bar{\zeta}^- \gamma^0 \psi^- - \frac{1}{\ell} \bar{\epsilon}^+ \gamma^0 \phi^+ \\
& - \frac{1}{\ell} \bar{\zeta}^+ \gamma^0 \psi^+ - \frac{1}{\ell} \bar{\eta} \gamma^0 \rho, \\
\delta u_1 &= -\bar{\epsilon}^+ \gamma^0 \psi^+, \\
\delta u_2 &= +\bar{\epsilon}^- \gamma^0 \psi^- - \bar{\epsilon}^+ \gamma^0 \rho - \bar{\rho} \gamma^0 \psi^+, \\
\delta u_3 &= +\bar{\epsilon}^- \gamma^0 \phi^- + \bar{\zeta}^- \gamma^0 \psi^- - \bar{\epsilon}^+ \gamma^0 \phi^+ \\
& - \bar{\zeta}^+ \gamma^0 \psi^+ - \bar{\eta} \gamma^0 \rho, \quad (B.1)
\end{aligned}$$

along with

$$\begin{aligned}
\delta\psi^+ &= d\epsilon^+ + \frac{1}{2} \omega \gamma_0 \epsilon^+ + \frac{1}{2\ell} \tau \gamma_0 \epsilon^+ - \frac{1}{2} y_1 \gamma_0 \epsilon^+ \\
\delta\psi^- &= d\epsilon^- + \frac{1}{2} \omega \gamma_0 \epsilon^- + \frac{1}{2\ell} \tau \gamma_0 \epsilon^- + \frac{1}{2} \omega^a \gamma_a \epsilon^+ \\
& + \frac{1}{2\ell} e^a \gamma_a \epsilon^+ + \frac{1}{2} y_1 \gamma_0 \epsilon^- \\
\delta\rho &= d\eta + \frac{1}{2} \omega \gamma_0 \eta + \frac{1}{2} s \gamma_0 \epsilon^+ + \frac{1}{2} \omega^a \gamma_a \epsilon^- + \frac{1}{2\ell} \tau \gamma_0 \eta \\
& + \frac{1}{2\ell} m \gamma_0 \epsilon^+ + \frac{1}{2\ell} e^a \gamma_a \epsilon^- \\
& - \frac{1}{2} y_1 \gamma_0 \eta - \frac{1}{2} y_2 \gamma_0 \epsilon^+, \\
\delta\phi^+ &= d\zeta^+ + \frac{1}{2} \omega \gamma_0 \epsilon^+ + \frac{1}{2} s \gamma_0 \eta + \frac{1}{2} \omega^a \gamma_a \zeta^- \\
& + \frac{1}{2} b^a \gamma_a \epsilon^- + \frac{1}{2\ell} \tau \gamma_0 \zeta^+ + \frac{1}{2\ell} m \gamma_0 \eta \\
& + \frac{1}{2\ell} e^a \gamma_a \zeta^- + \frac{1}{2\ell} t^a \gamma_a \epsilon^- - \frac{1}{2} y_1 \gamma_0 \zeta^+ \\
& - \frac{1}{2} y_2 \gamma_0 \eta - \frac{1}{2} y_3 \gamma_0 \epsilon^+, \\
\delta\phi^- &= d\zeta^- + \frac{1}{2} \omega \gamma_0 \zeta^- + \frac{1}{2} s \gamma_0 \epsilon^- + \frac{1}{2} \omega^a \gamma_a \eta \\
& + \frac{1}{2} b^a \gamma_a \epsilon^+ + \frac{1}{2\ell} \tau \gamma_0 \zeta^- + \frac{1}{2\ell} m \gamma_0 \epsilon^- \\
& + \frac{1}{2\ell} e^a \gamma_a \eta + \frac{1}{2\ell} t^a \gamma_a \epsilon^+ + \frac{1}{2} y_1 \gamma_0 \zeta^- + \frac{1}{2} y_2 \gamma_0 \epsilon^- \\
& + \frac{1}{2} z \gamma_0 \epsilon^+ + \frac{1}{2\ell} y \gamma_0 \epsilon^+, \quad (B.2)
\end{aligned}$$

where ϵ^\pm , ζ^\pm and η are the respective fermionic gauge parameters related to the fermionic generators Q^\pm , W^\pm and R . One can notice that in the vanishing cosmological constant limit, $\ell \rightarrow \infty$, we recover the supersymmetry transformations laws under which the curvature two-forms of the

extended Newtonian superalgebra transform in a covariant way [28].

References

1. D. Son, Toward an AdS/cold atoms correspondence: A Geometric realization of the Schrodinger symmetry. *Phys. Rev. D* **78**, 046003 (2008). [arXiv:0804.3972](#)
2. K. Balasubramanian, J. McGreevy, Gravity duals for non-relativistic CFTs. *Phys. Rev. Lett.* **101**, 061601 (2008). [arXiv:0804.4053](#)
3. S. Kachru, X. Liu, M. Mulligan, Gravity duals of Lifshitz-like fixed points. *Phys. Rev. D* **78**, 106005 (2008). [arXiv:0808.1725](#)
4. A. Bagchi, R. Gopakumar, Galilean Conformal Algebras and AdS/CFT. *JHEP* **07**, 037 (2009). [arXiv:0902.1385](#)
5. A. Bagchi, R. Gopakumar, I. Mandal, A. Miwa, GCA in 2d. *JHEP* **08**, 004 (2010). [arXiv:0912.1090](#)
6. M.H. Christensen, J. Hartong, N.A. Obers, B. Rollier, Torsional Newton–Cartan Geometry and Lifshitz Holography. *Phys. Rev. D* **89**, 061901 (2014). [arXiv:1311.4794](#)
7. M.H. Christensen, J. Hartong, N.A. Obers, B. Rollier, Boundary stress-energy tensor and Newton–Cartan geometry in Lifshitz holography. *JHEP* **01**, 057 (2014). [arXiv:1311.6471](#)
8. J. Hartong, E. Kiritsis, N.A. Obers, Lifshitz space-times for Schrödinger holography. *Phys. Lett. B* **746**, 318–324 (2015). [arXiv:1409.1519](#)
9. J. Hartong, E. Kiritsis, N.A. Obers, Schrödinger Invariance from Lifshitz Isometries in Holography and Field Theory. *Phys. Rev. D* **92**, 066003 (2015). [arXiv:1409.1522](#)
10. J. Hartong, E. Kiritsis, N.A. Obers, Field theory on Newton–Cartan backgrounds and symmetries of the Lifshitz vacuum. *JHEP* **08**, 006 (2015). [arXiv:1502.00228](#)
11. M. Taylor, Lifshitz holography. *Class. Quantum Gravity* **33**(3), 033001 (2016). [arXiv:1512.03554](#)
12. C. Hoyos, D.T. Son, Hall viscosity and electromagnetic response. *Phys. Rev. Lett.* **108**, 066805 (2012). [arXiv:1109.2651](#)
13. D.T. Son, Newton–Cartan geometry and the quantum hall effect (2013). [arXiv:1306.0638](#)
14. A.G. Abanov, A. Gromov, Electromagnetic and gravitational responses of two-dimensional noninteracting electrons in a background magnetic field. *Phys. Rev. B* **90**(1), 014435 (2014). [arXiv:1401.3703](#)
15. M. Geracie, K. Prabhu, M.M. Roberts, Curved non-relativistic spacetimes, Newtonian gravitation and massive matter. *J. Math. Phys.* **56**(10), 103505 (2015). [arXiv:1503.02682](#)
16. A. Gromov, K. Jensen, A.G. Abanov, Boundary effective action for quantum Hall states. *Phys. Rev. Lett.* **116**(12), 126802 (2016). [arXiv:1506.07171](#)
17. E. Cartan, Sur les variétés à connexion affine et la théorie de la relativité généralisée (première partie). *Ann. Ecole Norm. Sup.* **40**, 325–412 (1923)
18. E. Cartan, Sur les variétés à connexion affine et la théorie de la relativité généralisée (première partie) (Suite). *Ann. Ecole Norm. Sup.* **41**, 1–25 (1924)
19. D. Hansen, J. Hartong, N.A. Obers, Action principle for Newtonian gravity. *Phys. Rev. Lett.* **122**(6), 061106 (2019). [arXiv:1807.04765](#)
20. J.M. Lévy-Leblond, *Galilei group and Galilean invariance. Group Theory and its Applications*, ed. by E.M. Loebl, vol. II (Academic, New York, 1971), p. 221
21. D. Grigore, The projective unitary irreducible representations of the Galilei group in (1+2)-dimensions. *J. Math. Phys.* **37**, 460–473 (1996). [arXiv:hep-th/9312048](#)
22. S. Bose, The Galilean group in (2+1) space-times and its central extension. *Commun. Math. Phys.* **169**, 385–396 (1995)
23. C. Duval, P. Horvathy, The ‘Peierls substitution’ and the exotic Galilei group. *Phys. Lett. B* **479**, 284–290 (2000). [arXiv:hep-th/0002233](#)
24. R. Jackiw, V. Nair, Anyon spin and the exotic central extension of the planar Galilei group. *Phys. Lett. B* **480**, 237–238 (2000). [arXiv:hep-th/0003130](#)
25. G. Papageorgiou, B.J. Schroers, A Chern–Simons approach to Galilean quantum gravity in 2+1 dimensions. *JHEP* **11**, 009 (2009). [arXiv:0907.2880](#)
26. R. Grassie, Generalised Bargmann superalgebras (2020). [arXiv:2010.01894](#)
27. E.A. Bergshoeff, J. Rosseel, Three-dimensional extended Bargmann supergravity. *Phys. Rev. Lett.* **116**(25), 251601 (2016). [arXiv:1604.08042](#)
28. N. Ozdemir, M. Ozkan, O. Tunca, U. Zorba, Three-dimensional extended Newtonian (Super)gravity. *JHEP* **05**, 130 (2019). [arXiv:1903.09377](#)
29. A. Edery, Non-singular vortices with positive mass in 2+1 dimensional Einstein gravity with AdS₃ and Minkowski background. *JHEP* **01**, 166 (2021). [arXiv:2004.09295](#)
30. H. Bacry, J. Levy-Leblond, Possible kinematics. *J. Math. Phys.* **9**, 1605–1614 (1968)
31. R. Aldrovandi, A. Barbosa, L. Crispino, J. Pereira, Non-relativistic spacetimes with cosmological constant. *Class. Quantum Gravity* **16**, 495–506 (1999). [arXiv:gr-qc/9801100](#)
32. G. Gibbons, C. Patricot, Newton–Hooke space-times, Hpp waves and the cosmological constant. *Class. Quantum Gravity* **20**, 5225 (2003). [arXiv:hep-th/0308200](#)
33. J. Bruges, J. Gomis, K. Kamimura, Newton–Hooke algebras, non-relativistic branes and generalized pp-wave metrics. *Phys. Rev. D* **73**, 085011 (2006). [arXiv:hep-th/0603023](#)
34. P.D. Alvarez, J. Gomis, K. Kamimura, M.S. Plyushchay, (2+1)D Exotic Newton–Hooke symmetry. Duality and projective phase. *Ann. Phys.* **322**, 1556–1586 (2007). [arXiv:hep-th/0702014](#)
35. C. Duval, P. Horvathy, Conformal Galilei groups, Veronese curves, and Newton–Hooke spacetimes. *J. Phys. A* **44**, 335203 (2011). [arXiv:1104.1502](#)
36. C. Duval, G. Gibbons, P. Horvathy, Conformal and projective symmetries in Newtonian cosmology. *J. Geom. Phys.* **112**, 197–209 (2017). [arXiv:1605.00231](#)
37. G. Papageorgiou, B.J. Schroers, Galilean quantum gravity with cosmological constant and the extended q -Heisenberg algebra. *JHEP* **11**, 020 (2010). [arXiv:1008.0279](#)
38. J. Hartong, Y. Lei, N.A. Obers, Nonrelativistic Chern–Simons theories and three-dimensional Hořava–Lifshitz gravity. *Phys. Rev. D* **94**(6), 065027 (2016). [arXiv:1604.08054](#)
39. P. Concha, L. Ravera, E. Rodríguez, Three-dimensional exotic Newtonian gravity with cosmological constant. *Phys. Lett. B* **804**, 135392 (2020). [arXiv:1912.02836](#)
40. E. Bergshoeff, J. Gomis, P. Salgado-Rebolledo, Non-relativistic limits and three-dimensional coadjoint Poincaré gravity. *Proc. R. Soc. Lond. A* **476**(2240), 20200106 (2020). [arXiv:2001.11790](#)
41. R. Andringa, E.A. Bergshoeff, J. Rosseel, E. Sezgin, 3D Newton–Cartan supergravity. *Class. Quantum Gravity* **30**, 205005 (2013). [arXiv:1305.6737](#)
42. E. Bergshoeff, J. Rosseel, T. Zojer, Newton–Cartan supergravity with torsion and Schrödinger supergravity. *JHEP* **11**, 180 (2015). [arXiv:1509.04527](#)
43. J.A. de Azcárraga, D. Gútiérrez, J.M. Izquierdo, Extended $D = 3$ Bargmann supergravity from a Lie algebra expansion (2019). [arXiv:1904.12786](#)
44. N. Ozdemir, M. Ozkan, U. Zorba, Three-dimensional extended Lifshitz, Schrödinger and Newton–Hooke supergravity. *JHEP* **11**, 052 (2019). [arXiv:1909.10745](#)

45. P. Concha, L. Ravera, E. Rodríguez, Three-dimensional Maxwellian extended Bargmann supergravity. *JHEP* **04**, 051 (2020). [arXiv:1912.09477](#)
46. P. Concha, L. Ravera, E. Rodríguez, Three-dimensional non-relativistic extended supergravity with cosmological constant. *Eur. Phys. J. C* **80**(12), 1105 (2020). [arXiv:2008.08655](#)
47. P. Concha, M. Ipinza, L. Ravera, E. Rodríguez, Non-relativistic three-dimensional supergravity theories and semigroup expansion method. *JHEP* **02**, 094 (2021). [arXiv:2010.01216](#)
48. J.A. de Azcarraga, J.M. Izquierdo, M. Picon, O. Varela, Generating Lie and gauge free differential (super)algebras by expanding Maurer–Cartan forms and Chern–Simons supergravity. *Nucl. Phys. B* **662**, 185–219 (2003). [arXiv:hep-th/0212347](#)
49. F. Izaurieta, E. Rodríguez, P. Salgado, Expanding Lie (super)algebras through Abelian semigroups. *J. Math. Phys.* **47**, 123512 (2006). [arXiv:hep-th/0606215](#)
50. P. Concha, L. Ravera, E. Rodríguez, G. Rubio, Three-dimensional Maxwellian extended Newtonian gravity and flat limit. *JHEP* **10**, 181 (2020). [arXiv:2006.13128](#)
51. J. Gomis, A. Kleinschmidt, J. Palmkvist, P. Salgado-Rebolledo, Newton–Hooke/Carrollian expansions of (A)dS and Chern–Simons gravity. *JHEP* **02**, 009 (2020). [arXiv:1912.07564](#)
52. E. Witten, (2+1)-Dimensional gravity as an exactly soluble system. *Nucl. Phys. B* **311**, 46 (1988)
53. M. Hatsuda, M. Sakaguchi, Wess–Zumino term for the AdS superstring and generalized Inonu–Wigner contraction. *Prog. Theor. Phys.* **109**, 853–867 (2003). [arXiv:hep-th/0106114](#)
54. R. Caroca, I. Kondrashuk, N. Merino, F. Nadal, Bianchi spaces and their three-dimensional isometries as S-expansions of two-dimensional isometries. *J. Phys. A* **46**, 225201 (2013). [arXiv:1104.3541](#)
55. L. Andrianopoli, N. Merino, F. Nadal, M. Trigiante, General properties of the expansion methods of Lie algebras. *J. Phys. A* **46**, 365204 (2013). [arXiv:1308.4832](#)
56. M. Artebani, R. Caroca, M. Ipinza, D. Peñafiel, P. Salgado, Geometrical aspects of the Lie algebra S-expansion procedure. *J. Math. Phys.* **57**(2), 023516 (2016). [arXiv:1602.04525](#)
57. M. Ipinza, F. Lingua, D. Peñafiel, L. Ravera, An Analytic Method for S-Expansion involving Resonance and Reduction. *Fortsch. Phys.* **64**(11–12), 854–880 (2016). [arXiv:1609.05042](#)
58. D. Peñafiel, L. Ravera, Infinite S-expansion with ideal subtraction and some applications. *J. Math. Phys.* **58**(8), 081701 (2017). [arXiv:1611.05812](#)
59. C. Inostroza, I. Kondrashuk, N. Merino, F. Nadal, On the algorithm to find S-related Lie algebras. *J. Phys. Conf. Ser.* **1085**(5), 052011 (2018). [arXiv:1802.05765](#)
60. F. Izaurieta, E. Rodríguez, P. Salgado, Eleven-dimensional gauge theory for the M algebra as an Abelian semigroup expansion of osp(32|1). *Eur. Phys. J. C* **54**, 675–684 (2008). [arXiv:hep-th/0606225](#)
61. J.D. Edelstein, M. Hassaine, R. Troncoso, J. Zanelli, Lie-algebra expansions, Chern–Simons theories and the Einstein–Hilbert Lagrangian. *Phys. Lett. B* **640**, 278–284 (2006). [arXiv:hep-th/0605174](#)
62. J. de Azcarraga, J. Izquierdo, M. Picon, O. Varela, Expansions of algebras and superalgebras and some applications. *Int. J. Theor. Phys.* **46**, 2738–2752 (2007). [arXiv:hep-th/0703017](#)
63. F. Izaurieta, E. Rodríguez, P. Minning, P. Salgado, A. Perez, Standard general relativity from Chern–Simons gravity. *Phys. Lett. B* **678**, 213–217 (2009). [arXiv:0905.2187](#)
64. P. Concha, D. Peñafiel, E. Rodríguez, P. Salgado, Even-dimensional general relativity from Born–Infeld gravity. *Phys. Lett. B* **725**, 419–424 (2013). [arXiv:1309.0062](#)
65. P. Salgado, R.J. Szabo, O. Valdivia, Topological gravity and transgression holography. *Phys. Rev. D* **89**(8), 084077 (2014). [arXiv:1401.3653](#)
66. P. Concha, D. Penafiel, E. Rodríguez, P. Salgado, Chern–Simons and Born–Infeld gravity theories and Maxwell algebras type. *Eur. Phys. J. C* **74**, 2741 (2014). [arXiv:1402.0023](#)
67. P. Concha, E. Rodríguez, N = 1 Supergravity and Maxwell superalgebras. *JHEP* **09**, 090 (2014). [arXiv:1407.4635](#)
68. R. Durka, Resonant algebras and gravity. *J. Phys. A* **50**(14), 145202 (2017). [arXiv:1605.00059](#)
69. D.M. Peñafiel, L. Ravera, On the hidden Maxwell Superalgebra underlying D=4 Supergravity. *Fortsch. Phys.* **65**(9), 1700005 (2017). [arXiv:1701.04234](#)
70. R. Caroca, P. Concha, O. Fierro, E. Rodríguez, P. Salgado-Rebolledo, Generalized Chern–Simons higher-spin gravity theories in three dimensions. *Nucl. Phys. B* **934**, 240–264 (2018). [arXiv:1712.09975](#)
71. R. Caroca, P. Concha, E. Rodríguez, P. Salgado-Rebolledo, Generalizing the \mathfrak{bms}_3 and 2D-conformal algebras by expanding the Virasoro algebra. *Eur. Phys. J. C* **78**(3), 262 (2018). [arXiv:1707.07209](#)
72. D.M. Peñafiel, L. Ravera, Generalized cosmological term in $D = 4$ supergravity from a new AdS–Lorentz superalgebra. *Eur. Phys. J. C* **78**(11), 945 (2018). [arXiv:1807.07673](#)
73. R. Caroca, P. Concha, O. Fierro, E. Rodríguez, On the supersymmetric extension of asymptotic symmetries in three spacetime dimensions. *Eur. Phys. J. C* **80**(1), 29 (2020). [arXiv:1908.09150](#)
74. E. Bergshoeff, J.M. Izquierdo, T. Ortín, L. Romano, Lie algebra expansions and actions for non-relativistic gravity. *JHEP* **08**, 048 (2019). [arXiv:1904.08304](#)
75. L. Romano, Non-relativistic four dimensional p-Brane supersymmetric theories and Lie algebra expansion (2019). [arXiv:1906.08220](#)
76. A. Fontanella, L. Romano, Lie Algebra expansion and integrability in superstring sigma-models. *JHEP* **20**, 083 (2020). [arXiv:2005.01736](#)
77. P. Concha, E. Rodríguez, Non-relativistic gravity theory based on an enlargement of the extended Bargmann algebra. *JHEP* **07**, 085 (2019). [arXiv:1906.00086](#)
78. D.M. Peñafiel, P. Salgado-Rebolledo, Non-relativistic symmetries in three space-time dimensions and the Nappi–Witten algebra. *Phys. Lett. B* **798**, 135005 (2019). [arXiv:1906.02161](#)
79. O. Kasikci, N. Ozdemir, M. Ozkan, U. Zorba, Three-dimensional higher-order Schrödinger algebras and Lie algebra expansions. *JHEP* **04**, 067 (2020). [arXiv:2002.03558](#)
80. P. Concha, M. Ipinza, E. Rodríguez, Generalized Maxwellian exotic Bargmann gravity theory in three spacetime dimensions. *Phys. Lett. B* **807**, 135593 (2020). [arXiv:2004.01203](#)
81. P.S. Howe, J. Izquierdo, G. Papadopoulos, P. Townsend, New supergravities with central charges and Killing spinors in (2+1)-dimensions. *Nucl. Phys. B* **467**, 183–214 (1996). [arXiv:hep-th/9505032](#)
82. L. Andrianopoli, R. D’Auria, L. Ravera, Hidden gauge structure of supersymmetric free differential algebras. *JHEP* **08**, 095 (2016). [arXiv:1606.07328](#)
83. L. Andrianopoli, R. D’Auria, L. Ravera, More on the hidden symmetries of 11D supergravity. *Phys. Lett. B* **772**, 578–585 (2017). [arXiv:1705.06251](#)
84. L. Ravera, Hidden role of Maxwell superalgebras in the free differential algebras of $D = 4$ and $D = 11$ supergravity. *Eur. Phys. J. C* **78**(3), 211 (2018). [arXiv:1801.08860](#)
85. A. Banaudi, L. Ravera, Generalized AdS–Lorentz deformed supergravity on a manifold with boundary. *Eur. Phys. J. Plus* **133**(12), 514 (2018). [arXiv:1803.08738](#)
86. A. Achúcarro, P. Townsend, Extended supergravities in $d = (2+1)$ as Chern–Simons theories. *Phys. Lett. B* **229**, 383–387 (1989)
87. A. Giacomini, R. Troncoso, S. Willison, Three-dimensional supergravity reloaded. *Class. Quantum Gravity* **24**, 2845–2860 (2007). [arXiv:hep-th/0610077](#)

88. H.R. Afshar, E.A. Bergshoeff, A. Mehra, P. Parekh, B. Rollier, A Schrödinger approach to Newton-Cartan and Hořava–Lifshitz gravities. *JHEP* **04**, 145 (2016). [arXiv:1512.06277](#)
89. G. Festuccia, N. Seiberg, Rigid supersymmetric theories in curved superspace. *JHEP* **06**, 114 (2011). [arXiv:1105.0689](#)
90. V. Pestun, Localization of gauge theory on a four-sphere and supersymmetric Wilson loops. *Commun. Math. Phys.* **313**, 71–129 (2012). <http://arxiv.org/abs/0712.2824>
91. M. Mariño, Lectures on non-perturbative effects in large N gauge theories, matrix models and strings. *Fortsch. Phys.* **62**, 455–540 (2014). [arXiv:1206.6272](#)