

Three-dimensional Maxwellian Carroll gravity theory and the cosmological constant

Original

Three-dimensional Maxwellian Carroll gravity theory and the cosmological constant / Concha, P.; Penafiel, D.; Ravera, L.; Rodriguez, E.. - In: PHYSICS LETTERS. SECTION B. - ISSN 0370-2693. - STAMPA. - 823:136735(2021), pp. 1-10. [10.1016/j.physletb.2021.136735]

Availability:

This version is available at: 11583/2949104 since: 2022-01-12T12:06:30Z

Publisher:

Elsevier B.V.

Published

DOI:10.1016/j.physletb.2021.136735

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Three-dimensional Maxwellian Carroll gravity theory and the cosmological constant

Patrick Concha^{a,*}, Diego Peñafiel^b, Lucrezia Ravera^{c,d}, Evelyn Rodríguez^e

^a Departamento de Matemática y Física Aplicadas, Universidad Católica de la Santísima Concepción, Alonso de Ribera 2850, Concepción, Chile

^b Facultad de Ciencias, Universidad Arturo Prat, Iquique, Chile

^c DISAT, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy

^d INFN, Sezione di Torino, Via P. Giuria 1, 10125 Torino, Italy

^e Instituto de Matemática (INSTMAT), Universidad de Talca, Casilla 747, Talca 3460000, Chile

ARTICLE INFO

Article history:

Received 21 July 2021

Received in revised form 15 October 2021

Accepted 18 October 2021

Available online 20 October 2021

Editor: N. Lambert

ABSTRACT

In this work, we present the three-dimensional Maxwell Carroll gravity by considering the ultra-relativistic limit of the Maxwell Chern-Simons gravity theory defined in three spacetime dimensions. We show that an extension of the Maxwellian Carroll symmetry is necessary in order for the invariant tensor of the ultra-relativistic Maxwellian algebra to be non-degenerate. Consequently, we discuss the origin of the aforementioned algebra and theory as a flat limit. We show that the theoretical setup with cosmological constant yielding the extended Maxwellian Carroll Chern-Simons gravity in the vanishing cosmological constant limit is based on a new enlarged extended version of the Carroll symmetry. Indeed, the latter exhibits a non-degenerate invariant tensor allowing the proper construction of a Chern-Simons gravity theory which reproduces the extended Maxwellian Carroll gravity in the flat limit.

© 2021 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Symmetries in Physics play a fundamental role in the analysis and understanding of theories. In this perspective, ultra-relativistic (UR) symmetries, also known as Carroll symmetries, have attracted some interest over recent years. The Carroll symmetries arise in the UR limit, $c \rightarrow 0$, being c the speed of light [1,2]. The Carroll group introduced by Lévy-Leblond emerged as the UR contraction of the Poincaré group, dual to the non-relativistic (NR) contraction (i.e., $c \rightarrow \infty$) leading to the Galilean group.

Models with Carroll symmetries appear, for instance, in the context of high-energy Physics in the study of tachyon condensation [3], warped conformal field theories [4], and tensionless strings [5–9]. Moreover, further works on Carroll algebras in the context of electrodynamics and brane dynamics can be found for instance in [10–12]. On the other hand, the exploration of the Carroll limit corresponding to M2- as well as M3-branes propagating over $D = 11$ supergravity backgrounds in M-theory has been considered in [13]. In [14], flat and Anti-de Sitter (AdS) Carroll spaces were investigated, in the bosonic and in the supersymmetric case, studying the symmetries of a particle moving in such spaces. Models of Carrollian gravity have been developed in [15–18]. In particular, in [16], the respective NR and UR limits of the spin-3 Chern-Simons (CS) gravity in three spacetime dimensions were presented. Subsequently, the AdS Carroll CS gravity theory and its supersymmetric extension were discussed in [18] and in [19,20], respectively. More recently, the Carrollian version of the Jackiw-Teitelboim gravity was explored in [21,22].

On the other hand, there has been a growing interest, at the relativistic level, in exploring symmetries in three spacetime dimensions beyond the Poincaré and AdS ones. It is well assumed that a gravitational theory in three spacetime dimensions can be described by a CS action as a gauge theory, providing a useful setup to approach higher-dimensional models. A non-trivial extension of the Poincaré algebra is given by the Maxwell algebra which, in any dimension D , is characterized by the commutator

$$[\tilde{P}_A, \tilde{P}_B] = \tilde{Z}_{AB}. \quad (1.1)$$

* Corresponding author.

E-mail addresses: patrick.concha@ucsc.cl (P. Concha), dimolina@unap.cl (D. Peñafiel), lucrezia.ravera@polito.it (L. Ravera), evelyn.rodriguez@utalca.cl (E. Rodríguez).

The Maxwell algebra has been first introduced in four spacetime dimensions to describe a constant Minkowski spacetime in the presence of an electromagnetic background [23–26]. In arbitrary spacetime dimensions, the Maxwell algebra and generalizations have been useful to relate General Relativity with CS and Born-Infeld (BI) gravity theories [27–30]. In three spacetime dimensions, the Maxwell symmetries and extensions to supersymmetry and higher-spin, have been explored using the CS formalism with diverse physical implications [31–41].¹ Interestingly, the inclusion of a cosmological constant to the Maxwell CS gravity theory can be done considering the $so(2, 2) \times so(2, 1)$ algebra, also denoted as AdS-Lorentz (AdS- \mathcal{L}) algebra [53–55]. Applications of the AdS- \mathcal{L} algebra and generalizations can be found in [56,57], where the higher-dimensional pure Lovelock gravity is recovered as a particular limit of CS gravity theories based on the AdS- \mathcal{L} symmetries.² Besides, in [61–63], deformations of the Maxwell algebra and their dynamics through non-linear realizations were studied.

At the NR level, the study of a NR version of the three-dimensional CS Maxwell gravity theory was first considered in [64]. As was discussed in [64], a finite and non-degenerate NR CS action required to consider the NR limit of a particular U(1) enlargement of the Maxwell symmetry leading to a Maxwellian extended Bargmann (MEB) algebra. Interestingly, such symmetry can alternatively be recovered as a vanishing cosmological limit of an enlarged extended Bargmann (EEB) symmetry [65] and as a semigroup expansion of the Nappi-Witten algebra [65,66]. The generalization of the MEB symmetry and its supersymmetric extension have been subsequently approached in [67,68] and [69,70], respectively.

On the other hand, although the UR version of the Maxwell symmetry has been partially studied in [71], the corresponding three-dimensional CS action is unknown. Motivated by the prominent role of Maxwellian algebras in the context of (super)gravity and by the fact that, to our knowledge, the Maxwellian UR gravity model remains unexplored, we present a Maxwellian generalization of the three-dimensional CS Carroll gravity theory. In particular, we would like to study the effect of including, in the UR regime, a covariantly constant electromagnetic field in the three-dimensional CS gravity, without introducing a cosmological constant. In this work, we first show that the Maxwell Carroll symmetry can be obtained as an UR limit of the Maxwell algebra. However, similarly to the NR case, an extended version of the Maxwell Carroll is required. Such extension is necessary in order for the invariant tensor to be non-degenerate and thus allow the proper construction of a CS action. Subsequently, we include a cosmological constant to the theory by considering an enlarged Carroll symmetry which can be seen as the corresponding UR counterpart of the AdS- \mathcal{L} symmetry. We show that the model with cosmological constant is based on a novel enlarged extended version of the Carroll symmetry exhibiting a non-degenerate invariant tensor and therefore allowing the proper construction of a CS gravity theory. Let us emphasize that the UR CS gravity theories constructed here are new and, as their relativistic counterpart, they are related through a vanishing cosmological constant limit $\ell \rightarrow \infty$.

The paper is organized as follows: In Section 2 we briefly review the relativistic Maxwell and AdS- \mathcal{L} CS gravity theories in three spacetime dimensions. Sections 3, 4 and 5 contain our main results. In Section 3 we derive the Maxwellian Carroll algebra by considering the UR limit of the relativistic Maxwell algebra and we write an UR CS gravity action based on the Maxwellian Carroll algebra. In Section 4 we consider an extended version of the Maxwellian Carroll symmetry which exhibits a non-degenerate invariant bilinear form and therefore allows the proper construction of a CS Maxwellian extended Carroll gravity theory. In Section 5 we show that a cosmological constant can be incorporated to the extended Maxwellian Carroll CS gravity model by considering an enlarged Carroll algebra. We introduce additional bosonic generators to the enlarged Carroll algebra, obtaining an enlarged extended Carroll algebra, in order to have a non-degenerate invariant tensor allowing the proper construction of an UR CS gravity action. In the flat limit, the latter precisely boils down to the Maxwellian extended Carroll CS gravity theory. Section 6 is devoted to some concluding remarks and possible future developments.

2. Three-dimensional Maxwell and AdS-Lorentz Chern-Simons gravity

In this section, we will briefly review the three-dimensional AdS- \mathcal{L} CS gravity theory [53–55], along with the CS gravity theory obtained after considering its flat limit, known as the Maxwell gravity [33–36].

The AdS- \mathcal{L} algebra can be seen as a semi-simple enlargement of the Poincaré symmetry [72], allowing to incorporate a cosmological constant to the Maxwell gravity. Such enlarged algebra is spanned by the set of generators $\{\tilde{J}_A, \tilde{P}_A, \tilde{Z}_A\}$, which satisfy the non-vanishing commutation relations

$$\begin{aligned} [\tilde{J}_A, \tilde{J}_B] &= \epsilon_{ABC} \tilde{J}^C, & [\tilde{J}_A, \tilde{P}_B] &= \epsilon_{ABC} \tilde{P}^C, \\ [\tilde{J}_A, \tilde{Z}_B] &= \epsilon_{ABC} \tilde{Z}^C, & [\tilde{P}_A, \tilde{P}_B] &= \epsilon_{ABC} \tilde{Z}^C, \\ [\tilde{Z}_A, \tilde{Z}_B] &= \frac{1}{\ell^2} \epsilon_{ABC} \tilde{Z}^C, & [\tilde{P}_A, \tilde{Z}_B] &= \frac{1}{\ell^2} \epsilon_{ABC} \tilde{P}^C, \end{aligned} \quad (2.1)$$

where ℓ is a length parameter related to the (negative) cosmological constant through $\Lambda \propto -\frac{1}{\ell^2}$, the Lorentz indices $A, B, C = 0, 1, 2$ are lowered and raised with the Minkowski metric $\eta_{AB} = (-1, 1, 1)$ and ϵ_{ABC} is the three-dimensional Levi Civita tensor which satisfies $\epsilon_{012} = -\epsilon^{012} = 1$. Note that the flat limit $\ell \rightarrow \infty$ applied to the AdS- \mathcal{L} algebra (2.1) reproduces the Maxwell symmetry.

The AdS- \mathcal{L} algebra admits the following non-vanishing components of an invariant tensor of rank two:

$$\begin{aligned} \langle \tilde{J}_A \tilde{J}_B \rangle &= \tilde{\alpha}_0 \eta_{AB}, & \langle \tilde{J}_A \tilde{P}_B \rangle &= \tilde{\alpha}_1 \eta_{AB}, \\ \langle \tilde{J}_A \tilde{Z}_B \rangle &= \tilde{\alpha}_2 \eta_{AB}, & \langle \tilde{P}_A \tilde{P}_B \rangle &= \tilde{\alpha}_2 \eta_{AB}, \\ \langle \tilde{Z}_A \tilde{P}_B \rangle &= \frac{\tilde{\alpha}_1}{\ell^2} \eta_{AB}, & \langle \tilde{Z}_A \tilde{Z}_B \rangle &= \frac{\tilde{\alpha}_2}{\ell^2} \eta_{AB}, \end{aligned} \quad (2.2)$$

¹ For other relevant applications of the Maxwell algebras, together with supersymmetric extensions of the latter, we refer the interested reader to, e.g., [42–52].

² The interested reader can also find applications of supersymmetric extensions of the AdS- \mathcal{L} algebra in four spacetime dimensions in the context of supergravity theories formulated à la MacDowell-Mansouri in [58–60].

where $\tilde{\alpha}_0$, $\tilde{\alpha}_1$ and $\tilde{\alpha}_2$ are arbitrary constants. One can see that the invariant tensor given by (2.2) reduces to the invariant tensor for the Maxwell algebra when $\ell \rightarrow \infty$. Let us notice that both the Maxwellian and AdS- \mathcal{L} invariant tensors are non-degenerate, assuring the vanishing of the curvature two-forms when one considers the field equations of the associated CS gravity theory.

The gauge connection one-form A for the AdS- \mathcal{L} algebra reads

$$A = \tilde{W}^A \tilde{J}_A + \tilde{E}^A \tilde{P}_A + \tilde{K}^A \tilde{Z}_A, \tag{2.3}$$

where \tilde{W}^A denotes the spin-connection, \tilde{E}^A is the dreibein and \tilde{K}^A is the gauge field associated with the Maxwellian \tilde{Z}_A generator. The corresponding curvature two-form is given by

$$F = \tilde{R}^A(\tilde{W}) \tilde{J}_A + \tilde{R}^A(\tilde{E}) \tilde{P}_A + \tilde{R}^A(\tilde{K}) \tilde{Z}_A, \tag{2.4}$$

with

$$\begin{aligned} \tilde{R}^A(\tilde{W}) &:= d\tilde{W}^A + \frac{1}{2}\epsilon^{ABC}\tilde{W}_B\tilde{W}_C, \\ \tilde{R}^A(\tilde{E}) &:= d\tilde{E}^A + \epsilon^{ABC}\tilde{W}_B\tilde{E}_C + \frac{1}{\ell^2}\epsilon^{ABC}\tilde{K}_B\tilde{E}_C, \\ \tilde{R}^A(\tilde{K}) &:= d\tilde{K}^A + \epsilon^{ABC}\tilde{W}_B\tilde{K}_C + \frac{1}{2\ell^2}\epsilon^{ABC}\tilde{K}_B\tilde{K}_C + \frac{1}{2}\epsilon^{ABC}\tilde{E}_B\tilde{E}_C. \end{aligned} \tag{2.5}$$

Considering the gauge connection one-form (2.3) and the non-vanishing components of the invariant tensor (2.2) in the three-dimensional CS expression,

$$I = \frac{k}{4\pi} \int \left\langle AdA + \frac{2}{3}A^3 \right\rangle, \tag{2.6}$$

with k being the CS level of the theory related to the gravitational constant G , we find the following CS gravity action for the AdS- \mathcal{L} algebra [55]:

$$\begin{aligned} I_{\text{AdS-}\mathcal{L}} &= \frac{k}{4\pi} \int \left[\tilde{\alpha}_0 \left(\tilde{W}^A d\tilde{W}_A + \frac{1}{3}\epsilon^{ABC}\tilde{W}_A\tilde{W}_B\tilde{W}_C \right) \right. \\ &+ \tilde{\alpha}_1 \left(2\tilde{E}_A\tilde{R}^A(\tilde{W}) + \frac{2}{\ell^2}\tilde{E}_A\tilde{F}^A(\tilde{K}) + \frac{1}{3\ell^2}\epsilon^{ABC}\tilde{E}_A\tilde{E}_B\tilde{E}_C \right) \\ &\left. + \tilde{\alpha}_2 \left(\tilde{T}^A\tilde{E}_A + \frac{1}{\ell^2}\epsilon^{ABC}\tilde{E}_A\tilde{K}_B\tilde{E}_C + 2\tilde{K}_A\tilde{R}^A(\tilde{W}) + \frac{1}{\ell^2}\tilde{K}_A D_{\tilde{W}}\tilde{K}^A + \frac{1}{3\ell^4}\epsilon^{ABC}\tilde{K}_A\tilde{K}_B\tilde{K}_C \right) \right], \end{aligned} \tag{2.7}$$

where we have defined the curvature $\tilde{F}^A(\tilde{K}) := D_{\tilde{W}}\tilde{K}^A + \frac{1}{2\ell^2}\epsilon^{ABC}\tilde{K}_B\tilde{K}_C$, being $D_{\tilde{W}}$ the Lorentz covariant derivative, $D_{\tilde{W}}\Theta^A = d\Theta^A + \epsilon^{ABC}\tilde{W}_B\Theta_C$. Furthermore, we have that $\tilde{T}^A := d\tilde{E}^A + \epsilon^{ABC}\tilde{W}_B\tilde{E}_C$ is the usual torsion two-form.

From the CS action we can see that it is split in three independent different sectors. The piece along $\tilde{\alpha}_0$ corresponds to the gravitational (or exotic) Lagrangian [73], while the Einstein-Hilbert term, a cosmological constant term and a contribution involving the gauge field \tilde{K}^A appear along the $\tilde{\alpha}_1$ constant. Furthermore, the term related to the $\tilde{\alpha}_2$ constant contains a torsional term and also terms involving the additional field \tilde{K}^A . Naturally, the flat limit $\ell \rightarrow \infty$ of the relativistic CS action (2.7) leads to the Maxwell CS gravity action [33–36]. As it was shown in [36,55], the presence of the additional gauge field \tilde{K}^A modifies not only the asymptotic sector but also the vacuum of the Maxwell and AdS- \mathcal{L} CS gravity theories, respectively.

Non-relativistic versions of the Maxwell and the AdS- \mathcal{L} algebras have been recently considered in [64–69] and have required U(1) enlargements in order to have finite and non-degenerate NR CS gravity actions after the non-relativistic limit. The corresponding NR algebras are, as their relativistic counterparts, related through the flat limit $\ell \rightarrow \infty$.

In what follows, we will study diverse UR versions of the previously introduced three-dimensional CS gravity theories. First, we will show that a UR limit can be applied to obtain the Carrollian version of the Maxwell and AdS- \mathcal{L} gravity theories. To this aim, we decompose the A -index as follows:

$$A \rightarrow (0, a), \quad a = 1, 2. \tag{2.8}$$

Then, we will apply a particular redefinition to the aforementioned relativistic Maxwell and AdS- \mathcal{L} algebras and we will take the UR limit in order to get the Carrollian version of these algebras. We shall call them as Maxwellian and enlarged Carroll algebras, respectively. We will also consider the contraction at the level of the invariant tensors in order to construct the corresponding UR CS gravity actions. Nevertheless, although the UR symmetries are well-defined in the limit, the respective UR invariant bilinear forms obtained by performing the Carrollian contraction starting from the AdS- \mathcal{L} invariant tensor are degenerate. As we shall see, such drawback can be overcome by considering extensions of the Maxwellian and enlarged Carroll symmetries assuring the proper development of UR CS gravity theories.

3. Maxwellian Carroll Chern-Simons gravity theory

Let us consider now the Carrollian version of the Maxwell algebra. Let us recall that the relativistic Maxwell symmetry can be obtained by taking the vanishing cosmological constant limit $\ell \rightarrow \infty$ of (2.1).

The Maxwellian Carroll algebra is obtained by performing the indices decomposition (2.8) in the Maxwell algebra, and subsequently performing the Carroll contraction, in view of which we introduce the σ parameter by the redefinition

$$\tilde{J}_a \rightarrow \sigma K_a, \quad \tilde{P}_0 \rightarrow \sigma H, \quad \tilde{Z}_a \rightarrow \sigma Z_a, \quad \tilde{J}_0 \rightarrow J, \quad \tilde{P}_a \rightarrow P_a, \quad \tilde{Z}_0 \rightarrow Z. \quad (3.1)$$

Then, considering the limit $\sigma \rightarrow \infty$,³ we obtain the algebra generated by the set of generators $\{J, K_a, H, P_a, Z, Z_a\}$, satisfying the non-vanishing commutation relations

$$\begin{aligned} [J, K_a] &= \epsilon_{ab} K_b, & [J, P_a] &= \epsilon_{ab} P_b, & [K_a, P_b] &= -\epsilon_{ab} H, \\ [J, Z_a] &= \epsilon_{ab} Z_b, & [P_a, P_b] &= -\epsilon_{ab} Z, & [H, P_a] &= \epsilon_{ab} Z_b, \\ [K_a, Z] &= -\epsilon_{ab} Z_b, \end{aligned} \quad (3.2)$$

where $a = 1, 2$, $\epsilon_{ab} \equiv \epsilon_{0ab}$, $\epsilon^{ab} \equiv \epsilon^{0ab}$. Here $\{J, K_a, H, P_a, Z, Z_a\}$ are spacial rotations, Carrollian boosts, time translations, space translations and the UR Maxwellian generators Z and Z_a , respectively. Note that the three commutators appearing in the first line of (3.2) define the Carroll algebra [1,2], which is obtained by a Carroll contraction of the Poincaré algebra and appears by suppressing the Maxwell generators Z and Z_a . Analogously to the Maxwellian extended Bargmann symmetry [64], one can notice that the Z generator present in this Maxwellian generalization of the Carroll algebra is not a central charge and thus does not appear as a central extension of the Carroll algebra. On the other hand, the Maxwellian Carroll algebra presented here is contained as a particular case of the infinite-dimensional Carrollian Maxwell algebra introduced in [71], which was obtained applying the S -expansion procedure [74]. Here, we have shown that the Maxwellian Carroll symmetry can be alternatively recovered as a UR limit of the Maxwell algebra. Now, we extend the study of [71] to the explicit construction of a CS action for the UR Maxwell algebra.

To this end, let us first consider the corresponding gauge connection one-form

$$A = \tau H + e^a P_a + \omega J + \omega^a K_a + k Z + k^a Z_a. \quad (3.3)$$

The curvature two-form $F = dA + \frac{1}{2}[A, A]$ is given by

$$F = R(\tau) H + R^a(e^b) P_a + R(\omega) J + R^a(\omega^b) K_a + R(k) Z + R^a(k^b) Z_a, \quad (3.4)$$

where the UR curvatures associated with the Maxwellian Carroll algebra are defined as follows:

$$\begin{aligned} R(\tau) &= d\tau + \epsilon^{ab} \omega_a e_b, & R^a(e^b) &= de^a + \epsilon^{ac} \omega e_c, \\ R(\omega) &= d\omega, & R^a(\omega^b) &= d\omega^a + \epsilon^{ac} \omega \omega_c, \\ R(k) &= dk + \frac{1}{2} \epsilon^{ab} e_a e_b, & R^a(k^b) &= dk^a + \epsilon^{ac} \omega k_c + \epsilon^{ac} \tau e_c + \epsilon^{ac} k \omega_c. \end{aligned} \quad (3.5)$$

Now, in order to construct the CS action invariant under the Maxwellian Carroll algebra, we also require the non-vanishing components of the invariant bilinear form. These can be derived as the Carroll contraction of the non-vanishing components of the invariant tensor for the Maxwell algebra. In particular, let us note that the relativistic invariant tensor for the Maxwell symmetry appears from (2.2) considering the vanishing cosmological constant limit $\ell \rightarrow \infty$. Note that, in order to end up with a non-trivial invariant tensor for the contracted algebra, we also have to rescale the relativistic constants appearing in (2.2) as follows:

$$\tilde{\alpha}_0 \rightarrow \alpha_0, \quad \tilde{\alpha}_1 \rightarrow \sigma \alpha_1, \quad \tilde{\alpha}_2 \rightarrow \alpha_2. \quad (3.6)$$

Then, after taking the limit $\sigma \rightarrow \infty$, we obtain the following non-vanishing components of the invariant bilinear form for the Maxwellian Carroll algebra:

$$\begin{aligned} \langle JJ \rangle &= -\alpha_0, & \langle JH \rangle &= -\alpha_1, & \langle JZ \rangle &= -\alpha_2, \\ \langle K_a P_b \rangle &= \alpha_1 \delta_{ab}, & \langle P_a P_b \rangle &= \alpha_2 \delta_{ab}. \end{aligned} \quad (3.7)$$

Then, considering the gauge connection one-form for the Maxwellian Carroll algebra (3.3) and the non-vanishing components (3.7) in the general expression of the CS action (2.6), we find the following UR CS action for the Maxwellian Carroll algebra:

$$I_{\text{Maxwell Carroll}} = \frac{k}{4\pi} \int \left\{ -\alpha_0 \omega R(\omega) + \alpha_1 \left[-2\tau R(\omega) + 2e_a R^a(\omega^b) \right] + \alpha_2 \left[-2k R(\omega) + e_a R^a(e^b) \right] \right\}. \quad (3.8)$$

This novel UR CS gravity action is defined by three different independent sectors, each one being invariant under the Maxwellian Carroll algebra. The first term along the α_0 constant corresponds to the UR version of the exotic Lagrangian, while the term proportional to α_1 is the usual Carrollian gravity [16]. On the other hand, the term along α_2 corresponds to the novel Maxwellian Carroll contribution. Let us notice that the UR CS gravity action (3.8) can also be recovered from the relativistic Maxwell CS action. Let us recall that the relativistic Maxwell CS action is obtained after applying the vanishing cosmological constant limit $\ell \rightarrow \infty$ in (2.7). Indeed, the Maxwell Carroll CS action (3.8) appears by performing the following redefinition of the one-form gauge fields:

$$\tilde{W}^a \rightarrow \frac{1}{\sigma} \omega^a, \quad \tilde{E}^0 \rightarrow \frac{1}{\sigma} \tau, \quad \tilde{K}^a \rightarrow \frac{1}{\sigma} k^a, \quad \tilde{W}^0 \rightarrow \omega, \quad \tilde{E}^a \rightarrow e^a, \quad \tilde{K}^0 \rightarrow k, \quad (3.9)$$

together with (3.6), and consequently taking the limit $\sigma \rightarrow \infty$ directly at the level of the action.

³ The σ parameter is related to the speed of light c as $\sigma \rightarrow 1/c$ such that the limit $\sigma \rightarrow \infty$ corresponds to the ultra-relativistic limit, $c \rightarrow 0$.

Table 1
Generators of UR and NR versions of the Maxwell algebra.

	Maxwell Carroll	MEC	MEB	GMEB
Time generators	J, H, Z	J, H, Z, S	$J, H, Z,$	J, H, Z, N
Spatial generators	K_a, P_a, Z_a	K_a, P_a, Z_a, L_a	G_a, P_a, Z_a	G_a, P_a, Z_a, N_a
Central charges		T	S, M, T	S, M, T, V
Amount of generators	9	13	12	16

Although the Carroll contraction of the Maxwell algebra allows the construction of a finite UR CS gravity action, it does not avoid the degeneracy problem. Indeed, the Maxwellian Carroll algebra (3.2) only allows for a degenerate invariant bilinear form. In particular, in our construction the absence of the k_a gauge field, and therefore of its CS kinetic term, in the CS action (3.8) is due to the degeneracy of the invariant tensor. In the present case the Carrollian limit procedure led to an algebra with a degenerate bilinear form that does not reproduce a well-defined CS action, as (3.8) does not contain a kinetic term for each gauge field. Then, as was mentioned in [64], guaranteeing finiteness of the action in the contraction process does not guarantee a non-degenerate bilinear form. Then, a natural question is if there exist another Carrollian version of the Maxwell algebra allowing for a non-degenerate invariant tensor. The answer to this question will be given in the next section.

4. Maxwellian extended Carroll Chern-Simons gravity theory

A different Carrollian version of the Maxwell algebra is contained as a particular case in the infinite-dimensional Carrollian Maxwell algebra introduced in [71]. The algebra is generated by the set of generators $\{J, K_a, H, P_a, Z, Z_a, S, L_a, T\}$, satisfying the non-vanishing commutation relations

$$\begin{aligned}
 [J, K_a] &= \epsilon_{ab} K_b, & [J, P_a] &= \epsilon_{ab} P_b, & [K_a, P_b] &= -\epsilon_{ab} H, \\
 [J, Z_a] &= \epsilon_{ab} Z_b, & [P_a, P_b] &= -\epsilon_{ab} Z, & [H, P_a] &= \epsilon_{ab} Z_b, \\
 [K_a, Z] &= -\epsilon_{ab} Z_b, & [K_a, K_b] &= -\epsilon_{ab} S, & [J, L_a] &= \epsilon_{ab} L_b, \\
 [S, P_a] &= \epsilon_{ab} L_b, & [K_a, Z_b] &= -\epsilon_{ab} T, & [P_a, L_b] &= -\epsilon_{ab} T, \\
 [K_a, H] &= -\epsilon_{ab} L_b.
 \end{aligned} \tag{4.1}$$

We shall call the algebra (4.1) as Maxwellian Extended Carroll (MEC) algebra.⁴ The MEC algebra is characterized by the presence of the S and L_a generators along with a central charge T . Note that it reduces to the Maxwellian Carroll algebra when $S = T = L_a = 0$.

Here, we present a novel UR CS gravity action based on the MEC algebra (4.1). The latter admits the following non-vanishing components of a non-degenerate invariant tensor:

$$\begin{aligned}
 \langle JJ \rangle &= \langle JS \rangle = -\alpha_0, & \langle K_a K_b \rangle &= \alpha_0 \delta_{ab}, \\
 \langle JH \rangle &= -\alpha_1, & \langle K_a P_b \rangle &= \alpha_1 \delta_{ab}, \\
 \langle JZ \rangle = \langle JT \rangle = \langle SZ \rangle = \langle HH \rangle &= -\alpha_2, & \langle P_a P_b \rangle = \langle P_a L_b \rangle = \langle K_a Z_b \rangle &= \alpha_2 \delta_{ab},
 \end{aligned} \tag{4.2}$$

where α_0, α_1 and α_2 are arbitrary constants. Let us note that here non-degeneracy requires a Maxwellian extension of the Carroll algebra analogously to the non-relativistic case in which a Maxwellian extension of the Bargmann (MEB) algebra assures the non-degeneracy of the invariant tensor [64]. Nevertheless, unlike Carroll and Bargmann algebra, there is no duality between the MEC and the MEB algebras since they differ in the amount of generators. Although one could see some similarity with the generalized Maxwellian extended Bargmann (GMEB) algebra introduced in [67], they are also quite different at the level of amount of generators. Table 1 summarizes the generators content of the previously mentioned UR and non-relativistic versions of the Maxwell algebra.

To construct an UR CS action gauge-invariant under the MEC algebra (4.1), let us consider the following gauge connection one-form:

$$A = \tau H + e^a P_a + \omega J + \omega^a K_a + k Z + k^a Z_a + s S + t T + l^a L_a. \tag{4.3}$$

The corresponding curvature two-form $F = dA + \frac{1}{2} [A, A]$ is given by

$$F = R(\tau) H + R^a(e^b) P_a + R(\omega) J + R^a(\omega^b) K_a + R(k) Z + R^a(k^b) Z_a + R(s) S + R(t) T + R^a(l^b) L_a, \tag{4.4}$$

where the respective field-strengths of the gauge fields dual to the generators of the MEC algebra are given by (3.5) along with

$$\begin{aligned}
 R(s) &= ds + \frac{1}{2} \epsilon^{ab} \omega_a \omega_b, \\
 R(t) &= dt + \epsilon^{ab} \omega_a k_b + \epsilon^{ab} e_a l_b, \\
 R^a(l^b) &= dl^a + \epsilon^{ab} \omega l_b + \epsilon^{ab} s e_b + \epsilon^{ab} \tau \omega_b.
 \end{aligned} \tag{4.5}$$

Considering the gauge connection one-form for the MEC algebra (4.3) and the non-vanishing components of the invariant tensor (4.2) in the general expression of the CS action (2.6), we find the following UR CS action for the MEC algebra,

⁴ The MEC algebra can be obtained from the relativistic Maxwell algebra by performing a particular expansion (see [71], where the MEC algebra corresponds to a finite sub-case of the infinite-dimensional Maxwellian UR algebra constructed).

$$I_{\text{MEC}} = \frac{k}{4\pi} \int \left\{ \alpha_0 \left[-\omega R(\omega) - 2sR(\omega) + \omega_a R^a(\omega^b) \right] + \alpha_1 \left[-2\tau R(\omega) + 2e_a R^a(\omega^b) \right] \right. \\ \left. + \alpha_2 \left[-2kR(\omega) + e_a R^a(e^b) - 2sR(k) - \tau R(\tau) - 2tR(\omega) \right. \right. \\ \left. \left. + 2l_a R^a(e^b) + \omega_a R^a(k^b) + k_a R^a(\omega^b) \right] \right\}. \quad (4.6)$$

This novel UR CS gravity action is defined by three different independent sectors, each one being invariant under the MEC algebra. Note that, as in the previous case, the term proportional to α_1 is the usual Carrollian gravity [16]. However, unlike the Maxwell Carroll CS gravity, the k^a gauge field appears explicitly along the α_2 constant. The term along α_2 describes a new UR Maxwell gravity which requires the introduction of an additional gauge field l^a and two one-form gauge fields s and t , respectively dual to the generators L_a , S and T of the UR algebra (4.1). Note that the MEC algebra allows for a non-degenerate invariant bilinear form which implies that in the action there is a kinetic term of each gauge field and that the field equations of the theory correspond to the vanishing of the MEC field-strengths. Indeed, the equations of motion are given by

$$\begin{aligned} \delta\omega_a: & \quad \alpha_0 R^a(\omega^b) + \alpha_1 R^a(e^b) + \alpha_2 R^a(k^b) = 0, \\ \delta\omega: & \quad \alpha_0 [R(\omega) + R(s)] + \alpha_1 R(\tau) + \alpha_2 [R(k) + R(t)] = 0, \\ \delta e_a: & \quad \alpha_1 R^a(\omega^b) + \alpha_2 [R^a(e^b) + R^a(l^b)] = 0, \\ \delta\tau: & \quad \alpha_1 R(\omega) + \alpha_2 R(\tau) = 0, \\ \delta k_a: & \quad \alpha_2 R^a(\omega^b) = 0, \\ \delta k: & \quad \alpha_2 [R(\omega) + R(s)] = 0, \\ \delta s: & \quad \alpha_0 R(\omega) + \alpha_2 R(k) = 0, \\ \delta t: & \quad \alpha_2 R(\omega) = 0, \\ \delta l_a: & \quad \alpha_2 R^a(e^b) = 0. \end{aligned} \quad (4.7)$$

As the non-degeneracy of the invariant tensor (4.2) requires $\alpha_2 \neq 0$, the field equations (4.7) are equivalent to the vanishing of the curvature two-forms (3.5) along with (4.5).

5. Enlarged Carroll Chern-Simons gravity theory and extension

Analogously to the non-relativistic Maxwell gravity, the MEC CS gravity theory does not contain a cosmological constant. The inclusion of a cosmological constant in the UR CS gravity theory can be done considering the AdS-Carroll symmetry [18]. Here, we show that a cosmological constant can be incorporated to the MEC CS gravity model by considering a new enlarged Carroll algebra. Nevertheless, as in the Maxwell case, an extended UR symmetry is required in order to avoid degeneracy. For completeness, we first explore the enlarged Carroll CS gravity along with its flat limit.

5.1. Enlarged Carroll Chern-Simons gravity

An enlarged Carroll algebra can be derived applying the Carroll contraction to the AdS-Lorentz algebra (2.1). Indeed, performing the redefinition of the generators as in (3.1) and after taking the limit $\sigma \rightarrow \infty$, we get

$$\begin{aligned} [J, K_a] &= \epsilon_{ab} K_b, & [J, P_a] &= \epsilon_{ab} P_b, & [K_a, P_b] &= -\epsilon_{ab} H, \\ [J, Z_a] &= \epsilon_{ab} Z_b, & [P_a, P_b] &= -\epsilon_{ab} Z, & [H, P_a] &= \epsilon_{ab} Z_b, \\ [K_a, Z] &= -\epsilon_{ab} Z_b, & [P_a, Z] &= -\frac{1}{\ell^2} \epsilon_{ab} P_b, & [P_a, Z_b] &= -\frac{1}{\ell^2} \epsilon_{ab} H, \\ [Z, Z_a] &= \frac{1}{\ell^2} \epsilon_{ab} Z_b, \end{aligned} \quad (5.1)$$

where ℓ is a length parameter related to the inverse of the cosmological constant Λ . The present symmetry corresponds to the three-dimensional version of the UR symmetry denoted as $\mathcal{C}\mathcal{L}_4$ in [75]. Let us note that, as its relativistic counterpart, the Maxwell Carroll algebra (3.2) appears in the vanishing cosmological constant limit $\ell \rightarrow \infty$.

The gauge connection one-form for the enlarged Carroll symmetry reads

$$A = \tau H + e^a P_a + \omega J + \omega^a K_a + kZ + k^a Z_a. \quad (5.2)$$

The corresponding curvature two-form is given by

$$F = \hat{R}(\tau) H + \hat{R}^a(e^b) P_a + \hat{R}(\omega) J + \hat{R}^a(\omega^b) K_a + \hat{R}(k) Z + \hat{R}^a(k^b) Z_a, \quad (5.3)$$

where

$$\begin{aligned}
 \hat{R}(\tau) &= d\tau + \epsilon^{ab}\omega_a e_b + \frac{1}{\ell^2}\epsilon^{ab}e_a k_b, & \hat{R}^a(e^b) &= de^a + \epsilon^{ac}\omega e_c + \frac{1}{\ell^2}\epsilon^{ac}k e_c, \\
 \hat{R}(\omega) &= d\omega, & \hat{R}^a(\omega^b) &= d\omega^a + \epsilon^{ac}\omega\omega_c, \\
 \hat{R}(k) &= dk + \frac{1}{2}\epsilon^{ab}e_a e_b, \\
 \hat{R}^a(k^b) &= dk^a + \epsilon^{ac}\omega k_c + \epsilon^{ac}\tau e_c + \epsilon^{ac}k\omega_c + \frac{1}{\ell^2}\epsilon^{ac}k k_c.
 \end{aligned} \tag{5.4}$$

Let us note that the enlarged Carroll curvatures coincide with the Maxwell Carroll ones (3.5) in the flat limit $\ell \rightarrow \infty$. On the other hand, one can show that the enlarged Carroll algebra (5.1) admits the following non-vanishing components of an invariant tensor:

$$\begin{aligned}
 \langle JJ \rangle &= -\alpha_0, & \langle JH \rangle &= -\alpha_1, \\
 \langle K_a P_b \rangle &= \alpha_1 \delta_{ab}, & \langle JZ \rangle &= -\alpha_2, \\
 \langle P_a P_b \rangle &= \alpha_2 \delta_{ab}, & \langle ZH \rangle &= -\frac{\alpha_1}{\ell^2}, \\
 \langle Z_a P_b \rangle &= \frac{\alpha_1}{\ell^2} \delta_{ab}, & \langle ZZ \rangle &= -\frac{\alpha_2}{\ell^2},
 \end{aligned} \tag{5.5}$$

where the constants α_0 , α_1 and α_2 have been obtained by applying the rescaling (3.6) and the UR limit to the ones appearing in the relativistic invariant tensor (2.2). Then, considering the gauge connection one-form for the enlarged Carroll algebra (5.1) and the non-vanishing components of the invariant tensor (5.5) in the general expression of the CS action (2.6), we find the following UR CS action:

$$\begin{aligned}
 I_{\text{Enlarged Carroll}} &= \frac{k}{4\pi} \int \left\{ -\alpha_0 \omega \hat{R}(\omega) + 2\alpha_1 \left[-\tau \hat{R}(\omega) + e_a \hat{R}^a(\omega^b) + \frac{1}{\ell^2} e_a F^a(k^b) - \frac{1}{\ell^2} \tau \hat{R}(k) \right] \right. \\
 &\quad \left. + \alpha_2 \left[-2k \hat{R}(\omega) + e_a \hat{R}^a(e^b) - \frac{1}{\ell^2} k dk \right] \right\},
 \end{aligned} \tag{5.6}$$

where $F^a(k^b) \equiv dk^a + \epsilon^{ac}\omega k_c + \epsilon^{ac}k\omega_c + \frac{1}{\ell^2}\epsilon^{ac}k k_c$. The CS action (5.6) is invariant under the enlarged Carroll algebra (5.1) and reproduces the Maxwellian Carroll CS action (3.8) in the vanishing cosmological constant limit $\ell \rightarrow \infty$. Such CS action can alternatively be obtained from the relativistic AdS-Lorentz CS action (2.7) by considering the redefinition (3.9), together with the rescaling (3.6), and taking the UR limit $\sigma \rightarrow \infty$. Although the Carroll limit is well-defined and the CS action is finite, the enlarged Carroll symmetry only admits a degenerate invariant bilinear form. As in the Maxwellian case, the degeneracy problem can be overcome by considering an extended version of the enlarged Carroll algebra.

5.2. Enlarged extended Carroll Chern-Simons gravity action and flat limit

The inclusion of a cosmological constant to the non-degenerate UR version of the Maxwell CS gravity theory can be done introducing additional bosonic generators to the enlarged Carroll algebra (5.1). Let us consider the set of generators $\{J, K_a, H, P_a, Z, Z_a, S, L_a, T\}$ which satisfy the commutation relations of the MEC algebra (4.1) along with

$$\begin{aligned}
 [H, Z_a] &= \frac{1}{\ell^2} \epsilon_{ab} L_b, & [Z, P_a] &= \frac{1}{\ell^2} \epsilon_{ab} P_b, & [P_a, Z_b] &= -\frac{1}{\ell^2} \epsilon_{ab} H, \\
 [Z, Z_a] &= \frac{1}{\ell^2} \epsilon_{ab} Z_b, & [T, P_a] &= \frac{1}{\ell^2} \epsilon_{ab} L_b, & [Z_a, Z_b] &= -\frac{1}{\ell^2} \epsilon_{ab} T, \\
 [Z, L_a] &= \frac{1}{\ell^2} \epsilon_{ab} L_b,
 \end{aligned} \tag{5.7}$$

where ℓ is related to the inverse of the cosmological constant Λ . We denote the UR algebra (5.7) as the Enlarged Extended Carroll (EEC) algebra. Naturally, the EEC algebra reproduces the enlarged Carroll one by setting $S = T = L_a = 0$. Moreover, the MEC algebra is recovered in the vanishing cosmological constant limit $\ell \rightarrow \infty$. Observe also that, in (5.7), T is not a central charge, while it reduces to a central generator in the $\ell \rightarrow \infty$ limit. One can notice that the EEC algebra admits a non-degenerate invariant tensor whose non-vanishing components are given by (4.2) along with

$$\begin{aligned}
 \langle ZH \rangle &= -\frac{\alpha_1}{\ell^2}, & \langle Z_a P_b \rangle &= \frac{\alpha_1}{\ell^2} \delta_{ab}, \\
 \langle ZZ \rangle &= \langle ZT \rangle = -\frac{\alpha_2}{\ell^2}, & \langle Z_a Z_b \rangle &= \frac{\alpha_2}{\ell^2} \delta_{ab},
 \end{aligned} \tag{5.8}$$

where the non-degeneracy requires $\alpha_2 \neq 0$ and $\alpha_0 \neq \alpha_2 \ell^2$. On the other hand, the EEC gauge connection one-form reads

$$A = \tau H + e^a P_a + \omega J + \omega^a K_a + kZ + k^a Z_a + sS + tT + l^a L_a. \tag{5.9}$$

The curvature two-form is given by

$$F = \hat{R}(\tau) H + \hat{R}^a(e^b) P_a + \hat{R}(\omega) J + \hat{R}^a(\omega^b) K_a + \hat{R}(k) Z + \hat{R}^a(k^b) Z_a + \hat{R}(s) S + \hat{R}(t) T + \hat{R}^a(l^b) L_a, \tag{5.10}$$

where the respective field-strengths are given by (5.4) and

$$\begin{aligned} \hat{R}(s) &= ds + \frac{1}{2}\epsilon^{ab}\omega_a\omega_b, & \hat{R}(t) &= dt + \epsilon^{ab}\omega_a k_b + \epsilon^{ab}e_a l_b + \frac{1}{2\ell^2}\epsilon^{ab}k_a k_b, \\ \hat{R}^a(l^b) &= dl^a + \epsilon^{ab}\omega_l b + \epsilon^{ab}s e_b + \epsilon^{ab}\tau\omega_b + \frac{1}{\ell^2}\epsilon^{ab}\tau k_b + \frac{1}{\ell^2}\epsilon^{ab}t e_b + \frac{1}{\ell^2}\epsilon^{ab}k_l b. \end{aligned} \tag{5.11}$$

Naturally, the MEC curvatures (4.5) are recovered in the flat limit $\ell \rightarrow \infty$. The UR CS action based on the EEC algebra is obtained by taking into account the non-vanishing components of the invariant tensor (5.8) and the gauge connection one-form (5.9) into the general CS expression (2.6). By doing so, we find the following UR CS gravity action invariant under the EEC algebra:

$$\begin{aligned} I_{\text{EEC}} &= \frac{k}{4\pi} \int \left\{ \alpha_0 \left[-\omega \hat{R}(\omega) - 2s \hat{R}(\omega) + \omega_a \hat{R}^a(\omega^b) \right] \right. \\ &\quad + 2\alpha_1 \left[-\tau \hat{R}(\omega) + e_a \hat{R}^a(\omega^b) + \frac{1}{\ell^2} e_a F^a(k^b) - \frac{1}{\ell^2} \tau \hat{R}(k) \right] \\ &\quad + \alpha_2 \left[-2k \hat{R}(\omega) + e_a \hat{R}^a(e^b) - 2s \hat{R}(k) - \tau \hat{R}(\tau) - 2t \hat{R}(\omega) + 2l_a \hat{R}^a(e^b) \right. \\ &\quad \left. \left. + \omega_a \hat{R}^a(k^b) + k_a \hat{R}^a(\omega^b) - \frac{1}{\ell^2} k d k + \frac{1}{\ell^2} k_a \hat{R}^a(k^b) - \frac{2}{\ell^2} t \hat{R}(k) \right] \right\}, \end{aligned} \tag{5.12}$$

where $F^a(k^b) \equiv dk^a + \epsilon^{ac}\omega_c k^b + \epsilon^{ac}k\omega_c + \frac{1}{\ell^2}\epsilon^{ac}kk_c$. As in the enlarged Carroll CS action (5.6), the UR Maxwellian gauge field k and k_a appear in both the α_1 and α_2 sectors. Nevertheless, it is the very presence of the additional gauge fields which allows to avoid the degeneracy problem present in the enlarged Carroll case. Indeed, for $\alpha_2 \neq 0$ and $\alpha_0 \neq \alpha_2 \ell^2$, the field equations exactly correspond to the vanishing of the curvature two-forms (5.10). Furthermore, in the vanishing cosmological constant limit $\ell \rightarrow \infty$, the CS action reduces to the non-degenerate MEC gravity action (4.6).

It is important to mention that, unlike the enlarged Carroll symmetry, the EEC algebra has not been obtained as a Carrollian limit of a relativistic symmetry. It would be worth it to explore if, following the method used in [71], the present UR algebra and its non-degenerate invariant tensor can be alternatively recovered as an UR expansion of a relativistic algebra and its invariant tensor, respectively.

6. Conclusions

In this work we have presented novel UR gravity theories being Maxwellian generalizations and enlargements of the Carroll gravity. In the Maxwellian case, we have shown that an extended version of the Maxwell Carroll algebra, which we have denoted as MEC algebra, is required in order to avoid degeneracy. Such extension of the UR Maxwell algebra involves the presence of extra bosonic generators (S, L_a, T , where in particular T is a central charge), with respect to the Maxwellian Carroll algebra, necessary in order to produce a non-degenerate UR invariant tensor. The CS actions for the Maxwell Carroll and the MEC algebras have been then constructed. In particular, the non-degeneracy of the invariant trace in the MEC case ensures that the field equations arising from the MEC gravity theory are given by the vanishing of the curvature two-forms. Consequently, we have studied an alternative origin of the Maxwell Carroll and the MEC gravity theories as a flat limit, showing that both can be recovered in the vanishing cosmological constant limit (namely for $\ell \rightarrow \infty$) of a novel enlarged Carroll and enlarged extended Carroll (EEC) gravity theories, respectively. Although the enlarged Carroll algebra has a well-defined vanishing cosmological constant limit, it does not admit a non-degenerate invariant tensor. Remarkably, the EEC algebra admits a non-degenerate invariant bilinear form and therefore allows the proper construction of a CS gravity theory whose flat limit is also non-degenerate.

Let us note that the Maxwell Carroll and the MEC algebras correspond to particular sub-cases of the infinite-dimensional Carrollian Maxwell algebra defined in [71]. Here, we have extended the study of [71] to the explicit construction of a CS action for both UR Maxwell algebras. Such construction allowed us to elucidate which UR version of the infinite-dimensional Carrollian Maxwell algebra is a good candidate to construct a non-degenerate UR CS action. To our knowledge, the enlarged Carroll and the EEC algebras obtained here do not belong to a generalized family of UR infinite-dimensional algebras. It would be interesting to explore if, analogously to the Maxwellian case, the enlarged Carroll and its extended version can be seen as particular sub-cases of an infinite family of UR generalizations. In particular, one could expect that such generalization can be related to the UR expansion of the Maxwell algebra introduced in [71] through a vanishing cosmological constant limit $\ell \rightarrow \infty$.

On the other hand, as it has been shown that UR symmetry groups play a remarkable role in the contexts of flat holography (i.e., holography of flat space) and fluid/gravity correspondence [76–84], it would be intriguing to study the holographic implications of the MEC and EEC symmetries, in particular in connection with the fluid/gravity correspondence. Furthermore, it would be worth it to study supersymmetric extensions of the UR algebras and CS theories developed in the present paper, following the lines of what was done in, e.g., [19,20] in the case of AdS Carroll CS supergravity and flat limit. From the results we have obtained at the purely bosonic level, in the supersymmetric case we expect non-trivial bosonic and, in particular, fermionic enlargement and extensions to play a prominent role in the CS construction.

It would be interesting to go further in the study of the Maxwell and AdS- \mathcal{L} symmetries. In particular, one could explore the complete schematic “cube” summarizing the sequential contractions starting from the Maxwell and AdS- \mathcal{L} algebras. The NR version of both symmetries was recently introduced in [64,65]. In this paper, we have presented the respective UR counterparts with a detailed analysis of the non-degeneracy of the invariant tensor. It would be worth it to elucidate the respective Static Hopf and para-Galilei version of the Maxwell and AdS- \mathcal{L} algebras.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This work was funded by the National Agency for Research and Development ANID (ex-CONICYT) - PAI grant No. 77190078 (P.C.) and FONDECYT grants No. 1211077 (P.C.). This work was supported by the Research project Code DIREG_09/2020 (P.C.) of the Universidad Católica de la Santísima Concepción, Chile. P.C. would like to thank to the Dirección de Investigación and Vice-rectoría de Investigación of the Universidad Católica de la Santísima Concepción, Chile, for their constant support. L.R. would like to thank the Department of Applied Science and Technology of the Polytechnic University of Turin, and in particular Laura Andrianopoli and Francesco Raffa, for financial support.

References

- [1] J. Lévy-Leblond, Une nouvelle limite non-relativiste du groupe de poincaré, *Ann. Inst. Henri Poincaré A, Phys. Théor.* 3 (1965) 1–12.
- [2] H. Bacry, J. Levy-Leblond, Possible kinematics, *J. Math. Phys.* 9 (1968) 1605–1614.
- [3] G. Gibbons, K. Hashimoto, P. Yi, Tachyon condensates, Carrollian contraction of Lorentz group, and fundamental strings, *J. High Energy Phys.* 09 (2002) 061, arXiv:hep-th/0209034.
- [4] D.M. Hofman, B. Rollier, Warped conformal field theory as lower spin gravity, *Nucl. Phys. B* 897 (2015) 1–38, arXiv:1411.0672.
- [5] A. Bagchi, Tensionless strings and Galilean conformal algebra, *J. High Energy Phys.* 05 (2013) 141, arXiv:1303.0291.
- [6] A. Bagchi, S. Chakraborty, P. Parekh, Tensionless strings from worldsheet symmetries, *J. High Energy Phys.* 01 (2016) 158, arXiv:1507.04361.
- [7] A. Bagchi, S. Chakraborty, P. Parekh, Tensionless superstrings: view from the worldsheet, *J. High Energy Phys.* 10 (2016) 113, arXiv:1606.09628.
- [8] A. Bagchi, A. Banerjee, S. Chakraborty, P. Parekh, Inhomogeneous tensionless superstrings, *J. High Energy Phys.* 02 (2018) 065, arXiv:1710.03482.
- [9] A. Bagchi, A. Banerjee, S. Chakraborty, P. Parekh, Exotic origins of tensionless superstrings, *Phys. Lett. B* 801 (2020) 135139, arXiv:1811.10877.
- [10] C. Duval, G.W. Gibbons, P.A. Horvathy, P.M. Zhang, Carroll versus Newton and Galilei: two dual non-Einsteinian concepts of time, *Class. Quantum Gravity* 31 (2014) 085016, arXiv:1402.0657.
- [11] T.E. Clark, T. ter Veldhuis, AdS-Carroll branes, *J. Math. Phys.* 57 (11) (2016) 112303, arXiv:1605.05484.
- [12] R. Basu, U.N. Chowdhury, Dynamical structure of Carrollian electrodynamics, *J. High Energy Phys.* 04 (2018) 111, arXiv:1802.09366.
- [13] D. Roychowdhury, Carroll membranes, *J. High Energy Phys.* 10 (2019) 258, arXiv:1908.07280.
- [14] E. Bergshoeff, J. Gomis, L. Parra, The symmetries of the Carroll superparticle, *J. Phys. A* 49 (18) (2016) 185402, arXiv:1503.06083.
- [15] J. Hartong, Gauging the Carroll algebra and ultra-relativistic gravity, *J. High Energy Phys.* 08 (2015) 069, arXiv:1505.05011.
- [16] E. Bergshoeff, D. Grumiller, S. Prohazka, J. Rosseel, Three-dimensional spin-3 theories based on general kinematical algebras, *J. High Energy Phys.* 01 (2017) 114, arXiv:1612.02277.
- [17] E. Bergshoeff, J. Gomis, B. Rollier, J. Rosseel, T. ter Veldhuis, Carroll versus Galilei gravity, *J. High Energy Phys.* 03 (2017) 165, arXiv:1701.06156.
- [18] J. Matulich, S. Prohazka, J. Salzer, Limits of three-dimensional gravity and metric kinematical Lie algebras in any dimension, *J. High Energy Phys.* 07 (2019) 118, arXiv:1903.09165.
- [19] L. Ravera, AdS Carroll Chern-Simons supergravity in $2 + 1$ dimensions and its flat limit, *Phys. Lett. B* 795 (2019) 331–338, arXiv:1905.00766.
- [20] F. Ali, L. Ravera, \mathcal{N} -extended Chern-Simons Carrollian supergravities in $2 + 1$ spacetime dimensions, *J. High Energy Phys.* 02 (2020) 128, arXiv:1912.04172.
- [21] D. Grumiller, J. Hartong, S. Prohazka, J. Salzer, Limits of JT gravity, *J. High Energy Phys.* 02 (2021) 134, arXiv:2011.13870.
- [22] J. Gomis, D. Hidalgo, P. Salgado-Rebolledo, Non-relativistic and Carrollian limits of Jackiw-Teitelboim gravity, *J. High Energy Phys.* 05 (2021) 162, arXiv:2011.15053.
- [23] R. Schrader, The Maxwell group and the quantum theory of particles in classical homogeneous electromagnetic fields, *Fortschr. Phys.* 20 (1972) 701–734.
- [24] H. Bacry, P. Combe, J. Richard, Group-theoretical analysis of elementary particles in an external electromagnetic field. 1. the relativistic particle in a constant and uniform field, *Nuovo Cimento A* 67 (1970) 267–299.
- [25] H. Bacry, P. Combe, J.L. Richard, Group-theoretical analysis of elementary particles in an external electromagnetic field. 2. the nonrelativistic particle in a constant and uniform field, *Nuovo Cimento A* 70 (1970) 289–312.
- [26] J. Gomis, A. Kleinschmidt, On free Lie algebras and particles in electro-magnetic fields, *J. High Energy Phys.* 07 (2017) 085, arXiv:1705.05854.
- [27] J.D. Edelstein, M. Hassaine, R. Troncoso, J. Zanelli, Lie-algebra expansions, Chern-Simons theories and the Einstein-Hilbert Lagrangian, *Phys. Lett. B* 640 (2006) 278–284, arXiv:hep-th/0605174.
- [28] F. Izaurieta, E. Rodríguez, P. Minning, P. Salgado, A. Perez, Standard general relativity from Chern-Simons gravity, *Phys. Lett. B* 678 (2009) 213–217, arXiv:0905.2187.
- [29] P. Concha, D. Peñafiel, E. Rodríguez, P. Salgado, Even-dimensional general relativity from Born-Infeld gravity, *Phys. Lett. B* 725 (2013) 419–424, arXiv:1309.0062.
- [30] P. Concha, D. Penafiel, E. Rodríguez, P. Salgado, Chern-Simons and Born-Infeld gravity theories and Maxwell algebras type, *Eur. Phys. J. C* 74 (2014) 2741, arXiv:1402.0023.
- [31] D. Cangemi, One formulation for both lineal gravities through a dimensional reduction, *Phys. Lett. B* 297 (1992) 261–265, arXiv:gr-qc/9207004.
- [32] C. Duval, Z. Horvath, P.A. Horvathy, Chern-Simons gravity, based on a non-semisimple group, arXiv:0807.0977.
- [33] P. Salgado, R.J. Szabo, O. Valdivia, Topological gravity and transgression holography, *Phys. Rev. D* 89 (8) (2014) 084077, arXiv:1401.3653.
- [34] S. Hosenzadeh, A. Rezaei-Aghdam, $(2+1)$ -dimensional gravity from Maxwell and semisimple extension of the Poincaré gauge symmetric models, *Phys. Rev. D* 90 (8) (2014) 084008, arXiv:1402.0320.
- [35] R. Caroca, P. Concha, O. Fierro, E. Rodríguez, P. Salgado-Rebolledo, Generalized Chern-Simons higher-spin gravity theories in three dimensions, *Nucl. Phys. B* 934 (2018) 240–264, arXiv:1712.09975.
- [36] P. Concha, N. Merino, O. Miskovic, E. Rodríguez, P. Salgado-Rebolledo, O. Valdivia, Asymptotic symmetries of three-dimensional Chern-Simons gravity for the Maxwell algebra, *J. High Energy Phys.* 10 (2018) 079, arXiv:1805.08834.
- [37] P. Concha, D.M. Peñafiel, E. Rodríguez, On the Maxwell supergravity and flat limit in $2 + 1$ dimensions, *Phys. Lett. B* 785 (2018) 247–253, arXiv:1807.00194.
- [38] P. Concha, N -extended Maxwell supergravities as Chern-Simons theories in three spacetime dimensions, *Phys. Lett. B* 792 (2019) 290–297, arXiv:1903.03081.
- [39] D. Chernyavsky, N.S. Deger, D. Sorokin, Spontaneously broken $3d$ Hietarinta/Maxwell Chern-Simons theory and minimal massive gravity, *Eur. Phys. J. C* 80 (6) (2020) 556, arXiv:2002.07592.
- [40] H. Adami, P. Concha, E. Rodríguez, H. Safari, Asymptotic symmetries of Maxwell Chern-Simons gravity with torsion, *Eur. Phys. J. C* 80 (10) (2020) 967, arXiv:2005.07690.
- [41] R. Caroca, P. Concha, J. Matulich, E. Rodríguez, D. Tempo, Hypersymmetric extensions of Maxwell Chern-Simons gravity in $(2+1)$ dimensions, arXiv:2105.12243.
- [42] J.A. de Azcarraga, K. Kamimura, J. Lukierski, Generalized cosmological term from Maxwell symmetries, *Phys. Rev. D* 83 (2011) 124036, arXiv:1012.4402.
- [43] R. Durka, J. Kowalski-Glikman, M. Szczachor, Gauged AdS-Maxwell algebra and gravity, *Mod. Phys. Lett. A* 26 (2011) 2689–2696, arXiv:1107.4728.
- [44] J.A. de Azcarraga, K. Kamimura, J. Lukierski, Maxwell symmetries and some applications, *Int. J. Mod. Phys. Conf. Ser.* 23 (2013) 01160, arXiv:1201.2850.
- [45] P. Concha, E. Rodríguez, $N = 1$ Supergravity and Maxwell superalgebras, *J. High Energy Phys.* 09 (2014) 090, arXiv:1407.4635.
- [46] D.M. Peñafiel, L. Ravera, On the hidden Maxwell superalgebra underlying $D=4$ supergravity, *Fortschr. Phys.* 65 (9) (2017) 1700005, arXiv:1701.04234.
- [47] L. Ravera, Hidden role of Maxwell superalgebras in the free differential algebras of $D = 4$ and $D = 11$ supergravity, *Eur. Phys. J. C* 78 (3) (2018) 211, arXiv:1801.08860.
- [48] P. Concha, L. Ravera, E. Rodríguez, On the supersymmetry invariance of flat supergravity with boundary, *J. High Energy Phys.* 01 (2019) 192, arXiv:1809.07871.
- [49] P. Salgado-Rebolledo, The Maxwell group in $2+1$ dimensions and its infinite-dimensional enhancements, *J. High Energy Phys.* 10 (2019) 039, arXiv:1905.09421.
- [50] P. Concha, H. Safari, On stabilization of Maxwell-BMS algebra, *J. High Energy Phys.* 04 (2020) 073, arXiv:1909.12827.
- [51] S. Kibaroglu, O. Cebecioglu, Gauge theory of the Maxwell and semi-simple extended (anti) de Sitter algebra, arXiv:2007.14795.
- [52] O. Cebecioglu, S. Kibaroglu, Maxwell-modified metric affine gravity, arXiv:2104.12670.
- [53] J. Diaz, O. Fierro, F. Izaurieta, N. Merino, E. Rodríguez, P. Salgado, O. Valdivia, A generalized action for $(2 + 1)$ -dimensional Chern-Simons gravity, *J. Phys. A* 45 (2012) 255207, arXiv:1311.2215.
- [54] O. Fierro, F. Izaurieta, P. Salgado, O. Valdivia, Minimal AdS-Lorentz supergravity in three-dimensions, *Phys. Lett. B* 788 (2019) 198–205, arXiv:1401.3697.
- [55] P. Concha, N. Merino, E. Rodríguez, P. Salgado-Rebolledo, O. Valdivia, Semi-simple enlargement of the \mathfrak{bms}_3 algebra from a $\mathfrak{so}(2, 2) \oplus \mathfrak{so}(2, 1)$ Chern-Simons theory, *J. High Energy Phys.* 02 (2019) 002, arXiv:1810.12256.

- [56] P. Concha, R. Durka, C. Inostroza, N. Merino, E. Rodríguez, Pure Lovelock gravity and Chern-Simons theory, *Phys. Rev. D* 94 (2) (2016) 024055, arXiv:1603.09424.
- [57] P. Concha, E. Rodríguez, Generalized pure Lovelock gravity, *Phys. Lett. B* 774 (2017) 616–622, arXiv:1708.08827.
- [58] M. Ipinza, P. Concha, L. Ravera, E. Rodríguez, On the supersymmetric extension of Gauss-Bonnet like gravity, *J. High Energy Phys.* 09 (2016) 007, arXiv:1607.00373.
- [59] D.M. Peñafiel, L. Ravera, Generalized cosmological term in $D = 4$ supergravity from a new AdS–Lorentz superalgebra, *Eur. Phys. J. C* 78 (11) (2018) 945, arXiv:1807.07673.
- [60] A. Banaudi, L. Ravera, Generalized AdS-Lorentz deformed supergravity on a manifold with boundary, *Eur. Phys. J. Plus* 133 (12) (2018) 514, arXiv:1803.08738.
- [61] S. Bonanos, J. Gomis, Infinite sequence of Poincare group extensions: structure and dynamics, *J. Phys. A* 43 (2010) 015201, arXiv:0812.4140.
- [62] S. Bonanos, J. Gomis, A note on the Chevalley-Eilenberg cohomology for the Galilei and Poincare algebras, *J. Phys. A* 42 (2009) 145206, arXiv:0808.2243.
- [63] J. Gomis, K. Kamimura, J. Lukierski, Deformations of Maxwell algebra and their dynamical realizations, *J. High Energy Phys.* 08 (2009) 039, arXiv:0906.4464.
- [64] L. Avilés, E. Frodden, J. Gomis, D. Hidalgo, J. Zanelli, Non-relativistic Maxwell Chern-Simons gravity, *J. High Energy Phys.* 05 (2018) 047, arXiv:1802.08453.
- [65] P. Concha, E. Rodríguez, Non-relativistic gravity theory based on an enlargement of the extended Bargmann algebra, *J. High Energy Phys.* 07 (2019) 085, arXiv:1906.00086.
- [66] D.M. Peñafiel, P. Salgado-Rebolledo, Non-relativistic symmetries in three space-time dimensions and the Nappi-Witten algebra, *Phys. Lett. B* 798 (2019) 135005, arXiv:1906.02161.
- [67] P. Concha, M. Ipinza, E. Rodríguez, Generalized Maxwellian exotic Bargmann gravity theory in three spacetime dimensions, *Phys. Lett. B* 807 (2020) 135593, arXiv:2004.01203.
- [68] P. Concha, L. Ravera, E. Rodríguez, G. Rubio, Three-dimensional Maxwellian extended Newtonian gravity and flat limit, *J. High Energy Phys.* 10 (2020) 181, arXiv:2006.13128.
- [69] P. Concha, L. Ravera, E. Rodríguez, Three-dimensional Maxwellian extended Bargmann supergravity, *J. High Energy Phys.* 04 (2020) 051, arXiv:1912.09477.
- [70] P. Concha, M. Ipinza, L. Ravera, E. Rodríguez, Non-relativistic three-dimensional supergravity theories and semigroup expansion method, arXiv:2010.01216.
- [71] J. Gomis, A. Kleinschmidt, J. Palmkvist, P. Salgado-Rebolledo, Newton-Hooke/Carrollian expansions of (A)dS and Chern-Simons gravity, *J. High Energy Phys.* 02 (2020) 009, arXiv:1912.07564.
- [72] D.V. Soroka, V.A. Soroka, Semi-simple extension of the (super)Poincare algebra, *Adv. High Energy Phys.* 2009 (2009) 234147, arXiv:hep-th/0605251.
- [73] E. Witten, (2+1)-dimensional gravity as an exactly soluble system, *Nucl. Phys. B* 311 (1988) 46.
- [74] F. Izaurieta, E. Rodríguez, P. Salgado, Expanding Lie (super)algebras through Abelian semigroups, *J. Math. Phys.* 47 (2006) 123512, arXiv:hep-th/0606215.
- [75] G. Rubio, Gravedad newtoniana en teorías de gauge extendidas, PhD thesis, Concepcion U, 2020, 11.
- [76] A. Bagchi, Correspondence between asymptotically flat spacetimes and nonrelativistic conformal field theories, *Phys. Rev. Lett.* 105 (2010) 171601, arXiv:1006.3354.
- [77] A. Bagchi, R. Fareghbal, BMS/GCA redux: towards flatspace holography from non-relativistic symmetries, *J. High Energy Phys.* 10 (2012) 092, arXiv:1203.5795.
- [78] A. Bagchi, R. Basu, A. Kakkar, A. Mehra, Flat holography: aspects of the dual field theory, *J. High Energy Phys.* 12 (2016) 147, arXiv:1609.06203.
- [79] I. Lodato, W. Merbis, Super-BMS₃ algebras from $\mathcal{N} = 2$ flat supergravities, *J. High Energy Phys.* 11 (2016) 150, arXiv:1610.07506.
- [80] A. Bagchi, A. Mehra, P. Nandi, Field theories with conformal Carrollian symmetry, *J. High Energy Phys.* 05 (2019) 108, arXiv:1901.10147.
- [81] L. Ciambelli, C. Marteau, A.C. Petkou, P.M. Petropoulos, K. Siampos, Covariant Galilean versus Carrollian hydrodynamics from relativistic fluids, *Class. Quantum Gravity* 35 (16) (2018) 165001, arXiv:1802.05286.
- [82] L. Ciambelli, C. Marteau, A.C. Petkou, P.M. Petropoulos, K. Siampos, Flat holography and Carrollian fluids, *J. High Energy Phys.* 07 (2018) 165, arXiv:1802.06809.
- [83] L. Ciambelli, C. Marteau, Carrollian conservation laws and Ricci-flat gravity, *Class. Quantum Gravity* 36 (8) (2019) 085004, arXiv:1810.11037.
- [84] A. Campoleoni, L. Ciambelli, C. Marteau, P.M. Petropoulos, K. Siampos, Two-dimensional fluids and their holographic duals, *Nucl. Phys. B* 946 (2019) 114692, arXiv:1812.04019.