

Potential and mass matrix in gauged N=4 supergravity

Original

Potential and mass matrix in gauged N=4 supergravity / M., DE ROO; D. B., Westra; Sudhakar, Panda; Trigiante, Mario.
- In: JOURNAL OF HIGH ENERGY PHYSICS. - ISSN 1029-8479. - 2003:11(2003). [10.1088/1126-6708/2003/11/022]

Availability:

This version is available at: 11583/1723510 since: 2024-07-09T07:16:33Z

Publisher:

IOP publishing

Published

DOI:10.1088/1126-6708/2003/11/022

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Potential and mass-matrix in gauged $N = 4$ supergravity

To cite this article: Mees de Roo *et al* JHEP11(2003)022

View the [article online](#) for updates and enhancements.

You may also like

- [Group manifold reduction of dual \$N = 1\$ \$d = 10\$ supergravity](#)
Mees de Roo, Martijn G.C. Eenink, Dennis B. Westra et al.
- [Cosmological solutions from 4D \$N = 4\$ matter-coupled supergravity](#)
H L Dao
- [Options for gauge groups in five-dimensional supergravity](#)
John Ellis, Murat Günaydin and Marco Zagermann

Potential and mass-matrix in gauged $N = 4$ supergravity

Mess de Roo and Dennis B. Westra

*Institute for Theoretical Physics
Nijenborgh 4, 9747 AG Groningen, The Netherlands
E-mail: m.de.roo@phys.rug.nl, d.b.westra@phys.rug.nl*

Sudhakar Panda

*Harish-Chandra Research Institute
Chatnag Road, Jhusi, Allahabad 211019, India
E-mail: panda@mri.ernet.in*

Mario Trigiante

*Spinoza Institute
Leuvenlaan 4, 3508 TD Utrecht, The Netherlands
E-mail: m.trigiante@phys.uu.nl*

ABSTRACT: We discuss the potential and mass-matrix of gauged $N = 4$ matter coupled supergravity for the case of six matter multiplets, extending previous work by considering the dependence on all scalars. We consider all semi-simple gauge groups and analyse the potential and its first and second derivatives in the origin of the scalar manifold. Although we find in a number of cases an extremum with a positive cosmological constant, these are not stable under fluctuations of all scalar fields.

KEYWORDS: Extended Supersymmetry, Supergravity Models, Cosmology of Theories beyond the SM.

Contents

1. Introduction	1
2. The potential, its derivatives, and the mass-matrix	3
2.1 The potential	3
2.2 The first derivatives of the potential	6
2.3 The second derivatives of the potential	8
2.4 Masses of scalar and vector fields	9
3. Semi-simple gauge groups	10
3.1 $SO(2,1)_+^3 \otimes SO(3)_+$	11
3.2 $SO(2,1)_+^2 \otimes SO(2,1)_-^2$	13
3.3 $SO(3,1)_+ \otimes SO(2,1)_+ \otimes SO(2,1)_-$	13
3.4 $SO(3,1)_+ \otimes SO(3,1)_+$	14
3.5 $SO(3)_-^2 \otimes SO(3)_+^2$	15
3.6 $SO(3,1)_- \otimes SO(3)_- \otimes SO(3)_+$	15
3.7 $SO(3,1)_- \otimes SO(3,1)_-$	16
3.8 $SL(3, \mathbb{R})_- \otimes SO(3)_-$	16
3.9 $SU(2,1)_+ \otimes SO(2,1)_+$	17
4. Conclusions	17
A. $SU(1,1)$ scalars and angles	20
B. Generators and structure constants	22
C. The parameters P	24

1. Introduction

Recent astronomical observations [1, 2] have led to the conclusion that our universe is presently in a state of accelerating expansion, and that its cosmological properties are best described by assuming the presence of a large amount of dark matter of unknown origin, as well as an even larger amount of dark energy, which one considers as due to a cosmological constant. This situation has renewed interest in fundamental theories with scalar potentials, as these may have nonzero extremal values, thus presenting a possibility to explain the cosmological constant. Motivated by string and M theory, the search for the appropriate fundamental theory concentrates on the supergravity theories which arise as low energy limit of compactified string theories. A major problem in this respect is that the sign of the cosmological constant should be positive to explain the expansion, a

property which is hard to reconcile with theories which have their origin in string theory ([3, 4, 5], for a recent discussion see [6]).

In this paper we take the point of view that one should first obtain a solution of a four-dimensional supergravity theory with the appropriate properties (de Sitter, stability), and consider the connection with string theory as a second step. We will concentrate on the properties of gauged $N = 4$ supergravity theories coupled to additional matter multiplets. It is well-known that these theories do allow extrema with a positive cosmological constant [7]. In a previous paper [8] a situation where the scalar manifold is truncated to four scalar fields was discussed, and it was shown that a positive, stable extremum of the potential is possible when the scalar fluctuations are limited to the four directions which survive the truncation. In the present paper we will extend the scope of these investigations to include more general gauge groups, and also consider the fluctuations in all scalars present in the model.

A further motivation for this investigation arises from [9], where a stable solution in de Sitter space was constructed in gauged $N = 2$ supergravity. There is no obvious connection of this solution with string theory, but the authors indicated a possible relation with $N = 4$. Since $N = 4$ is a step closer to the maximal ten and eleven dimensional supergravity theories, the connection with string theory might be easier to obtain once the $N = 2$ case is raised to $N = 4$. In section 4 we will discuss the $N = 2$ aspect of our work in more detail.

We leave for later work the question of how these theories arise from the higher dimensional low-energy limit of string theory. It is well known that ungauged $N = 4$ supergravity theories in $d = 4$ can be obtained by toroidal compactification from $d = 10$ supergravity [10]. Gauged $N = 4$ supergravity obtained from Scherk-Schwarz compactifications [11] has been considered in the past [12, 13, 14], and also more recently in combination with flux compactification [15]–[20]. Nevertheless, to our knowledge the $SU(1, 1)$ duality angles [21], which play an essential role in obtaining extrema with a positive cosmological constant, have not been given a higher-dimensional origin.

Throughout this paper we will use the notation of and results from [8]. We will discuss the value of the potential and its derivatives, as well as the definition of the bosonic mass-matrices, in section 2. Nine semi-simple gauge groups satisfy the conditions that (i) the potential has an extremum for all scalar fields, (ii) the value of the potential in the extremum is positive. These groups are introduced in section 2.2 and their properties are discussed in detail in the different subsections of section 3.

To complete this Introduction we will review some basic properties of the parametrisation of the scalar sector of matter coupled $N = 4$ supergravity [22, 21]. We consider gauged $N = 4$ supergravity coupled to n vector multiplets. The scalar fields of the theory are¹ Z_a^R (real) and ϕ_α (complex), satisfying the constraints

$$\phi^\alpha \phi_\alpha = 1, \tag{1.1}$$

¹The indices α, β, \dots take on values 1 and 2, indices $R, S \dots$ the values $1, \dots, 6 + n$, and the indices a, b, \dots the values $1, \dots, 6$. The metric η_{RS} can be chosen as $\text{diag}(-1, -1, -1, -1, -1, -1, +1, \dots, +1)$, with n positive entries. In comparison to [21] we have replaced the complex scalars ϕ_{ij}^R by real scalars Z_a^R : $\phi_{ij}^R = \frac{1}{2} Z_a^R (G^a)_{ij}$, where the G^a are six matrices which ensure that Z_a^R transforms as a vector under $SO(6)$.

$$\eta_{RS} Z_a^R Z_b^S = -\delta_{ab}. \tag{1.2}$$

The scalars ϕ_α ($\phi^1 = (\phi_1)^*$, $\phi^2 = -(\phi_2)^*$) transform under global $SU(1,1)$ and local $U(1)$, the Z_a^R transform under local $SO(6) \times SO(n)$, and under global $SO(6,n)$. The constraints and the local symmetry restrict the scalars to the cosets $SU(1,1)/U(1)$ (two physical scalars) and $SO(6,n)/SO(6) \times SO(n)$ ($6n$ physical scalars).

There is a certain freedom in coupling the vector multiplets: for each multiplet, labeled by R , we can introduce an $SU(1,1)$ element, of which only a single angle α_R turns out to be important. These angles α_R can be reinterpreted as a modification of the $SU(1,1)$ scalars coupling to the multiplet R in the form

$$\phi_{(R)}^1 = e^{i\alpha_R} \phi^1, \quad \phi_{(R)}^2 = e^{-i\alpha_R} \phi^2, \quad \Phi_{(R)} = e^{i\alpha_R} \phi^1 + e^{-i\alpha_R} \phi^2. \tag{1.3}$$

The gauge group has to be a subgroup of $SO(6,n)$. For a semi-simple gauge group the α_R (called $SU(1,1)$ angles in this paper) have to be the same for all R belonging to the same factor of the gauge group. This gauging breaks the global $SO(6,n)$ symmetry of the ungauged theory. In the remainder of this paper we will limit ourselves to $n = 6$. The reason for the choice $n = 6$ is, besides its relative simplicity, that this case follows by toroidal compactification from $d = 10$ $N = 1$ supergravity without additional matter.

2. The potential, its derivatives, and the mass-matrix

2.1 The potential

The scalar potential $V(\phi, Z)$ can be written in the form

$$V = \sum_{i,j} (R^{(ij)}(\phi) V_{ij}(Z) + I^{(ij)} W_{ij}(Z)). \tag{2.1}$$

The indices i, j, \dots label the different factors in the gauge group G , which we will take to be semi-simple. $R^{(ij)}$ and $I^{(ij)}$ contain the $SU(1,1)$ scalars and depend on the gauge coupling constants and the $SU(1,1)$ angles, V_{ij} and W_{ij} contain the structure constants, depend on the matter fields, and are symmetric resp. antisymmetric in the indices i, j .

The extremum of the potential in the ϕ direction has been determined in [8]. For completeness we briefly review this analysis in appendix A. The conclusion is that in the extremum in the $SU(1,1)$ scalars the potential takes on the form

$$V_0 = \text{sgn} C_- \sqrt{\Delta} - T_-, \tag{2.2}$$

where (see [8])

$$C_- = \sum_{ij} g_i g_j \cos(\alpha_i - \alpha_j) V_{ij}, \tag{2.3}$$

$$T_- = \sum_{ij} a_{ij} W_{ij}, \tag{2.4}$$

$$\Delta = 2 \sum_{ij} \sum_{kl} V_{ij} V_{kl} a_{ik} a_{jl}, \tag{2.5}$$

Group	-	+	c	nc	Group	-	+	c	nc
SO(3) ₋	3	0	3	0	SO(3) ₊	0	3	3	0
SO(2,1) ₋	1	2	1	2	SO(2,1) ₊	2	1	1	2
SO(3,1) ₋	3 _c	3 _{nc}	3	3	SO(3,1) ₊	3 _{nc}	3 _c	3	3
SL(3,ℝ) ₋	3	5	3	5	SL(3,ℝ) ₊	5	3	3	5
SU(2,1) ₋	4 _c	4 _{nc}	4	4	SU(2,1) ₊	4 _{nc}	4 _c	4	4

Table 1: List of allowed simple groups. The first two columns indicate how the group is embedded in SO(6,6) with respect to the signs in the metric η_{RS} , the column c and nc indicate the number of compact and noncompact generators, respectively. The structure constants of these groups are presented in appendix B.

$$R^{(ij)} = \frac{\text{sgn}C_-}{\sqrt{\Delta}} \sum_{kl} V_{kl} (2a_{ik}a_{jl} - a_{ij}a_{kl}), \quad (2.6)$$

$$I^{(ij)} = -a_{ij}. \quad (2.7)$$

and

$$a_{ij} \equiv g_i g_j \sin(\alpha_i - \alpha_j). \quad (2.8)$$

The condition for this extremum is that $\Delta > 0$, which implies that at least two of the SU(1,1) angles must be different. This in turn implies that the gauge group must contain at least two simple subgroups.

Let us now make a list of possible simple subgroups of G , and discuss their embedding in SO(6,6). The metric η_{RS} is the invariant metric of the global symmetry group SO(6,6), which acts on the fields in the fundamental representation. The gauge group G acts in the adjoint representation, and has 12 or less generators (for $n = 6$). So the adjoint representation of G must fit into the fundamental representation of SO(6,6). The generators T_R of the simple subgroups of G in the fundamental representation are chosen in such a way that $g_{RS} \equiv \text{tr} T_R T_S = \pm 2\delta_{RS}$, with positive entries for the compact, and negative entries for the noncompact directions. The embedding of G in SO(6,6) is such that the metric g_{RS} coincides, up to an overall factor ± 2 , with η_{RS} . The factor 2 we absorb in the coupling constant of the corresponding gauge group (see appendix B for further properties of these groups). With these restrictions one can easily list all allowed simple factors G_i of G . These are presented in table 1. Note that groups of dimension 10 and higher are excluded since they do not leave enough room for a second nonabelian subgroup.

The starting point for the remaining analysis is then (2.1). The ingredients are

$$V_{ij} = \left(-\frac{1}{12} Z^{RU} Z^{SV} Z^{TW} + \frac{1}{4} Z^{RU} Z^{SV} (\eta^{TW} + Z^{TW}) \right) f^{(i)}_{RST} f^{(j)}_{UVW}, \quad (2.9)$$

$$W_{ij} = \frac{1}{36} \epsilon^{abcdef} Z_a^R Z_b^S Z_c^T Z_d^U Z_e^V Z_f^W f^{(i)}_{RST} f^{(j)}_{UVW}, \quad (2.10)$$

where $Z^{RS} = Z_a^R Z_a^S$. It is important to remember that the Z_a^R are not the independent scalars, due to the constraint (1.2). A useful parametrisation is given in terms of 6×6 matrices G (symmetric) and B (antisymmetric). We split the indices R, S, \dots of η_{RS} in

$A, B, \dots = 1, \dots, 6$, ($\eta_{AB} = -\delta_{AB}$) and $I, J, \dots = 7, \dots, 12$, ($\eta_{IJ} = +\delta_{IJ}$). The scalar constraint (1.2) then reads

$$XX^T - YY^T = \mathbb{1}_6, \tag{2.11}$$

where $X_a^A = Z_a^A$, $Y_a^{I-6} = Z_a^I$, which is solved by

$$X = \frac{1}{2} (G + G^{-1} + BG^{-1} - G^{-1}B - BG^{-1}B), \tag{2.12}$$

$$Y = \frac{1}{2} (G - G^{-1} - BG^{-1} - G^{-1}B - BG^{-1}B). \tag{2.13}$$

In [8] we limited ourselves to the case where

$$G = \begin{pmatrix} a\mathbb{1}_3 & 0 \\ 0 & a\mathbb{1}_3 \end{pmatrix} \quad (a > 0), \quad B = \begin{pmatrix} 0 & b\mathbb{1}_3 \\ -b\mathbb{1}_3 & 0 \end{pmatrix}, \tag{2.14}$$

in this paper we will use the complete G and B .

We will analyse the potential at the point Z_0 , the origin of the scalar manifold, given by $G = \mathbb{1}$, $B = 0$. We will see in section 2.2 that for many gauge groups the origin corresponds to an extremum of the potential in all directions. At $Z = Z_0$ we have

$$Z_{0a}^R = (\mathbb{1}_6 \ 0), \quad Z_0^{RS} = \begin{pmatrix} \mathbb{1}_6 & 0 \\ 0 & 0 \end{pmatrix}, \quad (\eta + Z)_0^{RS} = \begin{pmatrix} 0 & 0 \\ 0 & \mathbb{1}_6 \end{pmatrix}. \tag{2.15}$$

Consider V_{ij} , as given in (2.9). The first term contains a product of three Z_0^{RS} , which are diagonal and only non-vanishing if all of $RSTUVW$ are in the range $1 \dots 6$. The second term containing $\eta + Z$ is also diagonal with the two indices in the range $7 \dots 12$. Therefore in V_{ij} the indices of the structure constants of G_i and G_j are contracted, implying that they belong to the same factor in the product of groups. Therefore $V_{ij} = 0$ for $i \neq j$.

W_{ij} can only be nonzero if there are subgroups G_i and G_j of G , $i \neq j$, such that both have three generators in the range $1 \dots 6$ with structure constants f_{ABC} . These G_i must therefore be $\text{SO}(3)$ or $\text{SU}(2)$ subgroups of G_i and G_j .

Therefore we have:

$$\begin{aligned} V_{0ii} &= -\frac{1}{12} f^{(i)}_{ABC} f^{(i)}_{ABC} + \frac{1}{4} f^{(i)}_{ABI} f^{(i)}_{ABI}, \\ &= -\frac{1}{2} \quad \text{for } \text{SO}(3)_-, \text{SO}(3,1)_-, \text{SL}(3, \mathbb{R})_-, \\ &\quad -2 \quad \text{for } \text{SU}(2,1)_-, \\ &\quad \frac{1}{2} \quad \text{for } \text{SO}(2,1)_+, \\ &\quad \frac{3}{2} \quad \text{for } \text{SO}(3,1)_+, \\ &\quad \frac{15}{2} \quad \text{for } \text{SL}(3, \mathbb{R})_+, \\ &\quad 6 \quad \text{for } \text{SU}(2,1)_+, \\ &\quad 0 \quad \text{for } \text{SO}(2,1)_-, \text{SO}(3)_+, \\ V_{0ij} &= 0 \quad i \neq j, \end{aligned}$$

$$\begin{aligned}
 W_{0ij} &= 1 && \text{for subgroups } \text{SO}(3)_-, \text{SO}(3,1)_-, \text{SL}(3, \mathbb{R})_-, \\
 &= 0 && \text{otherwise.}
 \end{aligned} \tag{2.16}$$

Since $V_{0ij} = 0$ for $i \neq j$ we can simplify (2.2). We find

$$C_{0-} = \sum_i g_i^2 V_{0ii}, \tag{2.17}$$

$$T_{0-} = \sum_{ij} a_{ij} W_{0ij}, \tag{2.18}$$

$$\Delta_0 = 2 \sum_{ij} V_{0ii} V_{0jj} a_{ij}^2, \tag{2.19}$$

$$R^{(0ii)} = \frac{2 \text{sgn} C_{0-}}{\sqrt{\Delta_0}} \sum_j V_{0jj} a_{ij}^2. \tag{2.20}$$

The off-diagonal R^{ij} can also be nonzero, but they do not appear in the potential (or its first and second derivatives, as we shall see) in Z_0 because V_{0ij} is diagonal.

Further restrictions come from the value of V_{0ii} . To have $\Delta > 0$, we must have at least two subgroups G_i for which $V_{ii} \neq 0$. Note for instance that the groups $\text{SO}(3)_+$ and $\text{SO}(2,1)_-$ have zero V_{0ii} , and do not contribute to Δ in Z_0 . More restrictions will come from the requirement that Z_0 corresponds to an extremum of the potential.

There are two ways to make V_0 positive. One is to have $C_- > 0$, to which groups with positive V_{0ii} , such as $\text{SO}(2,1)_+$ or $\text{SO}(3,1)_+$, contribute. A further positive contribution can come from T_- , if W_{ij} is nonzero and the $\text{SU}(1,1)$ angles are appropriately chosen. This can be done for groups with $\text{SO}(3)_-$ as a subgroup.

2.2 The first derivatives of the potential

The unconstrained, independent scalar fields in the 12 vector multiplets are 21 components of the symmetric matrix G , and 15 components of the antisymmetric matrix B . It is convenient to introduce a single matrix $P = (G + B)$ for the 36 independent scalars. In appendix C we give the derivatives of X and Y in Z_0 with respect to these parameters. The first derivatives of V and W with respect to P are:

$$\frac{\partial V_{ij}}{\partial P} = \frac{1}{2} \left(\frac{\partial Z_f^R}{\partial P} Z_f^U + Z_f^R \frac{\partial Z_f^U}{\partial P} \right) Z^{SV} (\eta + Z)^{TW} f^{(i)}{}_{RST} f^{(j)}{}_{UVW}, \tag{2.21}$$

$$\begin{aligned}
 \frac{\partial W_{ij}}{\partial P} &= \frac{1}{12} \epsilon^{abcdef} \left(\frac{\partial Z_a^R}{\partial P} Z_b^S Z_c^T Z_d^U Z_e^V Z_f^W + \right. \\
 &\quad \left. + Z_a^R Z_b^S Z_c^T \frac{\partial Z_d^U}{\partial P} Z_e^V Z_f^W \right) f^{(i)}{}_{RST} f^{(j)}{}_{UVW}.
 \end{aligned} \tag{2.22}$$

Now we evaluate this in Z_0 , using the fact that the derivative of X with respect to P vanishes in Z_0 . The derivatives of V and W in Z_0 are then:

$$\left. \frac{\partial V_{ii}}{\partial P} \right|_0 = \left. \frac{\partial Z_A^I}{\partial P} \right|_0 f^{(i)}{}_{IBJ} f^{(i)}{}_{ABJ}, \quad \left. \frac{\partial V_{ij}}{\partial P} \right|_0 = 0 \text{ for } i \neq j, \tag{2.23}$$

Groups	Properties
$\text{SO}(3)_-^2 \otimes \text{SO}(3)_+^2$	
$\text{SO}(3)_- \otimes \text{SO}(2,1)_+ \otimes \text{SO}(2,1)_- \otimes \text{SO}(3)_+$	$\Delta_0 < 0$, no extremum
$\text{SO}(3)_- \otimes \text{SO}(2,1)_-^3$	$\Delta_0 = 0$
$\text{SO}(2,1)_+^3 \otimes \text{SO}(3)_+$	
$\text{SO}(2,1)_+^2 \otimes \text{SO}(2,1)_-^2$	
$\text{SO}(3,1)_+ \otimes \text{SO}(2,1)_+ \otimes \text{SO}(2,1)_-$	
$\text{SO}(3,1)_+ \otimes \text{SO}(3,1)_+$	
$\text{SO}(3,1)_+ \otimes \text{SO}(3)_- \otimes \text{SO}(3)_+$	$\Delta_0 < 0$, no extremum
$\text{SO}(3,1)_- \otimes \text{SO}(2,1)_+ \otimes \text{SO}(2,1)_-$	$\Delta_0 < 0$, no extremum
$\text{SO}(3,1)_- \otimes \text{SO}(3,1)_-$	
$\text{SO}(3,1)_- \otimes \text{SO}(3,1)_+$	$\Delta_0 < 0$, no extremum
$\text{SO}(3,1)_- \otimes \text{SO}(3)_- \otimes \text{SO}(3)_+$	
$\text{SL}(3, \mathbb{R})_+ \otimes \text{SO}(3)_+$	$\Delta_0 = 0$
$\text{SL}(3, \mathbb{R})_- \otimes \text{SO}(3)_-$	
$\text{SL}(3, \mathbb{R})_+ \otimes \text{SO}(2,1)_-$	$\Delta_0 = 0$
$\text{SL}(3, \mathbb{R})_- \otimes \text{SO}(2,1)_+$	$\Delta_0 < 0$, no extremum
$\text{SU}(2,1)_+ \otimes \text{SO}(2,1)_+$	
$\text{SU}(2,1)_+ \otimes \text{SO}(2,1)_-$	$\Delta_0 = 0$
$\text{SU}(2,1)_- \otimes \text{SO}(2,1)_+$	$\Delta_0 < 0$, no extremum
$\text{SU}(2,1)_- \otimes \text{SO}(2,1)_-$	$\Delta_0 = 0$

Table 2: List of possible gauge groups G . Nine groups have an extremum with respect to the matter scalars in Z_0 with positive Δ_0 .

$$\left. \frac{\partial W_{ij}}{\partial P} \right|_0 = \frac{1}{12} \epsilon^{ABCDEF} \left. \frac{\partial Z_A^I}{\partial P} \right|_0 f^{(i)}{}_{IBC} f^{(j)}{}_{DEF} - (i \leftrightarrow j). \quad (2.24)$$

In the derivatives of V_{ij} there are always contractions between the different groups, implying that the derivatives vanish for $i \neq j$. For the derivative of V_{ii} to be nonzero we see that G_i must have two indices AB and two indices IJ . This is not the case for the groups in table 1.

The derivative of W_{ij} is nonzero only if one of the groups G_j or G_i has an $\text{SO}(3)$ subgroup on the indices DEF , and the other has an $\text{SO}(2,1)$ subgroup for which the IBC structure constants are nonzero. Groups G with an $\text{SO}(3)_- \otimes \text{SO}(2,1)_+$ subgroup are therefore excluded.

If we require that Z_0 is an extremum of the potential, and that $\Delta > 0$ in the extremum is possible for a suitable choice of the parameters, the number of allowed groups becomes sufficiently small to make a complete analysis possible. The list of allowed groups is given in table 2.

To illustrate table 2, let's consider the group $\text{SO}(3,1)$. The commutation relations for the $\text{SO}(3,1)$ Lie algebra can be found in appendix B. The generators of the rotation subgroup are denoted by T , the boosts by K . To embed the adjoint of $\text{SO}(3,1)$ in $\text{SO}(6,6)$ we have two choices: either the three T correspond to the negative, and the boosts K to

the positive entries of η_{RS} ($\text{SO}(3,1)_-$), or the other way round ($\text{SO}(3,1)_+$). If we choose $\text{SO}(3)_- \otimes \text{SO}(3,1)_-$, the nonzero contributions to V_{0ii} (2.16) come from $\text{SO}(3)_-$ and from the rotation subgroup of $\text{SO}(3,1)_-$, both contribute $-\frac{1}{2}$, and Δ_0 (2.19) is positive. The structure constants f_{ABI} vanish, and do not contribute to V_0 in this case. For the same reason, the first derivatives of V and W (2.23), (2.24) vanish. So this case is interesting, and will appear as a subgroup of the groups considered in section 3.6 and 3.7. On the other hand, if we choose $\text{SO}(3)_- \otimes \text{SO}(3,1)_+$ then the boosts are in the ABC range of the indices, and structure constants f_{ABI} are nonzero. The potential V_{0ii} gets contributions from $\text{SO}(3)_-$, and now the second term in (2.16) will contribute positively. This will make Δ_0 negative. Also the first derivative of W will be nonzero, so this group cannot be used for our purposes.

2.3 The second derivatives of the potential

The full potential is given in (2.1), and its second derivatives are:

$$\frac{\partial^2 V}{\partial \phi^2} = \sum_{ij} \frac{\partial^2 R^{(ij)}}{\partial \phi^2} V_{ij}, \quad (2.25)$$

$$\frac{\partial^2 V}{\partial \phi \partial P} = \sum_{ij} \frac{\partial R^{(ij)}}{\partial \phi} \frac{\partial V_{ij}}{\partial P}, \quad (2.26)$$

$$\frac{\partial^2 V}{\partial P^2} = \sum_{ij} R^{(ij)} \frac{\partial^2 V_{ij}}{\partial P^2} + I^{(ij)} \frac{\partial^2 W_{ij}}{\partial P^2}. \quad (2.27)$$

The second derivatives (2.25) were studied in [8]. The sign of (2.25) depends on the sign of C_- . For positive (negative) C_- the extremum in the $\text{SU}(1,1)$ scalars is a minimum (maximum). The mixed second derivatives vanish if either the derivatives with respect to the $\text{SU}(1,1)$ scalars ϕ or with respect to the matter scalars vanishes. In this section we will evaluate the second derivatives (2.27) with respect to the matter scalars in Z_0 .

We therefore calculate the second derivatives of V_{ij} and W_{ij} with respect to the independent scalars P . They are:

$$\begin{aligned} \frac{\partial^2 V_{ij}}{\partial P_1 \partial P_2} = & \frac{1}{2} \left\{ \left(\frac{\partial^2 Z_f^R}{\partial P_1 \partial P_2} Z_f^U + Z_f^R \frac{\partial^2 Z_f^U}{\partial P_1 \partial P_2} \right) Z^{SV} (\eta + Z)^{TW} + \right. \\ & + \left(\frac{\partial Z_f^R}{\partial P_1} \frac{\partial Z_f^U}{\partial P_2} + \frac{\partial Z_f^R}{\partial P_2} \frac{\partial Z_f^U}{\partial P_1} \right) Z^{SV} (\eta + Z)^{TW} + \\ & + \left(\frac{\partial Z_f^R}{\partial P_1} Z_f^U + Z_f^R \frac{\partial Z_f^U}{\partial P_1} \right) \left(\frac{\partial Z_g^S}{\partial P_2} Z_g^V + Z_g^S \frac{\partial Z_g^V}{\partial P_2} \right) (\eta + Z)^{TW} + \\ & + \left. \left(\frac{\partial Z_f^R}{\partial P_1} Z_f^U + Z_f^R \frac{\partial Z_f^U}{\partial P_1} \right) Z^{SV} \left(\frac{\partial Z_g^T}{\partial P_2} Z_g^W + Z_g^T \frac{\partial Z_g^W}{\partial P_2} \right) \right\} \times \\ & \times f^{(i)}{}_{RST} f^{(j)}{}_{UVW}, \quad (2.28) \\ \frac{\partial^2 W_{ij}}{\partial P_1 \partial P_2} = & \frac{1}{12} \epsilon^{a_1 \dots a_6} \left\{ \frac{\partial^2 Z_{a_1}^R}{\partial P_1 \partial P_2} Z_{a_2}^S Z_{a_3}^T Z_{a_4}^U Z_{a_5}^V Z_{a_6}^W + \right. \\ & + 2 \frac{\partial Z_{a_1}^R}{\partial P_1} \frac{\partial Z_{a_2}^S}{\partial P_2} Z_{a_3}^T Z_{a_4}^U Z_{a_5}^V Z_{a_6}^W + \end{aligned}$$

$$\begin{aligned}
 & + 3 \frac{\partial Z_{a_1}^R}{\partial P_1} Z_{a_2}^S Z_{a_3}^T \frac{\partial Z_{a_4}^U}{\partial P_2} Z_{a_5}^V Z_{a_6}^W - \\
 & - (RST \leftrightarrow UVW) \left. \right\} f^{(i)}{}_{RST} f^{(j)}{}_{UVW}. \tag{2.29}
 \end{aligned}$$

We now evaluate both expressions for $Z = Z_0$. From (2.28) it is clear that in $Z = Z_0$ we will get contractions between the two structure constants, so that they must belong to the same subgroup. Therefore only the second derivatives of V_{ii} are nonzero in Z_0 . We find:

$$\begin{aligned}
 \frac{\partial^2 V_{ii}}{\partial P_1 \partial P_2} \Big|_0 &= \left\{ \frac{\partial^2 Z_A^R}{\partial P_1 \partial P_2} \Big|_0 f^{(i)}{}_{RBJ} f^{(i)}{}_{ABJ} + \frac{\partial Z_f^I}{\partial P_1} \Big|_0 \frac{\partial Z_f^J}{\partial P_2} \Big|_0 f^{(i)}{}_{IBK} f^{(i)}{}_{JBK} + \right. \\
 & + \left(\frac{\partial Z_A^I}{\partial P_1} \Big|_0 \frac{\partial Z_B^J}{\partial P_2} \Big|_0 f^{(i)}{}_{IJK} f^{(i)}{}_{ABK} + \frac{\partial Z_A^I}{\partial P_1} \Big|_0 \frac{\partial Z_B^J}{\partial P_2} \Big|_0 f^{(i)}{}_{AJK} f^{(i)}{}_{IBK} \right) \\
 & + \left. \left(\frac{\partial Z_A^I}{\partial P_1} \Big|_0 \frac{\partial Z_B^J}{\partial P_2} \Big|_0 f^{(i)}{}_{ICJ} f^{(i)}{}_{ACB} + \frac{\partial Z_A^I}{\partial P_1} \Big|_0 \frac{\partial Z_B^J}{\partial P_2} \Big|_0 f^{(i)}{}_{ICB} f^{(i)}{}_{ACJ} \right\}, \\
 \frac{\partial^2 W_{ij}}{\partial P_1 \partial P_2} \Big|_0 &= \frac{1}{12} \epsilon^{ABCDEF} \left(\frac{\partial^2 Z_A^R}{\partial P_1 \partial P_2} \Big|_0 f^{(i)}{}_{RBC} f^{(j)}{}_{DEF} + 2 \frac{\partial Z_A^I}{\partial P_1} \Big|_0 \frac{\partial Z_B^J}{\partial P_2} \Big|_0 f^{(i)}{}_{IJC} f^{(j)}{}_{DEF} + \right. \\
 & \left. + 3 \frac{\partial Z_A^I}{\partial P_1} \Big|_0 \frac{\partial Z_D^J}{\partial P_2} \Big|_0 f^{(i)}{}_{IBC} f^{(j)}{}_{JEF} - (i \leftrightarrow j) \right). \tag{2.30}
 \end{aligned}$$

The second derivatives of Z (X and Y) are given in appendix C.

2.4 Masses of scalar and vector fields

The mass-matrix should be normalised in relation to the kinetic terms. The kinetic term of the matter scalar fields Z is independent of the gauging - the gauge fields occur only in the covariantisations. The kinetic term of the Z_a^R reads (ignoring the gauge field contributions)

$$-\frac{1}{2} \eta_{RS} \partial_\mu Z_a^R \partial^\mu Z_a^S - \frac{1}{8} \eta_{RS} \eta_{TU} Z_a^R \overleftrightarrow{\partial}_\mu Z_b^S Z_a^T \overleftrightarrow{\partial}^\mu Z_b^U. \tag{2.31}$$

This should be evaluated in $Z = Z_0$ and expressed in terms of P_{ab} , giving

$$-\frac{1}{2} \partial_\mu P_{ab} \partial^\mu P_{ab}, \tag{2.32}$$

the standard normalization for scalar fields. Together with the contribution from the potential we therefore get

$$-\frac{1}{2} \partial_\mu P_{ab} \partial^\mu P_{ab} - V_0 - \frac{1}{2} (P_{ab} - \delta_{ab})(P_{cd} - \delta_{cd}) \frac{\partial^2 V}{\partial P_{ab} \partial P_{cd}} \Big|_0. \tag{2.33}$$

On shifting the scalar fields in the kinetic term by the constant δ_{ab} we see that the matrix of second derivatives we have calculated in the previous section is precisely the mass-matrix.

The mass-matrix for the $SU(1,1)$ scalars was given is given in A.10 in appendix A. The kinetic and mass terms for these scalars are:

$$-\frac{1}{2} (\partial_\mu x' \partial^\mu x' + \partial_\mu y' \partial^\mu y') - \frac{1}{2} \text{sgn} C_- \sqrt{\Delta_0} (x'^2 + y'^2). \tag{2.34}$$

In the case $C_- < 0$ we have two tachyons. The relation between $\phi_{1,2}$ and x and y is explained in appendix A.

The vector masses follow from the coupling of the vectors to the scalars Z . We have in the covariant derivative:

$$\mathcal{D}_\mu Z_a^R = \partial_\mu Z_a^R - V_{\mu ab} Z_b^R - A_\mu^S g f_{ST}^R Z_a^T. \quad (2.35)$$

After elimination of V the scalar kinetic term becomes:

$$-\frac{1}{2}\eta_{RS}D_\mu Z_a^R D^\mu Z_a^S - \frac{1}{2}\eta_{RS}\eta_{TU}Z_a^R Z_a^T D_\mu Z_b^S D^\mu Z_b^U, \quad (2.36)$$

where D contains the gauge field A only. Now substitute $Z = Z_0$ and isolate the A^2 terms. The result is, after writing out the indices R, S in terms of A, B and I, J :

$$-\frac{1}{2}g_i^2 A_\mu^A A^{\mu B} f^{(i)}_{ACI} f^{(i)}_{BCI} - g_i^2 A_\mu^A A^{\mu K} f^{(i)}_{ACI} f^{(i)}_{KCI} - \frac{1}{2}g_i^2 A_\mu^K A^{\mu L} f^{(i)}_{KCI} f^{(i)}_{LCI}. \quad (2.37)$$

We see that there are vector masses only for noncompact groups. That is to be expected, since these are the noncompact generators do not leave Z_0 invariant. The second term vanishes for all groups considered. The first term get contributions from gauge groups with an $\text{SO}(2,1)_+$ subgroups, the last one from $\text{SO}(2,1)_-$ subgroups. The masses are proportional to the corresponding g_i^2 , and independent of the $\text{SU}(1,1)$ angles.

3. Semi-simple gauge groups

Table 2 left us with nine allowed groups. In this section we will discuss these nine cases separately. In table 3 we give a list of the allowed groups, with their basic properties. The sign of C_- is important, because it determines the behaviour of the $\text{SU}(1,1)$ scalars: if $C_- < 0$, the $\text{SU}(1,1)$ scalars are at a maximum, if $C_- > 0$ at a minimum. Clearly a stable minimum in all 38 scalar directions requires $C_- > 0$. We now discuss the nine groups in the order of table 3.

Groups	C_-	V_0
$\text{SO}(2,1)_+^3 \otimes \text{SO}(3)_+$	$\frac{1}{2}(g_1^2 + g_2^2 + g_3^2)$	$\sqrt{a_{12}^2 + a_{13}^2 + a_{23}^2}$
$\text{SO}(2,1)_+^2 \otimes \text{SO}(2,1)_-^2$	$\frac{1}{2}(g_1^2 + g_2^2)$	$ a_{12} $
$\text{SO}(3,1)_+ \otimes \text{SO}(2,1)_+ \otimes \text{SO}(2,1)_-$	$\frac{1}{2}(3g_1^2 + g_2^2)$	$\sqrt{3} a_{12} $
$\text{SO}(3,1)_+ \otimes \text{SO}(3,1)_+$	$\frac{3}{2}(g_1^2 + g_2^2)$	$3 a_{12} $
$\text{SO}(3)_-^2 \otimes \text{SO}(3)_+^2$	$-\frac{1}{2}(g_1^2 + g_1^2)$	$- a_{12} - 2a_{12}$
$\text{SO}(3)_- \otimes \text{SO}(3,1)_- \otimes \text{SO}(3)_+$	$-\frac{1}{2}(g_1^2 + g_1^2)$	$- a_{12} - 2a_{12}$
$\text{SO}(3,1)_- \otimes \text{SO}(3,1)_-$	$-\frac{1}{2}(g_1^2 + g_2^2)$	$- a_{12} - 2a_{12}$
$\text{SL}(3, \mathbb{R})_- \otimes \text{SO}(3)_-$	$-\frac{1}{2}(g_1^2 + g_2^2)$	$- a_{12} - 2a_{12}$
$\text{SU}(2,1)_+ \otimes \text{SO}(2,1)_+$	$6g_1^2 + \frac{1}{2}g_2^2$	$2\sqrt{3} a_{12} $

Table 3: List of possible gauge groups G . V_0 is the value of the potential in $Z = Z_0$. $C_- > 0$ (< 0) implies that the $\text{SU}(1,1)$ scalars are at a minimum (maximum).

In the following sections 3.1–3.9 we will discuss the nine gauge groups which have $\Delta_0 > 0$ and an extremum for the matter scalars in the origin of the scalar manifold. We present for each case the mass-matrix for the matter scalars. The masses of the $SU(1, 1)$ scalars are always given by $\text{sgn}C_- \sqrt{\Delta_0}$, indicating that they are always both tachyonic or both positive.

For the first case, the group $SO(2, 1)_+^3 \otimes SO(3)_+$ presented in section 3.1, we will give full details of the analysis. For the other cases the procedure should then be clear, and we will limit ourselves to the presentation of the results.

3.1 $SO(2, 1)_+^3 \otimes SO(3)_+$

In this case the groups have to be arranged as follows:

$$R, S, \dots = \overbrace{1\ 2\ 7}^{i=1} \overbrace{3\ 4\ 8}^{i=2} \overbrace{5\ 6\ 9}^{i=3} \overbrace{10\ 11\ 12}^{i=4}. \quad (3.1)$$

Then all $V_{0ii} = 1/2$ for $i = 1, 2, 3$, while $V_{044} = 0$, and we have:

$$C_{0-} = \frac{1}{2}(g_1^2 + g_2^2 + g_3^2), \quad (3.2)$$

$$T_{0-} = 0, \quad (3.3)$$

$$\Delta_0 = a_{12}^2 + a_{13}^2 + a_{23}^2, \quad (3.4)$$

$$R^{(0ii)} = \frac{1}{\sqrt{\Delta_0}} (a_{i1}^2 + a_{i2}^2 + a_{i3}^2). \quad (3.5)$$

giving

$$V_0 = \sqrt{\Delta_0}, \quad (3.6)$$

$$\left. \frac{\partial V}{\partial P_{ab}} \right|_0 = 0, \quad (3.7)$$

$$\left. \frac{\partial^2 V}{\partial P_{ab} \partial P_{cd}} \right|_0 = \frac{1}{\sqrt{\Delta_0}} \sum_{i=1}^3 (a_{i1}^2 + a_{i2}^2 + a_{i3}^2) \left. \frac{\partial^2 V_{ii}}{\partial P_{ab} \partial P_{cd}} \right|_0 - \sum_{i < j} 2a_{ij} \left. \frac{\partial^2 W_{ij}}{\partial P_{ab} \partial P_{cd}} \right|_0. \quad (3.8)$$

Notice that the group $SO(3)_+$ does not contribute to the second derivatives, structure constants with indices JK appear in (2.30) in the third term, but in combination with a structure constant f_{ABK} , which $SO(3)_+$ does not have. The non-vanishing second derivative matrices required for calculating the mass-matrix are:

$$\begin{aligned} \left. \frac{\partial^2 V_{11}}{\partial P_{ab} \partial P_{cd}} \right|_0 &= (\delta_{ac} - \delta_{a1} \delta_{c1})(\delta_{b1} \delta_{d1} + \delta_{b2} \delta_{d2}), \\ \left. \frac{\partial^2 V_{22}}{\partial P_{ab} \partial P_{cd}} \right|_0 &= (\delta_{ac} - \delta_{a2} \delta_{c2})(\delta_{b3} \delta_{d3} + \delta_{b4} \delta_{d4}), \\ \left. \frac{\partial^2 V_{33}}{\partial P_{ab} \partial P_{cd}} \right|_0 &= (\delta_{ac} - \delta_{a3} \delta_{c3})(\delta_{b5} \delta_{d5} + \delta_{b6} \delta_{d6}), \\ \left. \frac{\partial^2 W_{12}}{\partial P_{ab} \partial P_{cd}} \right|_0 &= (\delta_{a1} \delta_{c2} - \delta_{a2} \delta_{c1})(\delta_{b5} \delta_{d6} - \delta_{b6} \delta_{d5}), \end{aligned}$$

$$\begin{aligned}
 \left. \frac{\partial^2 W_{13}}{\partial P_{ab} \partial P_{cd}} \right|_0 &= (\delta_{a1} \delta_{c3} - \delta_{a3} \delta_{c1})(\delta_{b3} \delta_{d4} - \delta_{b4} \delta_{d3}), \\
 \left. \frac{\partial^2 W_{23}}{\partial P_{ab} \partial P_{cd}} \right|_0 &= (\delta_{a2} \delta_{c3} - \delta_{a3} \delta_{c2})(\delta_{b1} \delta_{d2} - \delta_{b2} \delta_{d1}).
 \end{aligned}
 \tag{3.9}$$

The resulting eigenvalues of $\left. \frac{\partial^2 V}{\partial P_{ab} \partial P_{cd}} \right|_0$ are then

$$\begin{aligned}
 0 \ (6 \times), \quad & \frac{1}{\sqrt{\Delta_0}}(a_{12}^2 + a_{13}^2) \ (6 \times), \quad \frac{1}{\sqrt{\Delta_0}}(a_{13}^2 + a_{23}^2) \ (6 \times), \quad \frac{1}{\sqrt{\Delta_0}}(a_{12}^2 + a_{23}^2) \ (6 \times), \\
 & \frac{1}{\sqrt{\Delta_0}}(a_{12}^2 + a_{13}^2) \pm 2a_{23} \ (2 \times), \quad \frac{1}{\sqrt{\Delta_0}}(a_{13}^2 + a_{23}^2) \pm 2a_{12} \ (2 \times), \\
 & \frac{1}{\sqrt{\Delta_0}}(a_{12}^2 + a_{23}^2) \pm 2a_{13} \ (2 \times).
 \end{aligned}
 \tag{3.10}$$

One can obtain these eigenvalues as follows. The second derivatives of V_{ii} are diagonal in the parameters P , in the sense that always $a = c$, $b = d$. The list of elements of P giving a nonzero second derivative of V_{ii} is:²

$$\begin{aligned}
 V_{11} : & \frac{1}{\sqrt{\Delta_0}}(a_{12}^2 + a_{13}^2) [21, 22, 31, 32, 41, 42, 51, 52, 61, 62], \\
 V_{22} : & \frac{1}{\sqrt{\Delta_0}}(a_{12}^2 + a_{23}^2) [13, 14, 33, 34, 43, 44, 53, 54, 63, 64], \\
 V_{33} : & \frac{1}{\sqrt{\Delta_0}}(a_{13}^2 + a_{23}^2) [15, 16, 25, 26, 45, 46, 55, 56, 65, 66].
 \end{aligned}$$

On the other hand, W_{ij} has a non-vanishing second derivative for the following pairs of elements of P :

$$\begin{aligned}
 W_{12} : & -2a_{12}[(15, 26), (16, 25), (25, 16), (26, 15)], \\
 W_{13} : & -2a_{13}[(13, 34), (14, 33), (33, 14), (34, 13)], \\
 W_{23} : & -2a_{23}[(21, 32), (22, 31), (31, 22), (32, 21)].
 \end{aligned}$$

The six eigenvalues $(a_{13}^2 + a_{23}^2)/\sqrt{\Delta_0}$ are associated with the six diagonal elements of the second derivative matrix coming from V_{11} : 41, 42, 51, 52, 61, 62 are the corresponding elements of P . The six eigenvalues $(a_{12}^2 + a_{23}^2)/\sqrt{\Delta_0}$ and $(a_{12}^2 + a_{13}^2)/\sqrt{\Delta_0}$ arise in a similar way from the derivatives of V_{22} and V_{33} , respectively. The remaining diagonal contributions from V_{ii} combine with the corresponding derivatives of W_{ij} : V_{11} with W_{23} , etc. These elements of P give rise to 2×2 submatrices in the matrix of second derivatives, with the eigenvalues as indicated. The zero eigenvalues of the second derivative matrix are associated with the elements 11, 12, 23, 24, 35, 36 of P which do not occur anywhere in the second derivatives.

The six zero eigenvalues indicate that the solution $Z = Z_0$ breaks the gauge symmetry. In each of the $SO(2, 1)_+$ groups the $SO(2, 1)$ symmetry is broken to $U(1)$. The six massless scalars give masses to the gauge vectors, as we have seen in section 2.4.

²We use the notation ab to indicate the second derivative with respect to P_{ab} (diagonal elements of the 36×36 second derivative matrix), and (ab, cd) for the second derivative with respect to P_{ab} and P_{cd} (off-diagonal elements of the second derivative matrix).

The extremum we have obtained in this case is not stable. Assume that all 36 eigenvalues are nonnegative. If $(a_{12}^2 + a_{13}^2)/\sqrt{\Delta_0} \pm 2a_{23}$ has to be positive then necessarily

$$|a_{23}| < \sqrt{\Delta_0}(\sqrt{2} - 1). \tag{3.11}$$

This has to be true then for a_{13} and a_{23} as well, implying

$$\Delta_0 < 3\Delta_0(\sqrt{2} - 1)^2 < \Delta_0. \tag{3.12}$$

Therefore our assumption that all eigenvalues are positive must be false. In fact, there are two, four or six negative eigenvalues. To have six negative eigenvalues choose $a_{12} = a_{23} = a_{13} = \sqrt{\Delta_0/3}$.

3.2 $SO(2, 1)_+^2 \otimes SO(2, 1)_-^2$

The embedding of the subgroups is as follows:

$$R, S, \dots = \overbrace{1\ 2\ 7}^{i=1} \overbrace{3\ 4\ 8}^{i=2} \overbrace{5\ 9\ 10}^{i=3} \overbrace{6\ 11\ 12}^{i=4}. \tag{3.13}$$

Although the V_{0ii} vanish for each $SO(2, 1)_-$ factor, the second derivatives don't. We now have:

$$C_{0-} = \frac{1}{2}(g_1^2 + g_2^2), \tag{3.14}$$

$$T_{0-} = 0, \tag{3.14}$$

$$\Delta_0 = a_{12}^2, \tag{3.15}$$

$$R^{(0ii)} = \frac{1}{|a_{12}|} (a_{i1}^2 + a_{i2}^2), \tag{3.16}$$

$$V_0 = |a_{12}|. \tag{3.17}$$

Now we find the following eigenvalues:

$$0\ (8 \times), \quad |a_{12}|\ (4 \times), \quad |a_{12}| + R^{(033)}\ (8 \times), \quad |a_{12}| + R^{(044)}\ (8 \times), \\ R^{(033)}\ (2 \times), \quad R^{(044)}\ (2 \times), \quad 2a_{12}\ (2 \times), \quad -2a_{12}\ (2 \times). \tag{3.18}$$

$Z = Z_0$ breaks the gauge symmetry to $U(1)^4$, so the eight zero eigenvalues correspond to the Goldstone bosons that produce the masses of the gauge fields.

There are two negative eigenvalues, proportional to a_{12} . It does not help to set $\alpha_1 = \alpha_2$ to eliminate them, since this would make $\Delta = 0$ and invalidate the analysis of the $SU(1, 1)$ scalars.

3.3 $SO(3, 1)_+ \otimes SO(2, 1)_+ \otimes SO(2, 1)_-$

In this case the groups are arranged as follows:

$$R, S, \dots = \overbrace{1\ 2\ 3\ 10\ 11\ 12}^{i=1} \overbrace{4\ 5\ 7}^{i=2} \overbrace{6\ 8\ 9}^{i=3}. \tag{3.19}$$

The rotation subgroup of the $SO(3,1)$ subgroup is embedded on the indices $10 \dots 12$, the boosts on the indices $1 \dots 3$. Here we have:

$$C_{0-} = \frac{1}{2}(3g_1^2 + g_2^2), \quad (3.20)$$

$$T_{0-} = 0, \quad (3.21)$$

$$\Delta_0 = 3a_{12}^2, \quad (3.22)$$

$$R^{(0ii)} = \frac{1}{\sqrt{3}|a_{12}|} (3a_{i1}^2 + a_{i2}^2), \quad (3.23)$$

$$V_0 = \sqrt{3}|a_{12}|. \quad (3.24)$$

The complete list of eigenvalues in this case is:

$$\begin{aligned} & 0 \ (7 \times), \quad \frac{2|a_{12}|}{\sqrt{3}} \ (5 \times), \quad \sqrt{3}|a_{12}| \ (6 \times), \\ & \frac{|a_{12}|(1 + \sqrt{13})}{\sqrt{3}} \ (3 \times), \quad \frac{|a_{12}|(1 - \sqrt{13})}{\sqrt{3}} \ (3 \times), \\ & \frac{|a_{12}|(1 + \sqrt{37})}{\sqrt{3}} \ (1 \times), \quad \frac{|a_{12}|(1 - \sqrt{37})}{\sqrt{3}} \ (1 \times), \\ & \frac{1}{\sqrt{3}|a_{12}|} (3a_{12}^2 + 3a_{13}^2 + a_{23}^2) \ (4 \times), \quad \frac{1}{\sqrt{3}|a_{12}|} (2a_{12}^2 + 3a_{13}^2 + a_{23}^2) \ (6 \times). \end{aligned} \quad (3.25)$$

Altogether then we find that this group gives rise to four negative eigenvalues of the mass-matrix. The zero eigenvalues can again correspond to the Goldstone bosons of the broken noncompact gauge symmetries. The negative eigenvalues are proportional to $|a_{12}|$, which however we cannot set to zero because then also $\Delta_0 = 0$.

3.4 $SO(3,1)_+ \otimes SO(3,1)_+$

In this case the groups have to be arranged as follows:

$$R, S, \dots = \overbrace{1 \ 2 \ 3 \ 10 \ 11 \ 12}^{i=1} \overbrace{4 \ 5 \ 6 \ 7 \ 8 \ 9}^{i=2}. \quad (3.26)$$

In this case the rotation subgroup of the two $SO(3,1)$ subgroups is embedded on the indices $7 \dots 12$, the boosts on the indices $1 \dots 6$. Here V_{011} and V_{022} each are $\frac{3}{2}$, $W_{12} = 0$. and we have:

$$C_{0-} = \frac{3}{2}(g_1^2 + g_2^2), \quad (3.27)$$

$$T_{0-} = 0, \quad (3.28)$$

$$\Delta_0 = 9a_{12}^2, \quad (3.29)$$

$$R^{(0ii)} = \frac{1}{|a_{12}|} (a_{i1}^2 + a_{i2}^2), \quad (3.30)$$

$$V_0 = 3|a_{12}|. \quad (3.31)$$

The eigenvalues are:

$$0 \ (15 \times), \quad 2|a_{12}| \ (10 \times), \quad 4|a_{12}| \ (9 \times), \quad 8|a_{12}| \ (1 \times), \quad -4|a_{12}| \ (1 \times). \quad (3.32)$$

In this case there is a single negative eigenvalue. There are now more zero eigenvalues than the number required by the breaking of gauge invariance.

3.5 $SO(3)_-^2 \otimes SO(3)_+^2$

In the case of four $SO(3)$ groups these are arranged over the index values $R, S = 1, \dots, 12$ as follows:

$$R, S, \dots = \overbrace{1\ 2\ 3}^{i=1} \overbrace{4\ 5\ 6}^{i=2} \overbrace{7\ 8\ 9}^{i=3} \overbrace{10\ 11\ 12}^{i=4}. \quad (3.33)$$

This simplifies considerably the expressions derived in the section 2 for the potential, and its first and second derivatives in the point Z_0 . The only terms which contribute are those for which the indices on the structure constants are either ABC (in the range $1 \dots 6$) or IJK (in the range $7 \dots 12$). We find

$$C_{0-} = -\frac{1}{2}(g_1^2 + g_2^2), \quad (3.34)$$

$$T_{0-} = 2a_{12}, \quad (3.35)$$

$$\Delta_0 = a_{12}^2, \quad (3.36)$$

$$R^{(0ii)} = \frac{1}{|a_{12}|} \sum_i (a_{i1}^2 + a_{i2}^2), \quad (3.37)$$

$$V_0 = -|a_{12}| - 2a_{12}. \quad (3.38)$$

The eigenvalues of the mass-matrix are

$$-2a_{12} \ (36 \times). \quad (3.39)$$

If the $SU(1,1)$ angles are chosen such that V_0 is positive (de Sitter) then the eigenvalues of the mass-matrix are also positive for all 36 scalars.

In the present case $C_- < 0$ in Z_0 , which implies that for the $SU(1,1)$ scalars we have a maximum. So in this example there are two tachyons in the $SU(1,1)$ sector. In [8] we showed in the truncated model with two scalars that V_{11} and V_{22} , and therefore C_- , change sign on a circle around Z_0 . This turns the maximum for the $SU(1,1)$ scalars into a minimum. It will be interesting to see if this phenomenon also holds when all 36 matter scalars are taken into account.

3.6 $SO(3,1)_- \otimes SO(3)_- \otimes SO(3)_+$

In this case the groups have to be arranged as follows:

$$R, S, \dots = \overbrace{1\ 2\ 3\ 10\ 11\ 12}^{i=1} \overbrace{4\ 5\ 6}^{i=2} \overbrace{7\ 8\ 9}^{i=3}. \quad (3.40)$$

In this case the rotation subgroup of the $SO(3,1)$ subgroup is embedded on the indices $1 \dots 3$, the boosts on the indices $10 \dots 12$. Here we have:

$$C_{0-} = -\frac{1}{2}(g_1^2 + g_2^2), \quad (3.41)$$

$$T_{0-} = 2a_{12}, \quad (3.42)$$

$$\Delta_0 = a_{12}^2, \quad (3.43)$$

$$R^{(0ii)} = \frac{1}{|a_{12}|} (a_{i1}^2 + a_{i2}^2), \quad (3.44)$$

$$V_0 = -|a_{12}| - 2a_{12}. \quad (3.45)$$

We have the following eigenvalues:

$$\begin{aligned} 0 \ (3 \times), \quad -2a_{12} \ (18 \times), \quad 2|a_{12}| - 2a_{12} \ (9 \times), \\ 2|a_{12}| - 4a_{12} \ (5 \times), \quad 2|a_{12}| + 2a_{12} \ (1 \times). \end{aligned} \quad (3.46)$$

For $a_{12} < 0$ (de Sitter) there are no negative eigenvalues, and four zero eigenvalues. However, since $\text{sgn}C_- < 0$, the $SU(1,1)$ scalars produce the instability.

3.7 $SO(3,1)_- \otimes SO(3,1)_-$

In this case the groups have to be arranged as follows:

$$R, S, \dots = \overbrace{1 \ 2 \ 3 \ 10 \ 11 \ 12}^{i=1} \overbrace{4 \ 5 \ 6 \ 7 \ 8 \ 9}^{i=2}. \quad (3.47)$$

In this case the rotation subgroup of the two $SO(3,1)$ subgroups is embedded on the indices $1 \dots 6$, the boosts on the indices $7 \dots 12$. Here V_{011} and V_{022} each are $-\frac{1}{2}$, $W_{12} = 1$, and we have:

$$C_{0-} = -\frac{1}{2}(g_1^2 + g_2^2), \quad (3.48)$$

$$T_{0-} = 2a_{12}, \quad (3.49)$$

$$\Delta_0 = a_{12}^2, \quad (3.50)$$

$$R^{(0ii)} = \frac{1}{|a_{12}|} (a_{i1}^2 + a_{i2}^2), \quad (3.51)$$

$$V_0 = -|a_{12}| - 2a_{12}. \quad (3.52)$$

To make V_0 positive we have to choose $a_{12} < 0$. The eigenvalues of the mass-matrix are:

$$\begin{aligned} 0 \ (6 \times), \quad 2|a_{12}| - 2a_{12} \ (18 \times), \\ 2|a_{12}| - 4a_{12} \ (10 \times), \quad 2|a_{12}| + 2a_{12} \ (2 \times). \end{aligned} \quad (3.53)$$

For $a_{12} < 0$ there are no negative eigenvalues, and eight zero eigenvalues. However, since $C_- < 0$, the $SU(1,1)$ scalars produce the instability.

3.8 $SL(3, \mathbb{R})_- \otimes SO(3)_-$

In this case the groups are arranged as follows:

$$\overbrace{1 \ 2 \ 3 \ 4 \ 5 \ 7 \ 8 \ 9}^{i=1} \overbrace{4 \ 5 \ 6}^{i=2}. \quad (3.54)$$

The rotation subgroup of $SL(3, \mathbb{R})$ is placed on the indices $7, \dots, 9$, the noncompact generators on $1, \dots, 5$. In this case we have

$$C_{0-} = -\frac{1}{2}(g_1^2 + g_2^2), \quad (3.55)$$

$$T_{0-} = 2a_{12}, \quad (3.56)$$

$$\Delta_0 = a_{12}^2, \tag{3.57}$$

$$R^{(0ii)} = \frac{1}{|a_{12}|} (a_{i1}^2 + a_{i2}^2), \tag{3.58}$$

$$V_0 = -|a_{12}| - 2a_{12}. \tag{3.59}$$

The eigenvalues are:

$$0 \ (5 \times), \quad -2a_{12} \ (6 \times), \quad 6|a_{12}| + 4a_{12} \ (3 \times), \quad 6|a_{12}| - 2a_{12} \ (15 \times), \\ 6|a_{12}| - 6a_{12} \ (7 \times), \tag{3.60}$$

In the de Sitter case ($a_{12} < 0$) there are no negative eigenvalues, but of course the $SU(1, 1)$ scalars are tachyons because $C_- < 0$. The five zero eigenvalues are associated to the Goldstone fields which give mass to the gauge fields corresponding to the non-compact gauge generators.

In the AdS case ($a_{12} > 0$) the potential in the extremum is $V_0 = -3|a_{12}|$, in that case there are seven additional zero eigenvalues.

3.9 $SU(2, 1)_+ \otimes SO(2, 1)_+$

The gauge generators are arranged as follows:

$$\overbrace{1 \ 2 \ 3 \ 4 \ 7 \ 8 \ 9 \ 10}^{i=1} \ \overbrace{5 \ 6 \ 11}^{i=2} . \tag{3.61}$$

The compact generators of $SU(2, 1)$ are placed on the indices $7, \dots, 10$, the noncompact ones on $1, \dots, 4$. In this case we have:

$$C_{0-} = 6g_1^2 + \frac{1}{2}g_2^2, \tag{3.62}$$

$$T_{0-} = 0, \tag{3.63}$$

$$\Delta_0 = 12a_{12}^2, \tag{3.64}$$

$$R^{(0ii)} = \frac{1}{\sqrt{3}|a_{12}|} (6a_{i1}^2 + \frac{1}{2} a_{i2}^2), \tag{3.65}$$

$$V_0 = 2\sqrt{3}|a_{12}| > 0. \tag{3.66}$$

Again we find a de Sitter vacuum. The mass of the $SU(1, 1)$ scalars is $2\sqrt{3}|a_{12}|$ while in the matter sector the mass eigenvalues are:

$$0 \ (6 \times), \quad \sqrt{3}|a_{12}| \ (12 \times), \quad 2\sqrt{3}|a_{12}| \ (10 \times), \quad \sqrt{3}|a_{12}|(1 \pm 2\sqrt{2}) \ (4 \times). \tag{3.67}$$

The spectrum contains four tachyonic modes. The six zero modes correspond to the Goldstone bosons of the broken noncompact generators.

4. Conclusions

In this paper we have searched in gauged $N = 4$ supergravity with semi-simple gauge groups for examples that give a positive cosmological constant and a non-negative mass-matrix for all scalar fields. In all examples considered the scalar potential does allow a

positive extremum, but we have always found tachyons. In our search we limited ourselves to six vector multiplets, and therefore 36 matter scalars.

We found that there are two classes of gauge groups in the nine that were considered. Five have a positive extremum for all values of the parameters in the problem (coupling constants, $SU(1,1)$ angles), have tachyonic modes in the matter sector, and positive m^2 for the $SU(1,1)$ scalars. In four cases we find that the sign of potential in the extremum depends on the choice of parameters, that the matter scalars all have positive m^2 (if the parameters are chosen such that the extremum occurs for positive potential), and the two $SU(1,1)$ scalars are the tachyons. These last four are precisely all the cases with an $SO(3)_-$ subgroup. This distinction is not yet understood.

Certain features of the mass spectrum are clear. We always find the appropriate number of massless modes to provide for the Goldstone bosons of the broken gauge symmetries. In a number of cases we have more zero modes than Goldstone bosons. This might be an indication that such models can be embedded in a gauged supergravity with $N > 4$, and that these extra zero modes are related to Goldstone bosons that occur in the larger supergravity theory.

Another feature of gauged supergravity theories, remarked in [24], is the fact that the mass spectrum of gauged supergravity is often such that $3m^2/V_0$ is an integer. This is an interesting observation, since it makes such models unsuitable as candidates for slow-roll inflationary scenarios. We find the same property for the ratio of mass and potential in cases where the gauge group is a product of two simple groups, with one exception in the $SU(2,1)_+ \otimes SO(2,1)_+$ gauging in section 3.9, where this ratio contains a factor $\sqrt{2}$. For groups that have three simple factors and positive cosmological constant, the ratio becomes parameter dependent for some of the masses. However, also in these cases one does not have enough freedom to tune the parameters such that all tachyon masses become small.

As far as supersymmetry breaking is concerned the analysis of [21] can be applied. For the groups in sections 3.1–3.4 and 3.9 we always have $V_0 > 0$, and supersymmetry is completely broken. In sections 3.5–3.8 V_0 depends on the sign of a_{12} , and one finds that for $V_0 > 0$ supersymmetry is completely broken, for $V_0 \leq 0$ $N = 4$ supersymmetry is preserved. The supersymmetry variations of the fermions in these last three cases are proportional to $g_1\Phi_{(1)} - ig_2\Phi_{(2)}$, for which

$$|g_1\Phi_{(1)} - ig_2\Phi_{(2)}|^2 = R^{(011)} + R^{(022)} + 2I^{(012)} = 2(|a_{12}| - a_{12}). \quad (4.1)$$

This vanishes for $a_{12} > 0$, leading to unbroken supersymmetry in AdS spacetime. In these cases the potential in the AdS extremum is $V_0 = -3|a_{12}|$. This value follows also from the integrability condition arising from the supersymmetry variation of the gravitinos.

The relation of our $N = 4$ work with the $N = 2$ results of [9] remains intriguing. There are three cases presented in [9]. The one which seems most directly related to $N = 4$ supergravity has five vector multiplets, gauging, with the graviphoton, a six-dimensional $SO(2,1) \otimes SO(3)$ group. In addition, the model has two hyper-multiplets, giving a total of 18 scalar fields. The scalar manifold is

$$\left[\frac{SU(1,1)}{U(1)} \times \frac{SO(2,4)}{SO(2) \times SO(4)} \right] \times \left[\frac{SO(4,2)}{SO(4) \times SO(2)} \right], \quad (4.2)$$

where the last factor corresponds to the hyper-multiplets. The two $SU(1,1)$ scalars play a similar role as in $N = 4$ and allow the introduction of $SU(1,1)$ mixing angles in the coupling to the vectors. The gauge group is embedded in both $SO(4,2)$ groups, and it was shown in [9] how to obtain the manifold (4.2) from a truncation of

$$\frac{SU(1,1)}{U(1)} \times \frac{SO(6,6)}{SO(6) \times SO(6)}. \quad (4.3)$$

In the $N = 4$ case the gauge group would be $[SO(2,1) \times SO(3)]^2$, the $N = 2$ group being the diagonal subgroup. The only way to embed this group in $N = 4$ is as $SO(3)_- \times SO(2,1)_+ \times SO(2,1)_- \times SO(3)_+$, which we have in table 2 in section 2.2 but which does not have an extremum in the matter and $SU(1,1)$ scalars. We may restrict our analysis to points in the moduli space preserving the diagonal compact subgroup $SO(2)_D \times SO(3)_D$ of the gauge group (in this analysis we set the $g_1 = g_4, g_2 = g_3, \alpha_1 = \alpha_4, \alpha_2 = \alpha_3$). To this end it suffices to study the behavior of the scalar potential as a function of the $SO(2)_D \times SO(3)_D$ singlets only (indeed, the scalar potential being invariant in particular under $SO(2)_D \times SO(3)_D$, its dependence on scalar fields which are not singlets with respect to this group will be at least quadratic, and therefore in order to analyse the critical points of the potential exhibiting $SO(2)_D \times SO(3)_D$ symmetry, we may set these fields to zero). These singlets in the matter sector are four. This can be seen by first considering the action of the compact subgroup $SO(3)_- \times SO(2)_+ \times SO(2)_- \times SO(3)_+$ under which the matter scalar fields transform as follows:

$$(\mathbf{3} + \mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{1}) + (\mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{3} + \mathbf{1}) + (\mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1}) + (\mathbf{3} + \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{3} + \mathbf{1}). \quad (4.4)$$

The scalars in $(\mathbf{3} + \mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{1})$ and $(\mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{3} + \mathbf{1})$ correspond to the 16 scalars in the $SO(2,4)/SO(2) \times SO(4)$ and $SO(4,2)/SO(4) \times SO(2)$ cosets, those in $(\mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1})$ and $(\mathbf{3} + \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{3} + \mathbf{1})$ will be projected out by the $N = 4 \rightarrow N = 2$ truncation. Branching the above representations with respect to $SO(2)_D \times SO(3)_D$ we obtain two singlets $\varphi_{1,2}$ from $(\mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1})$, one singlet ξ from $(\mathbf{3}, \mathbf{1}, \mathbf{1}, \mathbf{3})$ and finally the $(\mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{1})$ singlet ξ_0 . A non vanishing value for ξ would therefore break $SO(3)_- \times SO(3)_+$ to $SO(3)_D$. Although the potential does not have critical points in the $SU(1,1)$ and $\varphi_{1,2}, \xi_0, \xi$ scalars, a numerical analysis shows that it can be extremized with respect to all the scalars except ξ , for large enough values of $|\xi|$ (at the origin of the matter sector the potential does not have an extremum with respect to the $SU(1,1)$ scalars). In other words there is an infinite range of values for ξ for which all the other scalars can be fixed so that the potential has a run-away behavior in ξ field only. In these points the potential is positive. It seems therefore that in lifting the $N = 2$ model with vector and hyper-multiplets to $N = 4$ new scalar fields emerge which destabilize the $N = 2$ de Sitter vacuum.

The program presented in this paper can be extended in many directions. One possibility could be to consider contractions (*CSO* groups) of the gauge groups studied here. Also the present analysis can be generalized to include Peccei-Quinn symmetries. These symmetries naturally appear in Scherk-Schwarz reductions [11, 15, 17], therefore such an investigation could also elucidate the relation between $N = 4$ supergravity with nonzero $SU(1,1)$ angles and string theory. Gaugings related to Scherk-Schwarz reductions from

$D = 5$ could also provide new ways for obtaining some of the $N = 2$ models with stable de Sitter vacua [9] as effective realizations of a larger gauged $N = 4$ theory (although a definite statement about this possibility would require an analysis that we postpone to future work). As an example we could consider the $N = 4$ supergravity coupled to six matter multiplets and gauge a group of the form $G_{S-S} \times \text{SO}(2, 1) \times \text{SO}(3)$ where G_{S-S} is a non-semi-simple gauge group á la Scherk–Schwarz. It was shown indeed that, for a certain choice of the gauge parameters, the effect of G_{S-S} alone amounts to a partial supersymmetry breaking from $N = 4$ to $N = 2$, in which the final effective supergravity is coupled to five vector multiplets and no hyper-multiplet. This is the ungauged version of one of the models considered in [9]. Therefore if we gauge in the $N = 4$ theory the group $G_{S-S} \times \text{SO}(2, 1) \times \text{SO}(3)$ and introduce $\text{SU}(1, 1)$ angles for each simple factor, we would be left at the level of the $N = 2$ effective model with a surviving $\text{SO}(2, 1) \times \text{SO}(3)$ gauge group. However, in order to recover in this framework one of the models without hyper-multiplets constructed in [9] a crucial ingredient would be the presence of the Fayet–Iliopoulos term corresponding to the $\text{SO}(3)$ factor, whose $N = 4$ origin is still unclear.

There are (at least) two aspects of this program which remain to be elucidated. The first is the existence or non-existence of stable de Sitter vacua in $N = 4$ supergravity, the second is the relation of gauged $N = 4$ supergravity with $\text{SU}(1, 1)$ angles with ten and/or eleven dimensions. If such a relation could be established, the no-go theorem of [3, 4, 5] would come into play, and one would know that to solve the first problem would require flux reduction or hyperbolic reduction and/or other ways around the no-go theorem [6].

Acknowledgments

MT would like to thank S. Ferrara for useful remarks. The work of DBW is part of the research programme of the “Stichting voor Fundamenteel Onderzoek van de Materie” (FOM). This work is supported in part by the European Commission RTN programme HPRN-CT-2000-00131, in which MdR and DBW are associated to the University of Utrecht. The work of MT is supported by a European Community Marie Curie Fellowship under contract HPRN-CT-2001-01276. SP thanks the ICTP in Trieste and the Centre for Theoretical Physics in Groningen for their hospitality.

A. $\text{SU}(1, 1)$ scalars and angles

We parametrise the scalars of the $\text{SU}(1, 1)/\text{U}(1)$ coset in a suitable $\text{U}(1)$ gauge as

$$\phi_1 = \frac{1}{\sqrt{1-r^2}}, \quad \phi_2 = \frac{re^{i\varphi}}{\sqrt{1-r^2}}. \tag{A.1}$$

The scalars r and φ then appear in the potential (2.1) through

$$\begin{aligned} R^{(ij)} &= \frac{g_i g_j}{2} (\Phi_i^* \Phi_j + \Phi_j^* \Phi_i) \\ &= g_i g_j \left(\cos(\alpha_i - \alpha_j) \frac{1+r^2}{1-r^2} - \frac{2r}{1-r^2} \cos(\alpha_i + \alpha_j + \varphi) \right), \end{aligned} \tag{A.2}$$

$$\begin{aligned}
 I^{(ij)} &= \frac{g_i g_j}{2i} (\Phi_i^* \Phi_j - \Phi_j^* \Phi_i) \\
 &= -g_i g_j \sin(\alpha_i - \alpha_j).
 \end{aligned}
 \tag{A.3}$$

Introducing

$$C_{\pm} = \sum_{ij} g_i g_j \cos(\alpha_i \pm \alpha_j) V_{ij}, \quad S_+ = \sum_{ij} g_i g_j \sin(\alpha_i + \alpha_j) V_{ij},
 \tag{A.4}$$

$$T_- = \sum_{ij} g_i g_j \sin(\alpha_i - \alpha_j) W_{ij},
 \tag{A.5}$$

we rewrite the potential as

$$V = C_- \frac{1+r^2}{1-r^2} - \frac{2r}{1-r^2} (C_+ \cos \varphi - S_+ \sin \varphi) - T_-.
 \tag{A.6}$$

This extremum in r and φ takes on the form

$$\begin{aligned}
 \cos \varphi_0 &= \frac{s_1 C_+}{\sqrt{C_+^2 + S_+^2}}, & \sin \varphi_0 &= -\frac{s_1 S_+}{\sqrt{C_+^2 + S_+^2}}, \\
 r_0 &= \frac{1}{\sqrt{C_+^2 + S_+^2}} (s_1 C_- + s_2 \sqrt{\Delta}), & \Delta &\equiv C_-^2 - C_+^2 - S_+^2,
 \end{aligned}
 \tag{A.7}$$

where s_1 and s_2 are signs. These are determined by requiring $r_0 \geq 0$ and $r_0 < 1$, this gives $s_1 = \text{sgn} C_-$ and $s_2 = -1$. Substitution of r_0 and φ_0 in V leads to

$$V_0 = \text{sgn} C_- \sqrt{\Delta} - T_-.
 \tag{A.8}$$

In the case that all $SU(1,1)$ angles α_i vanish, $S_+ = T_- = 0$ and $C_- = C_+$, and one finds $r_0 = 1$ and $\Delta = 0$. This is a singular point of the parametrisation, which we will exclude. It is generalisation of the Freedman-Schwarz potential [23] to the case of general matter coupling.

For the kinetic term and mass-matrix of the $SU(1,1)$ scalars we introduce:

$$\begin{aligned}
 x' &= \frac{2}{(1-r_0)^2} (r \cos \varphi - r_0 \cos \varphi_0), \\
 y' &= \frac{2}{(1-r_0)^2} (r \sin \varphi - r_0 \sin \varphi_0).
 \end{aligned}
 \tag{A.9}$$

In these variables we find

$$\mathcal{L}_\phi = -\frac{1}{2} \left(\frac{1-r_0^2}{1-r^2} \right)^2 (\partial_\mu x' \partial^\mu x' + \partial_\mu y' \partial^\mu y') - V_0 - \frac{1}{2} \text{sgn} C_- \sqrt{\Delta} (x'^2 + y'^2) + \dots
 \tag{A.10}$$

It is clear that we have two tachyons for $\text{sgn} C_- < 0$, and two positive mass scalars for $\text{sgn} C_- > 0$.

It is useful to also analyse the kinetic term of the vectors, since positivity³ of these terms might give further constraints on the $SU(1,1)$ scalars, to which the vectors couple.

³In [8] it was incorrectly stated that the kinetic terms of the vectors acquire the wrong sign. The discussion below should clarify this point.

In $Z = Z_0$ we have

$$\begin{aligned} \mathcal{L}_{\text{kin},A} = & -\frac{1}{4}F_{\mu\nu}{}^{+A}F^{\mu\nu}{}_{+A} \left(-\frac{\phi^1{}_{(A)} - \phi^2{}_{(A)}}{\Phi_{(A)}} + \frac{2}{|\Phi_{(A)}|^2} \right) - \\ & -\frac{1}{4}F_{\mu\nu}{}^{+I}F^{\mu\nu}{}_{+I} \left(\frac{\phi^1{}_{(I)} - \phi^2{}_{(I)}}{\Phi_{(I)}} \right) + \text{h.c.} \end{aligned} \quad (\text{A.11})$$

Here $\Phi_{(R)} \equiv e^{i\alpha_R}\phi^1 + e^{-i\alpha_R}\phi^2$. We find after some algebra:

$$-\frac{1}{4}F_{\mu\nu}{}^{+A}F^{\mu\nu}{}_{+A}S^{(A)} - \frac{1}{4}F_{\mu\nu}{}^{+I}F^{\mu\nu}{}_{+I}S^{*(I)} + \text{h.c.} \quad (\text{A.12})$$

Here $S^{(R)}$ is given by

$$S^{(R)} = \frac{1 + r e^{i(\varphi+2\alpha_R)}}{1 - r e^{i(\varphi+2\alpha_R)}}. \quad (\text{A.13})$$

The imaginary part of S gives a total derivative in the kinetic term. Therefore the kinetic terms are determined by

$$\text{Re}S^{(R)} = \frac{1 - r^2}{1 + r^2 - 2r \cos(\varphi + 2\alpha_R)}, \quad (\text{A.14})$$

showing that the domain of positivity is $r < 1$. In the extremum for the $SU(1,1)$ scalars, and in the origin of the matter scalar manifold, this becomes for the i 'th factor of the gauge group

$$\text{Re}S^{(i)} = \frac{g_i^2}{R^{(0ii)}}, \quad (\text{A.15})$$

where R^{0ii} is given in (2.20). Indeed, we find that always $R^{0ii} > 0$.

B. Generators and structure constants

Our conventions for the structure constant are the following:

$$[T_R, T_S] = i f_{RS}{}^U T_U, \quad f_{RST} \equiv f_{RS}{}^U \eta_{TU}, \quad (\text{B.1})$$

where the structure constants f are real. The structure constants f_{RST} are completely antisymmetric, since η_{RS} is proportional to the Cartan-Killing metric $\text{tr} T_R T_S$ of the gauge group. The factor 2 in this proportionality we absorb in a redefinition of the coupling constants g_i . Our choice for the generators is based on the Gell-Mann matrices (extended

Group	Generators (L; K)
SO(3)	$L_\alpha = \lambda_7, -\lambda_5, \lambda_2$
SO(2, 1)	$L_3 = \lambda_2, K_{1,2} = i\lambda_6, i\lambda_4,$
SO(3, 1)	$L_\alpha = \lambda_7, -\lambda_5, \lambda_2, K_\alpha = i\lambda_9, i\lambda_{10}, i\lambda_{11}$
SL(3, \mathbb{R})	$L_\alpha = \lambda_7, -\lambda_5, \lambda_2, K_\alpha = i\lambda_6, i\lambda_4, i\lambda_1, K_4 = i\lambda_3, K_5 = i\lambda_8$
SU(2, 1)	$L_\alpha = \lambda_1, \lambda_2, \lambda_3, L_4 = \lambda_8; K_1 = i\lambda_4, K_2 = i\lambda_5, K_3 = i\lambda_6, K_4 = i\lambda_7$

Table 4: Generators for the simple groups used in this paper. We always have $\alpha, \beta, \gamma = 1, \dots, 3$.

to 4×4 matrices to treat SO(3, 1) in the same context):

$$\begin{aligned}
 \lambda_1 &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \lambda_2 &= \begin{pmatrix} 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \lambda_3 &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\
 \lambda_4 &= \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \lambda_5 &= \begin{pmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \lambda_6 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\
 \lambda_7 &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \lambda_8 &= \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & & \\
 \lambda_9 &= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, & \lambda_{10} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, & \lambda_{11} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}. \tag{B.2}
 \end{aligned}$$

The groups we consider have compact generators, which in the following we will denote by L_R , and noncompact generators, denoted by K_R . We will specify the structure constants with three L 's, and with two K 's and one L . The choice of the generators of the simple groups used in this paper are summarized in table 4.

The structure constants for the SO(3) subgroups are in all cases

$$[L_\alpha, L_\beta] = i\epsilon_{\alpha\beta\gamma}L_\gamma, \quad (\alpha, \beta, \gamma = 1, 2, 3). \tag{B.3}$$

For SO(2, 1) we have

$$[L_3, K_\beta] = -i\epsilon_{3\beta\gamma}K_\gamma, \quad (\beta, \gamma = 1, 2). \tag{B.4}$$

In the case of SO(3, 1) the commutation relation involving noncompact generators are

$$[L_\alpha, K_\beta] = i\epsilon_{\alpha\beta\gamma}K_\gamma, \quad (\alpha, \beta, \gamma = 1, 2, 3). \tag{B.5}$$

For SL(3, \mathbb{R}) the relations between L and K are:

$$[L_\alpha, K_\beta] = -i\epsilon_{\alpha\beta\gamma}K_\gamma, \quad (\alpha \neq \beta, \quad \alpha, \beta, \gamma = 1, 2, 3),$$

	$K_1 = i\lambda_4$	$K_2 = i\lambda_6$	$K_3 = i\lambda_5$	$K_4 = i\lambda_7$
$L_1 = \lambda_1$	K_4	K_3	$-K_2$	$-K_1$
$L_2 = \lambda_2$	K_2	$-K_1$	K_4	$-K_3$
$L_3 = \lambda_3$	K_3	$-K_4$	$-K_1$	K_2
$L_4 = \lambda_8$	$\sqrt{3}K_3$	$\sqrt{3}K_4$	$-\sqrt{3}K_1$	$-\sqrt{3}K_2$

Table 5: Commutation relations between compact and noncompact generators in $SU(2,1)$. The table reads $[L_1, K_1] = iK_4$, etc.

$$[L_1, K_1] = i(K_4 - \sqrt{3}K_5), \quad [L_2, K_2] = i(K_4 + \sqrt{3}K_5), \quad [L_3, K_3] = -2iK_4. \quad (\text{B.6})$$

Finally, for $SU(2,1)$ we have

$$[L_\alpha, L_\beta] = 2i\epsilon_{\alpha\beta\gamma}L_\gamma, \quad (\alpha, \beta, \gamma = 1, 2, 3), \quad [L_\alpha, L_4] = 0. \quad (\text{B.7})$$

The commutation relation between the remaining generators are given in table 5. The structure constants presented for the different groups are the f_{RS}^T . For use in the calculation of second derivatives etc. the index T has to be lowered by η_{TU} , which may give a sign depending on the embedding of the group in $SO(6,6)$. Other commutation relations, such as $[K_\alpha, K_\beta]$ follow from the antisymmetry of f_{RST} .

C. The parameters P

The independent scalars are contained in G and B . Define $G_\pm \equiv (G \pm B)$, and $P \equiv G_+$. Then $G = \frac{1}{2}(P + P^T)$, $B = \frac{1}{2}(P - P^T)$, and we find

$$\begin{aligned} \frac{\partial G_{cd}}{\partial P_{ab}} &= \frac{1}{2}(\delta_{ac}\delta_{bd} + \delta_{ad}\delta_{bc}), & \frac{\partial B_{cd}}{\partial P_{ab}} &= \frac{1}{2}(\delta_{ac}\delta_{bd} - \delta_{ad}\delta_{bc}), \\ \frac{\partial G^{-1}_{cd}}{\partial P_{ab}} &= -\frac{1}{2} \left(G^{-1}_{ca}G^{-1}_{bd} + G^{-1}_{cb}G^{-1}_{ad} \right). \end{aligned} \quad (\text{C.1})$$

In the study of the potential we need the first and second derivatives of X and Y with respect to P . We find:

$$\begin{aligned} \left. \frac{\partial X_{cd}}{\partial P_{ab}} \right|_0 &= 0, \\ \left. \frac{\partial Y_{cd}}{\partial P_{ab}} \right|_0 &= \delta_{ad}\delta_{bc}, \\ \left. \frac{\partial^2 X_{ef}}{\partial P_{ab}\partial P_{cd}} \right|_0 &= \frac{1}{2}\delta_{ac}(\delta_{de}\delta_{bf} + \delta_{be}\delta_{df}), \\ \left. \frac{\partial^2 Y_{ef}}{\partial P_{ab}\partial P_{cd}} \right|_0 &= -\frac{1}{2}(\delta_{bc}\delta_{de}\delta_{af} + \delta_{ad}\delta_{be}\delta_{cf}), \end{aligned} \quad (\text{C.2})$$

References

- [1] C.L. Bennett et al., *First year Wilkinson microwave anisotropy probe (WMAP) observations: preliminary maps and basic results*, *Astrophys. J. Suppl.* **148** (2003) 1 [[astro-ph/0302207](#)].

- [2] SUPERNOVA SEARCH TEAM collaboration, A.G. Riess et al., *Observational evidence from supernovae for an accelerating universe and a cosmological constant*, *Astron. J.* **116** (1998) 1009–1038 [[astro-ph/9805201](#)];
SUPERNOVA COSMOLOGY PROJECT collaboration, S. Perlmutter et al., *Measurements of omega and lambda from 42 high-redshift supernovae*, *Astrophys. J.* **517** (1999) 565 [[astro-ph/9812133](#)].
- [3] G.W. Gibbons, *Aspects of supergravity theories*, in *Supersymmetry, supergravity and related topics*, F. Del Aguila, J. A. de Azcárraga and L. E. Ibáñez eds., World Scientific, Singapore 1985.
- [4] B. de Wit, D.J. Smit and N.D. Hari Dass, *Residual supersymmetry of compactified $D = 10$ supergravity*, *Nucl. Phys.* **B 283** (1987) 165.
- [5] J.M. Maldacena and C. Núñez, *Supergravity description of field theories on curved manifolds and a no go theorem*, *Int. J. Mod. Phys.* **A 16** (2001) 822 [[hep-th/0007018](#)].
- [6] P.K. Townsend, *Cosmic acceleration and M-theory*, [hep-th/0308149](#).
- [7] S.J. Gates Jr. and B. Zwiebach, *Gauged $N = 4$ supergravity theory with a new scalar potential*, *Phys. Lett.* **B 123** (1983) 200.
- [8] M. de Roo, D.B. Westra and S. Panda, *de Sitter solutions in $N = 4$ matter coupled supergravity*, *J. High Energy Phys.* **02** (2003) 003 [[hep-th/0212216](#)].
- [9] P. Fre, M. Trigiante and A. Van Proeyen, *Stable de Sitter vacua from $N = 2$ supergravity*, *Class. and Quant. Grav.* **19** (2002) 4167 [[hep-th/0205119](#)].
- [10] A.H. Chamseddine, *$N=4$ supergravity coupled to $N = 4$ matter*, *Nucl. Phys.* **B 185** (1981) 403.
- [11] J. Scherk and J.H. Schwarz, *Spontaneous breaking of supersymmetry through dimensional reduction*, *Phys. Lett.* **B 153** (1979) 61; *How to get masses from extra dimensions*, *Nucl. Phys.* **B 153** (1979) 61.
- [12] A.H. Chamseddine, *Interacting supergravity in ten-dimensions: the role of the six - index gauge field*, *Phys. Rev.* **D 24** (1981) 3065.
- [13] S. Thomas and P.C. West, *Dimensional reduction generates finiteness preserving soft terms*, *Nucl. Phys.* **B 245** (1984) 45.
- [14] M. Porrati and F. Zwirner, *Supersymmetry breaking in string derived supergravities*, *Nucl. Phys.* **B 326** (1989) 162.
- [15] E. Bergshoeff, M. de Roo and E. Eyras, *Gauged supergravity from dimensional reduction*, *Phys. Lett.* **B 413** (1997) 70 [[hep-th/9707130](#)].
- [16] R. D’Auria, S. Ferrara and S. Vaula, *$N = 4$ gauged supergravity and a IIB orientifold with fluxes*, *New J. Phys.* **4** (2002) 71 [[hep-th/0206241](#)].
- [17] L. Andrianopoli, R. D’Auria, S. Ferrara and M.A. Lledo, *Gauging of flat groups in four dimensional supergravity*, *J. High Energy Phys.* **07** (2002) 010 [[hep-th/0203206](#)]; *Duality and spontaneously broken supergravity in flat backgrounds*, *Nucl. Phys.* **B 640** (2002) 63 [[hep-th/0204145](#)].
- [18] R. D’Auria, S. Ferrara, M.A. Lledo and S. Vaula, *No-scale $N = 4$ supergravity coupled to Yang-Mills: the scalar potential and super Higgs effect*, *Phys. Lett.* **B 557** (2003) 278 [[hep-th/0211027](#)].

- [19] R. D'Auria, S. Ferrara, F. Gargiulo, M. Trigiante and S. Vaula, *$N = 4$ supergravity lagrangian for type-IIB on T^6/\mathbb{Z}_2 in presence of fluxes and d3-branes*, *J. High Energy Phys.* **06** (2003) 045 [[hep-th/0303049](#)].
- [20] C. Angelantonj, S. Ferrara and M. Trigiante, *New $D = 4$ gauged supergravities from $N = 4$ orientifolds with fluxes*, *J. High Energy Phys.* **10** (2003) 015 [[hep-th/0306185](#)]; *Unusual gauged supergravities from type-IIA and type-IIB orientifolds*, [hep-th/0310136](#).
- [21] M. de Roo and P. Wagemans, *Gauge matter coupling in $N = 4$ supergravity*, *Nucl. Phys.* **B 262** (1985) 644.
- [22] M. de Roo, *Matter coupling in $N = 4$ supergravity*, *Nucl. Phys.* **B 255** (1985) 515.
- [23] D.Z. Freedman and J.H. Schwarz, *$N = 4$ supergravity theory with local $SU(2) \times SU(2)$ invariance*, *Nucl. Phys.* **B 137** (1978) 333.
- [24] R. Kallosh, A.D. Linde, S. Prokushkin and M. Shmakova, *Gauged supergravities, de Sitter space and cosmology*, *Phys. Rev.* **D 65** (2002) 105016 [[hep-th/0110089](#)].