

Interval extropy and weighted interval extropy

Original

Interval extropy and weighted interval extropy / Buono, Francesco; Kamari, Osman; Longobardi, Maria. - In: RICERCHE DI MATEMATICA. - ISSN 1827-3491. - 72:1(2023), pp. 283-298. [10.1007/s11587-021-00678-x]

Availability:

This version is available at: 11583/2994632 since: 2024-11-25T11:49:59Z

Publisher:

Springer

Published

DOI:10.1007/s11587-021-00678-x

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Springer postprint/Author's Accepted Manuscript

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: <http://dx.doi.org/10.1007/s11587-021-00678-x>

(Article begins on next page)

Interval Extropy and Weighted Interval Extropy

Francesco Buono^{1*†}, Osman Kamari^{2†} and Maria
Longobardi^{3†}

¹Dipartimento di Matematica e Applicazioni “Renato Caccioppoli”, Università degli Studi di Napoli Federico II, Naples, Italy.

²Department of Business Management, University of Human Development, Sulaymaniyah, Iraq.

³Dipartimento di Biologia, Università degli Studi di Napoli Federico II, Naples, Italy.

*Corresponding author(s). E-mail(s): francesco.buono3@unina.it;
Contributing authors: osman.kamari@uhd.edu.iq;
maria.longobardi@unina.it;

†These authors contributed equally to this work.

Abstract

Recently, Extropy was introduced by Lad, Sanfilippo and Agrò as a complement dual of Shannon Entropy. In this paper, we propose dynamic versions of Extropy for doubly truncated random variables as measures of uncertainty called Interval Extropy and Weighted Interval Extropy. Some characterizations of random variables related to these new measures are given. Several examples are shown. These measures are evaluated under the effect of linear transformations and, finally, some bounds for them are presented.

Keywords: Uncertainty, Extropy, Weighted Extropy, Characterization

MSC Classification: 62N05 , 94A17

1 Introduction

The concept of Shannon entropy as a basic measure of uncertainty for a random variable was introduced by Shannon [20]. Let X be an absolutely continuous non-negative random variable having probability density function (pdf) f and cumulative distribution function (cdf) F . In Reliability Theory, X represents the lifetime of a system or a living organism. Shannon entropy for this kind of random variables is named differential entropy and is defined as follows:

$$H(X) = - \int_0^{+\infty} f(x) \log f(x) dx,$$

where \log denotes the natural logarithm. Recently, another measure of uncertainty, known as extropy, was proposed by Lad et al. [13] as a dual measure of Shannon entropy. For a non-negative random variable X the extropy is defined as below:

$$J(X) = -\frac{1}{2} \int_0^{+\infty} f^2(x) dx. \quad (1)$$

The concept of extropy is useful in many fields: for instance, it is applied in automatic speech recognition [4]. In particular, the extropy of a network output with respect to the training set can be obtained in order to compute a kind of transformed cross entropy. Moreover, extropy is a measure better than entropy in some scenarios in statistical mechanics and thermodynamics [14]. More recently, some applications of extropy have been done in pattern recognition [3, 10].

Qiu et al. [19] defined the extropy for residual lifetime $X_t = (X - t \mid X > t)$ whose pdf is $f_{X_t}(x) = \frac{f(x+t)}{\bar{F}(t)}$ and survival function $\bar{F}_{X_t}(x) = \frac{\bar{F}(x+t)}{\bar{F}(t)}$, $x > 0$, called the residual extropy (REx) at time t and defined as

$$J(X_t) = -\frac{1}{2(\bar{F}(t))^2} \int_t^{+\infty} f^2(x) dx,$$

where $\bar{F}(t) = \mathbb{P}(X > t) = 1 - F(t)$ is the survival (reliability) function of X .

Krishnan et al. [12] and Kamari and Buono [9] studied the dual measure of residual extropy for past lifetime ${}_tX = (X \mid X \leq t)$, whose pdf is $f_{{}_tX}(x) = \frac{f(x)}{F(t)}$ and cumulative distribution function $F_{{}_tX}(x) = \frac{F(x)}{F(t)}$, $0 < x < t$, called past extropy (PEX) and defined as follows:

$$J({}_tX) = -\frac{1}{2(F(t))^2} \int_0^t f^2(x) dx.$$

Recently, there has been growing attention to study uncertainty measures for doubly truncated random variable which is widely applied in many fields such

as survival analysis and reliability engineering. In survival analysis, if the lifetime of the item falls in an interval (t_1, t_2) , information about lifetime between these two points (also named doubly truncated failure time) is studied, see, for instance, Betensky and Martin [5], Khorashadizadeh et al. [11] and Poursaeed and Nematollahi [18]. Then, the random variable $(X | t_1 < X < t_2)$ is introduced with pdf $f_{X_{t_1, t_2}}(x) = \frac{f(x)}{F(t_2) - F(t_1)}$ and cdf $F_{X_{t_1, t_2}}(x) = \frac{F(x) - F(t_1)}{F(t_2) - F(t_1)}$, $t_1 < x < t_2$. With this motivation, Sunoj et.al. [21] introduced Interval Entropy to measure uncertainty in truncated random variable $(X | t_1 < X < t_2)$ that is defined as follows:

$$H(t_1, t_2) = - \int_{t_1}^{t_2} \frac{f(x)}{F(t_2) - F(t_1)} \log \frac{f(x)}{F(t_2) - F(t_1)} dx, \quad (2)$$

which is an extension of Shannon Entropy. If $t_2 \rightarrow +\infty$, then $H(t_1, t_2)$ tends to residual entropy which was introduced by Ebrahimi [8]. Moreover, if $t_1 \rightarrow 0$, then $H(t_1, t_2)$ tends to the past entropy defined by Di Crescenzo and Longobardi [6]. Several other properties of the interval entropy were studied by Misagh and Yari [16].

Di Crescenzo and Longobardi [7] defined weighted entropy, weighted residual entropy and weighted past entropy, which are respectively given by

$$\begin{aligned} H^w(X) &= - \int_0^{+\infty} x f(x) \log f(x) dx, \\ H^w(X_t) &= - \int_0^{+\infty} x \frac{f(x)}{F(t)} \log \frac{f(x)}{F(t)} dx, \\ H^w({}_tX) &= - \int_0^{+\infty} x \frac{f(x)}{F(t)} \log \frac{f(x)}{F(t)} dx. \end{aligned}$$

Balakrishnan et al. [2] introduced weighted Entropy and its dynamic versions as Weighted Residual Entropy and Weighted Past Entropy for residual and past lifetime as below:

$$\begin{aligned} J^w(X) &= -\frac{1}{2} \int_0^{+\infty} x f^2(x) dx, \\ J^w(X_t) &= -\frac{1}{2(F(t))^2} \int_t^{+\infty} x f^2(x) dx, \\ J^w({}_tX) &= -\frac{1}{2(F(t))^2} \int_0^t x f^2(x) dx. \end{aligned}$$

Weighted Interval Entropy was introduced by Misagh and Yari [15] for doubly truncated random variable $(X | t_1 < X < t_2)$ as follows:

$$IH^w(t_1, t_2) = - \int_{t_1}^{t_2} x \frac{f(x)}{F(t_2) - F(t_1)} \log \frac{f(x)}{F(t_2) - F(t_1)} dx.$$

In analogy with the novel measures of uncertainty (i.e., Interval Entropy and Weighted Interval Entropy), here we introduce the concepts of Interval Entropy and Weighted Interval Entropy for doubly truncated random variables.

2 Interval Entropy

Let us suppose that the random variable $(X \mid t_1 < X < t_2)$ represents the lifetime of a unit which fails between t_1 and t_2 where $(t_1, t_2) \in D = \{(u, v) \in \mathbb{R}_+^2 : F(u) < F(v)\}$, the Entropy for the doubly truncated random variable is defined as follows:

$$IJ(t_1, t_2) = IJ(X \mid t_1 < X < t_2) = - \frac{1}{2(F(t_2) - F(t_1))^2} \int_{t_1}^{t_2} f^2(x) dx, \quad (3)$$

which is an extension of Entropy and is called Interval Entropy (IE_X). In (3) we have omitted the dependence of X in the expression $IJ(t_1, t_2)$, but when it is necessary we denote by $IJ_X(t_1, t_2)$ the interval entropy of X to distinguish it from the interval entropy of another random variable.

Remark 1 It is clear that $IJ(0, t_2) = J(t_2 X)$, $IJ(t_1, +\infty) = J(X_{t_1})$ and $IJ(0, +\infty) = J(X)$ are Past Entropy, Residual Entropy and Entropy, respectively.

Example 1 Let $X \sim \text{Exp}(\lambda)$, $\lambda > 0$ and support $(0, +\infty)$. Based on (3), we evaluate the interval entropy of X for $0 < t_1 < t_2 < +\infty$ and we obtain

$$\begin{aligned} IJ(t_1, t_2) &= \frac{-1}{2(e^{-\lambda t_1} - e^{-\lambda t_2})^2} \int_{t_1}^{t_2} \lambda^2 e^{-2\lambda x} dx \\ &= -\frac{\lambda}{4} \cdot \frac{e^{-\lambda t_2} + e^{-\lambda t_1}}{e^{-\lambda t_1} - e^{-\lambda t_2}}. \end{aligned}$$

In Figure 1, we plot the interval entropy as a function of t_1 for fixed t_2 (Figure 1(a)) and vice versa (Figure 1(b)) for $\lambda = 1$.

Example 2 Let X follow the Weibull distribution, $W2(\alpha, \lambda)$, with parameters $\alpha = \lambda = 2$, $X \sim W2(2, 2)$. The cdf and the pdf of X are expressed as

$$F(x) = 1 - \exp(-2x^2), \quad f(x) = 4x \exp(-2x^2), \quad x \in (0, +\infty).$$

Since the expression of the interval entropy is not given in terms of elementary functions, in Figure 2, we plot the interval entropy as a function of t_1 for fixed t_2 (Figure 2(a)) and vice versa (Figure 2(b)). From Figure 2(b) we observe an asymptotic behavior of the interval entropy as $t_2 \rightarrow +\infty$ towards $-t_1$, i.e., when the interval entropy




Fig. 1: Plot of IJ in Example 1 as a function of t_1 or t_2 fixing the other one with $t_i = 2$ (blue), 3 (red), 4 (yellow) and 5 (violet), $i = 1, 2$.




Fig. 2: Plot of IJ in Example 2 as a function of t_1 or t_2 fixing the other one with $t_i = 1$ (violet), 2 (blue), 3 (red) and 4 (yellow), $i = 1, 2$.

$IJ(t_1, t_2)$ reduces to the residual entropy $J(X_{t_1})$. In fact, the residual entropy of X in t can be derived as

$$\begin{aligned}
 J(X_t) &= -\frac{1}{2\exp(-4t^2)} \int_t^{+\infty} 16x^2 \exp(-4x^2) dx \\
 &= -t - \frac{1}{\exp(-4t^2)} \int_t^{+\infty} \exp(-4x^2) dx \\
 &= -t - \frac{1}{2\sqrt{2}\exp(-4t^2)} \int_{2\sqrt{2}t}^{+\infty} \exp\left(-\frac{y^2}{2}\right) dy = -t - \frac{\sqrt{\pi}}{2} \frac{\bar{F}_Z(2\sqrt{2}t)}{\exp(-4t^2)},
 \end{aligned}$$

where $\bar{F}_Z(\cdot)$ is the survival function of $Z \sim N(0, 1)$.

Example 3 Let X follow the Lognormal distribution, $Lognormal(\mu, \sigma^2)$, with parameters $\mu = 0$, $\sigma^2 = 1$, $X \sim Lognormal(0, 1)$. The pdf of X is expressed

6 Interval Entropy and Weighted Interval Entropy




Fig. 3: Plot of IJ in Example 3 as a function of t_1 or t_2 fixing the other one with $t_i = 1$ (violet), 2 (blue), 3 (red) and 4 (yellow), $i = 1, 2$.

as

$$f(x) = \frac{1}{x\sqrt{2\pi}} \exp\left(-\frac{\log^2 x}{2}\right), \quad x \in (0, +\infty).$$

Since the expression of the interval entropy is not given in terms of elementary functions, in Figure 3, we plot the interval entropy as a function of t_1 for fixed t_2 (Figure 3(a)) and vice versa (Figure 3(b)).

Based on the above examples, it could seem that the interval entropy is always decreasing with respect to t_1 and always increasing with respect to t_2 . In the following, we provide two counterexamples to show that the interval entropy can be non monotonous with respect to t_1 and t_2 .

Example 4 Let X be a random variable with support $S = (a, +\infty)$, $a > 0$, whose cdf is defined as $F(x) = 1 - \left(\frac{a}{x}\right)^b$, $b > 0$. The interval entropy of X can be obtained as follows

$$\begin{aligned} IJ(t_1, t_2) &= -\frac{1}{2\left[\left(\frac{a}{t_1}\right)^b - \left(\frac{a}{t_2}\right)^b\right]^2} \int_{t_1}^{t_2} \frac{b^2 a^{2b}}{x^{2b+2}} dx \\ &= \frac{1}{2\left[\frac{1}{t_1^b} - \frac{1}{t_2^b}\right]^2} \frac{b^2}{2b+1} \left[\frac{1}{t_2^{2b+1}} - \frac{1}{t_1^{2b+1}}\right] \\ &= \frac{b^2 (t_1^{2b+1} - t_2^{2b+1})}{2(2b+1)t_1 t_2 (t_2^b - t_1^b)^2}. \end{aligned}$$

Let us focus on the case $a = 1$ and $b = 10$. In Figure 4 we have plotted the interval entropy of X as a function of t_1 for fixed different values of t_2 and we can observe that it is initially increasing and then decreasing with respect to t_1 .




Fig. 4: Plot of IJ in Example 4 as a function of t_1 fixing $t_2 = 2$ (blue), 3 (red), 4 (yellow) and 5 (violet).




Fig. 5: Plot of IJ in Example 5 as a function of $t_2 \in (1, 2)$ fixing $t_1 = 0.1$.

Example 5 Let X be a random variable with cdf and pdf respectively defined as

$$F(x) = \begin{cases} \exp\left(-\frac{1}{2} - \frac{1}{x}\right), & \text{if } x \in (0, 1] \\ \exp\left(-2 + \frac{x^2}{2}\right), & \text{if } x \in [1, 2) \end{cases}$$

$$f(x) = \begin{cases} \exp\left(-\frac{1}{2} - \frac{1}{x}\right) \frac{1}{x^2}, & \text{if } x \in (0, 1] \\ \exp\left(-2 + \frac{x^2}{2}\right) x, & \text{if } x \in [1, 2). \end{cases}$$

In Figure 5 we have plotted the interval entropy as a function of $t_2 \in (1, 2)$ with fixed $t_1 = 0.1$ and we can observe a non monotonic behavior.

In the following theorem, we show the connection among Entropy and its dynamic versions.

8 Interval Entropy and Weighted Interval Entropy

Theorem 1 Let X be a random variable denoting the lifetime of a component. For all $0 < t_1 < t_2 < +\infty$ the entropy can be decomposed as follows:

$$J(X) = F^2(t_1)J(t_1, X) + (F(t_2) - F(t_1))^2 IJ(t_1, t_2) + \bar{F}^2(t_2)J(X_{t_2}), \quad (4)$$

i.e., the Entropy is a function of PEx, REx and IEx.

Proof From the definition of entropy (1), we can write

$$\begin{aligned} J(X) &= -\frac{1}{2} \int_0^{+\infty} f^2(x) dx \\ &= -\frac{1}{2} \int_0^{t_1} f^2(x) dx - \frac{1}{2} \int_{t_1}^{t_2} f^2(x) dx - \frac{1}{2} \int_{t_2}^{+\infty} f^2(x) dx. \end{aligned} \quad (5)$$

Now, we observe that the terms in the RHS of (5) are related to past entropy, interval entropy and residual entropy as

$$\begin{aligned} -\frac{1}{2} \int_0^{t_1} f^2(x) dx &= F^2(t_1)J(t_1, X), \\ -\frac{1}{2} \int_{t_1}^{t_2} f^2(x) dx &= (F(t_2) - F(t_1))^2 IJ(t_1, t_2), \\ -\frac{1}{2} \int_{t_2}^{+\infty} f^2(x) dx &= \bar{F}^2(t_2)J(X_{t_2}), \end{aligned}$$

and then we obtain the stated result. \square

Relation (4) shows that the uncertainty about the failure time of a component consists of 3 parts:

- 1st. The uncertainty of the failure time in $(0, t_1)$;
- 2nd. The uncertainty of the failure time in (t_1, t_2) ;
- 3rd. The uncertainty about the failure time in $(t_2, +\infty)$.

The corresponding aging classes are defined as follows.

Definition 1 The random variable X is said to have decreasing IJ property if and only if for any fixed t_2 , $IJ(t_1, t_2)$ is decreasing respect to t_1 .

Definition 2 The random variable X is said to have increasing IJ property if and only if for any fixed t_1 , $IJ(t_1, t_2)$ is increasing respect to t_2 .

Remark 2 As seen in Example 1, the exponential distribution satisfies both the conditions in Definitions 1 and 2.

Definition 3 Let X be a non-negative and absolutely continuous random variable with cdf F and pdf f . The Generalized Failure Rate (GFR) functions of X in t_1 and t_2 (with $F(t_2) - F(t_1) > 0$) are defined in [17] as

$$h_i(t_1, t_2) = \frac{f(t_i)}{F(t_2) - F(t_1)}, \quad i = 1, 2. \quad (6)$$

An upper bound in terms of GFR is obtained for Interval Entropy in the following theorem.

Theorem 2 *Let X be an absolutely continuous non-negative random variable. If IJ is increasing in t_2 , then*

$$IJ(t_1, t_2) \leq -\frac{h_2(t_1, t_2)}{4}. \quad (7)$$

Proof By differentiating IEx with respect to t_2 , we have

$$\frac{\partial IJ(t_1, t_2)}{\partial t_2} = -\frac{h_2^2(t_1, t_2)}{2} - 2h_2(t_1, t_2)IJ(t_1, t_2). \quad (8)$$

If $IJ(t_1, t_2)$ is increasing in t_2 then (8) implies (7). \square

In the following proposition, we analyze the effect of a linear transformation on the interval entropy.

Proposition 3 *Let X be a non negative and absolutely continuous random variable and let $Y = aX + b$ where $a > 0$ and $b \geq 0$. The interval entropy of Y is given in terms of the interval entropy of X as*

$$IJ_Y(t_1, t_2) = \frac{1}{a} IJ_X\left(\frac{t_1 - b}{a}, \frac{t_2 - b}{a}\right), \quad (9)$$

where $t_1, t_2 \in S_Y$.

Proof The cdf and the pdf of Y can be expressed in terms of F_X and f_X as

$$F_Y(x) = F_X\left(\frac{x - b}{a}\right), \quad f_Y(x) = \frac{1}{a} f_X\left(\frac{x - b}{a}\right). \quad (10)$$

Hence, the interval entropy of Y can be expressed as

$$\begin{aligned} IJ_Y(t_1, t_2) &= -\frac{1}{2\left(F_X\left(\frac{t_2 - b}{a}\right) - F_X\left(\frac{t_1 - b}{a}\right)\right)^2} \int_{t_1}^{t_2} \frac{1}{a^2} f_X^2\left(\frac{x - b}{a}\right) dx \\ &= -\frac{1}{2\left(F_X\left(\frac{t_2 - b}{a}\right) - F_X\left(\frac{t_1 - b}{a}\right)\right)^2} \int_{\frac{t_1 - b}{a}}^{\frac{t_2 - b}{a}} \frac{1}{a} f_X^2(x) dx \\ &= \frac{1}{a} IJ_X\left(\frac{t_1 - b}{a}, \frac{t_2 - b}{a}\right), \end{aligned}$$

which completes the proof. \square

In the following theorem, we give a characterization of the exponential distribution based on the interval entropy.

Theorem 4 Let X be a random variable with support $(0, +\infty)$, differentiable and strictly positive pdf f and cdf F . Then, X is exponentially distributed if, and only if, for all (t_1, t_2) such that $0 < t_1 < t_2 < +\infty$, the following relation holds

$$IJ(t_1, t_2) = -\frac{1}{4} [h_1(t_1, t_2) + h_2(t_1, t_2)]. \quad (11)$$

Proof Let us suppose $X \sim \text{Exp}(\lambda)$. In Example 1, we have evaluated the interval entropy that is given by

$$IJ(t_1, t_2) = -\frac{\lambda}{4} \cdot \frac{e^{-\lambda t_2} + e^{-\lambda t_1}}{e^{-\lambda t_1} - e^{-\lambda t_2}}.$$

Moreover, about GFR functions of X , we have

$$h_1(t_1, t_2) = \frac{\lambda e^{-\lambda t_1}}{e^{-\lambda t_1} - e^{-\lambda t_2}},$$

$$h_2(t_1, t_2) = \frac{\lambda e^{-\lambda t_2}}{e^{-\lambda t_1} - e^{-\lambda t_2}},$$

and then the first part of the proof is completed.

Conversely, let us suppose (11) holds. Then, by making explicit the interval entropy and GFR functions, we obtain

$$-\frac{1}{2(F(t_2) - F(t_1))^2} \int_{t_1}^{t_2} f^2(x) dx = -\frac{f(t_1) + f(t_2)}{4(F(t_2) - F(t_1))}.$$

From the above equation, we have

$$\int_{t_1}^{t_2} f^2(x) dx = \frac{1}{2}(F(t_2) - F(t_1))(f(t_1) + f(t_2)). \quad (12)$$

By differentiating both sides of (12) with respect to t_1 , we get

$$-f^2(t_1) = -\frac{1}{2}f(t_1)(f(t_1) + f(t_2)) + \frac{1}{2}f'(t_1)(F(t_2) - F(t_1)),$$

which reduces to

$$-f^2(t_1) + f(t_1)f(t_2) = f'(t_1)(F(t_2) - F(t_1)). \quad (13)$$

By differentiating both sides of (13) with respect to t_2 , we get

$$f(t_1)f'(t_2) = f'(t_1)f(t_2),$$

which is equivalent to

$$\frac{f'(t_1)}{f(t_1)} = \frac{f'(t_2)}{f(t_2)},$$

i.e., the ratio is constant for $x > 0$,

$$\frac{f'(x)}{f(x)} = A. \quad (14)$$

Hence, by integrating both sides of (14) from 0 to t , we get

$$f(t) = f(0) e^{At},$$

and in order to satisfy the condition of normalization for the pdf f , we need $A = -f(0)$, i.e., f is the pdf of an exponential distribution. \square

Remark 3 The equilibrium random variable Y associated to a renewal process with lifetime X is a random variable of primary interest in the context of reliability theory, as pointed out in Barlow and Proschan [1]. The survival function and the probability density function of Y are expressed as

$$\bar{F}_Y(t) = \frac{1}{\mathbb{E}(X)} \int_t^{+\infty} \bar{F}_X(x) dx, \quad f_Y(t) = \frac{\bar{F}_X(t)}{\mathbb{E}(X)}$$

where $\mathbb{E}(X)$ is the expectation of X . We can define Entropy and its interval version of Y as follows:

$$J(Y) = -\frac{1}{2\mathbb{E}^2(X)} \int_0^\infty \bar{F}_X^2(x) dx,$$

$$IJ_Y(t_1, t_2) = -\frac{1}{2} \frac{\int_{t_1}^{t_2} \bar{F}_X^2(x) dx}{\left(\int_{t_1}^{t_2} \bar{F}_X(x) dx\right)^2}.$$

3 Weighted Interval Entropy

In order to give importance to the value assumed by the random variable, it is significant to introduce weighted versions of the measures of uncertainty. In fact, most of the well-known measure of discrimination are position-free, in the sense that they assume the same value for X and $X + b$ for any $b \in \mathbb{R}$. In Proposition 3, we have showed that the interval entropy does not change under translations since, for $Y = X + b$, we have $IJ_Y(t_1 + b, t_2 + b) = IJ_X(t_1, t_2)$. In this section, we will introduce and study the weighted version of the interval entropy and we will show that it is not invariant under translations.

Suppose X be a non-negative absolutely continuous random variable. For all t_1 and t_2 such that $(t_1, t_2) \in D = \{(u, v) \in \mathbb{R}_+^2 : F(u) < F(v)\}$ we define the Weighted Interval Entropy (WIE_x) of X as

$$IJ^w(t_1, t_2) = -\frac{1}{2(F(t_2) - F(t_1))^2} \int_{t_1}^{t_2} x f^2(x) dx, \quad (15)$$

in the same way in which in [15] the weighted interval entropy has been defined.

Remark 4 We notice that

$$\lim_{t_1 \rightarrow 0} IJ^w(t_1, t_2) = IJ^w(t_2, X) \quad \text{and} \quad \lim_{t_2 \rightarrow +\infty} IJ^w(t_1, t_2) = IJ^w(X, t_1),$$

where $IJ^w(t_2, X)$ and $IJ^w(X, t_1)$ are Weighted Past Entropy at time t_2 and Weighted Residual Entropy at time t_1 , respectively.

Example 6 Let $X \sim \text{Exp}(\lambda)$, $\lambda > 0$. Based on (15), we evaluate the weighted interval entropy of X for $0 < t_1 < t_2 < +\infty$ and we obtain

$$IJ^w(t_1, t_2) = \frac{-1}{2(e^{-\lambda t_1} - e^{-\lambda t_2})^2} \int_{t_1}^{t_2} x \lambda^2 e^{-2\lambda x} dx$$




Fig. 6: Plot of IJ^w in Example 6 as a function of t_1 or t_2 fixing the other one with $t_i = 2$ (blue), 3 (red), 4 (yellow) and 5 (violet), $i = 1, 2$.

$$= -\frac{\lambda}{4} \cdot \frac{t_1 e^{-2\lambda t_1} - t_2 e^{-2\lambda t_2}}{(e^{-\lambda t_1} - e^{-\lambda t_2})^2} - \frac{1}{8} \cdot \frac{e^{-\lambda t_2} + e^{-\lambda t_1}}{e^{-\lambda t_1} - e^{-\lambda t_2}}.$$

In Figure 6, we plot the interval entropy as a function of t_1 for fixed t_2 (Figure 6(a)) and vice versa (Figure 6(b)) for $\lambda = 1$.

Example 7 Let X follow the Weibull distribution with parameters $\alpha = \lambda = 2$, $X \sim W2(2, 2)$. Based on (15), we evaluate the weighted interval entropy of X for $0 < t_1 < t_2 < +\infty$ and we obtain

$$\begin{aligned} IJ^w(t_1, t_2) &= \frac{-1}{2(\exp(-2t_1^2) - \exp(-2t_2^2))^2} \int_{t_1}^{t_2} 16x^3 \exp(-4x^2) dx \\ &= \frac{t_2^2 \exp(-4t_2^2) - t_1^2 \exp(-4t_1^2)}{(\exp(-2t_1^2) - \exp(-2t_2^2))^2} - \frac{1}{4} \cdot \frac{\exp(-2t_1^2) + \exp(-2t_2^2)}{\exp(-2t_1^2) - \exp(-2t_2^2)}. \end{aligned}$$

In Figure 7, we plot the interval entropy as a function of t_1 for fixed t_2 (Figure 7(a)) and vice versa (Figure 7(b)). From Figure 7(b) we observe an asymptotic behavior of the weighted interval entropy as $t_2 \rightarrow +\infty$, i.e., when the weighted interval entropy $IJ^w(t_1, t_2)$ reduces to the weighted residual entropy $J^w(X_{t_1})$. In fact, the weighted residual entropy of X in t can be expressed as

$$J(X_t) = -t^2 - \frac{1}{4}.$$

Example 8 Let X follow the Lognormal distribution with parameters $\mu = 0$, $\sigma^2 = 1$, $X \sim \text{Lognormal}(0, 1)$. Since the expression of the weighted interval entropy is not given in terms of elementary functions, in Figure 8, we plot the weighted interval entropy as a function of t_1 for fixed t_2 (Figure 8(a)) and vice versa (Figure 8(b)).

In the following theorem, we prove that the expression of the weighted entropy is composed of three terms in function of the weighted past, residual and interval entropies.




Fig. 7: Plot of IJ^w in Example 7 as a function of t_1 or t_2 fixing the other one with $t_i = 1$ (violet), 2 (blue), 3 (red) and 4 (yellow), $i = 1, 2$.




Fig. 8: Plot of IJ^w in Example 8 as a function of t_1 or t_2 fixing the other one with $t_i = 1$ (violet), 2 (blue), 3 (red) and 4 (yellow), $i = 1, 2$.

Theorem 5 Let X be a random variable denoting the lifetime of a component. For all $0 < t_1 < t_2 < +\infty$ the weighted entropy can be decomposed as follows:

$$J^w(t_1, t_2) = F^2(t_1)J^w(t_1, X) + (F(t_2) - F(t_1))^2 IJ^w(t_1, t_2) + \bar{F}^2(t_2)J^w(X, t_2),$$

i.e., Weighted Entropy is a function of WPE x , WRE x and WIE x .

Proof The proof is similar to the one of Theorem 1 and hence it is omitted. □

In the following proposition, we analyze the effect of a linear transformation on the weighted interval entropy.

Proposition 6 Let X be a non negative and absolutely continuous random variable and let $Y = aX + b$ where $a > 0$ and $b \geq 0$. The weighted interval entropy of Y is given in terms of the interval entropy and weighted interval entropy of X as

$$IJ_Y^w(t_1, t_2) = IJ_X^w\left(\frac{t_1 - b}{a}, \frac{t_2 - b}{a}\right) + \frac{b}{a} IJ_X\left(\frac{t_1 - b}{a}, \frac{t_2 - b}{a}\right), \quad (16)$$

where $t_1, t_2 \in S_Y$.

Proof By using the expressions of the cdf and the pdf of Y in terms of F_X and f_X obtained in (10), the weighted interval entropy of Y can be expressed as

$$\begin{aligned} IJ_Y^w(t_1, t_2) &= -\frac{1}{2\left(F_X\left(\frac{t_2-b}{a}\right) - F_X\left(\frac{t_1-b}{a}\right)\right)^2} \int_{t_1}^{t_2} \frac{x}{a^2} f_X^2\left(\frac{x-b}{a}\right) dx \\ &= -\frac{1}{2\left(F_X\left(\frac{t_2-b}{a}\right) - F_X\left(\frac{t_1-b}{a}\right)\right)^2} \int_{\frac{t_1-b}{a}}^{\frac{t_2-b}{a}} x f_X^2(x) dx \\ &\quad -\frac{1}{2\left(F_X\left(\frac{t_2-b}{a}\right) - F_X\left(\frac{t_1-b}{a}\right)\right)^2} \int_{\frac{t_1-b}{a}}^{\frac{t_2-b}{a}} \frac{b}{a} f_X^2(x) dx \\ &= IJ_X^w\left(\frac{t_1 - b}{a}, \frac{t_2 - b}{a}\right) + \frac{b}{a} IJ_X\left(\frac{t_1 - b}{a}, \frac{t_2 - b}{a}\right), \end{aligned}$$

which completes the proof. \square

Remark 5 The results given in Propositions 3 and 6 about linear transformations could be generalized to monotonic transformations but, in these cases, we do not obtain a formula of interest, in the sense that the interval entropy and the weighted interval entropy of the transformed random variable are not expressed in terms of the ones of the original random variable.

In the following theorem, we present an upper bound for the weighted interval entropy given in terms of the generalized failure rate function.

Theorem 7 For an absolutely continuous non-negative random variable X , if the WIE x is increasing in t_2 , then we have

$$IJ^w(t_1, t_2) \leq -\frac{t_2 h_2(t_1, t_2)}{4}. \quad (17)$$

Proof The proof follows in analogy with the one of Theorem 2 and hence it is omitted. \square

4 Conclusion

In this paper dynamic versions of entropy for double truncated random variables have been presented. Several examples are given. The behavior under linear transformations of these new measures has been studied. Some bounds for them have been found in relation with the Generalized Failure Rate.

Acknowledgments. Francesco Buono and Maria Longobardi are members of the research group GNAMPA of INDAM (Istituto Nazionale di Alta Matematica), are partially supported by MIUR - PRIN 2017, project “Stochastic Models for Complex Systems”, no. 2017 JFFHSH. The present work was developed within the activities of the project 000009_ALTRLCDA_75_2021_FRA_LINEA_B.SIMONELLI funded by “Programma per il finanziamento della ricerca di Ateneo - Linea B” of the University of Naples Federico II.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- [1] Barlow, R. E., and F.J. Proschan (1996). *Mathematical Theory of Reliability*. Philadelphia: Society for Industrial and Applied Mathematics.
- [2] Balakrishnan, N., Buono, F., Longobardi, M. (2020). On weighted entropies. *Communications in Statistics-Theory and Methods*, DOI: 10.1080/03610926.2020.1860222.
- [3] Balakrishnan, N., Buono, F., Longobardi, M. (2022). On Tsallis entropy with an application to pattern recognition. *Statistics and Probability Letters*, <https://doi.org/10.1016/j.spl.2021.109241>.
- [4] Becerra, A., de la Rosa, J.I., González, E. et al. Training deep neural networks with non-uniform frame-level cost function for automatic speech recognition. *Multimed Tools Appl* 77, 27231–27267 (2018). <https://doi.org/10.1007/s11042-018-5917-5>.
- [5] Betensky, R. A. and Martin, E. C. (2003). Commentary: Failure-rate functions for doubly truncated random variables, *IEEE Transactions on Reliability*, 52(1), 7–8.
- [6] Di Crescenzo, A., Longobardi, M. (2002). Entropy-based measure of uncertainty in past lifetime distributions. *J. Appl. Probab.* 39: 434–440.
- [7] Di Crescenzo, A., Longobardi, M. (2006). On weighted residual and past entropies. *Scientiae Mathematicae Japonicae*, 64 (2), 255–266.

- [8] Ebrahimi, N (1996). How to measure uncertainty about residual lifetime. *Sankhya A*, 58, 48–57.
- [9] Kamari, O., Buono, F. (2020). On extropy of past lifetime distribution. *Ricerche di Matematica*, DOI:10.1007/s11587-020-00488-7.
- [10] Kazemi, M.R., Tahmasebi, S., Buono, F., Longobardi, M. Fractional Deng Entropy and Extropy and Some Applications. *Entropy* 2021, 23, 623. <https://doi.org/10.3390/e23050623>.
- [11] Khorashadizadeh, M.; Rezaei Roknabadi, A. H. and Mohtashami Borzadaran, G. R. (2012). Characterizations of lifetime distributions based on doubly truncated mean residual life and mean past to failure, *Communications in Statistics - Theory and Methods*, 41(6), 1105–1115.
- [12] Krishnan, A. S., Sunoj, S. M., Nair, N. U. (2020). Some reliability properties of extropy for residual and past lifetime random variables. *Journal of the Korean Statistical Society*, 49(2), 457–474.
- [13] Lad, F., Sanfilippo, G., Agrò, G. (2015). Extropy: complementary dual of entropy. *Statistical Science*, 30(1), 40–58.
- [14] Martinas, K., Frankowicz, M. (2000). Extropy-reformulation of the entropy principle. *Periodica Polytechnica Chemical Engineering*, 44(1): 29–38.
- [15] Misagh, F., Yari, G. H. (2011). On weighted interval entropy. *Statistics and Probability Letters*, 81(2), 188–194.
- [16] Misagh, F., Yari, G. (2012). Interval entropy and informative distance. *Entropy*, 14(3), 480-490.
- [17] Navarro, J., Ruiz, J.M. (1996). Failure-rate function for doubly-truncated random variables. *IEEE Transactions on Reliability*, 4 , 685–690.
- [18] Poursaeed, M. H. and Nematollahi, A. R. (2008). On the mean past and the mean residual life under double monitoring, *Communications in Statistics - Theory and Methods*, 37(7), 1119–1133.
- [19] Qiu, G., Jia, K. (2018). The residual extropy of order statistics. *Stat. Probab. Lett.* 133, 15–22.
- [20] C. E. Shannon, (1948). A mathematical theory of communication. *Bell System Technical J* , 27, 379–423.
- [21] S. M. Sunoj, P. G. Sankaran, S. S. Maya (2009) Characterizations of Life Distributions Using Conditional Expectations of Doubly (Interval) Truncated Random Variables, *Communications in Statistics - Theory and*

