

Parallel Kahler submanifolds of Quaternionic Kahler Symmetric Spaces

*Original*

Parallel Kahler submanifolds of Quaternionic Kahler Symmetric Spaces / DI SCALA, ANTONIO JOSE'; DMITRI V., Alekseevsky; Stefano, Marchiafava. - In: TOHOKU MATHEMATICAL JOURNAL. - ISSN 0040-8735. - 57:(2005), pp. 521-540.

*Availability:*

This version is available at: 11583/1399747 since:

*Publisher:*

*Published*

DOI:

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

# Parallel Kähler submanifolds of quaternionic Kähler symmetric spaces

DMITRI VLADIMIR ALEKSEEVSKY <sup>\*</sup>, ANTONIO JOSÉ DI SCALA <sup>\*</sup>  
AND STEFANO MARCHIAFAVA <sup>\*</sup>

Post print (i.e. final draft post-refereeing) version of an article published on *Tohoku Math. J. (2) Volume 57, Number 4 (2005)*, 521–540. , doi:10.2748/tmj/1140727071 .

Beyond the journal formatting, please note that there could be minor changes from this document to the final published version. The final published version is accessible from here:

<http://projecteuclid.org/euclid.tmj/1140727071>

This document has made accessible through PORTO, the Open Access Repository of Politecnico di Torino (<http://porto.polito.it>) following the editors requirements.

## Abstract

The non totally geodesic parallel  $2n$ -dimensional Kähler submanifolds of the  $n$ -dimensional quaternionic projective space were classified by K. Tsukada. Here we give the complete classification of non totally geodesic immersions of parallel  $2m$ -dimensional Kähler submanifolds in a quaternionic Kähler symmetric space of non zero scalar curvature, i.e., in a Wolf space  $W$  or in its non compact dual. They are exhausted by parallel Kähler submanifolds of a totally geodesic submanifold which is either an Hermitian symmetric space or a quaternionic projective space.

## 1 Introduction.

Let  $(\widetilde{M}^{4n}, \widetilde{g}, Q)$  be a quaternionic Kähler manifold with metric  $\widetilde{g}$  and parallel quaternionic structure  $Q$ . A submanifold  $M^{2m}$  together with a section  $J_1 \in \Gamma(Q)|_M$  such that  $J_1^2 =$

---

2000 *Mathematics Subject Classification*. Primary: 53C26; Secondary: 53C55, 53C40.

*Key words and phrases*. quaternionic Kähler manifolds, Kähler submanifolds, totally complex submanifolds

<sup>\*</sup> Supported by a EPSRC Grant GR/R69174, by Italian MURST, under the national project "Geometria delle varietà reali e complesse", and by G.N.S.A.G.A. of C.N.R. (Italy).

The third author is a member of EDGE, Research Training Network HPRN-CT-2000-00101, supported by the European Human Potential Programme.

$-1$  and  $J_1 TM = TM$  is called *Kähler* if  $J_1$  is parallel with respect to the Levi-Civita connection of  $\tilde{g}$ . We will study parallel Kähler submanifolds of a quaternionic Kähler symmetric space  $\widetilde{M}$  of *non zero scalar curvature*, that is, Kähler submanifolds  $M$  with parallel second fundamental form  $h$  in a Wolf space or in its non compact dual. In the case when  $\dim(\widetilde{M}) = 2 \dim M$ , we prove that *any curvature invariant and intrinsically locally symmetric Kähler submanifold is parallel, and hence extrinsically symmetric*.

Any parallel submanifold  $M$  of a Riemannian manifold  $\widetilde{M}$  is curvature invariant. Furthermore, a curvature invariant, in particular a parallel, maximal Kähler submanifold of a quaternionic Kähler manifold is also normal curvature invariant. Using these properties, we derive the following result from Naitoh's theorem 2.6 in the next section.

**Theorem 1.1.** *Any curvature invariant (in particular, any parallel) Kähler submanifold  $M^{2n}$  of the maximal dimension  $2n$  of a quaternionic Kähler symmetric space  $\widetilde{M}^{4n}$  different from the  $n$ -dimensional quaternionic projective space  $\mathbb{H}P^n$ ,  $\widetilde{M}^{4n} \neq \mathbb{H}P^n$ , is totally geodesic.*

We recall that a submanifold  $M$  of a Riemannian manifold  $\widetilde{M}$  is called *full* if  $M$  is not contained in a proper totally geodesic submanifold  $\overline{M}$  of  $\widetilde{M}$  and is called *1-full* (according to Tsukada [Tsu1]) if the first normal bundle  $N^1 M = h(TM, TM)$  of  $M$  coincides with the normal bundle  $T^\perp M$  of  $M$  in  $\widetilde{M}$ .

We associate with a Kähler submanifold  $M^{2m}$  of  $\widetilde{M}^{4n}$ , of arbitrary dimension  $2m$ , a symmetric 3-form  $C$ , called the *shape tensor*, and prove the following theorem.

**Theorem 1.2.** *Let  $(M^{2m}, J)$  be a geodesically complete parallel Kähler submanifold of a quaternionic Kähler symmetric space  $\widetilde{M}^{4n}$  and  $\overline{M}$  the minimal totally geodesic submanifold of  $\widetilde{M}$  containing  $M$ .*

- 1) *If the shape tensor  $C$  of  $M$  vanishes at one point, then  $\overline{M}$  is an Hermitian symmetric space and  $M$  is a full parallel Kähler submanifold of  $\overline{M}$ .*
- 2) *If  $C \neq 0$ , then  $\overline{M} = \mathbb{H}P^m$  and  $(M^{2m}, J)$  is a Hermitian symmetric manifold with parallel cubic line bundle, that is a product  $Q_{m-1} \times \mathbb{C}P^1$  of the complex quadric  $Q_{m-1} \subset \mathbb{C}P^m$  and the projective line  $\mathbb{C}P^1$ , or one of the following exceptional Hermitian symmetric spaces:  $\mathbb{C}P^1 \times \mathbb{C}P^1$ ,  $\mathbb{C}P^1 \times \mathbb{C}P^1 \times \mathbb{C}P^1$ ,  $Sp_2/U_2 \times \mathbb{C}P^1$ ,  $\mathbb{C}P^1$ ,  $Sp_3/U_3$ ,  $SU_6/S(U_3 \times U_3)$ ,  $SO_{12}/U_6$ ,  $E_7/T^1 \cdot E_6$ , with the canonical Tsukada imbedding into  $\mathbb{H}P^m$  as described in [Tsu2].*

Thus, the classification of parallel Kähler submanifolds of type 1) in a quaternionic Kähler symmetric space reduces to a description of parallel Kähler submanifolds of Hermitian symmetric spaces.

The classification of parallel Kähler submanifolds of  $\mathbb{C}P^N$  was first obtained by Nakagawa and Takagi [NT].

**Theorem 1.3.** ([NT]) *The only full parallel Kähler submanifolds of a complex projective space are, up to isometries, the images of the Veronese imbedding of the projective space  $PV$  associated with  $V = \mathbb{C}^{n+1}$  into the projectivization  $PS^2V$  of the symmetric square  $S^2V$  defined by*

$$\begin{aligned}\varphi : \mathbb{C}P^n = PV &\rightarrow PS^2V \\ [v] = \mathbb{C}v &\mapsto [v \otimes v],\end{aligned}$$

*of the Segre imbedding defined by*

$$\begin{aligned}\psi : \mathbb{C}P^n \times \mathbb{C}P^{n'} = PV \times PV' &\rightarrow P(V \otimes V') \\ ([v], [v']) &\mapsto [v \otimes v'],\end{aligned}$$

*or of the first canonical imbedding of compact irreducible Hermitian symmetric spaces of rank 2, i.e.,  $Q_n$ ,  $Gr_2(\mathbb{C}^{n+2})$ ,  $SO_{10}/SU_5$  and  $E_6/Spin_{10} \cdot T$ .*

The classification of all parallel Kähler submanifolds of a Hermitian symmetric space was established by Tsukada [Tsu1]. He proved that any such submanifold is a product of Veronese submanifolds, Segre submanifolds, canonical Kaehler imbeddings of compact Hermitian symmetric spaces of rank two and trivial factors (defined by the identity map). The Theorem in [Tsu1, p.130] implies the following

**Theorem 1.4.** *There is no full parallel (proper) Kähler submanifold  $M$  in a Hermitian symmetric space  $\widetilde{M}$  having no factor isometric to  $\mathbb{C}P^N$ . Any full parallel Kähler submanifold of  $\mathbb{C}P^{n_1} \times \mathbb{C}P^{n_2}$  has the form  $\psi_1(M_1) \times \psi_2(M_2)$ , where  $\psi_i(M_i) \subset \mathbb{C}P^{n_i}$  is one of the immersions in Theorem 1.3.*

Tsukada [Tsu1] proved that any parallel Kähler submanifold of a Hermitian symmetric space of non compact type is totally geodesic.

These results together give the full classification of non totally geodesic parallel Kähler submanifolds in a quaternionic Kähler symmetric space. A classification of maximal totally geodesic Kähler submanifolds of Wolf spaces in term of Satake diagrams was given by Takeuchi [Tak]. See also Section 6.

## 2 Preliminaries.

### 2.1 Gauss, Codazzi-Mainardi and Ricci equations

Let  $M$  be a submanifold of a Riemannian manifold  $\widetilde{M}$ . We denote by  $h : TM \times TM \rightarrow T^\perp M$  the second fundamental form of  $M$ , and by  $A^\xi$  the shape operator in the direction

of a normal vector  $\xi \in T^\perp M$  such that

$$\begin{aligned}\widetilde{\nabla}_X Y &= \nabla_X Y + h(X, Y) , \\ \widetilde{\nabla}_X \xi &= \nabla_X^\perp \xi - A^\xi X ,\end{aligned}$$

where  $X \in TM, Y \in \Gamma TM$  and  $\xi \in T^\perp M$ . Here  $\widetilde{\nabla}, \nabla, \nabla^\perp$  are the Levi-Civita connection of  $\widetilde{M}$  and the induced connections in  $TM$  and  $T^\perp M$ , respectively.

For  $X, Y \in T_x M$  we decompose the curvature operator  $\widetilde{R}_{X,Y}$  as

$$\widetilde{R}_{XY} = R_{XY}^{TT} + R_{XY}^{\perp T} + R_{XY}^{T\perp} + R_{XY}^{\perp\perp} ,$$

according to the decomposition

$$\text{End}(T_x \widetilde{M}) = \text{End}(T_x M) + \text{Hom}(T_x M, T_x^\perp M) + \text{Hom}(T_x^\perp M, T_x M) + \text{End}(T_x^\perp M).$$

Then we have the following *Gauss-Codazzi equations*:

$$(\text{Gauss}) \quad R_{XY}^{\top\top} = R_{XY} - h_X h_Y^t + h_Y h_X^t = R_{XY} - \sum_i A^{\xi_i} X \wedge A^{\xi_i} Y ,$$

$$(\text{Codazzi-Mainardi}) \quad R_{XY}^{\perp\top} Z = (\nabla'_X h)(Y, Z) - (\nabla'_Y h)(X, Z) ,$$

$$(\text{Ricci}) \quad R_{XY}^{\perp\perp} \xi = R_{XY}^\perp \xi - \sum_i \langle X, [A^{\xi_i}, A^\xi] Y \rangle \xi_i ,$$

where  $\xi_i$  is an orthonormal basis of  $T^\perp M$ ,  $X, Y \in TM$ ,  $\xi \in T^\perp M$ ,  $R, R^\perp$  are the curvature tensors of the connections  $\nabla, \nabla^\perp$ , and  $\nabla'$  is the connection in  $T^\perp M \otimes S^2 TM$  induced by  $\nabla^\perp$  and  $\nabla$ , respectively. (We identify a bivector  $X \wedge Y$  with the skew-symmetric operator  $Z \mapsto \langle Y, Z \rangle X - \langle X, Z \rangle Y$ .)

## 2.2 Parallel submanifolds of symmetric spaces

**Definition 2.1.** A submanifold  $M$  of a Riemannian manifold  $\widetilde{M}$  is called *parallel* if it has parallel second fundamental form  $h$ , i.e.,  $\nabla' h = 0$ .

**Definition 2.2.** A subspace  $V \subset T_x \widetilde{M}$  of a tangent space of a Riemannian manifold  $\widetilde{M}$  is called *curvature invariant* if

$$\widetilde{R}(V, V)V \subset V .$$

A submanifold  $M$  of  $\widetilde{M}$  is called *curvature invariant* if each tangent space  $T_x M$  is curvature invariant and it is called *normal curvature invariant* if each normal space  $T_x^\perp M$  is curvature invariant.

It follows from Codazzi-Mainardi equation that any parallel submanifold  $M$  of a Riemannian manifold  $\widetilde{M}$  is curvature invariant.

**Definition 2.3.** A submanifold  $M$  of a Riemannian manifold  $\widetilde{M}$  is called *1-full* if the first normal bundle  $N^1M = h(TM, TM)$  coincides with the normal bundle  $T^\perp M$ .

**Definition 2.4.** Let  $\widetilde{M} = G/K$  be a homogeneous Riemannian manifold. Fix an orbit  $\mathcal{V}$  of the isometry group  $G$  in the Grassmann bundle  $\text{Gr}_k(T\widetilde{M})$  of tangent  $k$ -planes of  $\widetilde{M}$ . If a  $k$ -plane  $V \in \mathcal{V}$  (respectively, if the orthogonal plane  $V^\perp, V \in \mathcal{V}$ ) is curvature invariant, then  $\mathcal{V}$  is called *curvature invariant* (respectively, *normal curvature invariant*).

A  $k$ -dimensional submanifold  $M \subset \widetilde{M}$  is called a  $\mathcal{V}$ -submanifold if  $T_x M \in \mathcal{V}$  for any  $x \in M$ . Obviously, if  $\mathcal{V}$  is (normal) curvature invariant, then any  $\mathcal{V}$ -submanifold is (normal) curvature invariant.

**Definition 2.5.** A submanifold  $M$  of a Riemannian manifold  $\widetilde{M}$  is called *extrinsically symmetric* if for any point  $x \in M$  there exists an involutive isometry (symmetry)  $s_x$  of  $\widetilde{M}$  preserving  $M$  such that  $s_x(x) = x$  and its differential at  $x$  satisfies

$$(1) \quad (s_x)_*|_{T_x M} = -\text{Id}, \quad (s_x)_*|_{T_x^\perp M} = \text{Id}.$$

We recall the following theorem of Naitoh [Na2].

**Theorem 2.6** (H. Naitoh). *Let  $\widetilde{M}$  be a simply connected Riemannian symmetric space. A submanifold  $M$  of  $\widetilde{M}$  is parallel and normal curvature invariant if and only if it is extrinsically symmetric.*

*Proof.* Let  $M \subset \widetilde{M}$  be an extrinsically symmetric submanifold. Remark that the symmetry  $s_x$  acts as  $-\text{Id}$  on any tensor space  $T_x^{\otimes p} \otimes T_x^{\perp \otimes q}$ , where  $p$  is odd. On the other hand, it preserves the tensor  $\nabla' h \in T_x^{\otimes 3} \otimes T_x^\perp$  and the curvature tensor  $\widetilde{R}$  at  $x$ . This implies that an extrinsically symmetric submanifold is parallel and normal curvature invariant. Conversely, if  $M$  is parallel and normal curvature invariant, then the automorphism  $(s_x)_* \in \text{Gl}(T_x \widetilde{M})$  defined by (1) preserves the curvature tensor  $\widetilde{R}_x$ , and hence can be extended to an involutive isometry  $s$  of  $\widetilde{M}$ . Now the inverse statement follows from a remarkable theorem of Strübing [Str].

**Theorem 2.7** (W. Strübing). *Let  $M$  be a parallel submanifold of a Riemannian manifold  $\widetilde{M}$  and  $s$  an isometry of  $\widetilde{M}$  which preserves a point  $x \in M$  and satisfies (1). Then  $s$  preserves any geodesic  $\gamma = \gamma(t)$  of  $M$  with  $\gamma(0) = x$ :  $s(\gamma(t)) = \gamma(-t)$ .*

The proof follows from the Frenet formulas for the curve  $\gamma(t)$  considered as a curve in

$\widetilde{M}$ :

$$\widetilde{\nabla}_{\dot{\gamma}} \begin{pmatrix} E_1 \\ \cdot \\ \cdot \\ \cdot \\ E_r \end{pmatrix} = \begin{pmatrix} 0 & k_1 & 0 & \dots & 0 & 0 \\ -k_1 & 0 & k_2 & \dots & 0 & 0 \\ 0 & -k_2 & 0 & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 0 & k_{r-1} \\ 0 & 0 & 0 & \dots & -k_{r-1} & 0 \end{pmatrix} \begin{pmatrix} E_1 \\ \cdot \\ \cdot \\ \cdot \\ E_r \end{pmatrix},$$

where  $E_1, \dots, E_r$  is an orthonormal Frenet frame along  $\gamma$  obtained from the fields  $\dot{\gamma}, \ddot{\gamma}, \ddot{\ddot{\gamma}}, \dots$  by Gram-Schmidt process,  $k_1, \dots, k_{r-1}$  are constants ("curvatures") and, moreover,  $E_1 = \dot{\gamma}, E_3, E_5, \dots \in \Gamma(TM)|_{\gamma}$  are  $\nabla$ -parallel fields and  $E_2 = h(\dot{\gamma}, \dot{\gamma})/|h(\dot{\gamma}, \dot{\gamma})|, E_4, E_6, \dots \in \Gamma(T^{\perp}M)|_{\gamma}$  are  $\nabla^{\perp}$ -parallel fields along  $\gamma$ . Indeed, the Frenet frame along  $\gamma(-t)$  and  $s_*E_i(t)$  satisfy Frenet equations with the same initial conditions  $(-1)^i E_i(0)$ . q.e.d.

Now we state the following fundamental result by Naitoh, which shows that up to a short list of exceptions, a parallel normal curvature invariant (or, equivalently, extrinsically symmetric)  $\mathcal{V}$ -submanifold of a symmetric space is in fact totally geodesic.

**Theorem 2.8.** (H. Naitoh[Na3]) Let  $\widetilde{M} = G/K$  be a compact simply connected symmetric space with simple isometry group  $G$ , and  $\mathcal{V}$  is an orbit of  $G$  in  $\text{Gr}_k(\widetilde{TM})$  which is curvature invariant and normal curvature invariant. Then any  $\mathcal{V}$ -submanifold is totally geodesic with the exception of the following cases:

- (a)  $\widetilde{M} = S^n = SO(n+1)/SO(n), 1 \leq k < n,$
- (b)  $\widetilde{M} = \mathbb{C}P^n, \mathcal{V}$  is the set of complex  $2k$ -subspaces,
- (c)  $\widetilde{M} = \mathbb{C}P^n, \mathcal{V}$  is the set of totally real  $n$ -subspaces,
- (d)  $\widetilde{M} = \mathbb{H}P^n, \mathcal{V}$  is the set of totally complex  $2n$ -subspaces,
- (e)  $\widetilde{M} = G/K$  is an irreducible symmetric space and  $\mathcal{V} = GT$ , where  $T$  is the tangent space to an irreducible symmetric  $R$ -space (i.e., the geometries associated with irreducible symmetric  $R$ -spaces).

The statement remains true also for non compact dual of  $G/K$ , [BENT].

The following result will be used in Section 5.

**Theorem 2.9.** (H. Naitoh[Na4]) Let  $M$  be a parallel submanifold of a symmetric space  $\widetilde{M}$ . If the first osculating space  $O_x^1 M = T_x M + h(T_x, T_x)$  at some point  $x \in M$  is curvature invariant, then  $M$  is contained in the totally geodesic submanifold  $\overline{M} = \exp(O_x^1 M)$  of  $\widetilde{M}$  generated by  $O_x^1 M$ .

Obviously,  $M$  is full in  $\overline{M}$ .

### 3 Kähler submanifolds of quaternionic Kähler manifolds

Let  $(\widetilde{M}^{4n}, Q, \widetilde{g})$  be a quaternionic Kähler manifold, that is, a Riemannian manifold  $(\widetilde{M}^{4n}, \widetilde{g})$  with a  $\widetilde{\nabla}$ -parallel quaternionic structure  $Q$ , i.e., a rank 3 subbundle of  $\text{End}(T\widetilde{M})$  locally generated by 3 skew-symmetric almost complex structures  $J_1, J_2, J_3 = J_1 J_2 = -J_2 J_1$ . For  $n = 1$ , in the definition we assume that  $(\widetilde{M}^4, \widetilde{g})$  is an anti-self-dual Einstein manifold.

Recall that the curvature tensor  $\widetilde{R}$  of a quaternionic Kähler manifold has the form

$$\widetilde{R} = \nu R_{\mathbb{H}P^n} + \widetilde{W} ,$$

where  $\widetilde{W}$  is an  $\mathfrak{sp}_n$ -valued 2-form satisfying the Bianchi identities (the *quaternionic Weyl tensor*),  $\nu = K/4n(n+2)$  is the *reduced scalar curvature*, which is proportional to the scalar curvature  $K$ , and

$$R_{\mathbb{H}P^n}(X, Y) = \frac{1}{4}(X \wedge Y + \sum J_\alpha X \wedge J_\alpha Y - 2 \sum_\alpha \langle J_\alpha X, Y \rangle J_\alpha) ,$$

where  $\alpha = 1, 2, 3$  and  $\langle \cdot, \cdot \rangle = \widetilde{g}(\cdot, \cdot)$ .

We recall also that the following identities hold:

$$[\widetilde{R}(X, Y), J_\alpha] = -\nu(\langle J_\gamma X, Y \rangle J_\beta - \langle J_\beta X, Y \rangle J_\gamma) ,$$

where  $(\alpha, \beta, \gamma)$  is a cyclic permutation of  $(1, 2, 3)$ . They are equivalent to the following identities

$$(2) \quad \widetilde{R}(J_\alpha X, J_\alpha Y)Z = \widetilde{R}(X, Y)Z + \nu(\langle J_\beta X, Y \rangle J_\beta Z + \langle J_\gamma X, Y \rangle J_\gamma Z) ,$$

which we will need later on.

**Definition 3.1.** A submanifold  $M^{2m}$  of a quaternionic Kähler manifold  $(\widetilde{M}^{4n}, Q, \widetilde{g})$  together with a section  $J_1 \in \Gamma(Q)|_M$  such that  $J_1^2 = -Id$  and  $J_1(TM) = TM$  is called

- 1) a *Kähler submanifold* if  $J_1$  is  $\widetilde{\nabla}$ -parallel,
- 2) a *totally complex submanifold* if  $J_2(TM) \perp TM$ , where  $J_2 \in Q$  is a complex structure anticommuting with  $J_1$ .

The Kähler submanifold  $M^{2m}$  considered as a manifold with the induced Riemannian metric  $g = \widetilde{g}|_M$  and the almost complex structure  $J = J_1|_{TM}$  is a Kähler manifold.

Recall that if the scalar curvature of  $(\widetilde{M}, \widetilde{g})$  is not zero then a Kähler submanifold  $M^{2m}$ ,  $m > 1$ , is totally complex ([AM2]). In particular,  $m \leq n$ . A Kähler submanifold of maximal possible dimension  $2n$  is called *maximal*.



Let  $(M^{2m}, J_1)$  be a Kähler submanifold of a quaternionic Kähler manifold  $\widetilde{M}^{4n}$ . We fix a local section  $J_2 \in \Gamma(Q)|_M$  such that  $J_2^2 = -1$  and  $J_1 J_2 = -J_2 J_1$ . One can check that

$$(3) \quad \widetilde{\nabla}_V J_2 = \omega(V) J_3 ,$$

where  $J_3 = J_1 J_2$  and  $\omega$  is a local 1-form on  $M$ . As in [AM1], we associate with the second fundamental form  $h$ , a (local)  $(0, 3)$ -tensor field  $C$  on  $M$ , called the *shape tensor*, defined by

$$C(X, Y, Z) := \langle J_2 h(X, Y), Z \rangle .$$

It is symmetric with respect to  $X, Y, Z$  and satisfies the following identities:

$$C(X, Y, JZ) = C(JX, Y, Z) = C(X, JY, Z) ,$$

which means that the associated endomorphism  $C_X$ ,  $X \in TM$ , defined by

$$\langle C_X Y, Z \rangle = C(X, Y, Z)$$

anticommutes with  $J$ .

If  $J'_2 = \cos \theta J_2 + \sin \theta J_1 J_2$  is another section, then the associated shape tensor  $C'$  is related to  $C$  by

$$C'_X = \cos \theta C_X + \sin \theta J_1 \circ C_X .$$

This implies that the  $\mathfrak{u}_m$ -valued 2-form  $[C, C](X, Y) := [C_X, C_Y]$  is globally defined and satisfies the Bianchi identities.

We define the  $(0, 4)$ -tensor field  $P$  as follows:

$$P(V; X, Y, Z) = (\nabla_V C)(X, Y, Z) + \omega(V) C(X, Y, JZ),$$

which is symmetric with respect to  $X, Y, Z$ .

**Proposition 3.2.** *Let  $(M^{2m}, J_1)$  be a curvature invariant Kähler submanifold of a quaternionic Kähler symmetric space. Then*

- 1) *the tangential part  $R^{TT}$  of the curvature tensor  $\widetilde{R}$  of  $\widetilde{M}$  is parallel and the tensor  $[C, C]$  satisfies the second Bianchi identity:*

$$\nabla R^{TT} = 0 , \quad \text{cycl}(\nabla_Z [C, C])(X, Y) = 0 ,$$

- 2) *If  $M$  is parallel, then  $P \equiv 0$ .*

*Proof.* The proof is the same as for the case  $n = m$ , which was done in [AM1]. q.e.d.

The following Lemma describes the relation between the covariant derivative of  $C$  and the tensor  $P$ .

**Lemma 3.3.** *Let  $(M^{2m}, J_1)$  be a totally complex submanifold of a quaternionic Kähler manifold  $\widetilde{M}^{4n}$ . Then the covariant derivative of the shape tensor  $C$  is given by*

$$-(\nabla_V C)(X, Y, Z) = \langle (\nabla'_V h)(X, Y), J_2 Z \rangle + \omega(V)C(X, Y, JZ) + \langle h(X, Y), J_2 h(V, Z) \rangle$$

or, equivalently,

$$(4) \quad -P(V; X, Y, Z) = \langle (\nabla'_V h)(X, Y), J_2 Z \rangle + \langle h(X, Y), J_2 h(V, Z) \rangle$$

for any tangent vectors  $X, Y, Z, V$ .

*Proof.* We extend vectors  $X, Y, Z \in T_x M$  to local tangent vector fields on  $M$  such that  $\nabla_V X = \nabla_V Y = \nabla_V Z = 0$  at  $x \in M$ . Then we have

$$\begin{aligned} -(\nabla_V C)(X, Y, Z) &= -VC(X, Y, Z) = V\langle h(X, Y), J_2 Z \rangle \\ &= \langle \nabla_V^\perp h(X, Y), J_2 Z \rangle + \langle h(X, Y), \nabla_V^\perp J_2 Z \rangle \\ &= \langle (\nabla'_V h)(X, Y), J_2 Z \rangle + \langle h(X, Y), \widetilde{\nabla}_V J_2 Z \rangle \\ &= \langle (\nabla'_V h)(X, Y), J_2 Z \rangle + \langle h(X, Y), (\widetilde{\nabla}_V J_2)Z + J_2 \widetilde{\nabla}_V Z \rangle \\ &= \langle (\nabla'_V h)(X, Y), J_2 Z \rangle + \langle h(X, Y), \omega(V)J_3 Z + J_2 h(V, Z) \rangle \\ &= \langle (\nabla'_V h)(X, Y), J_2 Z \rangle + \langle h(X, Y), -\omega(V)J_2 J_1 Z + J_2 h(V, Z) \rangle \\ &= \langle (\nabla'_V h)(X, Y), J_2 Z \rangle + \omega(V)C(X, Y, JZ) + \langle h(X, Y), J_2 h(V, Z) \rangle. \end{aligned}$$

q.e.d.

**Corollary 3.4.** 1) Assume that at some point  $x \in M$  the subspace  $(\nabla'_{T_x M} h)(T_x M, T_x M)$  is orthogonal to  $J_2 T_x M$ . Then  $P_x = 0$  and the first normal space  $N_x^1 = h(T_x M, T_x M)$  is totally complex, i.e.,  $J_1 N_x^1 = N_x^1$  and  $J_2 N_x^1$  is orthogonal to  $N_{1x}$ .

2) Assume that  $M$  is curvature invariant and the first normal space  $N_x^1$  at some point  $x \in M$  is totally complex. Then  $P_x(V; X, Y, Z) = \langle (\nabla'_V h)_x(X, Y), J_2 Z \rangle$  is symmetric in all arguments.

*Proof.* 1) The first term on the right member of (4) vanishes. Hence  $P_x(V; X, Y, Z) = -\langle h(X, Y), J_2 h(V, Z) \rangle$  is symmetric in all arguments. Since  $P_x(X, X, X, X) = \langle -h(X, X), J_2 h(X, X) \rangle = 0$ , we get the conclusion.

2) By taking Codazzi-Mainardi equation into account, it is obvious. q.e.d.

**Theorem 3.5.** *Let  $(M^{2m}, J_1)$  be a totally complex submanifold of a quaternionic Kähler manifold  $\widetilde{M}^{4n}$ . Assume that  $\langle (\nabla'_V h)(X, Y), J_2 Z \rangle = 0$  for any  $X, Y, Z, V \in TM$ , which is true if  $M$  is parallel. Then the first normal bundle  $N^1 M = h(TM, TM)$  is totally complex, i.e.,  $\langle h(X, Y), J_2 h(V, Z) \rangle = 0$  and the tensor field  $P = 0$ .*

Assume moreover that the reduced scalar curvature  $\nu$  of  $\widetilde{M}^{4n}$  is not zero. Then there are two cases:

- 1)  $C = 0$  at some point and then  $C \equiv 0$ , which means that  $N^1M \perp J_2TM$ , or
- 2)  $C \neq 0$  and then  $M$  is a locally symmetric Hermitian manifold with parallel cubic line bundle of type  $\nu$  ([AM1]). More precisely,  $M$  is locally isometric to one of the symmetric spaces:  $S = Q_{n-1} \times \mathbb{C}P^1, \mathbb{C}P^1 \times \mathbb{C}P^1, \mathbb{C}P^1 \times \mathbb{C}P^1 \times \mathbb{C}P^1, Sp_2/U_2 \times \mathbb{C}P^1, \mathbb{C}P^1, Sp_3/U_3, SU_6/S(U_3 \times U_3), SO_{12}/U_6, E_7/T^1 \cdot E_6$  or its non compact dual.

*Proof.* By Corollary 3.4, the tensor  $P$  vanishes, that is,

$$P(X, Y, Z, V) = (\nabla_V C)(X, Y, Z) + \omega(V)C(X, Y, JZ) \equiv 0.$$

It was shown in [AM1] that if  $C \neq 0$  at least at one point, then this condition means that the tensor field  $C$  generates a parallel holomorphic line bundle in the space of cubic symmetric forms of type  $(3, 0)$  such that the induced connection has curvature  $R^L = i\nu g \circ J$  (parallel cubic line bundle of type  $\nu$ ). All such Kähler manifolds are locally symmetric and locally isometric to one of the symmetric spaces described in [AM1, Thm. 3.14]. q.e.d.

## 4 Characterization of maximal parallel Kähler submanifolds of a quaternionic Kähler symmetric space

In this section we give a characterization of maximal parallel Kähler submanifolds  $M^{2n}$  of a quaternionic Kähler symmetric space  $\widetilde{M}^{4n}$ , of non zero scalar curvature.

**Theorem 4.1.** *Let  $M^{2n} \subset \widetilde{M}^{4n}$  be a complete maximal Kähler submanifold of a quaternionic symmetric space  $\widetilde{M}^{4n}$  of non zero scalar curvature. Then the following properties are equivalent:*

- (i)  $M$  is curvature invariant and locally symmetric.
- (ii)  $M$  is parallel.
- (iii)  $M$  is extrinsically symmetric.

*Proof.* For proof we need the following lemma.

**Lemma 4.2.** ([AM1, Prop. 2.8]) *Any curvature invariant maximal Kähler submanifold  $(M^{2n}, J)$  of a quaternionic Kähler manifold  $\widetilde{M}^{4n}$  is normal curvature invariant.*

*Proof.* The proof follows from the following identity which implies that the curvature tensor  $\widetilde{R}$  is invariant under the automorphism  $J_2$ :

$$\langle \widetilde{R}(J_2X, J_2Y)J_2Z, J_2W \rangle = \langle \widetilde{R}(X, Y)Z, W \rangle$$

for all  $X, Y, Z, W \in T\widetilde{M}$ .

q.e.d.

*Proof of the Theorem 4.1.* The equivalence (ii)  $\Leftrightarrow$  (iii) follows from the Lemma and Theorem 2.6. (ii)  $\Rightarrow$  (i) is well-known.

Thus, it remains to prove that (i)  $\Rightarrow$  (ii). Assume that  $M$  is curvature invariant and locally symmetric. Then, by Proposition 2.13 in [AM1, page 887] the tensor field  $[C, C]$  is parallel, i.e.,  $\nabla[C, C] = 0$ . We associate to the shape operator  $A$  the tensor  $[A, A] \in \Gamma(\Lambda^2 T^\perp M \otimes \Lambda^2 TM)$  by  $[A, A](\xi, \eta) = [A^\xi, A^\eta]$  for  $\xi, \eta \in T^\perp M$ .

We need the following lemma.

**Lemma 4.3.** *Let  $M^{2n}$  be a maximal Kähler submanifold of a quaternionic symmetric space  $\widetilde{M}^{4n}$ ,  $\nu \neq 0$ , and  $A^\xi$  its shape operator. Then the following holds:*

$$(\nabla_Z[C, C])(J_2\xi, J_2\eta)W = (\nabla'_Z[A, A])(\xi, \eta)W \quad .$$

*Proof of Lemma.* For  $\xi, \eta \in J_2T_xM$  and  $Z, W \in T_xM$ , we have

$$(\nabla_Z[C, C])(J_2\xi, J_2\eta)W = \nabla_Z(C \circ C)(J_2\xi, J_2\eta)W - \nabla_Z(C \circ C)(J_2\eta, J_2\xi)W \quad .$$

We have

$$\nabla_Z(C \circ C)(J_2\xi, J_2\eta)W = ((\nabla_Z C) \circ C)(J_2\xi, J_2\eta)W + (C \circ (\nabla_Z C))(J_2\xi, J_2\eta)W \quad .$$

By definition it follows that

$$(\nabla_Z C)_V W = \nabla_Z C_V W - C_{\nabla_Z V} W - C_V \nabla_Z W \quad .$$

Hence we obtain

$$\begin{aligned} \nabla_Z(C \circ C)(J_2\xi, J_2\eta)W &= ((\nabla'_Z A)^\xi \circ A^\eta)W + (A^\xi \circ (\nabla'_Z A)^\eta)W \\ &\quad - (C_{(\nabla_Z J_2)\xi} \circ C_{J_2\eta})W - (C_{J_2\xi} \circ C_{(\nabla_Z J_2)\eta})W \end{aligned}$$

Since  $(\nabla_Z J_2) = \omega(Z)J_3$ , we get

$$(C_{(\nabla_Z J_2)\xi} \circ C_{J_2\eta})W + (C_{J_2\xi} \circ C_{(\nabla_Z J_2)\eta})W = 0 \quad .$$

Then,

$$\nabla_Z(C \circ C)(J_2\xi, J_2\eta, W) = ((\nabla'_Z A)^\xi \circ A^\eta)W + (A^\eta \circ (\nabla'_Z A)^\xi)W = \nabla'_Z(A \circ A)(\xi, \eta, W) \quad .$$

Now, the lemma follows from the above identity.

q.e.d.

By using this lemma, we see that (i) implies  $(\nabla'_Z[A, A])(\xi, \eta)W = 0$ . Since  $J_1$  is parallel, we obtain that  $(\nabla'_Z[A, A])(\xi, J_1\eta)W = 0$ . From these two identities we get

$$\nabla'_Z(A \circ A)(\xi, \eta)W = ((\nabla'_Z A)^\xi \circ A^\eta)W + (A^\xi \circ (\nabla'_Z A)^\eta)W = 0.$$

Also, we have

$$\nabla'_{J_1 Z}(A \circ A)(\xi, \eta)W = ((\nabla'_{J_1 Z} A)^\xi \circ A^\eta)W + (A^\xi \circ (\nabla'_{J_1 Z} A)^\eta)W = 0.$$

Since  $M$  is curvature invariant, it follows that  $(\nabla'_{J_1 Z} A)^\xi X = -J_1(\nabla'_Z A)^\xi X$ . By using this fact together with the last two identities, we obtain

$$((\nabla'_Z A)^\xi \circ A^\eta)W = (A^\xi \circ (\nabla'_Z A)^\eta)W = 0.$$

Now, the theorem is a consequence of the following lemma. q.e.d.

**Lemma 4.4.** *Let  $M$  be a submanifold of a Riemannian manifold and  $A$  its shape operator. If*

$$((\nabla_X A)^\xi \circ A^\eta)W = (A^\xi \circ (\nabla_X A)^\eta)W = 0$$

*then  $M$  is parallel, i.e.,  $\nabla' A = 0$ .*

*Proof.* We decompose  $TM = \mathcal{N} \oplus \mathcal{N}^\perp$ , where

$$\mathcal{N} = \bigcap_{\xi \in TM^\perp} \ker(A^\xi), \quad \mathcal{N}^\perp = \text{span}\left(\bigcup_{\xi \in TM^\perp} \text{Image}(A^\xi)\right).$$

So, if  $Z \in \mathcal{N}^\perp$ , it follows that  $(\nabla'_X A)(\xi, Z) = 0$ . Let  $Z \in \mathcal{N}$  be any section. Observe that  $(\nabla'_X A)^\xi Z \in \mathcal{N}$ . On the other hand, we have  $(\nabla'_X A)^\xi Z = -A^\xi \nabla_X Z$ . Thus,  $(\nabla'_X A)^\xi Z \in \mathcal{N}^\perp$  and then  $(\nabla'_X A)^\xi Z = 0$ , that is,  $A$  is parallel. q.e.d.

## 5 Parallel Kähler submanifolds of quaternionic Kähler manifolds.

### 5.1 Reduction to the case of 1-full parallel Kähler submanifolds

Note that the intersection of totally geodesic submanifolds of a Riemannian manifold  $\widetilde{M}$  is a totally geodesic submanifold. Hence we may consider the minimal totally geodesic submanifold  $\overline{M}$  containing a given submanifold  $M$ .

In this subsection we prove the following theorem which reduces the classification of parallel Kähler submanifolds of a quaternionic Kähler symmetric manifold to the classification of 1-full parallel Kähler submanifolds in Hermitian or quaternionic Kähler symmetric spaces.

**Theorem 5.1.** *Let  $(M^{2m}, J)$  be a parallel Kähler submanifold of a symmetric quaternionic Kähler manifold  $\widetilde{M}^{4n}$  of non zero scalar curvature and  $\overline{M}$  the minimal totally geodesic submanifold of  $\widetilde{M}^{4n}$  containing  $M^{2m}$ .*

- 1) *If the shape tensor  $C$  of  $(M^{2m}, J)$  vanishes, then  $\overline{M}$  is a totally geodesic Hermitian symmetric space and  $(M^{2m}, J)$  is a full parallel Kähler submanifold of  $\overline{M}$ .*
- 2) *If  $C \neq 0$ , and hence  $(M^{2m}, J)$  is a Kähler manifold with parallel cubic line bundle, then  $\overline{M}$  is a quaternionic symmetric space of dimension  $4m$  and  $(M^{2m}, J)$  is a full parallel Kähler submanifold of  $\overline{M}$ .*

*Proof.* We need the following Lemma.

**Definition 5.2.** A parallel Kähler submanifold of a symmetric quaternionic Kähler manifold  $\widetilde{M}^{4n}$  is called *of type 1)* if the shape tensor  $C = 0$  and *of type 2)* otherwise.

**Lemma 5.3.** *Let  $M$  be a parallel Kähler submanifold of a symmetric quaternionic Kähler manifold with non zero scalar curvature.*

- 1) *If is of type 1), then*

$$J_2 T_x M \perp N_x^1 \quad \text{for all } x \in M.$$

- 2) *If it is of type 2), then*

$$J_2 T_x M = N_x^1 \quad \text{for all } x \in M.$$

*Proof of Lemma 5.3.* 1) is obvious, by definition of  $C$ . Before considering the case 2) let state some facts which hold true for any parallel submanifold  $M$ . As before, we use Latin letters  $X, Y, Z, \dots$  for vector fields in  $TM$  and Greek letters  $\xi, \eta, \dots$  for vector fields in  $T^\perp M$ . By hypothesis  $\nabla' h = 0$  we have the identity

$$(5) \quad \nabla_X^\perp (h(Y, Z)) = h(\nabla_X Y, Z) + h(Y, \nabla_X Z).$$

and

$$(6) \quad \widetilde{R}(TM, TM)TM \subset TM$$

Moreover, by (2) of Lemma 13 of [Na1],

$$(7) \quad \widetilde{R}(TM, TM)N^1 \subset N^1.$$

(Naitoh proved (7) as follows: the Ricci equation of the parallel submanifold can be written as

$$\tilde{R}(X, Y)\xi = R^\perp(X, Y)\xi - h(X, A^\xi Y) + h(A^\xi X, Y),$$

and for  $\xi = h(Z, T)$ , by (5), it follows that

$$R^\perp(X, Y)h(Z, T) = h(R(X, Y)Z, T) + h(Z, R(X, Y)T).$$

The conclusion follows immediately).

The proof of the Lemma follows directly from the next two Sublemmas.

**Sublemma 5.4.** *For any parallel Kähler submanifold  $M$  one has*

$$(8) \quad \tilde{R}(TM, N^1)TM \subset N^1.$$

Moreover, if  $M$  is of type 2) then

$$(9) \quad J_2TM \subset N^1,$$

*Proof of Sublemma 5.4.* Since  $\widetilde{M}^{4n}$  is a symmetric space and the submanifold  $M$  is curvature invariant, we have  $(\tilde{\nabla}_X \tilde{R})(Z, U)Y = 0$ , which can be written as

$$\begin{aligned} \nabla_X(\tilde{R}(Z, U)Y) + h(\tilde{R}(Z, U)Y, X) \\ = \tilde{R}(\nabla_X Z, U)Y + \tilde{R}(Z, \nabla_X U)Y + \tilde{R}(Z, U)\nabla_X Y \\ + \tilde{R}(h(X, Z), U)Y + \tilde{R}(Z, h(X, U))Y + \tilde{R}(Z, U)h(X, Y). \end{aligned}$$

The projection onto  $T^\perp M$  of this identity gives

$$(10) \quad \tilde{R}(h(X, Z), U)Y + \tilde{R}(Z, h(X, U))Y = h(\tilde{R}(Z, U)Y, X) - \tilde{R}(Z, U)h(X, Y).$$

By comparing (10) with the identity obtained by changing  $X \rightarrow J_1 X$  and  $U \rightarrow J_1 U$ , and taking account of (2), we deduce the following identity:

$$(11) \quad \begin{aligned} \tilde{R}(h(X, Z), U)Y = & (1/2)[- \nu(\langle J_2 h(X, Z), U \rangle J_2 Y + \langle J_3 h(X, Z), U \rangle J_3 Y) \\ & + h(\tilde{R}(Z, U)Y, X) + h(\tilde{R}(Z, JU)Y, JX) \\ & - \tilde{R}(Z, U)h(X, Y) - \tilde{R}(Z, JU)h(JX, Y)]. \end{aligned}$$

If  $M$  is of type 1) then (8) follows from (11), (6), (7). Let now assume that  $M$  is of type 2). We use (11) to compute the first two terms of the Bianchi identity  $0 = \tilde{R}(h(X, Z), U)Y + \tilde{R}(Y, h(X, Z))U + \tilde{R}(U, Y)h(X, Z)$ . Taking account of (7), we get

$$(12) \quad \begin{aligned} -\langle J_2 h(X, Z), U \rangle J_2 Y - \langle J_3 h(X, Z), U \rangle J_3 Y \\ + \langle J_2 h(X, Z), Y \rangle J_2 U + \langle J_3 h(X, Z), Y \rangle J_3 U \in N^1. \end{aligned}$$

Let us assume that at a point  $x \in M$  there exists a vector  $Y$  such that  $J_2 Y \notin (N^1)^\perp$ . If  $U = J_1 Y$ , then (12) gives

$$(13) \quad \langle J_3 h(X, Z), Y \rangle J_2 Y - \langle J_2 h(X, Z), Y \rangle J_3 Y \in N^1$$

and, by changing  $X \rightarrow J_1X$ , we get

$$(14) \quad \langle J_2h(X, Z), Y \rangle J_2Y + \langle J_3h(X, Z), Y \rangle J_3Y \in N^1.$$

By assumption, there exist vectors  $X, Z \in T_xM$  such that

$$\langle J_2h(X, Z), Y \rangle^2 + \langle J_3h(X, Z), Y \rangle^2 \neq 0.$$

Then (13) and (14) imply that  $J_2Y, J_3Y \in N^1$ . Now, for any  $U \in TM$ , (12) gives

$$(15) \quad \langle J_2h(X, Z), Y \rangle J_2U + \langle J_3h(X, Z), Y \rangle J_3U \in N^1,$$

from which, by comparing with the identity where  $U$  is replaced with  $JU$ , it is easy to deduce that  $J_2U \in N^1$ , for any  $U \in TM$ . (8) follows from (11), (6), (9) and (7). q.e.d.

**Sublemma 5.5.** *If  $M$  is of type 2) then*

$$(16) \quad J_2N^1 \subset TM.$$

*Proof of Sublemma 5.5.* Let us assume that the vector field  $\xi \in N^1$ . Since  $\tilde{R}(Y, \xi)Z \in N^1$  by (8), the identity  $(\tilde{\nabla}_X \tilde{R})(Y, \xi)Z = 0$  can be rewritten as

$$\begin{aligned} & \nabla_X^\perp \tilde{R}(Y, \xi)Z - A^{\tilde{R}(Y, \xi)Z} X \\ &= \tilde{R}(\nabla_X Y, \xi)Z + \tilde{R}(Y, \nabla_X^\perp \xi)Z + \tilde{R}(Y, \xi)\nabla_X Z \\ & \quad + \tilde{R}(h(X, Y), \xi)Z - \tilde{R}(Y, A_X^\xi)Z + \tilde{R}(Y, \xi)h(X, Z). \end{aligned}$$

By using repeatedly (5), (7) and (8), we get

$$\tilde{R}(h(X, Y), \xi)Z + \tilde{R}(Y, \xi)h(X, Z) \in O^1$$

and, by changing  $Y \rightarrow JY$  and  $\xi \rightarrow J\xi$ ,

$$\tilde{R}(J_1h(X, Y), J_1\xi)Z + \tilde{R}(J_1Y, J_1\xi)h(X, Z) \in O_x^1 = T_xM + h(T_xM, T_xM).$$

The last two identities together with (2) imply that

$$\begin{aligned} & \nu(\langle J_2h(X, Y), \xi \rangle J_2Z + \langle J_3h(X, Y), \xi \rangle J_3Z \\ & \quad + \langle J_2Y, \xi \rangle J_2h(X, Z) + \langle J_3Y, \xi \rangle J_3h(X, Z)) \in O_x^1. \end{aligned}$$

Since  $J_2Z, J_3Z \in N^1$  by Lemma 5.4, we conclude that

$$(17) \quad \langle J_2Y, \xi \rangle J_2h(X, Z) + \langle J_3Y, \xi \rangle J_3h(X, Z) \in O_x^1.$$

Let us assume that there exists a vector  $Y \in T_xM$  such that  $\langle J_2Y, \xi \rangle^2 + \langle J_3Y, \xi \rangle^2 \neq 0$ . We deduce easily, by comparing (17) with the identity obtained by the change  $Y \rightarrow J_1Y$ , that



$$J_2h(X, Z), J_3h(X, Z) \in O^1, \text{ for any } X, Z \in T_xM.$$

On the other hand, by Corollary 3.4,  $J_2h(X, Z)$  is orthogonal to  $N^1$ . Hence

$$J_2h(X, Z), J_3h(X, Z) \in TM, \text{ for any } X, Z \in T_xM,$$

and (5.5) follows. This finish the proof of Sublemma 5.5 and hence Lemma 5.3. q.e.d.

Now we prove the following Proposition which, together with Lemma 5.3, implies Theorem 5.1.

**Proposition 5.6.** *Let  $(M^{2m}, J)$  be a parallel Kähler submanifold of a locally symmetric quaternionic Kähler manifold. Then the first osculating space  $O_x^1 = T_xM + N_x^1$  at any point  $x \in M$  is curvature invariant, i.e.,*

$$\tilde{R}(O^1, O^1)O^1 \subset O^1.$$

*Remark.* The proposition remains true if  $\tilde{M}$  is a locally symmetric Kähler manifold, whose proof is the same as in the quaternionic Kähler case.

*Proof.* The identity  $(\tilde{\nabla}\tilde{R})(Y, Z)\xi = 0$  can be rewritten as

$$\begin{aligned} & \nabla_X^\perp(\tilde{R}(Y, Z)\xi) - A^{\tilde{R}(Y, Z)\xi}X \\ &= \tilde{R}(\nabla_X Y, Z)\xi + \tilde{R}(Y, \nabla_X Z)\xi + \tilde{R}(Y, Z)\nabla_X^\perp\xi \\ & \quad + \tilde{R}(h(X, Y), Z)\xi + \tilde{R}(Y, h(X, Z))\xi - \tilde{R}(Y, Z)A^\xi X. \end{aligned}$$

For  $\xi \in N^1$ , by taking account of (6), (7) and (5), this gives

$$\tilde{R}(h(X, Y), Z)\xi + \tilde{R}(Y, h(X, Z))\xi \in O^1.$$

By changing  $X \rightarrow J_1X$  and  $Z \rightarrow J_1Z$ , we have

$$\tilde{R}(J_1h(X, Y), J_1Z)\xi - \tilde{R}(Y, h(X, Z))\xi \in O^1.$$

By (2), we also have

$$\begin{aligned} & \tilde{R}(J_1h(X, Y), J_1Z)\xi \\ &= \tilde{R}(h(X, Y), Z)\xi + \nu(\langle J_2h(X, Y), Z \rangle J_2\xi + \langle J_3h(X, Y), Z \rangle J_3\xi) \in TM \end{aligned}$$

which implies

$$(18) \quad \tilde{R}(N^1, TM)N^1 \subset O^1.$$

Now the Bianchi identity gives

$$(19) \quad \tilde{R}(N^1, N^1)TM \subset O^1.$$

We rewrite the identity  $(\widetilde{\nabla} \widetilde{R})(Y, \eta)\xi = 0$  for  $\eta, \xi \in N^1$  as follows:

$$\begin{aligned} \widetilde{\nabla}_X(\widetilde{R}(Y, \eta)\xi) &= \widetilde{R}(\nabla_X Y, \eta)\xi + \widetilde{R}(Y, \nabla_X^\perp \eta)\xi + \widetilde{R}(Y, \eta)\nabla_X^\perp \xi \\ &\quad + \widetilde{R}(h(X, Y), \eta)\xi - \widetilde{R}(Y, A^\eta X)\xi - \widetilde{R}(Y, \eta)A^\xi X. \end{aligned}$$

Since the bundle  $O^1$  is invariant under parallel transport, it follows that  $\widetilde{R}(h(X, Y), \eta)\xi \in O_x^1$ , and hence

$$(20) \quad \widetilde{R}(N_x^1, N_x^1)N_x^1 \subset O_x^1 \quad .$$

Formulas (6), (7), (8), (18), (19) and (20) then imply Proposition 5.6 . q.e.d.

We also obtain the following corollary, which was proved by Tsukada [Tsu2] in the case of quaternionic projective space.

**Corollary 5.7.** *A non totally geodesic parallel totally complex submanifold  $(M^{2m}, J_1)$  of a symmetric quaternionic Kähler manifold  $\widetilde{M}^{4n}$  is 1-full if and only if it has maximal dimension, i.e.,  $n = m$ .*

*Proof.* We have the following orthogonal decomposition:

$$T\widetilde{M} = TM + J_2(TM) + N(M),$$

where  $N(M)$  is a quaternionic subbundle. If we assume that  $M$  is 1-full, then it follows that  $T^\perp M = J_2 TM + NM = N^1 M$ . By 1) of Corollary 3.4,  $N^1 M$  is totally complex, and hence  $NM = 0$ . Vice versa, if  $M$  has maximal dimension  $n = m$ , then  $J_2 TM = T^\perp M$ . Since  $M$  is not totally geodesic,  $M$  has type 2) and by Lemma 5.3, we get  $N^1 M = J_2 TM = T^\perp M$ . q.e.d.

**Remark 5.8.** As a consequence of Proposition 5.6 and Naitoh's Theorem 2.9, it follows that the concept of being 1-full and that of being full are equivalent for a parallel Kähler submanifold of a locally symmetric quaternionic Kähler manifold.

Now we can prove Theorem 5.1. By Proposition 5.6 and Theorem 2.9, the Kähler submanifold  $M^{2m}$  is 1-full in the totally geodesic submanifold  $\overline{M} = \exp(O_x^1 M)$ . In the case 1),  $\overline{M}$  is a totally complex totally geodesic submanifold, and hence a Hermitan symmetric space. In the case 2),  $\overline{M}$  is a quaternionic Kähler submanifold. q.e.d.

## 6 Totally geodesic maximal Kähler submanifolds of Wolf spaces

All totally geodesic maximal Kähler submanifolds  $M^{2n}$  of a Wolf space  $W = G/K = \widetilde{M}^{4n}$  were classified by Takeuchi in terms of *Satake diagrams* [Tak]. Here we sketch

another approach based on a simple observation that there exists a natural one to one correspondence between such submanifolds and involutive automorphisms of the complex Lie algebra  $\mathfrak{g} = \text{Lie}(G)^\mathbb{C}$ , which preserve the canonical ideal  $\mathfrak{sp}_1$  of the stability Lie algebra  $\mathfrak{k}$  and act non trivially on it. Similar ideas can be found in [Wo].

## 6.1 Lie algebra description of Wolf spaces

Recall that any simple complex Lie algebra  $\mathfrak{g}$  determine the Wolf space as follows. Let

$$\mathfrak{g} = \mathfrak{h} + \sum_{\alpha \in R} \mathbb{C}E_\alpha$$

be the Cartan decomposition of the Lie algebra  $\mathfrak{g}$  with respect to a Cartan subalgebra  $\mathfrak{h}$  and  $\Pi = \{\alpha_1, \dots, \alpha_\ell\}$  a system of simple roots of the root system  $R$ .

We denote by  $\mu$  the maximal root of  $R$  and by  $H_\mu = 2/(\mu, \mu)B_\mu^{-1} = [E_\mu, E_{-\mu}]$  the corresponding element of  $\mathfrak{h}$  such that  $\{H_\mu, E_{\pm\mu}\}$  is the standard basis of the 3-dimensional subalgebra  $\mathfrak{a}_1 = \mathfrak{sp}_1^\mu(\mathbb{C})$ . Then  $\text{ad}_{H_\mu}$  has the eigenvalues  $\pm 2, \pm 1, 0$  and the corresponding eigenspace decomposition

$$(21) \quad \mathfrak{g} = \mathfrak{g}_{-2} + \mathfrak{g}_{-1} + \mathfrak{g}_0 + \mathfrak{g}_1 + \mathfrak{g}_2$$

gives rise to a gradation of the Lie algebra  $\mathfrak{g}$ . Moreover, we have

$$\mathfrak{g}_{\pm 2} = \mathbb{C}E_{\pm\mu}, \quad \mathfrak{g}_{\pm 1} = \sum_{\alpha \in \pm R_1} \mathfrak{g}_\alpha, \quad \mathfrak{g}_0 = \mathfrak{h} + \sum_{\alpha \in R_0} \mathbb{C}E_\alpha = \mathfrak{g}_0' \oplus \mathbb{C}H_\mu,$$

where

$$R_1 = \{\alpha \in R; \alpha(H_\mu) = \frac{2(\alpha, \mu)}{(\mu, \mu)} = 1\}, \quad R_0 = \{\alpha \in R; (\alpha, \mu) = 0\}.$$

We put  $\varphi_0 = \exp i\pi(\text{ad}_{H_\mu})$ , which is an involutive automorphism of  $\mathfrak{g}$  with eigenspace decomposition

$$\mathfrak{g} = \mathfrak{g}_{\text{ev}} + \mathfrak{g}_{\text{odd}} = (\mathfrak{g}_{-2} + \mathfrak{g}_0 + \mathfrak{g}_2) + (\mathfrak{g}_{-1} + \mathfrak{g}_1).$$

Since  $\varphi_0$  commutes with the standard antilinear involution  $\tau$  of  $\mathfrak{g}$  associated with the Cartan decomposition, which determines the compact real form  $\mathfrak{g}^\tau = \{X \in \mathfrak{g}; \tau(X) = X\}$ ,  $\varphi_0$  defines a symmetric decomposition

$$\mathfrak{g}^\tau = \mathfrak{g}_{\text{ev}}^\tau + \mathfrak{g}_{\text{odd}}^\tau = (\mathfrak{sp}_1^\mu + \mathfrak{g}_0')^\tau + (\mathfrak{g}_{-1} + \mathfrak{g}_1)^\tau = \mathfrak{k} + \mathfrak{m}$$

of the compact Lie algebra  $\mathfrak{g}^\tau$ . We denote by  $G$  the adjoint (compact) Lie group with the Lie algebra  $\mathfrak{g}^\tau$  and by  $K = N_G(\mathfrak{a}_1) = Sp_1^\mu \cdot K'$  the normalizer of the 3-dimensional subalgebra (which is the connected Lie group generated by the subalgebra  $\mathfrak{k} = \mathfrak{g}_{\text{ev}}^\tau$ ). Then  $W = G/K$  is a simply connected irreducible symmetric space  $W = G/K$  associated

with this symmetric decomposition. Moreover, it has a natural structure of quaternionic Kähler symmetric space, which is called the *Wolf space associated with the Lie algebra  $\mathfrak{g}$* . The quaternionic structure  $Q$  in the tangent space  $T_o W = \mathfrak{g}_{\text{odd}}^\tau$  is given by  $Q = \text{ad}_{\mathfrak{sp}_1^\mu}|_{\mathfrak{g}_{\text{odd}}^\tau}$ .

Remark that the pair  $(G, K)$  is determined by the grading element  $d = H_\mu$  of the gradation (21) and the antilinear involution  $\tau$  with  $\tau d = -d$ . Conversely, a pair  $(d, \tau)$ , where  $d$  is the grading element of a gradation (21) with  $\dim \mathfrak{g}_{\pm 2} = 1$  and  $\tau$  is an antilinear involution of  $\mathfrak{g}$  with  $\tau d = -d$ , defining a compact real form  $\mathfrak{g}^\tau$  of  $\mathfrak{g}$  defines a Wolf space  $W = G/K$ , and any such pairs are conjugated by an inner automorphism of  $\mathfrak{g}$ .

## 6.2 Totally geodesic extrinsically symmetric Kähler submanifolds of a Wolf space

Let  $W = G/K$  be a Wolf space associated with a complex simple Lie algebra  $\mathfrak{g}$  and  $(d = H_\mu, \tau)$  be the pair that determines  $(G, K)$  as above. Since the isotropy group  $K = Sp_1^\mu \cdot K'$  acts transitively on the unit sphere of all complex structures  $J \in Q = \text{ad}_{\mathfrak{a}_1}|_{\mathfrak{m}}$ , any totally geodesic Kähler submanifold  $M$  of  $W$  containing  $o = eK \in W$  is  $K$ -equivalent to a submanifold  $M' \ni o$ , whose tangent space  $T_o M$  is invariant under some fixed complex structure  $J_1 \in Q$ . We choose as  $J_1$  the complex structure  $J_1 = \text{ad}_{iH_\mu}|_{\mathfrak{g}_{\text{odd}}^\tau}$ . We will call a totally geodesic Kähler submanifold  $M$  of  $W$  *admissible* if it contains  $o$  and the tangent space  $T_o M$  is  $J_1$ -invariant.

**Theorem 6.1.** *Let  $W = G/K$  be a Wolf space associated with a complex simple Lie algebra  $\mathfrak{g}$ ,  $d = H_\mu$  be the grading element of the gradation (21) and  $\tau$  be the antilinear involution defining the compact real form  $\text{Lie } G = \mathfrak{g}^\tau$  of  $\mathfrak{g}$ .*

- 1) *There is a natural one to one correspondence between*
  - i) *involutive automorphisms  $\sigma$  of  $\mathfrak{g}$  which commute with  $\tau$  and satisfy condition  $\sigma(E_{\pm\mu}) = -E_{\pm\mu}$ , and*
  - ii) *(connected) admissible totally geodesic extrinsically symmetric Kähler submanifolds  $M(\sigma)$  of  $W = G/K$  given by  $M(\sigma) = W^{s_\sigma}$ , where  $W^{s_\sigma} \ni o$  is the connected component of the fixed points set of the symmetry  $s_\sigma : W \ni aK \mapsto \sigma(a)K$ . Moreover,  $\dim M(\sigma) = (1/2) \dim W$ .*
- 2) *Submanifolds  $M(\sigma)$  and  $M(\sigma_1)$  are  $G$ -equivalent if and only if the involutive automorphisms  $\sigma$  and  $\sigma_1$  are conjugated by an element of  $K$ .*
- 3) *For any submanifold  $M(\sigma)$  there is another canonically defined totally geodesic extrinsically symmetric Kähler submanifold  $M(\sigma')$  associated with the involutive automorphism  $\sigma' = \varphi_0 \circ \sigma$  such that one has the orthogonal decomposition  $T_o W = T_o M(\sigma) + T_o M(\sigma')$ .*

- 4) The pair of involutive automorphism  $\sigma$  and  $\sigma' = \varphi_0 \circ \sigma$  is determined by the restriction of  $\sigma$  to  $\mathfrak{g}'_0$ . Two automorphisms  $\sigma$  and  $\sigma_1$  define  $G$ -equivalent pairs  $(M(\sigma), M(\sigma'))$  and  $(M(\sigma_1), M(\sigma'_1))$  of submanifolds if and only if the automorphism  $\sigma|_{\mathfrak{g}'_0}$  is conjugated to  $\sigma_1|_{\mathfrak{g}'_0}$  or  $\sigma'_1|_{\mathfrak{g}'_0}$  in the group of automorphisms of  $\mathfrak{g}'_0$ .

*Proof of Theorem.* 1) Let  $M = L/L_0 = Lo$  be an admissible totally geodesic extrinsically symmetric Kähler submanifold of the Wolf space  $W = G/K$  and

$$\mathfrak{g} = \mathfrak{g}_{-2} + \mathfrak{g}_{-1} + \mathfrak{g}_0 + \mathfrak{g}_1 + \mathfrak{g}_2 = \mathfrak{g}_{\text{ev}} + \mathfrak{g}_{\text{odd}}$$

the  $\text{ad}_{H_\mu}$ -eigenspace decomposition of the complex Lie algebra  $\mathfrak{g}$ . We identify the complexified tangent space  $T_o^\mathbb{C}W$  with  $\mathfrak{p} = \mathfrak{g}_{\text{odd}} = \mathfrak{g}_{-1} + \mathfrak{g}_1$ .

The symmetry  $s_o$  of  $M$  at point  $o$  induces a complex linear involutive transformation  $s_{o*}$  of  $T_o^\mathbb{C}W = \mathfrak{p} = \mathfrak{g}_{-1} + \mathfrak{g}_1$ , which by assumption commutes with the complex structure  $J_1 = \text{ad}_{iH_\mu}|_{\mathfrak{p}}$ . This implies that the eigenspace decomposition of  $s_{o*}$  has the form

$$\mathfrak{p} = (\mathfrak{g}_{-1}^+ + \mathfrak{g}_1^+) + (\mathfrak{g}_{-1}^- + \mathfrak{g}_1^-),$$

where the  $+1$ -eigenspace  $\mathfrak{m}^\mathbb{C} = \mathfrak{g}_{-1}^+ + \mathfrak{g}_1^+$  is the complexification of the tangent space  $\mathfrak{m} = T_oM$  and  $\mathfrak{g}_{-1}^- + \mathfrak{g}_1^-$  is its orthogonal complement. The graded subspace  $\mathfrak{m}^\mathbb{C}$  generates a graded Lie subalgebra  $\ell = [\mathfrak{m}^\mathbb{C}, \mathfrak{m}^\mathbb{C}] + \mathfrak{m}^\mathbb{C}$  of  $\mathfrak{g}$ . Since  $[\mathfrak{m}^\mathbb{C}, \mathfrak{m}^\mathbb{C}]$  cannot contain the subalgebra  $\mathfrak{sp}_1^\mu(\mathbb{C})$ , it belongs to  $\mathfrak{g}_{-1} + \mathfrak{g}_0 + \mathfrak{g}_1$ . In particular,  $[\mathfrak{g}_1^+, \mathfrak{g}_1^+] = [\mathfrak{g}_{-1}^+, \mathfrak{g}_{-1}^+] = 0$ . On the other hand,  $\ell_0 = [\mathfrak{m}^\mathbb{C}, \mathfrak{m}^\mathbb{C}] \subset \mathfrak{g}_0$  contains  $H_\mu$ , since  $M = L/L_0$  is a Hermitian symmetric space.

We denote by  $\sigma$  the involutive automorphism of the group  $G$  and its Lie algebra  $\mathfrak{g}^\tau$  defined by conjugation with the symmetry  $s_o$ , and extend it to a complex linear automorphism  $\sigma$  of  $\mathfrak{g}$ , which commutes with  $\tau$ . Since the restriction  $\sigma|_{\mathfrak{p}} = s_o|_{\mathfrak{p}}$  commutes with  $J_1 = \text{ad}_{iH_\mu}|_{\mathfrak{p}}$ , we have  $\sigma(H_\mu) = H_\mu$ , that is,  $\sigma$  preserves the gradation of  $\mathfrak{g}$  defined by  $H_\mu$ . In particular,  $\sigma(E_{\pm\mu}) = \epsilon E_{\pm\mu}$ , where  $\epsilon = \pm 1$ . Assume that  $\epsilon = +1$ , i.e.,  $\sigma(E_{\pm\mu}) = E_{\pm\mu}$ . Then  $(s_o)_*|_{T_oW}$  commutes with the quaternionic structure  $Q = \text{ad}_{\mathfrak{sp}_1}^\mu(\mathbb{C})$ , which contradicts the assumption that  $M$  is totally complex. Hence  $\sigma(E_{\pm\mu}) = -E_{\pm\mu}$ . We have proven that the automorphism  $\sigma$  defined by the symmetry  $s_o$  satisfies all conditions of the theorem.

Now we remark that

$$[\mathfrak{g}_{\pm 1}^+, \mathfrak{g}_{\pm 1}^+] = [\mathfrak{g}_{\pm 1}^-, \mathfrak{g}_{\pm 1}^-] = 0,$$

since  $\sigma|_{\mathfrak{g}_{\pm 2}} = -\text{Id}$ . This means that  $\mathfrak{g}_{\pm 1} = \mathfrak{g}_{\pm 1}^+ + \mathfrak{g}_{\pm 1}^-$  is a decomposition of the complex symplectic vector space  $\mathfrak{g}_{\pm 1}$ , with the symplectic form  $\omega$  defined by  $[X, Y] = \omega(X, Y)E_{\pm\mu}$ , into direct sum of two Lagrangian subspaces. In particular,

$$\dim \mathfrak{g}_1^+ = \dim \mathfrak{g}_1^- = \dim \mathfrak{g}_{-1}^+ = \dim \mathfrak{g}_{-1}^- = \frac{1}{4} \dim W.$$

Conversely, let  $\sigma$  be an involutive automorphism commuting with  $\tau$  and acting as  $-\text{Id}$  on  $\mathfrak{g}_{-2} + \mathfrak{g}_2$ . Then it preserves  $H_\mu = [E_\mu, E_{-\mu}]$ . Hence its eigenspaces decomposition has the form

$$\mathfrak{g} = \mathfrak{g}_{-2} + \mathfrak{g}_{-1}^+ + \mathfrak{g}_{-1}^- + \mathfrak{g}_0^+ + \mathfrak{g}_0^- + \mathfrak{g}_1^+ + \mathfrak{g}_1^- + \mathfrak{g}_2.$$

Moreover,  $[\mathfrak{g}_{\pm 1}^+, \mathfrak{g}_{\pm 1}^+] = [\mathfrak{g}_{\pm 1}^-, \mathfrak{g}_{\pm 1}^-] = 0$  and the four spaces  $\mathfrak{g}_{\pm 1}^\pm$  have the same dimension. One can easily check that the subalgebras

$$\ell^- = \mathfrak{g}_{-1}^- + \mathfrak{g}_1^-, \quad \ell^+ = \mathfrak{g}_{-1}^+ + \mathfrak{g}_1^+$$

define two totally geodesic extrinsically symmetric Kähler submanifolds  $M^+ = M(\sigma)$  and  $M^- = M(\varphi_0 \circ \sigma)$  of the same dimension  $2n = (1/2) \dim W$ .

To prove that the correspondence between  $\sigma$  and  $M(\sigma)$  is a bijection, it is sufficient to show that two involutive automorphisms  $\sigma, \sigma'$  coincide if they have the same restriction to  $\mathfrak{g}_{-1} + \mathfrak{g}_1$  or, equivalently, that the fixed point set  $\mathfrak{g}^\sigma = \mathfrak{g}_{-1}^\sigma + \mathfrak{g}_0'^\sigma + \mathbb{C}H_\mu + \mathfrak{g}_1^\sigma$  can be reconstructed from  $\mathfrak{g}_{-1}^\sigma + \mathfrak{g}_1^\sigma$ . Since  $\mathfrak{g}_0' = [\mathfrak{g}_{-1}, \mathfrak{g}_1]$ , we have

$$\mathfrak{g}_0'^\sigma = [\mathfrak{g}_{-1}, \mathfrak{g}_1]^\sigma = [\mathfrak{g}_{-1}^\sigma, \mathfrak{g}_1^\sigma].$$

2) If  $M(\sigma)$  and  $M(\sigma_1)$  are  $G$ -equivalent, there exists an isometry  $k \in K$  such that  $kM(\sigma) = M(\sigma_1)$ . Then the conjugation by  $k$  transforms  $\sigma$  into  $\sigma'$ . The converse statement is also clear.

3) is obvious. To prove 4), it is sufficient to check that an automorphism  $\rho = \sigma^{-1} \circ \sigma'$  acting trivially on  $\mathfrak{g}_{\text{ev}} = \mathfrak{g}_{-2} + \mathfrak{g}_0 + \mathfrak{g}_2$  is either trivial or equal to  $\varphi_0$ . It follows from the fact that the isometry of  $W$  associated to  $\rho$  with the fixed point  $o$  commutes with the stability subgroup  $K$  acting irreducibly on  $T_o W$ . q.e.d.

It is not difficult to describe all automorphisms  $\sigma$  of  $\mathfrak{g}$  which correspond to totally geodesic extrinsically symmetric Kaehler submanifolds  $M(\sigma)$  in terms of Kac diagrams, see [GOV]. Here we state only a corollary which we use in the proof of Theorem 1.1.

**Corollary 6.2.** *Let  $W = G/K$  be a Wolf space or its non compact dual. Then, up to an isometry, there exist finitely many totally geodesic extrinsically symmetric Kähler submanifolds of  $W$ . Any one of them has dimension  $(1/2) \dim W$ .*

*Proof.* The claim for Wolf spaces follows from Theorem 6.1. It remains true for non compact dual  $W'$ , since totally geodesic Kähler extrinsically symmetric submanifolds can be characterized as totally geodesic Kähler submanifolds which are normal curvature invariant and the restriction of the natural one-to-one correspondence between totally geodesic submanifolds of  $W$  and  $W'$  gives a one-to-one correspondence between such submanifolds. q.e.d.

Remark that in a symmetric space  $M$  there could be even a continuous number of non equivalent totally geodesic submanifolds of given dimension, for example geodesics in a symmetric space of rank greater than 1.

## 7 Proof of Theorems 1.1 and 1.2

*Proof of Theorem 1.1.* Let  $M$  be a curvature invariant maximal Kähler submanifold of a Wolf space or its dual. By Lemma 4.2,  $M$  is also normal curvature invariant. Hence for any point  $x \in M$  there exists an involutive isometry  $s_o$  such that  $s_o|_{T_x M} = -\text{Id}$  and  $s_o|_{T_x^\perp M} = \text{Id}$ , see the proof of Theorem 2.6. This shows that the totally geodesic submanifold  $M(x) = \exp(T_x M)$  is an extrinsically symmetric maximal Kähler submanifold. Hence by 6.2, the tangent space  $T_x M$  belongs to one of the finitely many orbits  $\mathcal{V} = G(V) \subset \text{Gr}_{2n} T(G/K)$ . By continuity reason,  $M$  is a  $\mathcal{V}$ -submanifold, where  $\mathcal{V}$  is defined by one of the extrinsically symmetric Kähler submanifolds. Since  $\mathcal{V}$  is curvature and normal curvature invariant, by applying Naitoh's Theorem 2.8,  $M$  is totally geodesic if  $\widetilde{M} \neq \mathbb{H}P^n$  or the dual quaternionic hyperbolic space  $\mathbb{H}H^n$  (The last statement for  $\widetilde{M} \neq \mathbb{H}P^n$  can also be obtained directly by using Theorem 5.4 and Remark 5.5 of [Na2] for the Grassmannian  $G_2(\mathbb{C}^{n+2})$ . An elementary proof that  $G_2(\mathbb{C}^{n+2})$  does not contain non totally geodesic maximal Kähler submanifolds was given in [ADM]). It is known ([Tsu2]) that any parallel Kähler submanifold of  $\mathbb{H}H^n$  is totally geodesic. This proves Theorem 1.1.

*Proof of Theorem 1.2.* The first claim was proved in Theorem 5.1. Assume that the shape tensor  $C \neq 0$ . Then by Theorem 5.1,  $M^{2m}$  is a parallel maximal Kähler submanifold of a quaternionic Kähler symmetric space  $\widetilde{M}^{4m}$ . Theorem 1.1 then implies that  $\widetilde{M} = \mathbb{H}P^m$ . Now result follows from Tsukada's classification of parallel Kähler submanifolds of  $\mathbb{H}P^m$ . q.e.d.

ACKNOWLEDGEMENT. The authors like to thank heartily the referee for useful remarks.

## References

- [ADM] D.V. ALEKSEEVSKY, A.J. DI SCALA AND S. MARCHIAFAVA *Kähler submanifolds of Wolf spaces*, Proceedings of the 6th International Workshop on Differential Geometry and its Applications at Cluj-Napoca, Romania, September 2003 (to appear)
- [AM1] D.V. ALEKSEEVSKY AND S. MARCHIAFAVA *Hermitian and Kähler submanifolds of a quaternionic Kähler manifold*, Osaka J. Math. **38** (2001), 869–904.
- [AM2] D.V. ALEKSEEVSKY AND S. MARCHIAFAVA *A twistor construction of Kähler submanifolds of a quaternionic Kähler manifold*, Ann. Mat. Pura Appl. 183 (published on line: 28 January 2004)

- [B] J. BERNDT *Symmetric submanifolds of symmetric spaces*, Proceedings of The Seventh International Workshop on Differential Geometry **7** (Taegu, 2002), 1–15, Kyungpook Natl. Univ., Taegu, 2003.
- [BENT] J. BERNDT, J.-H. ESHENBURG, H. NAITOH AND K. TSUKADA *Symmetric submanifolds associated with the irreducible symmetric R-spaces* Preprint (2003)
- [BRS] F. BURSTALL, J. RAWNSLEY AND S. SALAMON *Stable harmonic 2-spheres in symmetric spaces*, Bull. Amer. Math. Soc. **16** (1987), 274–278.
- [GOV] V.V. GORBATSEVICH, A.L. ONISHCHIK AND E.B. VINBERG *Lie groups and Lie algebras*, III, Encyclopaedia of Math. Sci., **41** (1994), 113–121
- [Na1] H. NAITOH *Isotropic submanifolds with parallel second fundamental forms in symmetric spaces*, Osaka J. Math. **17** (1980), 95–110
- [Na2] H. NAITOH *Symmetric submanifolds of compact symmetric spaces*, Tsukuba J. Math. **10** (1986), 215–242.
- [Na3] H. NAITOH *Grassmann geometries on compact symmetric spaces of exceptional type*, Japan J. Math. **26** (2000), 157–206.
- [Na4] H. NAITOH *Parallel submanifolds of complex space forms I*, Nagoya Math. J. **90** (1983), 85–117.
- [NT] H. NAKAGAWA AND R. TAKAGI *On locally symmetric Kaehler submanifolds in a complex projective space*, J. Math. Soc. Japan **28** (1976), 638–667.
- [Str] W. STRÜBING *Symmetric submanifolds of Riemannian manifolds*, Math. Ann. **245** (1979), no. 1, 37–44.
- [Ta] H. TASAKI *Quaternionic submanifolds in quaternionic symmetric spaces*, Tohoku Math. Journ. **38** (1986), 513–538.
- [Tak] M. TAKEUCHI *Totally complex submanifolds of quaternionic symmetric spaces*, Japan. J. Math. **12** (1986), 161–189.
- [Tsu1] K. TSUKADA *Parallel Kaehler Submanifolds of Hermitian Symmetric Spaces*, Math. Z. **190** (1985), 129–150.
- [Tsu2] K. TSUKADA *Parallel submanifolds in a quaternion projective space*, Osaka J. Math. **22** (1985), 187–241.
- [Wo] J. WOLF *Complex forms of Quaternionic Symmetric Spaces*, arXiv:math.DG/0308283 v2 2 Oct 2003.



THE UNIVERSITY OF HULL  
COTTINGHAM ROAD  
HU6 7RX, HULL  
UNITED KINGDOM

*E-mail addresses:* D.V.Alekseevsky@maths.hull.ac.uk

DIPARTIMENTO DI MATEMATICA  
POLITECNICO DI TORINO,  
CORSO DUCA DEGLI ABRUZZI 24,  
10129 TORINO  
ITALY

*E-mail address:* antonio.discal@polito.it

DIPARTIMENTO DI MATEMATICA  
UNIVERSITA' DEGLI STUDI DI ROMA "LA SAPIENZA",  
P.LE A. MORO 2,  
00185 ROMA  
ITALY

*E-mail address:* marchiaf@mat.uniroma1.it